

Kuranishi homology and Kuranishi cohomology

Dominic Joyce

Abstract

A *Kuranishi space* is a topological space equipped with a *Kuranishi structure*, defined by Fukaya and Ono. Kuranishi structures occur naturally on many moduli spaces in differential geometry. If (M, ω) is a symplectic manifold with compatible almost complex structure J , then moduli spaces of stable J -holomorphic curves in M have Kuranishi structures. Kuranishi spaces are important in symplectic geometry, for defining Gromov–Witten invariants, Lagrangian Floer cohomology, and so on.

Let Y be an orbifold, and R a commutative ring or $\mathbb{Q}$ -algebra. We shall define two new homology and cohomology theories of Y : *Kuranishi (co)homology* $KH_*, KH^*(Y; R)$, for R a $\mathbb{Q}$ -algebra, and *effective Kuranishi (co)homology* $KH_*^{\text{ef}}, KH_{\text{ec}}^*(Y; R)$, for R a commutative ring, such as $R = \mathbb{Z}$. Our main result is that $KH_*, KH_*^{\text{ef}}(Y; R)$ are isomorphic to singular homology, and $KH^*, KH_{\text{ec}}^*(Y; R)$ are isomorphic to compactly-supported cohomology. We also define five different kinds of *Kuranishi (co)bordism* $KB_*, KB^*(Y; R)$. These are new topological invariants of Y which turn out to be very large, far larger than the (co)homology groups.

The (co)bordism theories $KB_*, KB^*(Y; R)$ are spanned over R by isomorphism classes $[X, \mathbf{f}]$, where X is a compact oriented Kuranishi space without boundary, and $\mathbf{f} : X \rightarrow Y$ is a strongly smooth map. The (co)homology theories are defined using (co)chain complexes $KC_*, \dots, KC_{\text{ec}}^*(Y; R)$ spanned by isomorphism classes $[X, \mathbf{f}, \mathbf{G}]$, where X is a compact oriented Kuranishi space with boundary and corners, $\mathbf{f} : X \rightarrow Y$ is strongly smooth, and $\mathbf{G}$ is some extra *gauge-fixing data* upon X . The main purpose of $\mathbf{G}$ is to ensure the automorphism groups $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ are finite, which is necessary to get a well-behaved (co)homology theory.

These theories are powerful new tools in symplectic geometry. We define new Gromov–Witten type invariants in Kuranishi bordism, containing more information than conventional Gromov–Witten invariants. In a sequel we hope to use these to prove the integrality conjecture for Gopakumar–Vafa invariants. We also explain how to reformulate Lagrangian Floer cohomology using Kuranishi cohomology rather than singular homology; this will lead to significant technical simplifications, and improved results.

The results of this book are briefly surveyed in [38].

Contents

1	Introduction	3
2	Kuranishi structures and Kuranishi spaces	8
2.1	Manifolds with corners and generalized corners	8
2.2	Orbifolds and orbibundles	16
2.3	Kuranishi neighbourhoods and coordinate changes	21
2.4	Kuranishi structures	24
2.5	Strongly smooth maps and strong submersions	25
2.6	Boundaries and fibre products	27
2.7	Orientations and orientation conventions	28
2.8	Coorientations	31
2.9	Almost complex and almost CR structures	33
2.10	Co-almost complex and co-almost CR structures	35
3	Gauge-fixing data and co-gauge-fixing data	36
3.1	Definition of (co-)gauge-fixing data	38
3.2	Existence of (co-)gauge-fixing data	42
3.3	Finiteness of automorphism groups	48
3.4	Quotients by finite groups	50
3.5	Restriction of (co-)gauge-fixing data to boundaries	51
3.6	Extension of (co-)gauge-fixing data from boundaries	52
3.7	Pushforwards and pullbacks of (co-)gauge-fixing data	54
3.8	Fibre products of (co-)gauge-fixing data	56
3.9	Effective gauge-fixing data and co-gauge-fixing data	69
4	Kuranishi homology and cohomology	78
4.1	Classical homology and cohomology	78
4.2	(Effective) Kuranishi homology	82
4.3	Morphisms between singular and Kuranishi homology	85
4.4	Kuranishi cohomology and Poincaré duality for $\partial Y = \emptyset$	88
4.5	The case $\partial Y \neq \emptyset$, and relative Kuranishi (co)homology	95
4.6	Effective Kuranishi cohomology	100
4.7	Products on (effective) Kuranishi (co)homology	102
4.8	Kuranishi (co)homology as a bivariant theory on (co)chains	108
4.9	Why we need (co-)gauge-fixing data	113
5	Kuranishi bordism and cobordism	115
5.1	Classical bordism and cobordism groups	115
5.2	Kuranishi bordism groups	117
5.3	Morphisms of Kuranishi bordism groups	119
5.4	Kuranishi cobordism and Poincaré duality	122
5.5	Products on Kuranishi (co)bordism	125
5.6	Orbifold strata and operators $\Pi^{\Gamma, \rho}$	128
5.7	How large are Kuranishi (co)bordism groups?	137

6	Applications in Symplectic Geometry	141
6.1	Moduli spaces of closed Riemann surfaces	141
6.2	Gromov–Witten invariants	148
6.3	Integrality properties of Gromov–Witten invariants	151
6.4	Moduli spaces of stable holomorphic discs	158
6.5	Choosing co-gauge-fixing data for the $\overline{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$	162
6.6	Lagrangian Floer cohomology for one Lagrangian	169
6.7	Open Gromov–Witten invariants	174
6.8	Conclusions	177
A	Tent functions	178
A.1	Tent functions on compact manifolds	180
A.2	Tent functions on orbifolds	199
A.3	Tent functions on Kuranishi chains $[X, \mathbf{f}, \mathbf{G}], [X, \mathbf{f}, \underline{\mathbf{G}}]$	203
B	Proof that $H_*^{\text{si}}(Y; R) \cong KH_*^{\text{ef}}(Y; R)$	217
B.1	Step 1: cutting X_a into X_{ac} with trivial stabilizers	222
B.2	Step 2: deforming X_{ac} to compact manifolds $\tilde{X}_{ac}$	225
B.3	Step 3: triangulating the $\tilde{X}_a$ with simplices $\sigma_{ac}(\Delta_k)$	235
B.4	Step 4: $\Pi_{\text{si}}^{\text{ef}} : H_k^{\text{si}}(Y; R) \rightarrow KH_k^{\text{ef}}(Y; R)$ is injective	238
C	Proof that $KH_*(Y; R) \cong KH_*^{\text{ef}}(Y; R)$	241
C.1	Step 1: cutting the $X_a, \tilde{X}_d$ into small pieces $X_{ac}, \tilde{X}_{df}$	250
C.2	Step 2: lifting to $\tilde{X}_{ac}$ with trivial stabilizers	262
C.3	Step 3: lifting to a cycle in $\tilde{K}C_k(Y; R)$	264
C.4	Step 4: lifting to a cycle in $\tilde{K}C_k^{\text{ef}}(Y; R)$	270
C.5	Step 5: $\Pi_{\text{ef}}^{\text{Kh}} : KH_*^{\text{ef}}(Y; R) \rightarrow KH_*(Y; R)$ is injective	280
D	Identification of products $\cup, \cap, \bullet$	283
	References	285

1 Introduction

The goal of this book is to develop new tools for use in areas of symplectic geometry concerned with ‘counting’ J -holomorphic curves — Gromov–Witten theory, Lagrangian Floer cohomology, contact homology, Symplectic Field Theory and so on — and elsewhere, such as in the ‘string topology’ of Chas and Sullivan [11]. Moduli spaces $\overline{\mathcal{M}}$ of (stable) J -holomorphic curves Σ in a symplectic manifold (M, ω) are generally not smooth manifolds, but *Kuranishi spaces*, topological spaces equipped with a *Kuranishi structure*, as in Fukaya and Ono [25]. To ‘count’ the number of curves in $\overline{\mathcal{M}}$ one generally perturbs $\overline{\mathcal{M}}$ to construct a singular homology class called a *virtual moduli cycle* or *chain*. Several versions of this virtual moduli cycle construction exist [24, 25, 37, 49, 52, 53, 55, 65, 70].

A great deal of the technical complexity of these areas is due to the uneasy interplay between Kuranishi spaces, the raw geometric data, and virtual moduli

cycles and chains in singular homology, the abstract perturbations. This is particularly true in areas like Lagrangian Floer cohomology in which the moduli spaces $\bar{\mathcal{M}}$ have boundaries and corners, and one must consider compatibility of virtual moduli chains at the boundaries. Anyone who has seriously tried to read the mammoth Fukaya et al. [24] will appreciate this.

Let Y be an orbifold, and R a commutative ring or $\mathbb{Q}$ -algebra. The basic idea of this book is to define a new homology theory, the *Kuranishi homology* $KH_*(Y; R)$ of Y with coefficients in R . There are two variations, *Kuranishi homology* $KH_*(Y; R)$ for R a $\mathbb{Q}$ -algebra, and *effective Kuranishi homology* $KH_*^{\text{ef}}(Y; R)$ for R a commutative ring, such as $R = \mathbb{Z}$. Kuranishi homology $KH_*(Y; R)$ is better behaved at the chain level. Both are isomorphic to singular homology $H_*^{\text{si}}(Y; R)$, so they are not new topological invariants; the point is to use them as a replacement for singular homology, in situations where one wants to construct virtual chains for moduli spaces.

Kuranishi homology is the homology of a chain complex $(KC_*(Y; R), \partial)$ of *Kuranishi chains*. Elements of $KC_k(Y; R)$ are finite sums $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$ for $\rho_a \in R$, where $[X_a, \mathbf{f}_a, \mathbf{G}_a]$ is the isomorphism class of a triple $(X_a, \mathbf{f}_a, \mathbf{G}_a)$, with X_a a compact oriented Kuranishi space with boundary and corners, of virtual dimension $\text{vdim } X_a = k$, $\mathbf{f}_a : X_a \rightarrow Y$ a strongly smooth map, and $\mathbf{G}_a$ some *gauge-fixing data* for $(X_a, \mathbf{f}_a)$. The main point of $\mathbf{G}_a$ is to ensure that $\text{Aut}(X_a, \mathbf{f}_a, \mathbf{G}_a)$ is finite, since it turns out that if we allow infinite automorphism groups then the resulting homology groups are always zero.

We also define Poincaré dual theories of (*effective*) *Kuranishi cohomology* $KH^*(Y; R)$, $KH_{\text{ec}}^*(Y; R)$, which are isomorphic to compactly-supported cohomology $H_{\text{cs}}^*(Y; R)$, using a cochain complex $(KC^*(Y; R), d)$ of *Kuranishi cochains*. Elements of $KC^k(Y; R)$ are finite sums $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{C}_a]$, where X_a is a compact Kuranishi space with $\text{vdim } X_a = \dim Y - k$, $\mathbf{f}_a : X \rightarrow Y$ is a *strong submersion* with a relative orientation (*coorientation*), and $\mathbf{C}_a$ is *co-gauge-fixing data* for $(X_a, \mathbf{f}_a)$.

Here is an important difference between manifolds and Kuranishi spaces. If $f : X \rightarrow Y$ is a submersion of manifolds, then $\dim X \geq \dim Y$. Thus, if we defined $KC^k(Y; R)$ using manifolds X_a we would have $KC^k(Y; R) = 0$ for $k > 0$, which would be no use at all. However, if X is a Kuranishi space and $\mathbf{f} : X \rightarrow Y$ is a strong submersion, we need not have $\text{vdim } X \geq \dim Y$. In fact, if X is a manifold or Kuranishi space and $\mathbf{f} : X \rightarrow Y$ is (strongly) smooth, we can always change the Kuranishi structure of X to get a strong submersion $\mathbf{f}^Y : X^Y \rightarrow Y$. So strong submersions are easy to produce.

If $X, \tilde{X}$ are Kuranishi spaces and $\mathbf{f} : X \rightarrow Y$, $\tilde{\mathbf{f}} : \tilde{X} \rightarrow Y$ are strong submersions, there is a *fibre product* Kuranishi space $X \times_Y \tilde{X}$, with a strong submersion $\pi_Y : X \times_Y \tilde{X} \rightarrow Y$. Using this we define a *cup product* on $KC^*(Y; R)$ by

$$[X, \mathbf{f}, \mathbf{C}] \cup [\tilde{X}, \tilde{\mathbf{f}}, \tilde{\mathbf{C}}] = [X \times_Y \tilde{X}, \pi_Y, \mathbf{C} \times_Y \tilde{\mathbf{C}}], \quad (1)$$

where $\mathbf{C} \times_Y \tilde{\mathbf{C}}$ is a fibre product of co-gauge-fixing data. Then $\cup$ is associative, supercommutative, and compatible with d , and induces $\cup$ on $KH^*(Y; R)$ which is identified with the usual $\cup$ on $H_{\text{cs}}^*(Y; R)$ under $H_{\text{cs}}^*(Y; R) \cong KH^*(Y; R)$. There is also a *cap product* $\cap : KC_*(Y; R) \times KC^*(Y; R) \rightarrow KC_*(Y; R)$.

This illustrates the fact that *Kuranishi homology and cohomology are very well behaved at the (co)chain level*. For comparison, if one works with chains in singular homology, to take the intersection of two singular chains one must perturb them so they are transverse, and then triangulate the intersection by simplices, making arbitrary choices. Working with Kuranishi cochains, we just take cup products in a canonical way. So, *using Kuranishi (co)chains can eliminate problems with transversality*.

Here is how all this helps in the formation of virtual cycles and chains. Consider for example the moduli space $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ of genus g stable J -holomorphic curves with m marked points in a symplectic manifold M . Then $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ is a compact oriented Kuranishi space without boundary, and has *evaluation maps* $\mathbf{ev}_i : \bar{\mathcal{M}}_{g,m}(M, J, \beta) \rightarrow M$ for $i = 1, \dots, m$, strong submersions sending a curve to its i^{th} marked point. We define a virtual cycle for $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ to be $[\bar{\mathcal{M}}_{g,m}(M, J, \beta), \mathbf{ev}_1 \times \dots \times \mathbf{ev}_m, \mathbf{C}]$ in $KC^*(M^m; \mathbb{Q})$, where $\mathbf{C}$ is any choice of co-gauge-fixing data for $(\bar{\mathcal{M}}_{g,m}(M, J, \beta), \mathbf{ev}_1 \times \dots \times \mathbf{ev}_m)$.

Thus, *the moduli space is its own virtual cycle*. There is no need to perturb the moduli space, or triangulate it with simplices; we only have to choose (co-)gauge-fixing data, which is always possible, and is a much milder and less disruptive process. Choosing (co-)gauge-fixing data to satisfy prescribed boundary conditions or other compatibilities is much easier than trying to do the same with perturbations and singular chains. As a result, we can generally translate relationships between moduli spaces into *exact algebraic identities between virtual chains, at the (co)chain level, using cup and cap products*. This will lead to huge simplifications in the theory of Lagrangian Floer cohomology [4].

We also define *Kuranishi bordism* $KB_*(Y; R)$, and Poincaré dual *Kuranishi cobordism* $KB^*(Y; R)$. Elements of $KB_k(Y; R)$ are finite sums $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a]$ for $\rho_a \in R$, where $[X_a, \mathbf{f}_a]$ is the isomorphism class of $(X_a, \mathbf{f}_a)$, for X_a a compact oriented Kuranishi space without boundary with $\text{vdim } X_a = k$, and $\mathbf{f}_a : X_a \rightarrow Y$ is strongly smooth. The relations in $KB_k(Y; R)$ are that $[X, \mathbf{f}] + [X', \mathbf{f}'] = [X \amalg X', \mathbf{f} \amalg \mathbf{f}']$ and $[\partial W, \mathbf{e}|_{\partial W}] = 0$, for W compact oriented Kuranishi space with boundary but without corners, with $\text{vdim } W = k + 1$, and $\mathbf{e} : W \rightarrow Y$ strongly smooth. The definition for $KB^k(Y; R)$ is the same except that $\text{vdim } X_a = \dim Y - k$ and $\mathbf{f}_a : X_a \rightarrow Y$, $\mathbf{e} : W \rightarrow Y$ are cooriented strong submersions. We define a *cup product* on $KB^*(Y; R)$ by

$$[X, \mathbf{f}] \cup [\tilde{X}, \tilde{\mathbf{f}}] = [X \times_Y \tilde{X}, \pi_Y], \quad (2)$$

as for (1). Generally, to define or prove something for Kuranishi (co)bordism, we just take the analogue in Kuranishi (co)homology, restrict to Kuranishi spaces without boundary, and omit (co-)gauge-fixing data.

In fact we define five variations on this idea, by imposing extra conditions on or adding extra structure to $X_a, \mathbf{f}_a$. One version, *trivial stabilizers Kuranishi (co)bordism* $KB_*^{\text{tr}}, KB_*^{\text{trc}}(Y; R)$ has $KB_*^{\text{tr}}(Y; \mathbb{Z}) \cong MSO_*(Y)$ and $KB_*^{\text{trc}}(Y; \mathbb{Z}) \cong MSO_{\text{cs}}^*(Y)$ for manifolds Y , where $MSO_*(Y)$ and $MSO_{\text{cs}}^*(Y)$ are the conventional bordism and compactly-supported cobordism groups defined by Atiyah [5] using algebraic topology. However, the other four Kuranishi

(co)bordism theories are new topological invariants of Y . We show that they are *huge*: $KB_k(Y; R)$ has infinite rank over R for all nonempty orbifolds Y , commutative rings R without torsion (such as $R = \mathbb{Z}$ or $\mathbb{Q}$), and even $k \in \mathbb{Z}$, for instance. So it is probably not feasible to compute them and write them down.

The author intends these Kuranishi (co)bordism theories as new tools for studying (closed) Gromov–Witten theory. Since moduli spaces of (closed) J -holomorphic curves are compact, oriented Kuranishi spaces without boundary, they define classes in Kuranishi (co)bordism groups. So we can define new Gromov–Witten type invariants $GW_{g,m}^{\text{Kb}}(M, \omega, \beta)$ in $KB_*(M^m; \mathbb{Z})$ or $KB_*(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Z})$, which lift conventional Gromov–Witten invariants $GW_{g,m}^{\text{si}}(M, \omega, \beta)$ in $H_*(M^m; \mathbb{Q})$ or $H_*(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Q})$ in singular homology up to Kuranishi bordism.

There are two important points about these new invariants $GW_{g,m}^{\text{Kb}}(M, \omega, \beta)$. Firstly, they take values in very large groups, so they *contain more information than conventional Gromov–Witten invariants*. Much of this extra information concerns the *orbifold strata* of the moduli spaces, that is, for each finite group Γ the $GW_{g,m}^{\text{Kb}}(M, \omega, \beta)$ include data which ‘counts’ J -holomorphic curves in M with symmetry group Γ . Secondly, they are defined in groups over $\mathbb{Z}$ rather than $\mathbb{Q}$. This makes them a good context for studying *integrality properties* of Gromov–Witten invariants. In §6.3 we will outline a method for using them to prove the Gopakumar–Vafa Integrality Conjecture for Gromov–Witten invariants of Calabi–Yau 3-folds, which the author hopes to complete in [41].

There will also be applications of Kuranishi (co)homology or Kuranishi (co)bordism which are not related to J -holomorphic curves. For example, in [40] we will exploit the particularly good behaviour of Kuranishi (co)chains to give a nice formulation of the *String Topology* programme of Chas and Sullivan [11], in which the String Topology operations satisfy the desired identities *at the chain level*. Also, we show in §5.4 that for compact manifolds Y , trivial stabilizers Kuranishi cobordism $KH_{\text{trc}}^*(Y; \mathbb{Z})$ is isomorphic to cobordism $MSO^*(Y)$. This is the only differential-geometric description of $MSO^*(Y)$ that the author knows, and it may be useful to somebody.

We begin in Chapter 2 by defining Kuranishi structures and Kuranishi spaces, operations on them such as strongly smooth maps and fibre products, and geometric structures such as (co)orientations and (co)almost complex structures. This is partly taken from Fukaya and Ono [24, 25], partly modifying [24, 25], and partly new. Chapter 3 discusses *gauge-fixing data* and *co-gauge-fixing data*, vital ingredients of our Kuranishi (co)homology theories. Their main purpose is to make automorphism groups finite. If X is a compact Kuranishi space, $\mathbf{f} : X \rightarrow Y$ is strongly smooth, and $\mathbf{G}$ is gauge-fixing data for $(X, \mathbf{f})$, then $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ is finite, although $\text{Aut}(X, \mathbf{f})$ may be infinite.

Chapter 4 studies *Kuranishi (co)homology* $KH_*, KH^*(Y; R)$ for R a $\mathbb{Q}$ -algebra, and *effective Kuranishi (co)homology* $KH_*^{\text{ef}}, KH_{\text{ec}}^*(Y; R)$ for R a commutative ring. We define pushforwards h_* on chains and homology, and pullbacks h^* on cochains and cohomology, and cup and cap products $\cup, \cap$ — all the usual package for homology and cohomology theories, and all work at the

(co)chain level. Our main results are Theorems 4.8 and 4.9, which show that $KH_*(Y; R)$ and $KH_*^{\text{ef}}(Y; R)$ are isomorphic to singular homology $H_*^{\text{si}}(Y; R)$. Their proofs, deferred to Appendices A–C, take up more than one third of the book. We deduce using Poincaré duality that $KH^*(Y; R)$ and $KH_{\text{ec}}^*(Y; R)$ are isomorphic to compactly-supported cohomology $H_{\text{cs}}^*(Y; R)$. We show by example that if we omitted (co-)gauge-fixing data $\mathbf{G}$ from the chains $[X, \mathbf{f}, \mathbf{G}]$ then we would have $KH_*(Y; R) = KH^*(Y; R) = \{0\}$ for all Y , because of problems when $\text{Aut}(X, \mathbf{f})$ is infinite.

Chapter 5 defines two kinds of classical bordism groups $B_*, B_*^{\text{eo}}(Y; R)$, five kinds of Kuranishi bordism groups $KB_*, KB_*^{\text{eb}}, KB_*^{\text{tr}}, KB_*^{\text{ac}}, KB_*^{\text{eac}}(Y; R)$, and their Poincaré dual cobordism groups $KB^*, KB_{\text{ecb}}^*, KB_{\text{trc}}^*, KB_{\text{ca}}^*, KB_{\text{eca}}^*(Y; R)$. We define pushforwards, cup and cap products, and morphisms between the various (co)bordism groups, and from Kuranishi (co)bordism to (effective) Kuranishi (co)homology. We also define the *orbifold strata* $X^{\Gamma, \rho}$ of Kuranishi space X . Restricting to orbifold strata induces morphisms $\Pi^{\Gamma, \rho} : KB_*(Y; R) \rightarrow KB_{*-\dim \rho}(Y; R)$ of Kuranishi (co)bordism groups, which change degree. Using these composed with projection to homology, we prove that Kuranishi (co)bordism groups are very large.

In Chapter 6 we discuss applications to symplectic geometry. Our aim is not to give a complete treatment, but only to demonstrate the potential of our theories with a few examples, which may perhaps inspire readers to learn about Kuranishi (co)homology or Kuranishi (co)bordism, and use it in their own research. Sections 6.1–6.3 discuss closed Gromov–Witten invariants using Kuranishi bordism, and outline a method of attack on the Gopakumar–Vafa Integrality Conjecture.

Sections 6.4–6.7 discuss moduli spaces of J -holomorphic curves in M with boundary in a Lagrangian L , and open Gromov–Witten theory, using Kuranishi cochains and Kuranishi cohomology. In §6.6 we associate a gapped filtered A_∞ algebra $(\widehat{KC}^*(L; \Lambda_{\text{nov}}^*[1], \mathfrak{m})$ to L in a much simpler way than Fukaya et al. [24], and so define *bounding cochains* b and *Lagrangian Floer cohomology* $HF^*(L, b; \Lambda_{\text{nov}}^*)$ for one Lagrangian L . Then §6.7 gives a new definition of *open Gromov–Witten invariants* $OGW(M, L, \lambda)$ for a Lagrangian L with $[L] = 0 \in H_n(M; \mathbb{Q})$, together with a choice of bounding cochain b . Both $HF^*(L, b; \Lambda_{\text{nov}}^*)$ and the $OGW(M, L, \lambda)$ are independent of the choice of almost complex structure J , in a certain sense.

Finally, Appendices A–D prove the main results of Chapter 4, Theorems 4.8, 4.9 and 4.34. The proofs of Theorems 4.8 and 4.9 in Appendices B and C draw some inspiration from results of Fukaya and Ono on multisections and virtual cycles [25, §3, §6], [24, §A1], and also from Fukaya et al. [24, §30], who consider a situation in which transverse multisections must be carefully chosen on a collection of Kuranishi spaces, with compatibility conditions at the boundaries.

Significant parts of this book may read like a kind of commentary on Fukaya, Ono et al. [24, 25], which includes suggesting improvements and correcting errors. Lest this create the wrong impression, I would like to record here my great admiration for [24, 25], and my gratitude to Fukaya, Oh, Ohta and Ono for the creativity and long labour involved in writing them.

In [38] we provide a short ‘User’s Guide’ for Kuranishi (co)homology and Kuranishi (co)bordism, consisting of the most important definitions and results from this book, but omitting the proofs. Some readers may find it helpful to begin with [38] rather than this book.

Acknowledgements. I am grateful to Mohammed Abouzaid, Manabu Akaho, Kenji Fukaya, Ezra Getzler, Shinichiroh Matsuo, Yong-Geun Oh, Hiroshi Ohta, Kauru Ono, Paul Seidel, Ivan Smith and Dennis Sullivan for useful conversations. This research was partially supported by EPSRC grant EP/D07763X/1.

2 Kuranishi structures and Kuranishi spaces

We begin by discussing *Kuranishi spaces*, that is, topological spaces with Kuranishi structures. Kuranishi structures were introduced by Fukaya and Ono [25] as a tool for studying moduli problems, and further developed by Fukaya, Oh, Ohta and Ono [24]. We take ideas freely from both, and also give some new definitions and (in the author’s opinion) some improvements of those in [24, 25]. Since Kuranishi structures are so important in what follows, and are not yet in widespread use, we give plenty of detail.

2.1 Manifolds with corners and generalized corners

We shall define four classes of manifolds. The first three are entirely standard: *manifolds without boundary*, which are locally modelled on $\mathbb{R}^n$, *manifolds with boundary*, locally modelled on $\mathbb{R}^n$ or $[0, \infty) \times \mathbb{R}^{n-1}$, and *manifolds with corners*, locally modelled on $[0, \infty)^k \times \mathbb{R}^{n-k}$ for $k = 0, \dots, n$.

The fourth class, which we call *manifolds with generalized corners*, or *g-corners* for short, is new. It is similar to, but more general than, the notion of *locally polygonal set* used by Fukaya et al. [24, Def. 35.4]. We begin by defining the local models for manifolds with g-corners in $\mathbb{R}^n$, which we call *regions with g-corners*.

Definition 2.1. Let W be a nonempty, connected, open set in $\mathbb{R}^n$, and $f_1, \dots, f_N : W \rightarrow \mathbb{R}$ be smooth maps, such that:

- (a) For every subset $\{i_1, \dots, i_l\}$ of $\{1, \dots, N\}$, the subset $S_{\{i_1, \dots, i_l\}} = \{\mathbf{x} \in W : f_{i_1}(\mathbf{x}) = \dots = f_{i_l}(\mathbf{x}) = 0\}$ is an embedded submanifold of $\mathbb{R}^n$, without boundary.
- (b) In (a), for all $\mathbf{x} \in S_{\{i_1, \dots, i_l\}}$, we have $\langle df_{i_1}|_{\mathbf{x}}, \dots, df_{i_l}|_{\mathbf{x}} \rangle_{\mathbb{R}} = \{\alpha \in (\mathbb{R}^n)^* : \alpha|_{T_{\mathbf{x}} S_{\{i_1, \dots, i_l\}}} \equiv 0\}$ as vector subspaces of $(\mathbb{R}^n)^*$.

Define $U = \{\mathbf{x} \in W : f_i(\mathbf{x}) \geq 0, i = 1, \dots, N\}$. Then U is a closed subset of W . Let U° be the *interior* of U in W , and $\overline{U^\circ}$ the closure of U° in W . Suppose $U = \overline{U^\circ}$. Then we call U a *region with generalized corners*, or *g-corners*, in $\mathbb{R}^n$.

Definition 2.2. Let X be a paracompact Hausdorff topological space.

- (i) An n -dimensional chart on X without boundary is a pair (U, ϕ) , where U is an open subset in $\mathbb{R}^n$, and $\phi : U \rightarrow X$ is a homeomorphism with a nonempty open set $\phi(U)$ in X .
- (ii) An n -dimensional chart on X with boundary for $n \geq 1$ is a pair (U, ϕ) , where U is an open subset in $\mathbb{R}^n$ or in $[0, \infty) \times \mathbb{R}^{n-1}$, and $\phi : U \rightarrow X$ is a homeomorphism with a nonempty open set $\phi(U)$.
- (iii) An n -dimensional chart on X with corners for $n \geq 1$ is a pair (U, ϕ) , where U is an open subset in $[0, \infty)^k \times \mathbb{R}^{n-k}$ for some $0 \leq k \leq n$, and $\phi : U \rightarrow X$ is a homeomorphism with a nonempty open set $\phi(U)$.
- (iv) An n -dimensional chart on X with generalized corners, or g -corners, is a pair (U, ϕ) , where U is a region with g -corners in $\mathbb{R}^n$, and $\phi : U \rightarrow X$ is a homeomorphism with a nonempty open set $\phi(U)$.

These are increasing order of generality, that is, (i) $\Rightarrow$ (ii) $\Rightarrow$ (iii) $\Rightarrow$ (iv).

Let A, B be subsets of $\mathbb{R}^n$ and $\alpha : A \rightarrow B$ be continuous. We call α *smooth* if it extends to a smooth map between open neighbourhoods of A, B , that is, if there exists an open subset A' of $\mathbb{R}^n$ with $A \subseteq A'$ and a smooth map $\alpha' : A' \rightarrow \mathbb{R}^n$ with $\alpha'|_A \equiv \alpha$. If A is open we can take $A' = A$ and $\alpha' = \alpha$. We call $\alpha : A \rightarrow B$ a *diffeomorphism* if it is a homeomorphism and $\alpha : A \rightarrow B$, $\alpha^{-1} : B \rightarrow A$ are smooth.

Let $(U, \phi), (V, \psi)$ be n -dimensional charts on X , which may be without boundary, or with boundary, or with corners, or with g -corners. We call (U, ϕ) and (V, ψ) *compatible* if $\psi^{-1} \circ \phi : \phi^{-1}(\phi(U) \cap \psi(V)) \rightarrow \psi^{-1}(\phi(U) \cap \psi(V))$ is a diffeomorphism between subsets of $\mathbb{R}^n$, in the sense above.

An n -dimensional manifold structure on X , which may be *without boundary*, or *with boundary*, or *with corners*, or *with generalized corners (with g -corners)*, is a system of pairwise compatible n -dimensional charts $\{(U^i, \phi^i) : i \in I\}$ on X with $X = \bigcup_{i \in I} \phi^i(U^i)$, where the (U^i, ϕ^i) are without boundary, or with boundary, or with corners, or with g -corners, respectively. When we just refer to a manifold, we will usually mean a manifold with g -corners.

Remark 2.3. (a) Note that in Definition 2.1(a) we do not prescribe the dimension of $S_{\{i_1, \dots, i_l\}}$, and in particular, we do not require $\dim S_{\{i_1, \dots, i_l\}} = n - l$. Definition 2.1(b) implies that $\dim S_{\{i_1, \dots, i_l\}} \geq n - l$. If $\dim S_{\{i_1, \dots, i_l\}} = n - l$ then $df_{i_1}, \dots, df_{i_l}$ are linearly independent near $S_{\{i_1, \dots, i_l\}}$, but if $\dim S_{\{i_1, \dots, i_l\}} > n - l$ then $df_{i_1}, \dots, df_{i_l}$ are linearly dependent along $S_{\{i_1, \dots, i_l\}}$.

If $\dim S_{\{i_1, \dots, i_l\}} = n - l$ for all $\{i_1, \dots, i_l\}$ then U is an n -manifold with corners (not g -corners), locally modelled on $[0, \infty)^l \times \mathbb{R}^{n-l}$. Thus, we only generalize manifolds with corners by allowing $\dim S_{\{i_1, \dots, i_l\}} > n - l$.

(b) The purpose of requiring $U = \overline{U}^\circ$ in Definition 2.1 is to ensure U has no pieces of dimension less than n . For example, we could take $n = N = 2$, $W = \mathbb{R}^2$ and $f_1(x_1, x_2) = x_1$, $f_2(x_1, x_2) = -x_1$, which gives $U = \{(0, x_2) : x_2 \in \mathbb{R}\}$, with dimension 1 rather than 2. But then $U^\circ = \emptyset$, so $\overline{U}^\circ = \emptyset$, and $U \neq \overline{U}^\circ$.

(c) An important class of examples of regions with g -corners is given by taking $W = \mathbb{R}^n$ and each $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ to be linear. Then Definition 2.1(a),(b) hold

automatically, with each $S_{\{i_1, \dots, i_l\}}$ a linear subspace of $\mathbb{R}^n$, and U is a *polyhedral cone* in $\mathbb{R}^n$, which satisfies $U = \overline{U^\circ}$ if and only if $U^\circ \neq \emptyset$.

More generally, we can take $W = \mathbb{R}^n$ and each $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ to be affine, that is, of the form $f_i(x_1, \dots, x_n) = a_1 x_1 + \dots + a_n x_n + b$. Then U is a *convex polyhedron* in $\mathbb{R}^n$, not necessarily compact, which satisfies $U = \overline{U^\circ}$ if and only if $U^\circ \neq \emptyset$, that is, if and only if $\dim U = n$.

For example, the solid octahedron in $\mathbb{R}^3$

$$O = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : |x_1| + |x_2| + |x_3| \leq 1\} \quad (3)$$

is a region with g-corners, which may be defined using the eight affine functions

$$\begin{aligned} f_1(x_1, x_2, x_3) &= x_1 + x_2 + x_3 + 1, & f_2(x_1, x_2, x_3) &= x_1 + x_2 - x_3 + 1, \\ f_3(x_1, x_2, x_3) &= x_1 - x_2 + x_3 + 1, & f_4(x_1, x_2, x_3) &= x_1 - x_2 - x_3 + 1, \\ f_5(x_1, x_2, x_3) &= -x_1 + x_2 + x_3 + 1, & f_6(x_1, x_2, x_3) &= -x_1 + x_2 - x_3 + 1, \\ f_7(x_1, x_2, x_3) &= -x_1 - x_2 + x_3 + 1, & f_8(x_1, x_2, x_3) &= -x_1 - x_2 - x_3 + 1. \end{aligned}$$

This is not a manifold with corners, since four 2-dimensional faces of O meet at the vertex $(1, 0, 0)$, but three 2-dimensional faces of $[0, \infty)^3$ meet at the vertex $(0, 0, 0)$, so O near $(1, 0, 0)$ is not locally modelled on $[0, \infty)^3$ near $(0, 0, 0)$.

(d) To a first approximation, regions with g-corners and manifolds with corners look locally like convex polyhedrons in $\mathbb{R}^n$. Let U be a region with g-corners defined using $W, f_1, \dots, f_N$ in Definition 2.1, and let $\mathbf{y} \in U$. Define $\hat{W} = \mathbb{R}^n$ and $\hat{f}_1, \dots, \hat{f}_N : \mathbb{R}^n \rightarrow \mathbb{R}$ by $\hat{f}_i(\mathbf{x}) = f_i(\mathbf{y}) + df_i|_{\mathbf{y}}(\mathbf{x} - \mathbf{y})$, so that $\hat{f}_1, \dots, \hat{f}_N$ are the unique affine functions on $\mathbb{R}^n$ with $\hat{f}_i(\mathbf{x}) = f_i(\mathbf{x}) + O(|\mathbf{x} - \mathbf{y}|^2)$.

Let $\hat{U}$ be the region with g-corners defined using $\hat{W}, \hat{f}_1, \dots, \hat{f}_N$. Then $\hat{U}$ is a convex polyhedron in $\mathbb{R}^n$ (not necessarily compact), and $U, \hat{U}$ are basically isomorphic near $\mathbf{y}$. In particular, from Definition 2.1(b) it follows that for all subsets $\{i_1, \dots, i_l\} \subseteq \{1, \dots, N\}$, the submanifolds $S_{\{i_1, \dots, i_l\}}$ from $f_{i_1}, \dots, f_{i_l}$ and $\hat{S}_{\{i_1, \dots, i_l\}}$ from $\hat{f}_{i_1}, \dots, \hat{f}_{i_l}$ have the same tangent space at $\mathbf{y}$, and the same dimension near $\mathbf{y}$, and this implies that U and $\hat{U}$ have the same stratification into faces and corners of different dimensions near $\mathbf{y}$.

(e) However, despite (d), regions with g-corners in $\mathbb{R}^n$ are a *more general class of local models* than convex polyhedrons in $\mathbb{R}^n$, and yield a more general notion of manifolds with g-corners than we would have obtained by taking charts to be modelled on open sets in convex polyhedra in $\mathbb{R}^n$.

To see this, note that polyhedral cones C in $\mathbb{R}^n$ are classified up to the action of $\text{GL}(n, \mathbb{R})$ by both *discrete parameters* — the number of k -dimensional faces for $0 < k < n$, and how they intersect — and *continuous parameters* — if C has at least $n+2$ codimension one faces then the family of polyhedral cones with the same discrete structure as C up to $\text{GL}(n, \mathbb{R})$ is in general positive-dimensional. Classifying polyhedral cones in $\mathbb{R}^n$ up to $\text{GL}(n, \mathbb{R})$ is important, as this is what is preserved by our notion of diffeomorphism of subsets of $\mathbb{R}^n$ in Definition 2.2.

If U is a convex polyhedron in $\mathbb{R}^n$, and F is a connected, open dimension l face of U , then U is modelled on a fixed polyhedral cone C_F in $\mathbb{R}^n$ near $\mathbf{y}$ for all

$\mathbf{y}$ in F . However, if U is a region with g-corners in $\mathbb{R}^n$, and F is a connected, open, dimension l face of U for $l > 0$, then U is modelled on a polyhedral cone $C_{\mathbf{y}}$ in $\mathbb{R}^n$ near $\mathbf{y}$ for each $\mathbf{y}$ in F , where the discrete parameters of $C_{\mathbf{y}}$ up to $\text{GL}(n, \mathbb{R})$ are independent of $\mathbf{y}$, but *the continuous parameters can vary with $\mathbf{y}$* .

This can only happen when $n \geq 4$, since the local models for manifolds with g-corners are unique near strata of codimension 1 and 2, so we need both $l \geq 3$ and $\dim F = n - l > 0$ to have nontrivial continuous variation of $C_{\mathbf{y}}$ along F . The simplest example would be a region with g-corners in $\mathbb{R}^4$ in which five 3-dimensional faces come together along a 1-dimensional edge.

The point of the rather general, non-explicit local models of Definition 2.1 is to allow this local variation of the polyhedral cone model.

(f) The only places in this book where the details of the definition of manifolds with g-corners matter is in the material on *tent functions* in Appendix A, and their application in Appendices B and C to prove Theorems 4.8 and 4.9. For the rest of the book, all we need is that manifolds with g-corners have a well-behaved notion of boundary.

Next we define the *boundary* ∂X of a manifold X with boundary or (g-)corners. To motivate the definition, consider $[0, \infty)^2$ in $\mathbb{R}^2$, regarded as a manifold with corners. If we took $\partial([0, \infty)^2)$ to be the subset $[0, \infty) \times \{0\} \cup \{0\} \times [0, \infty)$ of $[0, \infty)^2$, then $\partial([0, \infty)^2)$ would not be a manifold with corners near $(0, 0)$. Instead, we take $\partial([0, \infty)^2)$ to be the *disjoint union* of the two boundary strata $[0, \infty) \times \{0\}$ and $\{0\} \times [0, \infty)$. This is a manifold with boundary, but now $\partial([0, \infty)^2)$ is *not a subset of* $[0, \infty)^2$, since two points in $\partial([0, \infty)^2)$ correspond to the vertex $(0, 0)$ in $[0, \infty)^2$.

Definition 2.4. Let U be a region with g-corners in $\mathbb{R}^n$, defined using $W, f_1, \dots, f_N$ as in Definition 2.1. If $u \in U$, let $\{i_1, \dots, i_l\}$ be the subset of i in $\{1, \dots, N\}$ for which $f_i(u) = 0$, and let $S_{\{i_1, \dots, i_l\}}$ be as in Definition 2.1. Then $S_{\{i_1, \dots, i_l\}}$ is an embedded submanifold of W containing u . Define the *codimension of u in U* to be $n - \dim S_{\{i_1, \dots, i_l\}}$.

It is not difficult to see that if u has codimension m , then $n - m$ is the maximum dimension of embedded submanifolds S in $\mathbb{R}^n$ with $u \in S \subseteq U \subseteq \mathbb{R}^n$, and this maximum is achieved by $S_{\{i_1, \dots, i_l\}}$. Therefore the codimension of u depends only on u and the set U up to diffeomorphism (in the sense of Definition 2.2), and is independent of the choice of $W, f_1, \dots, f_N$ used to define U .

Let X be an n -manifold with g-corners. Then X is covered by a system of charts $\{(U^i, \phi^i) : i \in I\}$, with each U^i a region with g-corners in $\mathbb{R}^n$. Let $x \in X$, so that $x = \phi^i(u)$ for some $i \in I$ and $u \in U^i$. Define the *codimension of x in X* to be the codimension of u in U^i . This is independent of the choice of chart (U^i, ϕ^i) , since if $x = \phi^i(u) = \phi^j(u')$, then U^i near u is diffeomorphic to U^j near u' using $\phi_j \circ (\phi_i)^{-1}$. For each $k = 0, \dots, n$, define $S_k(X)$ to be the subset of $x \in X$ of codimension k , the *codimension k stratum* of X . Then $X = \coprod_{k=0}^n S_k(X)$. One can show that each $S_k(X)$ has the structure of an $(n - k)$ -manifold without boundary, with closure $\overline{S_k(X)} = \bigcup_{l=k}^n S_l(X)$.

Let $x \in \overline{S_1(X)} = \bigcup_{l=1}^n S_l(X)$. A *local boundary component B of X at*

x is a local choice of connected component of $S_1(X)$ near x . That is, for each sufficiently small open neighbourhood U of x in X , B gives a choice of connected component V of $U \cap S_1(X)$ with $x \in \bar{V}$, and any two such choices U, V and U', V' must be compatible in the sense that $x \in \overline{(V \cap V')}$. It is easy to see that if $x \in S_1(X)$ then x has a unique local boundary component B , and if $x \in S_2(X)$ then x has exactly two local boundary components B_1, B_2 , and if $x \in S_k(X)$ for $k \geq 2$ then x has at least k local boundary components.

As a set, define the *boundary* ∂X of X by

$$\partial X = \{(x, B) : x \in \overline{S_1(X)}, B \text{ is a local boundary component for } X \text{ at } x\}.$$

Then ∂X naturally has the structure of an $(n-1)$ -manifold with g -corners. To see this, let $(x, B) \in \partial X$, and suppose (U^i, ϕ^i) is a chart on X with $x \in \phi^i(U^i)$. Then U^i is a region with g -corners in $\mathbb{R}^n$ defined using some $W, f_1, \dots, f_N$, with $x = \phi^i(u)$ for some unique $u \in U^i$. Now $(\phi^i)^{-1}(B)$ is a local boundary component of U^i at u . From Definition 2.1 it follows that $(\phi^i)^{-1}(B)$ coincides with $f_j^{-1}(0)$ near u for some $j = 1, \dots, N$.

Definition 2.1(b) with $l = 1$ and $i_1 = j$ then implies that df_j is nonzero on $f_j^{-1}(0) = S_{\{j\}}$, so $f_j^{-1}(0)$ is a closed, embedded $(n-1)$ -submanifold of U^j containing u . Thus we can choose a diffeomorphism $\psi : \tilde{W} \rightarrow f_j^{-1}(0) \subset U^j$ from an open subset $\tilde{W} \subset \mathbb{R}^{n-1}$ to an open neighbourhood $\psi(\tilde{W})$ of u in $f_j^{-1}(0)$. Define functions $\tilde{f}_1, \dots, \tilde{f}_{N-1} : \tilde{W} \rightarrow \mathbb{R}$ to be $\tilde{f}_a = f_a \circ \psi$ for $a = 1, \dots, j-1$ and $\tilde{f}_a = f_{a+1} \circ \psi$ for $a = j, \dots, N-1$. Then $\tilde{W}, \tilde{f}_1, \dots, \tilde{f}_{N-1}$ satisfy Definition 2.1 and define a region with corners $\tilde{U}$ in $\mathbb{R}^{n-1}$, with $\psi(\tilde{U})$ an open neighbourhood of u in $U^i \cap f_j^{-1}(0)$. Make $\tilde{W}, \tilde{U}$ smaller if necessary to ensure that $\psi(\tilde{U}) \subseteq \overline{(\phi^i)^{-1}(B)}$. Define $\tilde{\phi} : \tilde{U} \rightarrow \partial X$ by $\tilde{\phi}(\tilde{u}) = (\phi^i \circ \psi(\tilde{u}), B)$. Then $(\tilde{U}, \tilde{\phi})$ is an $(n-1)$ -dimensional chart on ∂X with g -corners, with $(x, B) \in \tilde{\phi}(\tilde{U})$. Any two such charts are compatible, by compatibility of the (U^i, ϕ^i) . Thus, covering ∂X with a system of these charts makes it into an $(n-1)$ -manifold with g -corners.

If X is a manifold without boundary then $\partial X = \emptyset$, and if X is a manifold with boundary then ∂X is a manifold without boundary. In this case we may regard ∂X as a subset of X , since each $x \in S_1(X)$ has a unique local boundary component B , so $(x, B) \mapsto x$ is injective. If X has corners (not g -corners) then ∂X has corners (not g -corners). If X is compact then ∂X is compact.

We define *smooth maps* and *submersions* of manifolds with g -corners.

Definition 2.5. Let X, Y be manifolds with g -corners, with $\dim X = m$, $\dim Y = n$, and $f : X \rightarrow Y$ be a continuous map. We call f *smooth* if whenever $(U, \phi), (V, \psi)$ are charts on X, Y respectively compatible with the manifold structures on X, Y , where $U \subseteq \mathbb{R}^m$ and $V \subseteq \mathbb{R}^n$ are regions with g -corners, then

$$\psi^{-1} \circ f \circ \phi : (f \circ \phi)^{-1}(\psi(V)) \longrightarrow \psi^{-1}(f \circ \phi(U)) \quad (4)$$

is a smooth map from $(f \circ \phi)^{-1}(\psi(V)) \subseteq U \subseteq \mathbb{R}^m$ to $\psi^{-1}(f \circ \phi(U)) \subseteq V \subseteq \mathbb{R}^n$. That is, as in Definition 2.2, $\psi^{-1} \circ f \circ \phi$ should extend to a smooth map between open subsets of $\mathbb{R}^m, \mathbb{R}^n$ containing $(f \circ \phi)^{-1}(\psi(V))$ and $\psi^{-1}(f \circ \phi(U))$.

We will define when a smooth map $f : X \rightarrow Y$ is a *submersion*. When $\partial X = \emptyset = \partial Y$, the definition is well-known. But when X, Y have boundary and (g-)corners, we will impose complicated extra conditions on f over $\partial^k X$ and $\partial^l Y$ for $k, l \geq 0$. Suppose first that $f : X \rightarrow Y$ is a smooth map and $\partial X = \emptyset = \partial Y$. We call f a *submersion* if for all $(U, \phi), (V, \psi)$ as above and all $u \in (f \circ \phi)^{-1}(\psi(V))$, the linear map $d(\psi^{-1} \circ f \circ \phi)|_u : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is surjective. This implies $m \geq n$. More geometrically, X, Y have tangent bundles TX, TY , and the derivative df is a morphism of vector bundles $df : TX \rightarrow f^*(TY)$, and f is a submersion if df is surjective.

Next, suppose that $f : X \rightarrow Y$ is a smooth map and $\partial Y = \emptyset$, but do not assume that $\partial X = \emptyset$. We call f a *submersion* if for all $k \geq 0$, the map $f|_{\partial^k X} : \partial^k X \rightarrow Y$ satisfies $d(f|_{\partial^k X}) : T(\partial^k X) \rightarrow f|_{\partial^k X}^*(TY)$ is surjective. This implies that $\partial^k X = \emptyset$ for all $k \geq m - n$. Also, if $f : X \rightarrow Y$ is a submersion then $f|_{\partial X} : X \rightarrow Y$ is a submersion.

Finally, suppose that $f : X \rightarrow Y$ is a smooth map, but do not assume that $\partial X = \emptyset$ or $\partial Y = \emptyset$. We give an *inductive definition* of what it means for f to be a submersion. We call f a submersion if:

- (a) $df : TX \rightarrow f^*(TY)$ is a surjective morphism of vector bundles over X ;
- (b) there is a unique decomposition $\partial X = \partial_+^f X \amalg \partial_-^f X$, where $\partial_+^f X, \partial_-^f X$ are open and closed subsets of ∂X , such that $f_+ = f|_{\partial_+^f X}$ maps $(\partial_+^f X)^\circ \rightarrow Y^\circ$, and $f_- = f|_{\partial_-^f X} : \partial_-^f X \rightarrow Y$ factors uniquely as $f_- = f_- \circ \iota$, where $f_- : \partial_-^f X \rightarrow \partial Y$ is a smooth map and $\iota : \partial Y \rightarrow Y$ is the natural immersion, and f_- maps $(\partial_+^f X)^\circ \rightarrow (\partial Y)^\circ$; and
- (c) $f_+ : \partial_+^f X \rightarrow Y$ and $f_- : \partial_-^f X \rightarrow \partial Y$ are both submersions.

This definition is recursive, as the definition of f being a submersion involves $f_\pm$ being submersions in (c), but since $\dim \partial_\pm X = \dim X - 1$, the definition makes sense by induction on $\dim X = 0, 1, 2, \dots$.

The following example may help readers to understand the definition. Let W, Y be manifolds with g-corners, and set $X = W \times Y$, with $f : X \rightarrow Y$ the projection $W \times Y \rightarrow Y$. Then f is a submersion in the sense above. We have $\partial X = \partial(W \times Y) = (\partial W \times Y) \amalg (W \times \partial Y)$. Then $\partial_+^f X = \partial W \times Y$, with $f_+ : \partial_+^f X \rightarrow Y$ the projection $\partial W \times Y \rightarrow Y$, and $\partial_-^f X = W \times \partial Y$, with $f_- : \partial_-^f X \rightarrow \partial Y$ the projection $W \times \partial Y \rightarrow \partial Y$. In general, one can think of submersions $f : X \rightarrow Y$ as being *locally* like a projection $W \times Y \rightarrow Y$ for W, Y manifolds with g-corners. If Y is a manifold with g-corners then $\text{id}_Y : Y \rightarrow Y$ is a submersion, taking $\partial_+^{\text{id}_Y} Y = \emptyset$ and $\partial_-^{\text{id}_Y} Y = \partial Y$.

One justification for this complicated definition of submersions is that if $f : X \rightarrow Y, f' : X' \rightarrow Y$ are smooth maps of manifolds with g-corners with at least one of f, f' a submersion, there is a well-behaved notion of *fibre product* $X \times_{f, Y, f'} X'$, which is a manifold with g-corners.

Proposition 2.6. *Suppose X, X', Y are manifolds with g-corners and $f : X \rightarrow Y, f' : X' \rightarrow Y$ are smooth maps, with at least one of f, f' a submersion. Define*

the **fibre product** $X \times_{f,Y,f'} X'$ or $X \times_Y X'$ by

$$X \times_{f,Y,f'} X' = X \times_Y X' = \{(p, p') \in X \times X' : f(p) = f'(p')\}. \quad (5)$$

Then $X \times_Y X'$ is a closed, embedded submanifold of $X \times X'$, and so $X \times_Y X'$ is a manifold with g-corners, which is compact if X, X' are compact.

Define maps $\pi_X : X \times_Y X' \rightarrow X$, $\pi_{X'} : X \times_Y X' \rightarrow X'$, and $\pi_Y : X \times_Y X' \rightarrow Y$ by $\pi_X : (p, p') \mapsto p$, $\pi_{X'} : (p, p') \mapsto p'$, and $\pi_Y : (p, p') \mapsto f(p) = f'(p')$. Then $\pi_X, \pi_{X'}, \pi_Y$ are smooth, and π_X is a submersion if f' is a submersion, and $\pi_{X'}$ is a submersion if f is a submersion, and π_Y is a submersion if both f', f are submersions.

If $\partial Y = \emptyset$ then we have a natural diffeomorphism

$$\partial(X \times_{f,Y,f'} X') \cong (\partial X \times_{f|_{\partial X}, Y, f'} X') \amalg (X \times_{f,Y,f'|_{\partial X}} \partial X'), \quad (6)$$

where if f is a submersion then $f|_{\partial X} : \partial X \rightarrow Y$ is a submersion, and the same for f' , so the fibre products in (6) make sense.

In general, allowing $\partial Y \neq \emptyset$, if f is a submersion then

$$\partial(X \times_{f,Y,f'} X') \cong (\partial_+^f X \times_{f_+, Y, f'} X') \amalg (X \times_{f,Y,f'|_{\partial X}} \partial X'), \quad (7)$$

and if f' is a submersion then

$$\partial(X \times_{f,Y,f'} X') \cong (\partial X \times_{f|_{\partial X}, Y, f'_+} X') \amalg (X \times_{f,Y,f'_+} \partial_+^{f'} X'), \quad (8)$$

and if both f, f' are submersions then both (7)–(8) hold, and also

$$\begin{aligned} \partial(X \times_{f,Y,f'} X') &\cong (\partial_+^f X \times_{f_+, Y, f'} X') \amalg (X \times_{f,Y,f'_+} \partial_+^{f'} X') \\ &\quad \amalg (\partial_-^f X \times_{f_-, \partial Y, f'_-} \partial_-^{f'} X'), \end{aligned} \quad (9)$$

$$\text{with } \partial_+^{\pi_Y}(X \times_{f,Y,f'} X') \cong (\partial_+^f X \times_{f_+, Y, f'} X') \amalg (X \times_{f,Y,f'_+} \partial_+^{f'} X'), \quad (10)$$

$$\text{and } \partial_-^{\pi_Y}(X \times_{f,Y,f'} X') \cong \partial_-^f X \times_{f_-, \partial Y, f'_-} \partial_-^{f'} X'. \quad (11)$$

Note that Proposition 2.6 would not be true if in defining submersions $f : X \rightarrow Y$ we had not imposed conditions over $\partial^k X$ and $\partial^l Y$ for all $k, l \geq 0$, since otherwise $X \times_Y X'$ might fail to be a manifold with g-corners on strata coming from $\partial^k X$ or $\partial^{k'} X'$. Thus we do need the complex, inductive definition in Definition 2.5. The proof of Proposition 2.6 is straightforward. The main thing is to check that $X \times_Y X'$ is a *submanifold* of $X \times X'$, by working with coordinate charts on X, X' . The submersion conditions on $\partial^k X, \partial^{k'} X'$ ensure that $X \times_Y X'$ intersects the parts of the boundary and corners of $X \times X'$ coming from $\partial^k X, \partial^{k'} X'$ suitably transversely.

A manifold X with (g-)corners has a natural involution $\sigma : \partial^2 X \rightarrow \partial^2 X$.

Definition 2.7. Let X be an n -manifold with g-corners. Then $\partial^2 X = \partial(\partial X)$ is an $(n-2)$ -manifold with g-corners. Points of $\partial^2 X$ are of the form $((x, B_1), B')$, where $(x, B_1) \in \partial X$, so that $x \in X$ and B_1 is a local boundary component for X at x , and B' is a local boundary component for ∂X at (x, B_1) . We would like to understand B' in terms of local boundary components for X at x . Let U be a sufficiently small open neighbourhood of $x \in X$. Then B_1 determines a connected component V_1 of $U \cap S_1(X)$ with $x \in \bar{V}_1$, and B' determines a connected component V' of $U \cap \bar{V}_1 \cap S_2(X)$.

From above, if $y \in S_2(X)$ then there are exactly two local boundary components for X at y . Applying this to $y \in V'$ with the same U , we find that there exists a unique second connected component V_2 of $U \cap S_1(X)$ with $V_1 \neq V_2$ such that V' is a subset of $U \cap \bar{V}_2 \cap S_2(X)$. This V_2 determines a unique local boundary component B_2 of X at x , with $B_1 \neq B_2$. Conversely, B_1, B_2 determine B' , since V' is the unique connected component of $U \cap \bar{V}_1 \cap \bar{V}_2 \cap S_2(X)$ with $x \in \bar{V}'$. Let us write $B_1 \cap B_2$ for B' . Then each point in $\partial^2 X$ may be uniquely written as $((x, B_1), B_1 \cap B_2)$, where $x \in X$ and B_1, B_2 are distinct local boundary components of X at x which intersect in codimension 2 in X .

Define $\sigma : \partial^2 X \rightarrow \partial^2 X$ by $\sigma : ((x, B_1), B_1 \cap B_2) \mapsto ((x, B_2), B_2 \cap B_1)$. Then σ is a *natural, smooth, free involution* of $\partial^2 X$. If X is oriented, so that ∂X and $\partial^2 X$ are oriented, then σ is orientation-reversing.

These involutions σ are important in problems of extending data from ∂X to X . We call this the *Extension Principle*, and we state it for arbitrary maps, for continuous functions or sections, and for smooth functions and sections.

Principle 2.8. (Extension Principle)(a) Let X be a manifold with g-corners, P a nonempty set, and $H : \partial X \rightarrow P$ an arbitrary map. Then a necessary and sufficient condition for there to exist $G : X \rightarrow P$ with $G|_{\partial X} \equiv H$ is that $H|_{\partial^2 X}$ is invariant under $\sigma : \partial^2 X \rightarrow \partial^2 X$ in Definition 2.7, that is, $H|_{\partial^2 X} \circ \sigma \equiv H|_{\partial^2 X}$. **(b)** Let X be a manifold with g-corners, $E \rightarrow X$ a vector bundle, $\zeta : \partial X \rightarrow \mathbb{R}$ a continuous map, and t a continuous section of $E|_{\partial X}$. Then necessary and sufficient conditions for there to exist continuous $\eta : X \rightarrow \mathbb{R}$ with $\eta|_{\partial X} \equiv \zeta$, and a continuous section s of E with $s|_{\partial X} \equiv t$, are that $\zeta|_{\partial^2 X}$ and $t|_{\partial^2 X}$ should be invariant under $\sigma : \partial^2 X \rightarrow \partial^2 X$. Here $\sigma : \partial^2 X \rightarrow \partial^2 X$ has a natural lift $\hat{\sigma} : E|_{\partial^2 X} \rightarrow E|_{\partial^2 X}$, and $t|_{\partial^2 X}$ is invariant under σ means that $t|_{\partial^2 X} \circ \sigma \equiv \hat{\sigma} \circ t|_{\partial^2 X}$. **(c)** Let X be a manifold with corners (*not* g-corners), $E \rightarrow X$ a vector bundle, $\zeta : \partial X \rightarrow \mathbb{R}$ be smooth, and t a smooth section of $E|_{\partial X}$. Then necessary and sufficient conditions for there to exist smooth $\eta : X \rightarrow \mathbb{R}$ with $\eta|_{\partial X} \equiv \zeta$, and a smooth section s of E with $s|_{\partial X} \equiv t$, are that $\zeta|_{\partial^2 X}$ and $t|_{\partial^2 X}$ should be invariant under $\sigma : \partial^2 X \rightarrow \partial^2 X$.

The point is that for manifolds with (g-)corners, the ‘inclusion’ map $\iota : \partial X \rightarrow X$ taking $\iota : (x, B) \mapsto x$ is *not injective* over $S_k(X)$ for $k \geq 2$. By the ‘restriction’ $G|_{\partial X}$, we mean $G \circ \iota$. Clearly, in (a) there can only exist $G : X \rightarrow P$ with $G|_{\partial X} = H$ if whenever $p \neq q \in \partial X$ with $\iota(p) = \iota(q)$ we have $H(p) = H(q)$. Then p, q lie over $\partial^2 X$, and it is easy to show that $H(p) = H(q)$ for all such p, q if and only if $H|_{\partial^2 X}$ is σ -invariant.

This establishes necessity in (a), and for (b),(c) the argument is the same. Sufficiency in (a) is more-or-less trivial, we can take $G|_{X^\circ} \equiv p \in P$. For (b),(c), we prove local sufficiency near any point of X that lifts to ∂X , then join local choices of η, s with a partition of unity to get a global choice.

Note that if X has g-corners, not corners, then in (c) for $\zeta|_{\partial^2 X}, t|_{\partial^2 X}$ to be σ -invariant is a necessary but *not* sufficient for ζ, t to extend smoothly to X . As an example, take X to be the octahedron O in (3), and let $\zeta : \partial O \rightarrow \mathbb{R}$ be smooth with $\zeta|_{\partial^2 X} \equiv \zeta|_{\partial^2 X} \circ \sigma$. Then at the vertex $(1, 0, 0)$ of O , we can prescribe the derivatives of ζ along the four edges of O meeting at $(1, 0, 0)$ independently. However, if $\zeta = \eta|_{\partial O}$ with η smooth then these four derivatives would be determined by $d\eta|_{(1,0,0)}$, which has only three parameters, a contradiction.

That is, *the smooth Extension Principle fails for manifolds with g-corners*. This will cause problems in proofs in Appendices A–C. To get round them, we develop a notion of *piecewise smooth extensions* in §A.1.4, which do work for manifolds and orbifolds with g-corners.

Some very similar ideas can be found in Fukaya et al. [24], which is not surprising, as they also have to solve essentially the same problems that we need the Extension Principle for. They prove a result close to Principle 2.8(c) in [24, Lem. 30.121], and in [24, Rem. 30.122] they note that this result does not hold for polygons, which is their way of saying that the smooth Extension Principle fails for manifolds with g-corners, and that instead one can work with piecewise smooth multisections, just as we use piecewise smooth extensions for manifolds with g-corners.

2.2 Orbifolds and orbibundles

Next we introduce *orbifolds*. Some useful references are Adem, Leida and Ruan [1], the appendix in Chen and Ruan [12], and Satake's original paper [66].

Definition 2.9. Let X be a paracompact Hausdorff topological space. For orbifolds *without boundary*, an *n-dimensional orbifold chart on X* is a triple (U, Γ, ϕ) , where Γ is a finite group with a linear representation on $\mathbb{R}^n$, U is a Γ -invariant open subset in $\mathbb{R}^n$, and $\phi : U/\Gamma \rightarrow X$ is a homeomorphism with a nonempty open set $\phi(U/\Gamma)$ in X .

For orbifolds *with boundary* we also allow orbifold charts of the form (U, Γ, ϕ) , where Γ is a finite group with a linear representation on $\mathbb{R}^n$ preserving the subset $[0, \infty) \times \mathbb{R}^{n-1}$, U is a Γ -invariant open subset in $[0, \infty) \times \mathbb{R}^{n-1}$, and $\phi : U/\Gamma \rightarrow X$ is a homeomorphism with a nonempty open set $\phi(U/\Gamma)$ in X .

For orbifolds *with corners* we also allow orbifold charts (U, Γ, ϕ) , where Γ is a finite group with a linear representation on $\mathbb{R}^n$ preserving the subset $[0, \infty)^k \times \mathbb{R}^{n-k}$ for some $0 \leq k \leq n$, U is a Γ -invariant open subset in $[0, \infty)^k \times \mathbb{R}^{n-k}$, and $\phi : U/\Gamma \rightarrow X$ is a homeomorphism with a nonempty open set $\phi(U/\Gamma)$ in X .

For orbifolds *with generalized corners (g-corners)* we also allow orbifold charts (U, Γ, ϕ) , where Γ is a finite group with a linear representation on $\mathbb{R}^n$, U is a Γ -invariant region with g-corners in $\mathbb{R}^n$, and $\phi : U/\Gamma \rightarrow X$ is a homeomorphism with a nonempty open set $\phi(U/\Gamma)$ in X .

For such a chart (U, Γ, ϕ) , let $p \in \text{Im } \phi \subseteq X$, choose $u_p \in U$ with $\phi(\Gamma u_p) = p$, and define $\Gamma_p = \{\gamma \in \Gamma : \gamma u_p = u_p\}$. Choose a Γ_p -invariant open neighbourhood U_p of u_p in U such that if $\gamma \in \Gamma$ and both $u, \gamma u$ lie in U_p then $\gamma \in \Gamma_p$; this holds if U_p is sufficiently small. Then the map $U_p/\Gamma_p \rightarrow (\Gamma U_p)/\Gamma \subseteq U/\Gamma$ given by $\Gamma_p u \mapsto \Gamma u$ is a homeomorphism. Define $\phi_p : U_p/\Gamma_p \rightarrow X$ by $\phi_p(\Gamma_p u) = \phi(\Gamma u)$. Then ϕ_p is a homeomorphism with its image, so (U_p, Γ_p, ϕ_p) is an orbifold chart on X . We call (U_p, Γ_p, ϕ_p) an *induced chart* of (U, Γ, ϕ) for p .

Let (U, Γ, ϕ) and (V, Δ, ψ) be n -dimensional orbifold charts on X . We say (U, Γ, ϕ) , (V, Δ, ψ) are *compatible* if for each $p \in \phi(U/\Gamma) \cap \psi(V/\Delta)$ there exist induced charts (U_p, Γ_p, ϕ_p) and (V_p, Δ_p, ψ_p) for p , an isomorphism of finite groups $\rho_p : \Gamma_p \rightarrow \Delta_p$, and a ρ_p -equivariant diffeomorphism $\sigma_p : U_p \rightarrow V_p$, such that $\phi_p = \psi_p \circ (\sigma_p)_*$, where $(\sigma_p)_* : U_p/\Gamma_p \rightarrow V_p/\Delta_p$ is the induced homeomorphism.

An *n -dimensional orbifold structure* on X is a system of n -dimensional orbifold charts $\{(U^i, \Gamma^i, \phi^i) : i \in I\}$ on X such that $X = \bigcup_{i \in I} \phi^i(U^i/\Gamma^i)$ and $(U^i, \Gamma^i, \phi^i), (U^j, \Gamma^j, \phi^j)$ are pairwise compatible for all $i, j \in I$. We call X with its n -dimensional orbifold structure an *n -dimensional orbifold* (either *without boundary*, or *with boundary*, or *with corners*, or *with g-corners*, according to which classes of orbifold charts above we allow). When we just refer to an orbifold, we usually mean an orbifold with g-corners.

If X is an n -dimensional orbifold and $p \in X$ then for some orbifold chart (U, Γ, ϕ) on X we have $p = \phi(\Gamma u_p)$ for $u_p \in U$. Define the *stabilizer group* $\text{Stab}(p)$ of p to be the finite group $\{\gamma \in \Gamma : \gamma u_p = u_p\}$. Up to isomorphism, this is independent of choices of (U, Γ, ϕ) and u_p . The representation of $\text{Stab}(p)$ on $T_{u_p} U \cong \mathbb{R}^n$ is also independent of choices, up to isomorphisms of $\text{Stab}(p)$ and automorphisms of $\mathbb{R}^n$.

Generalizing Definition 2.4, we can define the *boundary* ∂X of an orbifold with (g-)corners, which is also an orbifold with (g-)corners. Following Definition 2.7 we can define a natural involution $\sigma : \partial^2 X \rightarrow \partial^2 X$. This σ may not be free in the orbifold case, but if X is oriented then σ is orientation-reversing. Principle 2.8 holds, with orbifolds in place of manifolds.

Remark 2.10. (a) In defining orbifolds, some authors require the group Γ to act *effectively* on $\mathbb{R}^n$ in orbifold charts (U, Γ, ϕ) , that is, every non-identity element in Γ must act nontrivially on $\mathbb{R}^n$. Our definition is more general, and allows orbifolds such as $\{0\}/\Gamma$ in which generic points have nontrivial stabilizer groups. We will call orbifolds in which Γ acts effectively in orbifold charts (U, Γ, ϕ) *effective orbifolds*. They will be discussed in §3.9, and will be important in defining effective Kuranishi (co)homology.

One reason for allowing non-effective orbifolds is that some important examples of moduli spaces are not effective. For example, the moduli space $\bar{\mathcal{M}}_{1,1}$ of Deligne–Mumford stable Riemann surfaces of genus one with one marked point, to be discussed in §6.1, is a non-effective orbifold, whose generic points have stabilizer group $\mathbb{Z}_2$.

(b) Suppose (U, Γ, ϕ) is an orbifold chart on X with corners or g-corners, and $u \in S_k(U)$ for $k \geq 2$ so that there are $l \geq k$ local boundary components $B_1, \dots, B_l$ of U at u , and $\Delta \subset \Gamma$ is the subgroup of Γ fixing u . Then we

allow Δ to permute $B_1, \dots, B_l$, we do not require Δ to fix each local boundary component of U at u . Thus, the stabilizer group of $\phi(\Gamma u)$ in X is Δ , but the stabilizer group of $\phi(\Gamma(u, B_i))$ in ∂X is the subgroup of Δ fixing B_i , for $i = 1, \dots, l$. So, under $\iota : \partial X \rightarrow X$ mapping $(x, B) \mapsto x$, the induced morphisms of stabilizer groups $\iota_* : \text{Stab}((x, B)) \rightarrow \text{Stab}(x)$ are injective, but need not be isomorphisms.

In Definition 5.15, for Γ a finite group and ρ an isomorphism class of non-trivial Γ -representations, we will define the *orbifold stratum* $X^{\Gamma, \rho}$ of an orbifold X . It is an immersed suborbifold of X , with points (x, λ) for $x \in X$ and $\lambda : \Gamma \rightarrow \text{Stab}_X(x)$ an injective group morphism. The fact above has the important consequence that *for orbifolds with corners or g-corners, restricting to orbifold strata does not commute with taking boundaries*, that is, $\partial(X^{\Gamma, \rho}) \neq (\partial X)^{\Gamma, \rho}$.

In Chapter 5 we will discuss *Kuranishi (co)bordism groups* $KB_*, KB^*(Y; R)$. We will show that $KB_*, KB^*(Y; R)$ are *very large*, as for ‘chains’ $[X, \mathbf{f}]$ there is invariant information in orbifold strata $X^{\Gamma, \rho}$ for all Γ, ρ with $|\Gamma|$ odd, which can be recovered from $KB_*, KB^*(Y; R)$. However, this does not apply to Kuranishi (co)homology. Since restricting to orbifold strata does not commute with boundaries for orbifolds and Kuranishi spaces with (g-)corners, for Kuranishi (co)chains $[X, \mathbf{f}, \mathbf{G}]$ or $[X, \mathbf{f}, \mathbf{C}]$, no information from orbifold strata X^Γ survives in Kuranishi (co)homology. This is implicit in the proof of the isomorphism of (effective) Kuranishi homology with singular homology in Appendices A–C, see in particular §A.2.

Definition 2.11. Let X, Y be orbifolds, and $f : X \rightarrow Y$ a continuous map. We call f *smooth* if for all $p \in X$ and $q = f(p)$ in Y there are induced orbifold charts $(U_p, \Gamma_p, \phi_p), (V_q, \Delta_q, \psi_q)$ on X, Y for p, q , a morphism of finite groups $\rho_{pq} : \Gamma_p \rightarrow \Delta_q$, and a ρ_{pq} -equivariant smooth map $\sigma_{pq} : U_p \rightarrow V_q$, such that $f \circ \phi_p(\Gamma_p u) = \psi_q(\Delta_q \sigma_{pq}(u))$ for all $u \in U_p$. (**Warning:** this definition is oversimplified and requires modification, as in Remark 2.12 below.)

We call f a *submersion* if all the σ_{pq} are submersions of manifolds, in the sense of Definition 2.5. We call f an *embedding* if f is injective, all the ρ_{pq} are isomorphisms, and all the σ_{pq} are embeddings. We call f a *diffeomorphism* if it is a homeomorphism, all the ρ_{pq} are isomorphisms, and all the σ_{pq} are diffeomorphisms with their images. If f is a diffeomorphism then f^{-1} is a diffeomorphism.

If X, Y, Z are orbifolds and $f : X \rightarrow Y, g : Y \rightarrow Z$ are smooth then $g \circ f : X \rightarrow Z$ is smooth. If f is smooth and $p \in X$ with $q = f(p)$ in Y there is a natural group homomorphism $f_* : \text{Stab}(p) \rightarrow \text{Stab}(q)$, unique up to conjugation in $\text{Stab}(p), \text{Stab}(q)$, given by $f_* = \rho_{pq}$ above under the isomorphisms $\text{Stab}(p) \cong \Gamma_p, \text{Stab}(q) \cong \Delta_q$. Embeddings and diffeomorphisms induce isomorphisms f_* on stabilizer groups.

Remark 2.12. This definition of smooth maps of orbifolds is based on Satake [66, p. 361], who introduced orbifolds (then called *V-manifolds*). However, there is a problem with it, discussed by Adem et al. [1, p. 23-4, p. 47-50]: smooth maps of orbifolds defined this way do not have all the properties one wants, in particular, if $f : X \rightarrow Y$ is a smooth map of orbifolds and $E \rightarrow Y$ is an orbifold

vector bundle (see Definition 2.13), then the pullback orbifold vector bundle $f^*(E) \rightarrow X$ may not be well-defined.

The problem is this: if X, Y, f, p, q and some choices of orbifold charts (U_p, Γ_p, ϕ_p) , (V_q, Δ_q, ψ_q) are as above, then the ρ_{pq} -equivariant smooth map $\sigma_{pq} : U_p \rightarrow V_q$ satisfying $f \circ \phi_p(\Gamma_p u) = \psi_q(\Delta_q \sigma_{pq}(u))$ for $u \in U_p$ may not be uniquely defined. For instance, if Δ_q is abelian and $\delta \in \Delta_q$ then $\sigma'_{pq} = \delta \circ \sigma_{pq}$ is also a ρ_{pq} -equivariant smooth map $\sigma'_{pq} : U_p \rightarrow V_q$ with $f \circ \phi_p(\Gamma_p u) = \psi_q(\Delta_q \sigma'_{pq}(u))$ for all $u \in U_p$, but we may have $\sigma'_{pq} \neq \sigma_{pq}$. Thus, if we write down some construction involving the σ_{pq} , and the result is not independent of the choice of σ_{pq} , then the construction is not well-defined. This holds for pullbacks $f^*(E)$ of orbifold vector bundles, and also for fibre products of orbifolds below.

There is a solution to this problem. Using their ‘groupoid’ point of view on orbifolds (for them, a *groupoid* is roughly speaking equivalent to an orbifold with a particular choice of atlas of charts (U, Γ, ϕ)), Moerdijk and Pronk [57, §5] define *strong maps* of orbifolds. Also, Chen and Ruan [12, §4.4] define *good maps* of orbifolds, also with choices of atlases. Both strong and good maps have well-behaved pullbacks of orbifold vector bundles, and fibre products. Lupercio and Uribe [54, Prop. 5.1.7] prove that these notions of strong maps and good maps coincide. Adem et al. [1, §2.4] call strong/good maps *morphisms of orbifolds*. One can also use the theory of *stacks* from algebraic geometry: as in [54, §8.4], an orbifold is basically a Deligne–Mumford stack over smooth manifolds, and then strong and good maps correspond to morphisms of stacks.

In fact the notion of smooth map of orbifolds we want to work with is that of strong maps [57], good maps [12] or orbifold morphisms [1], rather than the weaker notion of Satake [66]. But this is not a central issue for us, and to work in one of the frameworks of [1, 12, 57] would increase notation, and sacrifice clarity and geometric intuition. So we will have our cake and eat it, by making the following convention: strong/good maps are equivalent to Satake’s smooth maps together with some *extra data*, and this extra data enables us to make unique choices of σ_{pq} in Definition 2.11, in a functorial way. We will leave this extra data implicit, but use it to choose σ_{pq} uniquely whenever we need to, such as in Definition 2.14 below. This convention entitles us to ignore the problem of non-uniqueness of σ_{pq} .

Our definition of orbifold vector bundles is equivalent to the *good orbifold vector bundles* of Chen and Ruan [12, §4.3].

Definition 2.13. Let X be an orbifold. An *orbifold vector bundle*, or *orbibundle*, $E \rightarrow X$ with fibre $\mathbb{R}^k$ is an orbifold E called the *total space* and a smooth map $\pi : E \rightarrow X$ called the *projection*, such that:

- (a) for each orbifold chart (U, Γ, ϕ) on X there is an orbifold chart $(E_U, \Gamma, \hat{\phi})$ on E with $\hat{\phi}(E_U/\Gamma) = \pi^{-1}(\phi(U/\Gamma))$, and a Γ -equivariant submersion $\pi_U : E_U \rightarrow U$ with $\pi \circ \hat{\phi}(\Gamma \hat{e}) = \phi(\Gamma \pi_U(\hat{e}))$ for all $\hat{e} \in E_U$, such that E_U has the structure of a vector bundle over U with fibre $\mathbb{R}^k$ and projection $\pi_U : E_U \rightarrow U$, and the Γ -action on E_U preserves the vector bundle structure.

- (b) Let (U, Γ, ϕ) , (V, Δ, ψ) , $p \in \phi(U/\Gamma) \cap \psi(V/\Delta)$, (U_p, Γ_p, ϕ_p) , (V_p, Δ_p, ψ_p) , $\rho_p : \Gamma_p \rightarrow \Delta_p$ and $\sigma_p : U_p \rightarrow V_p$ be as in Definition 2.9, and (U, Γ, ϕ) , (V, Δ, ψ) lift to $(E_U, \Gamma, \hat{\phi})$, $(E_V, \Delta, \hat{\psi})$ on E with projections $\pi_U : E_U \rightarrow U$, $\pi_V : E_V \rightarrow V$ as in (a). Set $E_{U_p} = \pi_U^{-1}(U_p)$, $E_{V_p} = \pi_V^{-1}(V_p)$, so that E_{U_p}, E_{V_p} are vector bundles over U_p, V_p with projections $\pi_U : E_{U_p} \rightarrow U_p$, $\pi_V : E_{V_p} \rightarrow V_p$. Define $\hat{\phi}_p : E_{U_p}/\Gamma_p \rightarrow E$, $\hat{\psi}_p : E_{V_p}/\Delta_p \rightarrow E$ by $\hat{\phi}_p(\Gamma_p \hat{e}) = \hat{\phi}(\Gamma \hat{e})$, $\hat{\psi}_p(\Delta_p \hat{f}) = \hat{\psi}(\Delta \hat{f})$, so that $(E_{U_p}, \Gamma_p, \hat{\phi}_p)$, $(E_{V_p}, \Delta_p, \hat{\psi}_p)$ are induced charts for E . Then there must exist a ρ_p -equivariant isomorphism of vector bundles $\hat{\sigma}_p : E_{U_p} \rightarrow E_{V_p}$ over U_p, V_p , such that $\pi_V \circ \hat{\sigma}_p = \sigma_p \circ \pi_U$ and $\hat{\phi}_p = \hat{\psi}_p \circ (\hat{\sigma}_p)_*$.

A *section* s of an orbibundle $\pi : E \rightarrow X$ is a smooth map $s : X \rightarrow E$ with $\pi \circ s = \text{id}_X$, the identity on X . For any n -dimensional orbifold X we may define the *tangent bundle* TX , an orbibundle with fibre $\mathbb{R}^n$. Morphisms (or isomorphisms, embeddings, ...) of orbibundles may be defined in the obvious way, as smooth maps (or diffeomorphisms, embeddings, ...) of the total spaces of the bundles, which lift to morphisms (or isomorphisms, embeddings, ...) of vector bundles $E_U \rightarrow U$ on orbifold charts $(E_U, \Gamma, \hat{\phi})$.

In Proposition 2.6 we defined *fibre products* of manifolds. We now extend this to orbifolds. There are some subtleties to do with stabilizer groups which mean that even as a set, the orbifold fibre product $X \times_Y X'$ is not given by (5). For orbifolds without boundary our definition is equivalent to that of Moerdijk and Pronk [57, §5], but because of our convention in Remark 2.12 we can present it in a more explicit, differential-geometric way.

Definition 2.14. Let X, X', Y be orbifolds and $f : X \rightarrow Y$, $f' : X' \rightarrow Y$ be smooth maps, at least one of which is a submersion. Then for $p \in X$ and $p' \in X'$ with $f(p) = q = f'(p')$ in Y we have homomorphisms of stabilizer groups $f_* : \text{Stab}(p) \rightarrow \text{Stab}(q)$, $f'_* : \text{Stab}(p') \rightarrow \text{Stab}(q)$. Thus we can form the double coset space

$$\begin{aligned} & f_*(\text{Stab}(p)) \backslash \text{Stab}(q) / f'_*(\text{Stab}(p')) \\ &= \{ \{ f_*(\gamma) \delta f'_*(\gamma') : \gamma \in \text{Stab}(p), \gamma' \in \text{Stab}(p') \} : \delta \in \text{Stab}(q) \}. \end{aligned}$$

As a set, we define

$$\begin{aligned} X \times_Y X' &= \{ (p, p', \Delta) : p \in X, p' \in X', f(p) = f'(p'), \\ & \Delta \in f_*(\text{Stab}(p)) \backslash \text{Stab}(f(p)) / f'_*(\text{Stab}(p')) \}. \end{aligned} \quad (12)$$

We give this the structure of an orbifold as follows. Suppose $p \in X$, $p' \in X'$ and $q \in Y$ with $f(p) = y = f'(p')$, and let (U_p, Γ_p, ϕ_p) , $(U_{p'}, \Gamma_{p'}, \phi_{p'})$, (V_q, Δ_q, ψ_q) be induced orbifold charts, so that $\Gamma_p \cong \text{Stab}(p)$, $\Gamma_{p'} \cong \text{Stab}(p')$, $\Delta_q \cong \text{Stab}(q)$. Making $U_p, U_{p'}$ smaller if necessary, there exist homomorphisms $\rho_{pq} : \Gamma_p \rightarrow \Delta_q$ and $\rho'_{p'q} : \Gamma_{p'} \rightarrow \Delta_q$ identified with $f_* : \text{Stab}(p) \rightarrow \text{Stab}(q)$ and $f'_* : \text{Stab}(p') \rightarrow \text{Stab}(q)$, and $\rho_{pq}, \rho'_{p'q}$ -equivariant smooth maps $\sigma_{pq} : U_p \rightarrow V_q$

and $\sigma'_{p'q} : U'_{p'} \rightarrow V_q$ at least one of which is a submersion, inducing f, f' on $\text{Im } \phi_p$ and $\text{Im } \phi'_{p'}$. Define an orbifold chart on $X \times_Y X'$ by

$$((U_p \times U_{p'}) \times_{\sigma_{pq} \times \sigma'_{p'q}, V_p \times V_q, \pi_q} (V_q \times \Delta_q), \Gamma_p \times \Gamma'_{p'}, \psi_{pp'q}). \quad (13)$$

Here the first term in (13) is a fibre product of smooth manifolds, as in Proposition 2.6. Define $\pi_q : (V_q \times \Delta_q) \rightarrow V_q \times V_q$ by $\pi_q(v, \delta) \mapsto (v, \delta \cdot v)$. The action of $\Gamma_p \times \Gamma'_{p'}$ on the first term in (13) is induced by the obvious action on $U_p \times U_{p'}$, and the action on $V_q \times \Delta_q$ by $(\gamma, \gamma') : (v, \delta) \mapsto (\rho_{pq}(\gamma) \cdot v, \rho'_{p'q}(\gamma') \delta \rho_{pq}(\gamma)^{-1})$.

Defining the map $\psi_{pp'q}$ from the first term of (13) to the set (12) is tricky, because $\text{Stab}(p), \text{Stab}(p'), \text{Stab}(q)$ and the morphisms f_*, f'_* between them are only really defined up to conjugation. (Actually, one needs the ideas of Remark 2.12 at this point.) Let (u, u', v, δ) lie in the first term of (13), with $\phi_p(\Gamma_p u) = x$, $\phi'_{p'}(\Gamma'_{p'} u') = x'$ and $\psi_q(\Delta_q v) = y$. Then we have natural identifications $\text{Stab}(x) \cong \text{Stab}_{\Gamma_p}(u)$, $\text{Stab}(x') \cong \text{Stab}_{\Gamma'_{p'}}(u')$ and $\text{Stab}(y) \cong \text{Stab}_{\Delta_q}(v)$. But $f_* : \text{Stab}(x) \rightarrow \text{Stab}(y)$ is naturally identified with $\rho'_{p'q} : \text{Stab}_{\Gamma'_{p'}}(u') \rightarrow \text{Stab}_{\Delta_q}(\delta \cdot v)$. To identify $\text{Stab}_{\Delta_q}(\delta \cdot v)$ with our model $\text{Stab}_{\Delta_q}(v)$ for $\text{Stab}(y)$ we must choose $\epsilon \in \Delta_q$ such that $\epsilon \cdot v = \delta \cdot v$. (Note that just fixing $\epsilon = \delta$ is not a good choice; we want the choice of ϵ to depend only on the points $v, \delta \cdot v$, not on δ .) With these identifications, we can define $\psi_{pp'q}$ by

$$\begin{aligned} \psi_{pp'q} : (u, u', v, \delta) \mapsto \\ (\psi_p(\Gamma_p u), \psi'_{p'}(\Gamma'_{p'} u'), \rho_{pq}(\text{Stab}_{\Gamma_p}(u)) \delta \epsilon^{-1} \rho'_{p'q}(\text{Stab}_{\Gamma'_{p'}}(u'))). \end{aligned}$$

The orbifold charts (13) are compatible, and make $X \times_Y X'$ into an orbifold. The projections $\pi_X : X \times_Y X' \rightarrow X$, $\pi_{X'} : X \times_Y X' \rightarrow X'$ and $\pi_Y : X \times_Y X' \rightarrow Y$ are smooth maps, and the analogue of Proposition 2.6 holds. If Y is a manifold, or more generally if all the maps $f_* : \text{Stab}(x) \rightarrow \text{Stab}(f(x))$ or $f'_* : \text{Stab}(x') \rightarrow \text{Stab}(f'(x'))$ are surjective, then as a topological space $X \times_Y X'$ is the topological space fibre product of X, X' over Y in (5). But in general the orbifold and topological space fibre products do not coincide.

2.3 Kuranishi neighbourhoods and coordinate changes

Let X be a paracompact Hausdorff topological space.

Definition 2.15. A *Kuranishi neighbourhood* (V_p, E_p, s_p, ψ_p) of $p \in X$ satisfies:

- (i) V_p is an orbifold, which may have boundary, corners, or g-corners;
- (ii) $E_p \rightarrow V_p$ is an orbifold vector bundle over V_p ;
- (iii) $s_p : V_p \rightarrow E_p$ is a smooth section; and
- (iv) ψ_p is a homeomorphism from $s_p^{-1}(0)$ to an open neighbourhood of p in X , where $s_p^{-1}(0)$ is the subset of V_p where the section s_p is zero.

We call E_p the *obstruction bundle*, and s_p the *Kuranishi map*.

Remark 2.16. Definition 2.15 follows [25, Def. 5.1], save that in (i) we allow the V_p to have boundary or (g-)corners which Fukaya and Ono do not, though in [25, Def. 17.3] they allow the V_p to have boundary, and in (iii) following [24] we require s_p to be *smooth*, whereas Fukaya and Ono allow the s_p to be only *continuous*. We discuss smoothness of s_p in Remark 2.19 below.

In [24, Def. A1.1] Fukaya et al. define a Kuranishi neighbourhood to be $(V_p, E_p, \Gamma_p, \psi_p, s_p)$, where V_p is a manifold possibly with boundary or corners, $E_p \rightarrow V_p$ a trivial vector bundle, Γ_p a finite group acting effectively on V_p and E_p , s_p a smooth Γ_p -equivariant section of E_p , and $\psi_p : s_p^{-1}(0)/\Gamma_p \rightarrow X$ a homeomorphism with its image. Then $(V_p/\Gamma_p, E_p/\Gamma_p, s_p/\Gamma_p, \psi_p)$ is a Kuranishi neighbourhood in our sense. One reason why we opted for the (V_p, E_p, s_p, ψ_p) form in this book is that much of Chapter 3 would not work for Kuranishi neighbourhoods of the form $(V_p, E_p, \Gamma_p, \psi_p, s_p)$.

Definition 2.17. Let $(V_p, E_p, s_p, \psi_p), (\tilde{V}_p, \tilde{E}_p, \tilde{s}_p, \tilde{\psi}_p)$ be two Kuranishi neighbourhoods of $p \in X$. We call $(V_p, \dots, \psi_p)$ and $(\tilde{V}_p, \dots, \tilde{\psi}_p)$ *isomorphic* with *isomorphism* $(\alpha, \hat{\alpha})$ if there exist a diffeomorphism $\alpha : V_p \rightarrow \tilde{V}_p$ and an isomorphism of orbibundles $\hat{\alpha} : E_p \rightarrow \alpha^*(\tilde{E}_p)$, such that $\tilde{s}_p \circ \alpha \equiv \hat{\alpha} \circ s_p$ and $\tilde{\psi}_p \circ \alpha \equiv \psi_p$.

We call $(V_p, \dots, \psi_p), (\tilde{V}_p, \dots, \tilde{\psi}_p)$ *equivalent* if there exist open neighbourhoods $U_p \subseteq V_p, \tilde{U}_p \subseteq \tilde{V}_p$ of $\psi_p^{-1}(p), \tilde{\psi}_p^{-1}(p)$ such that $(U_p, E_p|_{U_p}, s_p|_{U_p}, \psi_p|_{U_p})$ and $(\tilde{U}_p, \tilde{E}_p|_{\tilde{U}_p}, \tilde{s}_p|_{\tilde{U}_p}, \tilde{\psi}_p|_{\tilde{U}_p})$ are isomorphic.

Definition 2.18. Let (V_p, E_p, s_p, ψ_p) and (V_q, E_q, s_q, ψ_q) be Kuranishi neighbourhoods of $p \in X$ and $q \in \psi_p(s_p^{-1}(0))$ respectively. We call a pair $(\phi_{pq}, \hat{\phi}_{pq})$ a *coordinate change* from $(V_q, \dots, \psi_q)$ to $(V_p, \dots, \psi_p)$ if:

- (a) $\phi_{pq} : V_q \rightarrow V_p$ is a smooth embedding of orbifolds. When V_p, V_q have boundaries or (g-)corners, ϕ_{pq} maps $S_k(V_q) \rightarrow S_k(V_p)$ in the notation of Definition 2.4, for all $k \geq 0$. Thus ϕ_{pq} is compatible with boundaries and (g-)corners, and induces embeddings $\phi_{pq} : \partial^k V_q \rightarrow \partial^k V_p$ for all $k \geq 0$.
- (b) $\hat{\phi}_{pq} : E_q \rightarrow \phi_{pq}^*(E_p)$ is an embedding of orbibundles over V_q ;
- (c) $\hat{\phi}_{pq} \circ s_q \equiv s_p \circ \phi_{pq}$;
- (d) $\psi_q \equiv \psi_p \circ \phi_{pq}$; and
- (e) Choose an open neighbourhood W_{pq} of $\phi_{pq}(V_q)$ in V_p , and an orbifold vector subbundle F_{pq} of $E_p|_{W_{pq}}$ with $\phi_{pq}^*(F_{pq}) = \hat{\phi}_{pq}(E_q)$, as orbifold vector subbundles of $\phi_{pq}^*(E_p)$ over V_q . Write $\hat{s}_p : W_{pq} \rightarrow E_p/F_{pq}$ for the projection of $s_p|_{W_{pq}}$ to the quotient bundle E_p/F_{pq} . Now $s_p|_{\phi_{pq}(V_q)}$ lies in F_{pq} by (c), so $\hat{s}_p|_{\phi_{pq}(V_q)} \equiv 0$. Thus there is a well-defined derivative

$$d\hat{s}_p : N_{\phi_{pq}(V_q)} V_p \rightarrow (E_p/F_{pq})|_{\phi_{pq}(V_q)},$$

where $N_{\phi_{pq}(V_q)} V_p$ is the normal orbifold vector bundle of $\phi_{pq}(V_q)$ in V_p . Pulling back to V_q using ϕ_{pq} , and noting that $\phi_{pq}^*(F_{pq}) = \hat{\phi}_{pq}(E_q)$, gives

a morphism of orbifold vector bundles over V_q :

$$d\hat{s}_p : \frac{\phi_{pq}^*(TV_p)}{(d\phi_{pq})(TV_q)} \longrightarrow \frac{\phi_{pq}^*(E_p)}{\hat{\phi}_{pq}(E_q)}. \quad (14)$$

We require that (14) should be an *isomorphism* over $s_q^{-1}(0)$, and hence on an open neighbourhood of $s_q^{-1}(0)$ in V_q . Note that this forces $\dim V_p - \text{rank } E_p = \dim V_q - \text{rank } E_q$. Now since $s_q \equiv 0$ on $s_q^{-1}(0)$, we have $s_p \equiv 0$ on $\phi_{pq}(s_q^{-1}(0))$ by (c), and because of this, the restriction of $d\hat{s}_p$ in (14) to $s_q^{-1}(0)$ is *independent* of the choices of W_{pq} and F_{pq} . So the condition that (14) is an isomorphism over $s_q^{-1}(0)$ is independent of choices.

Remark 2.19. Definition 2.18 is based on Fukaya, Ono et al. [25, Def. 5.3], [24, Def. A1.3], with two important modifications. Firstly, the condition in (a) on ϕ_{pq} mapping $S_k(V_q) \rightarrow S_k(V_p)$ is new; it is necessary for the notion of a boundary or codimension k corner point to be preserved by coordinate changes, so that boundaries of Kuranishi spaces are well-behaved. Secondly, (e) is also new, and it replaces the notion in [25, Def. 5.6], [24, Def. A1.14] that a Kuranishi structure *has a tangent bundle*.

There is a possibility of confusion in the literature here. As well as the 1999 paper of Fukaya and Ono [25], there are currently three different versions of Fukaya et al. [24] in circulation, dated 2000, 2006 and 2008. Each of these four uses different definitions of Kuranishi space. In this book, references are to the 2008 version of [24]. In [25] and the 2000 and 2006 versions of [24], by saying that a Kuranishi structure ‘has a tangent bundle’, Fukaya et al. mean that for each coordinate change $(\phi_{pq}, \hat{\phi}_{pq})$ in the Kuranishi structure, we are *given* an isomorphism of orbibundles over V_q

$$\chi_{pq} : \frac{\phi_{pq}^*(TV_p)}{(d\phi_{pq})(TV_q)} \longrightarrow \frac{\phi_{pq}^*(E_p)}{\hat{\phi}_{pq}(E_q)}, \quad (15)$$

satisfying compatibilities under composition of coordinate changes. But these χ_{pq} need have no particular relation to $d\hat{s}_p$ in (14).

In the 2008 version of [24], the definition [24, Def. A1.14] that a Kuranishi structure ‘has a tangent bundle’ is basically equivalent to Definition 2.18(e). That is, Fukaya et al. identify the maps χ_{pq} in (15) with $d\hat{s}_p$, which they write as $d_{\text{fiber}}s_p$. This change is necessary, and important. With the definitions of ‘Kuranishi space with a tangent bundle’ used in [25] and the 2000, 2006 versions of [24], in which χ_{pq} is unrelated to $d\hat{s}_p$, the construction of virtual cycles and virtual classes is invalid. This is remarked upon in the 2008 version [24, A1.7(iv)], which also gives an example [24, Ex. A1.64] of bad behaviour using the previous definition of tangent bundles.

Our Definition 2.18(e), and the corresponding definition [24, Def. A1.14] in the 2008 version of [24], require the Kuranishi maps s_p to be *smooth*, as in [24, Def. A1.1], or at least C^1 , rather than merely continuous as in [25, Def. 5.1]. This is a significant issue because in constructing Kuranishi structures on moduli

spaces of J -holomorphic curves, it is straightforward to make the s_p continuous at singular curves, but one has to do much more work to arrange for the s_p to be smooth. We will return to this issue in Remark 6.5 and in §6.8.

2.4 Kuranishi structures

We can now define *Kuranishi structures* on topological spaces.

Definition 2.20. Let X be a paracompact Hausdorff topological space. A *germ of Kuranishi neighbourhoods* at $p \in X$ is an equivalence class of Kuranishi neighbourhoods (V_p, E_p, s_p, ψ_p) of p , for equivalence as in Definition 2.17.

Suppose (V_p, E_p, s_p, ψ_p) lies in such a germ. Then for any open neighbourhood U_p of $\psi_p^{-1}(p)$ in V_p , $(U_p, E_p|_{U_p}, s_p|_{U_p}, \psi_p|_{U_p})$ also lies in the germ. As a shorthand, we say that some condition on the germ *holds for sufficiently small* $(V_p, \dots, \psi_p)$ if whenever $(V_p, \dots, \psi_p)$ lies in the germ, the condition holds for $(U_p, \dots, \psi_p|_{U_p})$ for all sufficiently small U_p as above.

A *Kuranishi structure* κ on X assigns a germ of Kuranishi neighbourhoods for each $p \in X$ and a *germ of coordinate changes* between them in the following sense: for each $p \in X$, for all sufficiently small $(V_p, \dots, \psi_p)$ in the germ at p , for all $q \in \text{Im } \psi_p$, and for all sufficiently small $(V_q, \dots, \psi_q)$ in the germ at q , we are given a coordinate change $(\phi_{pq}, \hat{\phi}_{pq})$ from $(V_q, \dots, \psi_q)$ to $(V_p, \dots, \psi_p)$. These coordinate changes should be compatible with equivalence in the germs at p, q in the obvious way, and satisfy:

- (i) $\dim V_p - \text{rank } E_p$ is independent of p ; and
- (ii) if $q \in \text{Im } \psi_p$ and $r \in \text{Im } \psi_q$ then $\phi_{pq} \circ \phi_{qr} = \phi_{pr}$ and $\hat{\phi}_{pq} \circ \hat{\phi}_{qr} = \hat{\phi}_{pr}$.

We call $\text{vdim } X = \dim V_p - \text{rank } E_p$ the *virtual dimension* of the Kuranishi structure. Note that Definition 2.18(e) already implies that $\dim V_p - \text{rank } E_p$ is invariant under coordinate changes, so it is locally constant on X ; part (i) says it is globally constant.

A *Kuranishi space* (X, κ) is a topological space X with a Kuranishi structure κ . Usually we will just refer to X as the Kuranishi space, suppressing κ , except when we need to distinguish between different Kuranishi structures $\kappa, \tilde{\kappa}$ on the same topological space X .

We say that a Kuranishi space (X, κ) is *without boundary*, or *with boundary*, or *with corners*, or *with g-corners*, if for all $p \in X$ and all sufficiently small $(V_p, \dots, \psi_p)$ in the germ at p in κ , V_p is an orbifold without boundary, or with boundary, or with corners, or with g-corners, respectively. When we just refer to a Kuranishi space, we usually mean a Kuranishi space with g-corners.

Remark 2.21. Definition 2.20 is based on [25, Def. 5.3], [24, Def. A1.5], but since our definition of coordinate change includes the extra condition Definition 2.18(e), as in Remark 2.19, our notion of Kuranishi structure is *stronger* than that of Fukaya et al., and roughly corresponds to their definition of Kuranishi structure *with a tangent bundle* [25, Def. 5.6], [24, Def. A1.14]. Note too that [25] and the 2000 and 2006 versions of [24] define Kuranishi structures in terms of

germs of Kuranishi neighbourhoods, as above, but the 2008 version of [24] does not use germs.

An orbifold X is equivalent to a Kuranishi space with E_p the zero vector bundle and $s_p \equiv 0$ in all Kuranishi neighbourhoods (V_p, E_p, s_p, ψ_p) , and can be covered by the single Kuranishi neighbourhood $(X, X, 0, \text{id}_X)$.

Definition 2.22. Let (X, κ) be a Kuranishi space. A Kuranishi neighbourhood (V, E, s, ψ) on X is called *compatible with the Kuranishi structure κ* if for all $p \in \text{Im } \psi$ and for all sufficiently small Kuranishi neighbourhoods $(V_p, \dots, \psi_p)$ in the germ at p in κ , there exists a coordinate change $(\phi_p, \hat{\phi}_p)$ from $(V_p, \dots, \psi_p)$ to (V, E, s, ψ) .

Actually, when we use compatible Kuranishi neighbourhoods, we shall usually implicitly suppose that for all sufficiently small $(V_p, \dots, \psi_p)$ we are given a *particular choice* of coordinate change $(\phi_p, \hat{\phi}_p)$ from $(V_p, \dots, \psi_p)$ to (V, E, s, ψ) , and that these choices are compatible with coordinate changes $(\phi_{pq}, \hat{\phi}_{pq})$ in the germ of coordinate changes in κ in the obvious way. So probably these choices of $(\phi_p, \hat{\phi}_p)$ should be part of the definition.

Definition 2.20 implies that for $p \in X$, all sufficiently small $(V_p, \dots, \psi_p)$ in the germ at p in κ are compatible with κ . Thus, every Kuranishi space can be covered by compatible Kuranishi neighbourhoods (with particular choices of coordinate changes $(\phi_p, \hat{\phi}_p)$ as above). If X is compact we can choose a finite cover by compatible Kuranishi neighbourhoods.

We can also define *stabilizer groups* of points in a Kuranishi space.

Definition 2.23. Let X be a Kuranishi space, $p \in X$, and $(V_p, \dots, \psi_p)$ be a Kuranishi neighbourhood in the germ at p . Define the *stabilizer group* $\text{Stab}(p)$ of p to be the stabilizer group $\text{Stab}(\psi_p^{-1}(p))$ of the point $\psi_p^{-1}(p)$ in the orbifold V_p . By definition of equivalence in Definition 2.17, up to isomorphism this is independent of the choice of $(V_p, \dots, \psi_p)$, and so is well-defined.

2.5 Strongly smooth maps and strong submersions

The next definition is based on Fukaya and Ono [25, Def. 6.6]. The equivalent definition in [24, Def. A1.13] instead uses *good coordinate systems*, as in §3.1, but we prefer not to choose a good coordinate system unnecessarily.

Definition 2.24. Let X be a Kuranishi space, and Y a topological space. A *strongly continuous map* $\mathbf{f} : X \rightarrow Y$ consists of, for all $p \in X$ and all sufficiently small (V_p, E_p, s_p, ψ_p) in the germ of Kuranishi neighbourhoods at p , a choice of continuous map $f_p : V_p \rightarrow Y$, such that for all $q \in \text{Im } \psi_p$ and sufficiently small $(V_q, \dots, \psi_q)$ in the germ at q with coordinate change $(\phi_{pq}, \hat{\phi}_{pq})$ from $(V_q, \dots, \psi_q)$ to $(V_p, \dots, \psi_p)$ in the germ of coordinate changes, we have $f_p \circ \phi_{pq} = f_q$. Then $\mathbf{f}$ induces a continuous map $f : X \rightarrow Y$ in the obvious way.

If Y is an orbifold and all f_p are smooth, we call $\mathbf{f}$ *strongly smooth*. We call $\mathbf{f}$ a *strong submersion* if all the f_p are submersions, in the sense of Definitions 2.5

and 2.11 which includes conditions on f_p over $\partial^k V_p$ and $\partial^l Y$ for $k, l \geq 0$. In fact, in this book we will usually restrict to the case $\partial Y = \emptyset$ for simplicity, except in §4.5. Note that Fukaya and Ono [24, 25] use the notation *weakly submersive*, rather than strong submersion.

We call two strongly continuous or strongly smooth maps $\mathbf{f} : X \rightarrow Y$, $\mathbf{f}' : X \rightarrow Y$ *equal* if for all $p \in X$ and all sufficiently small $(V_p, \dots, \psi_p)$ in the germ at p , the maps $f_p : V_p \rightarrow Y$ and $f'_p : V_p \rightarrow Y$ are the same. Note that this allows there to exist $(V_p, \dots, \psi_p)$ in the germ at p such that only one of $f_p : V_p \rightarrow Y$ and $f'_p : V_p \rightarrow Y$ are given, or both are given but are not equal; if both are given, the definition of ‘sufficiently small’ requires only that $f_p|_{U_p} \equiv f'_p|_{U_p}$ for some open $\psi_p^{-1}(p) \in U_p \subseteq V_p$. Equal $\mathbf{f} : X \rightarrow Y$, $\mathbf{f}' : X \rightarrow Y$ induce the same continuous maps $f : X \rightarrow Y$, $f' : X \rightarrow Y$.

We shall need to generalize this to the case when Y is a Kuranishi space. As far as the author knows, the next two definitions are new.

Definition 2.25. Let $(V_p, E_p, s_p, \psi_p), (W_r, F_r, t_r, \xi_r)$ be Kuranishi neighbourhoods in Kuranishi spaces X, Y . A *smooth map* from $(V_p, \dots, \psi_p)$ to $(W_r, \dots, \xi_r)$ is a pair $(f_{rp}, \hat{f}_{rp})$ such that

- (a) $f_{rp} : V_p \rightarrow W_r$ is a smooth map of orbifolds;
- (b) $\hat{f}_{rp} : E_p \rightarrow f_{rp}^*(F_r)$ is a morphism of orbibundles over V_p ; and
- (c) $\hat{f}_{rp} \circ s_p \equiv t_r \circ f_{rp}$.

We call $(f_{rp}, \hat{f}_{rp})$ a *submersion* if f_{rp} is a submersion in the sense of Definitions 2.5 and 2.11, and $\hat{f}_{rp} : E_p \rightarrow f_{rp}^*(F_r)$ is surjective. Following Definition 2.17, we call $(f_{rp}, \hat{f}_{rp})$ an *equivalence* if there exist open neighbourhoods V'_p of $\psi_p^{-1}(p)$ in V_p and W'_r of $\xi_r^{-1}(r)$ in W_r such that $f_{rp}|_{V'_p} : V'_p \rightarrow W'_r$ is a diffeomorphism, and $\hat{f}_{rp}|_{V'_p} : E_p|_{V'_p} \rightarrow f_{rp}^*(F_r|_{W'_r})$ is an isomorphism of orbibundles.

A *strongly smooth map* $\mathbf{f} : X \rightarrow Y$ inducing a given continuous map $f : X \rightarrow Y$ consists of, for all p in X with $r = f(p)$ in Y , for all sufficiently small (W_r, F_r, t_r, ξ_r) in the germ of Kuranishi neighbourhoods at r , for all sufficiently small (V_p, E_p, s_p, ψ_p) in the germ at p , a smooth map $(f_{rp}, \hat{f}_{rp})$ from $(V_p, \dots, \psi_p)$ to $(W_r, \dots, \xi_r)$, such that if $(\phi_{pq}, \hat{\phi}_{pq})$ lies in the germ of coordinate changes on X then $f_{rp} \circ \phi_{pq} = f_{rq}$ and $\hat{f}_{rp} \circ \hat{\phi}_{pq} = \hat{f}_{rq}$, and if $(\phi_{rs}, \hat{\phi}_{rs})$ lies in the germ of coordinate changes on Y then $\phi_{rs} \circ f_{sp} = f_{rp}$ and $\hat{\phi}_{rs} \circ \hat{f}_{sp} = \hat{f}_{rp}$.

We call $\mathbf{f}$ a *strong submersion* if all the $(f_{rp}, \hat{f}_{rp})$ are submersions. We call $\mathbf{f}$ a *strong diffeomorphism* if $f : X \rightarrow Y$ is a homeomorphism and all the $(f_{rp}, \hat{f}_{rp})$ are equivalences. We call two strongly maps $\mathbf{f} : X \rightarrow Y$, $\mathbf{f}' : X \rightarrow Y$ *equal* if they induce the same continuous maps $f : X \rightarrow Y$, $f' : X \rightarrow Y$, and for all $p \in X$ and $r = f(p)$ in Y and all sufficiently small $(V_p, \dots, \psi_p), (W_r, \dots, \xi_r)$ in the germs at p, r in X, Y , we have $(f_{rp}, \hat{f}_{rp}) = (f'_{rp}, \hat{f}'_{rp})$.

Definition 2.26. Let X, Y, Z be Kuranishi spaces, and $\mathbf{f} : X \rightarrow Y$, $\mathbf{g} : Y \rightarrow Z$ be strongly smooth maps inducing continuous maps $f : X \rightarrow Y$, $g : Y \rightarrow Z$.

Define the *composition* $\mathbf{g} \circ \mathbf{f} : X \rightarrow Z$ as follows: let $p \in X$, $q = f(p) \in Y$ and $r = g(q) \in Z$, let $(V_p, \dots, \psi_p)$, $(V'_q, \dots, \psi'_q)$, $(V''_r, \dots, \psi''_r)$ be sufficiently small Kuranishi neighbourhoods in the germs at p, q, r in X, Y, Z , let $(f_{qp}, \hat{f}_{qp})$ be the smooth map from $(V_p, \dots, \psi_p)$ to $(V'_q, \dots, \psi'_q)$ in the germ of $\mathbf{f}$, and $(g_{rq}, \hat{g}_{rq})$ the smooth map from $(V'_q, \dots, \psi'_q)$ to $(V''_r, \dots, \psi''_r)$ in the germ of $\mathbf{g}$. Then the composition $(g_{rq} \circ f_{qp}, \hat{g}_{rq} \circ \hat{f}_{qp})$ is a smooth map from $(V_p, \dots, \psi_p)$ to $(V''_r, \dots, \psi''_r)$. It is easy to verify that the germ of all such compositions forms a strongly smooth map $\mathbf{g} \circ \mathbf{f} : X \rightarrow Z$, which induces the continuous map $g \circ f : X \rightarrow Z$. Clearly, if $\mathbf{f}, \mathbf{g}$ are strong submersions then so is $\mathbf{g} \circ \mathbf{f}$, and composition is associative, $(\mathbf{h} \circ \mathbf{g}) \circ \mathbf{f} = \mathbf{h} \circ (\mathbf{g} \circ \mathbf{f})$.

If $\mathbf{f} : X \rightarrow Y$ is a strong diffeomorphism inducing the homeomorphism $f : X \rightarrow Y$, it is not difficult to show that there exists a strong diffeomorphism $\mathbf{f}^{-1} : Y \rightarrow X$ inducing $f^{-1} : Y \rightarrow X$, such that the composition $\mathbf{f}^{-1} \circ \mathbf{f} : X \rightarrow X$ equals the identity on X , and $\mathbf{f} \circ \mathbf{f}^{-1} : Y \rightarrow Y$ equals the identity on Y , using the notion of *equal* strongly smooth maps in Definition 2.25.

2.6 Boundaries and fibre products

We now define the *boundary* ∂X of a Kuranishi space X , which is itself a Kuranishi space of dimension $\text{vdim } X - 1$. Our definition extends [25, Lem. 17.8] to the case of Kuranishi spaces with (g-)corners, and is modelled on Definition 2.4.

Definition 2.27. Let X be a Kuranishi space with g-corners. We shall define a Kuranishi space ∂X called the *boundary* of X . The points of ∂X are equivalence classes $[p, (V_p, \dots, \psi_p), B]$ of triples $(p, (V_p, \dots, \psi_p), B)$, where $p \in X$, $(V_p, \dots, \psi_p)$ lies in the germ of Kuranishi neighbourhoods at p , and B is a local boundary component of V_p at $\psi_p^{-1}(p)$, in the sense of Definition 2.4.

Two triples $(p, (V_p, \dots, \psi_p), B), (q, (\tilde{V}_q, \dots, \tilde{\psi}_q), \tilde{B})$ are *equivalent* if $p = q$, and the Kuranishi neighbourhoods $(V_p, \dots, \psi_p), (\tilde{V}_q, \dots, \tilde{\psi}_q)$ are equivalent so that we are given an isomorphism $(\alpha, \hat{\alpha}) : (U_p, \dots, \psi_p|_{U_p}) \rightarrow (\tilde{U}_q, \dots, \tilde{\psi}_q|_{\tilde{U}_q})$ for open $\psi_p^{-1}(p) \in U_p \subseteq V_p$ and $\tilde{\psi}_q^{-1}(q) \in \tilde{U}_q \subseteq \tilde{V}_q$, and $\alpha_*(B) = \tilde{B}$ near $\tilde{\psi}_q^{-1}(q)$.

We can define a unique natural *topology* and *Kuranishi structure* on ∂X , such that $(\partial V_p, E_p|_{\partial V_p}, s_p|_{\partial V_p}, \psi'_p)$ is a Kuranishi neighbourhood on ∂X for each Kuranishi neighbourhood $(V_p, \dots, \psi_p)$ on X , where $\psi'_p : (s_p|_{\partial V_p})^{-1}(0) \rightarrow \partial X$ is given by $\psi'_p : (q, B) \mapsto [\psi_p(q), (V_p, \dots, \psi_p), B]$ for $(q, B) \in \partial V_p$ with $s_p(q) = 0$. Then $\text{vdim}(\partial X) = \text{vdim } X - 1$, and ∂X is compact if X is compact.

Note that since ∂V_p is not actually a subset of V_p , but a set of points (v, B) for $v \in V_p$, writing $E_p|_{\partial V_p}$ and $s_p|_{\partial V_p}$ here is an abuse of notation. A more accurate way to say it is that we have an immersion $\iota : \partial V_p \rightarrow V_p$ acting by $\iota : (v, B) \mapsto v$, and by $E_p|_{\partial V_p}$ we mean $\iota^*(E_p)$, and by $s_p|_{\partial V_p}$ we mean $s_p \circ \iota$.

If X is a manifold with g-corners then Definition 2.7 gave a natural smooth free involution $\sigma : \partial^2 X \rightarrow \partial^2 X$, and Definition 2.9 noted this also works for orbifolds, though σ may not be free. In the same way, if X is a Kuranishi space there is a natural strong diffeomorphism $\sigma : \partial^2 X \rightarrow \partial^2 X$ with $\sigma^2 = \text{id}_X$, such that if (V_p, E_p, s_p, ψ_p) is a Kuranishi neighbourhood on X and $(\partial^2 V_p, \dots, \psi_p|_{\partial^2 V_p})$

is the induced Kuranishi neighbourhood on $\partial^2 X$, then σ is represented by $(\sigma_p, \hat{\sigma}_p) : (\partial^2 V_p, \dots, \psi_p|_{\partial^2 V_p}) \rightarrow (\partial^2 V_p, \dots, \psi_p|_{\partial^2 V_p})$, where $\sigma_p : \partial^2 V_p \rightarrow \partial^2 V_p$ is the involution for V_p , and $\hat{\sigma}_p$ its natural lift to $E_p|_{\partial^2 V_p} = \partial^2 E_p$. If X is oriented in the sense of §2.7 below, then $\sigma : \partial^2 X \rightarrow \partial^2 X$ is orientation-reversing.

Fukaya et al. [24, Def. A1.37] define *fibre products* of Kuranishi spaces when $\mathbf{f}, \mathbf{f}'$ are strong submersions to a manifold Y . We generalize to orbifolds Y . One can also generalize to Kuranishi spaces Y , but we will not need to.

Definition 2.28. Let X, X' be Kuranishi spaces, Y be a smooth orbifold, and $\mathbf{f} : X \rightarrow Y, \mathbf{f}' : X' \rightarrow Y$ be strongly smooth maps inducing continuous maps $f : X \rightarrow Y$ and $f' : X' \rightarrow Y$. Suppose at least one of $\mathbf{f}, \mathbf{f}'$ is a strong submersion. We shall define the *fibre product* $X \times_Y X'$ or $X \times_{\mathbf{f}, Y, \mathbf{f}'} X'$, a Kuranishi space. As a set, the underlying topological space $X \times_Y X'$ is given by (12), as for fibre products of orbifolds. The definition is modelled on Definition 2.14.

Let $p \in X, p' \in X'$ and $q \in Y$ with $f(p) = q = f'(p')$. Let $(V_p, E_p, s_p, \psi_p), (V_{p'}, E_{p'}, s_{p'}, \psi_{p'})$ be sufficiently small Kuranishi neighbourhoods in the germs at p, p' in X, X' , and $f_p : V_p \rightarrow Y, f_{p'} : V_{p'} \rightarrow Y$ be smooth maps in the germs of $\mathbf{f}, \mathbf{f}'$ at p, p' respectively. Define a Kuranishi neighbourhood on $X \times_Y X'$ by

$$(V_p \times_{f_p, Y, f_{p'}} V_{p'}, \pi_{V_p}^*(E_p) \oplus \pi_{V_{p'}}^*(E_{p'}), (s_p \circ \pi_{V_p}) \oplus (s_{p'} \circ \pi_{V_{p'}}), (\psi_p \circ \pi_{V_p}) \times (\psi_{p'} \circ \pi_{V_{p'}}) \times \chi_{pp'}). \quad (16)$$

Here $V_p \times_{f_p, Y, f_{p'}} V_{p'}$ is the fibre product of orbifolds from Definition 2.14, defined as at least one of $f_p, f_{p'}$ is a submersion, and $\pi_{V_p}, \pi_{V_{p'}}$ are the projections from $V_p \times_Y V_{p'}$ to $V_p, V_{p'}$. The final term $\chi_{pp'}$ in (16) maps the biquotient terms in (12) for $V_p \times_Y V_{p'}$ to the same terms in (12) for the set $X \times_Y X'$, so that for $v \in V_p$ and $v' \in V_{p'}$ with $f_p(v) = f(\psi_p(v)) = y = f_{p'}(v') = f'(\psi_{p'}(v'))$ in Y we define

$$\chi_{pp'}|_{(v, v')} : (f_p)_*(\text{Stab}(v)) \backslash \text{Stab}(y) / (f_{p'})_*(\text{Stab}(v')) \longrightarrow f_*(\text{Stab}(\psi_p(v))) \backslash \text{Stab}(y) / f'_*(\text{Stab}(\psi_{p'}(v')))$$

to be the map induced by the isomorphisms $(\psi_p)_* : \text{Stab}(v) \rightarrow \text{Stab}(\psi_p(v))$ and $(\psi_{p'})_* : \text{Stab}(v') \rightarrow \text{Stab}(\psi_{p'}(v'))$.

It is easy to verify that coordinate changes between Kuranishi neighbourhoods in X and X' induce coordinate changes between neighbourhoods (16). So the systems of germs of Kuranishi neighbourhoods and coordinate changes on X, X' induce such systems on $X \times_Y X'$. This gives a *Kuranishi structure* on $X \times_Y X'$, making it into a *Kuranishi space*. Clearly $\text{vdim}(X \times_Y X') = \text{vdim } X + \text{vdim } X' - \dim Y$, and $X \times_Y X'$ is compact if X, X' are compact.

2.7 Orientations and orientation conventions

We can now discuss *orientations* of Kuranishi spaces X . Our definition is basically equivalent to Fukaya et al. [24, Def. A1.17], noting as in Remarks 2.19 and 2.21 that our notion of Kuranishi structure is roughly equivalent to that of Kuranishi structure with tangent bundles in the 2008 version of [24].

Definition 2.29. Let X be a Kuranishi space. An *orientation* on X assigns, for all $p \in X$ and all sufficiently small Kuranishi neighbourhoods (V_p, E_p, s_p, ψ_p) in the germ at p , orientations on the fibres of the orbibundle $TV_p \oplus E_p$ varying continuously over V_p . These must be compatible with coordinate changes, in the following sense. Let $q \in \text{Im } \psi_p$, $(V_q, \dots, \psi_q)$ be sufficiently small in the germ at q , and $(\phi_{pq}, \hat{\phi}_{pq})$ be the coordinate change from $(V_q, \dots, \psi_q)$ to $(V_p, \dots, \psi_p)$ in the germ. Define $d\hat{s}_p$ near $s_q^{-1}(0) \subseteq V_q$ as in (14).

Locally on V_q , choose any orientation for the fibres of $\phi_{pq}^*(TV_p)/(\text{d}\phi_{pq})(TV_q)$, and let $\phi_{pq}^*(E_p)/\hat{\phi}_{pq}(E_q)$ have the orientation induced from this by the isomorphism $d\hat{s}_p$ in (14). These induce an orientation on $\frac{\phi_{pq}^*(TV_p)}{(\text{d}\phi_{pq})(TV_q)} \oplus \frac{\phi_{pq}^*(E_p)}{\hat{\phi}_{pq}(E_q)}$, which is independent of the choice for $\phi_{pq}^*(TV_p)/(\text{d}\phi_{pq})(TV_q)$. Thus, these local choices induce a natural orientation on the orbibundle $\frac{\phi_{pq}^*(TV_p)}{(\text{d}\phi_{pq})(TV_q)} \oplus \frac{\phi_{pq}^*(E_p)}{\hat{\phi}_{pq}(E_q)}$ near $s_q^{-1}(0)$. We require that in oriented orbibundles over V_q near $s_q^{-1}(0)$, we have

$$\begin{aligned} \phi_{pq}^*[TV_p \oplus E_p] \cong (-1)^{\dim V_q(\dim V_p - \dim V_q)} [TV_q \oplus E_q] \oplus \\ \left[\frac{\phi_{pq}^*(TV_p)}{(\text{d}\phi_{pq})(TV_q)} \oplus \frac{\phi_{pq}^*(E_p)}{\hat{\phi}_{pq}(E_q)} \right], \end{aligned} \quad (17)$$

where $TV_p \oplus E_p$ and $TV_q \oplus E_q$ have the orientations assigned by the orientation on X . An *oriented Kuranishi space* is a Kuranishi space with an orientation.

Suppose X, X' are oriented Kuranishi spaces, Y is an oriented orbifold, and $f : X \rightarrow Y$, $f' : X' \rightarrow Y$ are strong submersions. Then by §2.6 we have Kuranishi spaces ∂X and $X \times_Y X'$. These can also be given orientations in a natural way. We follow the orientation conventions of Fukaya et al. [24, §45].

Convention 2.30. First, our conventions for manifolds:

- (a) Let X be an oriented manifold with boundary ∂X . Then we define the orientation on ∂X such that $TX|_{\partial X} = \mathbb{R}_{\text{out}} \oplus T(\partial X)$ is an isomorphism of oriented vector bundles, where $\mathbb{R}_{\text{out}}$ is oriented by an outward-pointing normal vector to ∂X .
- (b) Let X, X', Y be oriented manifolds, and $f : X \rightarrow Y$, $f' : X' \rightarrow Y$ be submersions. Then $df : TX \rightarrow f^*(TY)$ and $df' : TX' \rightarrow (f')^*(TY)$ are surjective maps of vector bundles over X, X' . Choosing Riemannian metrics on X, X' and identifying the orthogonal complement of $\text{Ker } df$ in TX with the image $f^*(TY)$ of df , and similarly for f' , we have isomorphisms of vector bundles over X, X' :

$$TX \cong f^*(TY) \oplus \text{Ker } df \quad \text{and} \quad TX' \cong (f')^*(TY) \oplus \text{Ker } df'. \quad (18)$$

Define orientations on the fibres of $\text{Ker } df$, $\text{Ker } df'$ over X, X' such that (18) are isomorphisms of oriented vector bundles, where TX, TX' are oriented by the orientations on X, X' , and $f^*(TY), (f')^*(TY)$ by the ori-

entation on Y . Then we define the orientation on $X \times_Y X'$ so that

$$\begin{aligned} T(X \times_Y X') &\cong (f \circ \pi_X)^*(TY) \oplus \pi_X^*(\text{Ker } df) \oplus \pi_{X'}^*(\text{Ker } df') \\ &\cong \pi_X^*(TX) \oplus \pi_{X'}^*(\text{Ker } df') \\ &\cong (-1)^{\dim Y(\dim X - \dim Y)} \pi_X^*(\text{Ker } df) \oplus \pi_{X'}^*(TX') \end{aligned} \quad (19)$$

are isomorphisms of oriented vector bundles. Here $\pi_X : X \times_Y X' \rightarrow X$ and $\pi_{X'} : X \times_Y X' \rightarrow X'$ are the natural projections, and $f \circ \pi_X \equiv f' \circ \pi_{X'}$.

Note that the second line of (19) makes sense if f' is a submersion but f is only smooth, and the third line makes sense if f is a submersion but f' is only smooth. Thus, our convention extends to fibre products $X \times_{f,Y,f'} X'$ in which only one of f, f' is a submersion.

These extend immediately to orbifolds. They also extend to the Kuranishi space versions in Definitions 2.27 and 2.28; for Definition 2.28 they are described in [24, Conv. 45.1(4)]. We will not give details, but here is an algorithm to deduce Kuranishi space orientation conventions from manifold ones.

Let M be an oriented manifold or orbifold. Then M is a Kuranishi space with a single Kuranishi neighbourhood $(V_q, E_q, s_q, \psi_q) = (M, M, 0, \text{id}_M)$. The orientation on M induces an orientation on the fibres of $TV_q \oplus E_q = TM$, so M is oriented in the Kuranishi space sense.

Let $E \rightarrow M$ be a vector bundle or orbibundle. Write $\bar{E}$ for the total space of E , as an orbifold, with projection $\pi : \bar{E} \rightarrow M$, and regard M as a suborbifold of $\bar{E}$, the zero section of E . Then we can define a second Kuranishi neighbourhood $(V_p, E_p, s_p, \psi_p) = (\bar{E}, \pi^*(E), \text{id}_E, \text{id}_M)$. Here $s_p = \text{id}_E : \bar{E} \rightarrow \pi^*(E)$ is just the identity map on E , so $s_p^{-1}(0)$ is $M \subset \bar{E}$, the zero section of E . There is an obvious coordinate change $(\phi_{pq}, \hat{\phi}_{pq}) = (\text{id}_M, 0)$.

Now given any operation requiring an orientation convention which makes sense for manifolds, such as boundaries or fibre products of smooth submersions above, we can define it for manifolds, and then use the example above and Definition 2.29 to deduce what the convention must be for Kuranishi neighbourhoods of the form $(\bar{E}, \pi^*(E), \text{id}_E, \text{id}_M)$. This is enough to define the convention for general Kuranishi neighbourhoods.

If X is an oriented Kuranishi space, we often write $-X$ for the same Kuranishi space with the opposite orientation. The next result comes from [24, Lem. 45.3], except the second line of (20), which is elementary. In (a) we assume $\partial Y = \emptyset$ to avoid having to split $\partial X_1 = \partial_+^{\mathbf{f}_1} X_1 \amalg \partial_-^{\mathbf{f}_1} X_1$, and so on, as in the manifolds case in Definition 2.5 and Proposition 2.6.

Proposition 2.31. *Let $X_1, X_2, \dots$ be oriented Kuranishi spaces, $Y, Y_1, \dots$ be oriented orbifolds, and $\mathbf{f}_1 : X_1 \rightarrow Y, \dots$ be strongly smooth maps, with at least one strong submersion in each fibre product below. Then the following hold, in oriented Kuranishi spaces:*

(a) *If $\partial Y = \emptyset$, for $\mathbf{f}_1 : X_1 \rightarrow Y$ and $\mathbf{f}_2 : X_2 \rightarrow Y$ we have*

$$\begin{aligned} \partial(X_1 \times_Y X_2) &\cong (\partial X_1) \times_Y X_2 \amalg (-1)^{\text{vdim } X_1 + \dim Y} X_1 \times_Y (\partial X_2) \\ \text{and } X_1 \times_Y X_2 &\cong (-1)^{(\text{vdim } X_1 - \dim Y)(\text{vdim } X_2 - \dim Y)} X_2 \times_Y X_1. \end{aligned} \quad (20)$$

(b) For $\mathbf{f}_1 : X_1 \rightarrow Y_1$, $\mathbf{f}_2 : X_2 \rightarrow Y_1 \times Y_2$ and $\mathbf{f}_3 : X_3 \rightarrow Y_2$, we have

$$(X_1 \times_{Y_1} X_2) \times_{Y_2} X_3 \cong X_1 \times_{Y_1} (X_2 \times_{Y_2} X_3).$$

(c) For $\mathbf{f}_1 \times \mathbf{f}_2 : X_1 \rightarrow Y_1 \times Y_2$, $\mathbf{f}_3 : X_2 \rightarrow Y_1$ and $\mathbf{f}_4 : X_3 \rightarrow Y_2$, we have

$$X_1 \times_{Y_1 \times Y_2} (X_2 \times X_3) \cong (-1)^{\dim Y_2 (\dim Y_1 + \dim X_2)} (X_1 \times_{Y_1} X_2) \times_{Y_2} X_3.$$

A useful special case of Proposition 2.31(a) with $X_1 = [0, 1]$, $X_2 = X$ and Y a point gives the decomposition in oriented Kuranishi spaces:

$$\partial([0, 1] \times X) \cong (\{1\} \times X) \amalg -(\{0\} \times X) \amalg -([0, 1] \times \partial X).$$

2.8 Coorientations

We will also need a notion of *relative orientation* for a strong submersion $\mathbf{f} : X \rightarrow Y$. We call it a *coorientation*, as we use orientations in Kuranishi homology, and coorientations in Kuranishi cohomology.

Definition 2.32. Let X be a Kuranishi space, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ a strong submersion. A *coorientation* for $(X, \mathbf{f})$ assigns, for all $p \in X$ and all sufficiently small Kuranishi neighbourhoods (V_p, E_p, s_p, ψ_p) in the germ at p with submersion $f_p : V_p \rightarrow Y$ representing $\mathbf{f}$, orientations on the fibres of the orbibundle $\text{Ker } df_p \oplus E_p$ varying continuously over V_p , where $df_p : TV_p \rightarrow f_p^*(TY)$ is the (surjective) derivative of f_p .

These must be compatible with coordinate changes, in the following sense. Let $q \in \text{Im } \psi_p$, $(V_q, \dots, \psi_q)$ be sufficiently small in the germ at q , let $f_q : V_q \rightarrow Y$ represent $\mathbf{f}$, and $(\phi_{pq}, \hat{\phi}_{pq})$ be the coordinate change from $(V_q, \dots, \psi_q)$ to $(V_p, \dots, \psi_p)$ in the germ. Then we require that in oriented orbibundles over V_q near $s_q^{-1}(0)$, we have

$$\begin{aligned} \phi_{pq}^* [\text{Ker } df_p \oplus E_p] &\cong (-1)^{\dim V_q (\dim V_p - \dim V_q)} [\text{Ker } df_q \oplus E_q] \oplus \\ &\quad \left[\frac{\phi_{pq}^*(TV_p)}{(d\phi_{pq})(TV_q)} \oplus \frac{\phi_{pq}^*(E_p)}{\phi_{pq}(E_q)} \right], \end{aligned} \quad (21)$$

by analogy with (17), where $\frac{\phi_{pq}^*(TV_p)}{(d\phi_{pq})(TV_q)} \oplus \frac{\phi_{pq}^*(E_p)}{\phi_{pq}(E_q)}$ is oriented as in Definition 2.29.

Suppose now that Y is oriented. Then an orientation on X is equivalent to a coorientation for $(X, \mathbf{f})$, since for all p , $(V_p, \dots, \psi_p)$, f_p as above, the isomorphism $TV_p \cong f_p^*(TY) \oplus \text{Ker } df_p$ induces isomorphisms of orbibundles over V_p :

$$(TV_p \oplus E_p) \cong f_p^*(TY) \oplus (\text{Ker } df_p \oplus E_p). \quad (22)$$

There is a 1-1 correspondence between orientations on X and coorientations for $(X, \mathbf{f})$ such that (22) holds in oriented orbibundles, where $TV_p \oplus E_p$ is oriented by the orientation on X , and $\text{Ker } df_p \oplus E_p$ by the coorientation for $(X, \mathbf{f})$, and $f_p^*(TY)$ by the orientation on Y . Taking the direct sum of $f_p^*(TY)$ with (21) and using (22) yields (17), so this is compatible with coordinate changes.

Convention 2.33. Here are our conventions for manifolds:

- (a) Let X, Y be manifolds with $\partial Y = \emptyset$ and $f : X \rightarrow Y$ a submersion. Suppose (X, f) is cooriented. Then we define a coorientation for $(\partial X, f|_{\partial X})$ such that $\text{Ker}(df)|_{\partial X} = (-1)^{\dim Y} \mathbb{R}_{\text{out}} \oplus \text{Ker} d(f|_{\partial X})$ is an isomorphism of oriented vector bundles, where $(df)|_{\partial X}$ maps $TX|_{\partial X} \rightarrow f^*(TY)|_{\partial X}$ and $d(f|_{\partial X})$ maps $T(\partial X) \rightarrow f^*(TY)|_{\partial X}$ and $\mathbb{R}_{\text{out}}$ is oriented by an outward-pointing normal vector to ∂X in $\text{Ker}(df)|_{\partial X}$.

Now drop the assumption that $\partial Y = \emptyset$. Then as in Definition 2.5 we have a decomposition $\partial X = \partial_+^f X \amalg \partial_-^f X$, and submersions $f_+ : \partial_+^f X \rightarrow Y$ and $f_- : \partial_-^f X \rightarrow \partial Y$. Define a coorientation for $(\partial_+^f X, f_+)$ such that $\text{Ker}(df)|_{\partial_+^f X} = (-1)^{\dim Y} \mathbb{R}_{\text{out}} \oplus \text{Ker} df_+$. Define a coorientation for $(\partial_-^f X, f_-)$ such that $\text{Ker}(df)|_{\partial_-^f X} = \text{Ker} df_-$.

- (b) Let X, X', Y be manifolds, $f : X \rightarrow Y$, $f' : X' \rightarrow Y$ be submersions, and $(X, f), (X', f')$ be cooriented. Define a coorientation on $(X \times_Y X', \pi_Y)$ such that

$$\text{Ker}(d\pi_Y : T(X \times_Y X') \rightarrow \pi_Y^*(TY)) \cong \pi_X^*(\text{Ker} df) \oplus \pi_{X'}^*(\text{Ker} df')$$

is an isomorphism of oriented vector bundles, where $\text{Ker} df$ and $\text{Ker} df'$ are oriented by the coorientations for $(X, f), (X', f')$. Note that we do not need Y oriented, in contrast to Convention 2.30(b).

- (c) We can also combine orientations and coorientations in fibre products, as follows. Let X, X', Y be manifolds, $f : X \rightarrow Y$ be smooth, $f' : X' \rightarrow Y$ be a submersion, and suppose X is oriented and (X', f') is cooriented. As in (19), define orientations on $X \times_Y X'$ and $X' \times_Y X$ such that

$$\begin{aligned} T(X \times_Y X') &\cong \pi_X^*(TX) \oplus \pi_{X'}^*(\text{Ker} df'), \\ T(X' \times_Y X) &\cong (-1)^{\dim Y (\dim X' - \dim Y)} \pi_{X'}^*(\text{Ker} df') \oplus \pi_X^*(TX) \end{aligned}$$

are isomorphism of oriented vector bundles, where $TX, \text{Ker} df'$ are oriented by the orientation on X and the coorientation on (X', f') . Note that we do not need Y oriented, nor f a submersion.

These extend to orbifolds and Kuranishi spaces as for Convention 2.30.

As in Definition 2.32, if $\mathbf{f} : X \rightarrow Y$ is a strong submersion and Y is oriented, there is a 1-1 correspondence between orientations on X and coorientations for $(X, \mathbf{f})$. The signs in Convention 2.33 then correspond with those in Convention 2.30. Therefore the analogue of Proposition 2.31 holds for coorientations. In particular, for strong submersions $\mathbf{f}_a : X_a \rightarrow Y$ with $(X_a, \mathbf{f}_a)$ cooriented for $a = 1, 2, 3$, taking $\partial Y = \emptyset$ in (23), we have

$$\begin{aligned} (\partial(X_1 \times_Y X_2), \pi_Y) &\cong ((\partial X_1) \times_Y X_2, \pi_Y) \amalg \\ &\quad (-1)^{\text{vdim } X_1 + \dim Y} (X_1 \times_Y (\partial X_2), \pi_Y), \end{aligned} \quad (23)$$

$$(X_1 \times_Y X_2, \pi_Y) \cong (-1)^{(\text{vdim } X_1 - \dim Y)(\text{vdim } X_2 - \dim Y)} (X_2 \times_Y X_1, \pi_Y), \quad (24)$$

$$((X_1 \times_Y X_2) \times_Y X_3, \pi_Y) \cong (X_1 \times_Y (X_2 \times_Y X_3), \pi_Y). \quad (25)$$

Similarly, if X_1 is oriented, $\mathbf{f}_1 : X_1 \rightarrow Y$ is strongly smooth, $\mathbf{f}_2 : X_2 \rightarrow Y$ is a cooriented strong submersion, and $\partial Y = \emptyset$ then

$$\partial(X_1 \times_Y X_2) \cong ((\partial X_1) \times_Y X_2) \amalg (-1)^{\text{vdim } X_1 + \dim Y} (X_1 \times_Y (\partial X_2)) \quad (26)$$

in oriented Kuranishi spaces, combining orientations and coorientations as in Convention 2.33(c).

Here is how to generalize equations (23) and (26) to the case $\partial Y \neq \emptyset$. If X is a Kuranishi space and $\mathbf{f} : X \rightarrow Y$ a strong submersion then as for submersions of manifolds in Definition 2.5, we can define a decomposition $\partial X = \partial_+^{\mathbf{f}} X \amalg \partial_-^{\mathbf{f}} X$, and strong submersions $\mathbf{f}_+ : \partial_+^{\mathbf{f}} X \rightarrow Y$ and $\mathbf{f}_- : \partial_-^{\mathbf{f}} X \rightarrow \partial Y$. When $\mathbf{f}_1 : X_1 \rightarrow Y$ and $\mathbf{f}_2 : X_2 \rightarrow Y$ are cooriented strong submersions, and we do not assume $\partial Y = \emptyset$, equation (23) should be replaced by

$$(\partial_+^{\pi_Y} (X_1 \times_Y X_2), \pi_Y) \cong ((\partial_+^{\mathbf{f}_1} X_1) \times_Y X_2, \pi_Y) \amalg (-1)^{\text{vdim } X_1 + \dim Y} (X_1 \times_Y (\partial_+^{\mathbf{f}_2} X_2), \pi_Y), \quad (27)$$

$$(\partial_-^{\pi_Y} (X_1 \times_Y X_2), \pi_{\partial Y}) \cong ((\partial_-^{\mathbf{f}_1} X_1) \times_{\partial Y} (\partial_-^{\mathbf{f}_2} X_2), \pi_{\partial Y}). \quad (28)$$

If X_1 is oriented, $\mathbf{f}_1 : X_1 \rightarrow Y$ is strongly smooth, and $\mathbf{f}_2 : X_2 \rightarrow Y$ is a cooriented strong submersion, then equation (26) should be replaced by

$$\partial(X_1 \times_Y X_2) \cong ((\partial X_1) \times_Y X_2) \amalg (-1)^{\text{vdim } X_1 + \dim Y} (X_1 \times_Y (\partial_+^{\mathbf{f}_2} X_2)). \quad (29)$$

2.9 Almost complex and almost CR structures

We now define *almost complex structures* on Kuranishi spaces. They are a substitute for Fukaya and Ono's notion of *stably almost complex* Kuranishi spaces [25, Def. 5.17]. We will use them to define almost complex Kuranishi bordism $KB_*^{\text{ac}}(Y; R)$ in Chapter 5, and in the author's approach to the Gopakumar–Vafa Integrality Conjecture in Chapter 6.

Definition 2.34. Let V be an orbifold without boundary. An *almost complex structure* J on V is a tensor $J = J_a^b$ in $C^\infty(TV \otimes T^*V)$ with $J^2 = -1$, that is, $J_a^b J_b^c = -\delta_a^c$ in index notation. Let $E \rightarrow V$ be an orbibundle. An *almost complex structure* K on the fibres of E is $K \in C^\infty(E^* \otimes E)$ with $K^2 = -1$.

Let X be a Kuranishi space without boundary. An *almost complex structure* (J, K) on X assigns for all $p \in X$ and all sufficiently small (V_p, E_p, s_p, ψ_p) in the germ at p , a choice of almost complex structure J_p on V_p , and a choice of almost complex structure K_p on the fibres of E_p . These choices must satisfy the following conditions. For all $p \in X$, for all $(V_p, \dots, \psi_p)$ sufficiently small in the germ at p with almost complex structures J_p, K_p , for all $q \in \text{Im } \psi_p$, and for all sufficiently small $(V_q, \dots, \psi_q)$ in the germ at q with almost complex structures J_q, K_q , if $(\phi_{pq}, \hat{\phi}_{pq})$ is the coordinate change from $(V_q, \dots, \psi_q)$ to $(V_p, \dots, \psi_p)$ in the germ of coordinate changes, then:

- (a) $d\phi_{pq} \circ J_q = \phi_{pq}^*(J_p) \circ d\phi_{pq}$ as morphisms of orbibundles $TV_q \rightarrow \phi_{pq}^*(TV_p)$, that is, ϕ_{pq} is a pseudoholomorphic map of almost complex orbifolds;

- (b) $\hat{\phi}_{pq} \circ K_q = \phi_{pq}^*(K_p) \circ \hat{\phi}_{pq}$ as morphisms of orbibundles $E_q \rightarrow \phi_{pq}^*(E_p)$; and
- (c) Parts (a) and (b) imply that the orbibundles $\phi_{pq}^*(TV_p)/(\mathrm{d}\phi_{pq})(TV_q)$ and $\phi_{pq}^*(E_p)/\hat{\phi}_{pq}(E_q)$ over V_q appearing in (14) have almost complex structures J_{pq}, K_{pq} on their fibres, by projection from $\phi_{pq}^*(J_p), \phi_{pq}^*(K_p)$. We require that $K_{pq} \circ \mathrm{d}\hat{s}_p = \mathrm{d}\hat{s}_p \circ J_{pq}$ over $s_q^{-1}(0)$ in V_q , for $\mathrm{d}\hat{s}_p$ as in (14).

If (J_p, K_p) exists for (V_p, E_p, s_p, ψ_p) as above then $\dim V_p$ and $\mathrm{rank} E_p$ are even, so $\mathrm{vdim} X = \dim V_p - \mathrm{rank} E_p$ is even. Thus X can admit an almost complex structure only if $\mathrm{vdim} X$ is even. Our next definition is an odd-dimensional analogue of almost complex structures, in the same way that contact structures are odd-dimensional analogues of symplectic structures.

Definition 2.35. Let V be an orbifold with boundary, but without (g-)corners. An *almost CR structure* (D, J) on V is an orbifold vector subbundle D of TV with $\dim V - \mathrm{rank} D = 1$, such that $D|_{\partial V} = T(\partial V) \subset TV|_{\partial V}$, and an almost complex structure J on the fibres of D .

Let X be a Kuranishi space with boundary, but without (g-)corners. An *almost CR structure* $(\mathbf{D}, \mathbf{J}, \mathbf{K})$ on X assigns for all $p \in X$ and all sufficiently small (V_p, E_p, s_p, ψ_p) in the germ at p , a choice of almost CR structure (D_p, J_p) on V_p , and a choice of almost complex structure K_p on the fibres of E_p . These choices must satisfy the following conditions. For all $p \in X$, for all $(V_p, \dots, \psi_p)$ sufficiently small in the germ at p with structures $(D_p, J_p), K_p$, for all $q \in \mathrm{Im} \psi_p$, and for all sufficiently small $(V_q, \dots, \psi_q)$ in the germ at q with structures $(D_q, J_q), K_q$, if $(\phi_{pq}, \hat{\phi}_{pq})$ is the coordinate change from $(V_q, \dots, \psi_q)$ to $(V_p, \dots, \psi_p)$ in the germ of coordinate changes, then:

- (a) $\mathrm{d}\phi_{pq}(D_q)$ is an orbisubbundle of $\phi_{pq}^*(D_p)$ in $\phi_{pq}^*(TV_p)$, and $\mathrm{d}\phi_{pq} \circ J_q = \phi_{pq}^*(J_p) \circ \mathrm{d}\phi_{pq}|_{D_q}$ as morphisms of orbibundles $D_q \rightarrow \phi_{pq}^*(D_p)$;
- (b) $\hat{\phi}_{pq} \circ K_q = \phi_{pq}^*(K_p) \circ \hat{\phi}_{pq}$ as morphisms of orbibundles $E_q \rightarrow \phi_{pq}^*(E_p)$; and
- (c) Part (a) and the fact that D_q, D_p both have codimension 1 yield an isomorphism $\phi_{pq}^*(D_p)/(\mathrm{d}\phi_{pq})(D_q) \cong \phi_{pq}^*(TV_p)/(\mathrm{d}\phi_{pq})(TV_q)$ of orbibundles over V_q . So projecting $\phi_{pq}^*(J_p)$ gives an almost complex structure J_{pq} on $\phi_{pq}^*(TV_p)/(\mathrm{d}\phi_{pq})(TV_q)$. Part (b) implies that projecting $\phi_{pq}^*(K_p)$ gives an almost complex structure K_{pq} on $\phi_{pq}^*(E_p)/\hat{\phi}_{pq}(E_q)$. We require that $K_{pq} \circ \mathrm{d}\hat{s}_p = \mathrm{d}\hat{s}_p \circ J_{pq}$ over $s_q^{-1}(0)$ in V_q , for $\mathrm{d}\hat{s}_p$ as in (14).

If V is an orbifold with boundary but without (g-)corners, and (D, J) is an almost CR structure on V , then ∂V is an orbifold without boundary, and $J|_{\partial V}$ is an almost complex structure on ∂V , as $T(\partial V) = D|_{\partial V}$. In the same way, if X is a Kuranishi space with boundary but without (g-)corners and $(\mathbf{D}, \mathbf{J}, \mathbf{K})$ is an almost CR structure on X then ∂X is a Kuranishi space without boundary, and $(\mathbf{J}, \mathbf{K})|_{\partial X}$ is an almost complex structure on ∂X .

2.10 Co-almost complex and co-almost CR structures

Just as we can generalize orientations on X in §2.7 to coorientations for $(X, \mathbf{f})$ for $\mathbf{f} : X \rightarrow Y$ a strong submersion in §2.8, so we can generalize §2.9 to give notions of *co-almost complex structures* and *co-almost CR structures* for $(X, \mathbf{f})$.

Definition 2.36. Let V, Y be orbifolds without boundary, and $f : V \rightarrow Y$ a submersion. A *co-almost complex structure* L for (V, f) is an almost complex structure L on the fibres of the orbibundle $\text{Ker}(df : TV \rightarrow f^*(TY))$ over V .

Let X be a Kuranishi space without boundary, and $\mathbf{f} : X \rightarrow Y$ a strong submersion. A *co-almost complex structure* $(\mathbf{L}, \mathbf{K})$ for $(X, \mathbf{f})$ assigns for all $p \in X$ and all sufficiently small (V_p, E_p, s_p, ψ_p) in the germ at p with submersion $f_p : V_p \rightarrow Y$, a co-almost complex structure L_p for (V_p, f_p) , and an almost complex structure K_p on the fibres of E_p , satisfying analogues of Definition 2.34(a)–(c).

Let V be an orbifold with boundary but without (g-)corners, and $f : V \rightarrow Y$ a submersion. A *co-almost CR structure* (D, L) for (V, f) is an orbifold vector subbundle D of $\text{Ker}(df : TV \rightarrow f^*(TY))$ with $\text{rank } D = \dim V - \dim Y - 1$, such that $D|_{\partial V} = \text{Ker}(d(f|_{\partial V}) : T(\partial V) \rightarrow f|_{\partial V}^*(TY))$, and an almost complex structure L on the fibres of D .

Let X be a Kuranishi space with boundary, but without (g-)corners. A *co-almost CR structure* $(\mathbf{D}, \mathbf{L}, \mathbf{K})$ for $(X, \mathbf{f})$ assigns a co-almost CR structure (D_p, L_p) for (V_p, f_p) , and an almost complex structure K_p on the fibres of E_p , for all $p, (V_p, \dots, \psi_p), f_p$ as above, satisfying analogues of Definition 2.35(a)–(c).

As in Definition 2.35, if (D, L) is a co-almost CR structure for (V, f) then $L|_{\partial V}$ is a co-almost complex structure for $(\partial V, f|_{\partial V})$, and if $(\mathbf{D}, \mathbf{L}, \mathbf{K})$ is a co-almost CR structure for $(X, \mathbf{f})$ then $(\mathbf{L}, \mathbf{K})|_{\partial X}$ is a co-almost complex structure for $(\partial X, \mathbf{f}|_{\partial X})$.

We can *combine co-almost complex structures on fibre products*. Let X, X' be Kuranishi spaces, Y an orbifold, $\mathbf{f} : X \rightarrow Y$, $\mathbf{f}' : X' \rightarrow Y$ strong submersions, and $(\mathbf{L}, \mathbf{K}), (\mathbf{L}', \mathbf{K}')$ co-almost structures for $(X, \mathbf{f}), (X', \mathbf{f}')$. Then we can define a natural co-almost complex structure $(\mathbf{L}, \mathbf{K}) \times_Y (\mathbf{L}', \mathbf{K}')$ for $(X \times_Y X', \pi_Y)$. Suppose $p \in X, p' \in X'$ with $f(p) = f'(p')$ in Y , and $(V_p, \dots, \psi_p), (V'_{p'}, \dots, \psi'_{p'})$ are sufficiently small Kuranishi neighbourhoods of p, p' in X, X' , and $f_p : V_p \rightarrow Y, f'_{p'} : V'_{p'} \rightarrow Y$ are submersions representing $\mathbf{f}, \mathbf{f}'$, and $(L_p, K_p), (L'_{p'}, K'_{p'})$ represent $(\mathbf{L}, \mathbf{K}), (\mathbf{L}', \mathbf{K}')$ on $(V_p, \dots, \psi_p), (V'_{p'}, \dots, \psi'_{p'})$.

Then $(V_{pp'}, \dots, \psi_{pp'})$ is a Kuranishi neighbourhood on $X \times_Y X'$, where $V_{pp'} = V_p \times_Y V'_{p'}$, and $E_{pp'} = \pi_{V_p}^*(E_p) \oplus \pi_{V'_{p'}}^*(E'_{p'})$, and $\pi_{pp'} = \pi_Y : V_{pp'} \rightarrow Y$ represents $\pi_Y : X \times_Y X' \rightarrow Y$. We have a natural isomorphism

$$\begin{aligned} \text{Ker}(d\pi_{pp'} : TV_{pp'} \rightarrow \pi_{pp'}^*(TY)) &\cong \pi_{V_p}^*(\text{Ker}(df_p : TV_p \rightarrow f_p^*(TY))) \oplus \\ &\pi_{V'_{p'}}^*(\text{Ker}(df'_{p'} : TV'_{p'} \rightarrow (f'_{p'})^*(TY))). \end{aligned} \quad (30)$$

Define an almost complex structure $L_{pp'}$ on the fibres of the l.h.s. of (30) to be $\pi_{V_p}^*(L_p) \oplus \pi_{V'_{p'}}^*(L'_{p'})$ on the r.h.s. of (30). Define an almost complex structure $K_{pp'}$ on $E_{pp'}$ to be $\pi_{V_p}^*(K_p) \oplus \pi_{V'_{p'}}^*(K'_{p'})$ on $\pi_{V_p}^*(E_p) \oplus \pi_{V'_{p'}}^*(E'_{p'})$. Then these

$(L_{pp'}, K_{pp'})$ for all p, p' induce a co-almost complex structure $(\mathbf{L}, \mathbf{K}) \times_Y (\mathbf{L}', \mathbf{K}')$ for $(X \times_Y X', \pi_Y)$.

Similarly, we can naturally define the fibre product $(\mathbf{L}, \mathbf{K}) \times_Y (\mathbf{D}', \mathbf{L}', \mathbf{K}')$ of a co-almost complex structure $(\mathbf{L}, \mathbf{K})$ for $(X, \mathbf{f})$ and a co-almost CR structure $(\mathbf{D}', \mathbf{L}', \mathbf{K}')$ for $(X', \mathbf{f}')$, which is a co-almost CR structure for $(X \times_Y X', \pi_Y)$.

Given $\mathbf{f} : X \rightarrow Y$ strongly smooth and $\mathbf{f}' : X' \rightarrow Y$ a strong submersion, it is also possible to define the fibre product of an almost complex structure $(\mathbf{J}, \mathbf{K})$ on X and a co-almost complex structure $(\mathbf{L}', \mathbf{K}')$ for $(X', \mathbf{f}')$, to get an almost complex structure $(\mathbf{J}, \mathbf{K}) \times_Y (\mathbf{L}', \mathbf{K}')$ for $X \times_{\mathbf{f}, Y, \mathbf{f}'} X'$. However, this construction is not quite natural. The issue is that if $p \in X$ with $(J_p, K_p), f_p$ representing $(\mathbf{J}, \mathbf{K}), \mathbf{f}$ on $(V_p, \dots, \psi_p)$, and $p' \in X'$ with $(L'_{p'}, K'_{p'}), f'_{p'}$ representing $(\mathbf{L}', \mathbf{K}'), \mathbf{f}'$ on $(V'_{p'}, \dots, \psi'_{p'})$, then J_p is an almost complex structure on the fibres of TV_p , and $L'_{p'}$ is an almost complex structure on the fibres of $\text{Ker}(df'_{p'} : TV'_{p'} \rightarrow (f')^*(TY))$. We can write

$$T(V_p \times_{f_p, Y, f'_{p'}} V'_{p'}) \cong \pi_{V_p}^*(TV_p) \oplus \pi_{V'_{p'}}^*(\text{Ker}(df'_{p'} : TV'_{p'} \rightarrow (f')^*(TY))), \quad (31)$$

but this splitting is not natural; only the short exact sequence

$$0 \rightarrow \pi_{V'_{p'}}^*(\text{Ker}(df'_{p'} : TV'_{p'} \rightarrow (f')^*(TY))) \rightarrow T(V_p \times_{f_p, Y, f'_{p'}} V'_{p'}) \rightarrow \pi_{V_p}^*(TV_p) \rightarrow 0$$

is natural. So to define $(\mathbf{J}, \mathbf{K}) \times_Y (\mathbf{L}', \mathbf{K}')$ we must choose splittings (31) for all p, p' with $f(p) = f'(p')$, compatible with coordinate changes $(\phi_{pq}, \hat{\phi}_{pq})$ and $(\phi'_{p'q'}, \hat{\phi}'_{p'q'})$, and then define $J_{pp'}$ to be $\pi_{V_p}^*(J_p) \oplus \pi_{V'_{p'}}^*(L'_{p'})$, using (31).

In the same way, after choosing splittings (31), we can define fibre products $(\mathbf{D}, \mathbf{J}, \mathbf{K}) \times_Y (\mathbf{L}', \mathbf{K}')$ of an almost CR structure $(\mathbf{D}, \mathbf{J}, \mathbf{K})$ on X and a co-almost complex structure $(\mathbf{L}', \mathbf{K}')$ for $(X', \mathbf{f}')$, and $(\mathbf{J}, \mathbf{K}) \times_Y (\mathbf{D}', \mathbf{L}', \mathbf{K}')$ of an almost complex structure $(\mathbf{J}, \mathbf{K})$ on X and a co-almost CR structure $(\mathbf{D}', \mathbf{L}', \mathbf{K}')$ for $(X', \mathbf{f}')$. Both are almost CR structures on $X \times_{\mathbf{f}, Y, \mathbf{f}'} X'$.

3 Gauge-fixing data and co-gauge-fixing data

The basic idea of this book is to define a homology theory $KH_*(Y; R)$ for an orbifold Y using as chains pairs $(X, \mathbf{f})$, where X is a compact Kuranishi space and $\mathbf{f} : X \rightarrow Y$ is a strongly smooth map. However, we show in §4.9 below that if we allow chains $(X, \mathbf{f})$ for which the *automorphism group* $\text{Aut}(X, \mathbf{f})$ of strong diffeomorphisms $\mathbf{a} : X \rightarrow X$ up to equality such that $\mathbf{f} \circ \mathbf{a} = \mathbf{f}$ is infinite, the resulting homology groups are always zero.

To deal with this problem, we add some *extra data* $\mathbf{G}$ to $(X, \mathbf{f})$ for which $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ is finite, and use as chains triples $(X, \mathbf{f}, \mathbf{G})$. Exactly what this extra data is does not matter very much, but here are the properties we would like it to satisfy for this book and in [4, 39–41]. Parts (a), (e) are not essential for well-behaved (co)homology theories, but will be needed in [4, 39–41]. In (e), we restrict to X with corners (not g-corners) because Principle 2.8 fails for smooth extensions from boundaries of general manifolds or orbifolds with g-corners.

Property 3.1. We want to define some kind of *extra data* $\mathbf{G}$ for pairs $(X, \mathbf{f})$, where X is a compact Kuranishi space with g-corners, Y an orbifold and $\mathbf{f} : X \rightarrow Y$ a strongly smooth map, such that:

- (a) Every pair $(X, \mathbf{f})$ should admit extra data $\mathbf{G}$. If Γ is a finite subgroup of $\text{Aut}(X, \mathbf{f})$, then there should exist Γ -invariant extra data for $(X, \mathbf{f})$.
- (b) The *automorphism group* $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ of isomorphisms $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \mathbf{G}) \rightarrow (X, \mathbf{f}, \mathbf{G})$ is finite, where $\mathbf{b}$ is a lift of $\mathbf{a}$ to $\mathbf{G}$.
- (c) Suppose Γ is a finite group acting on $(X, \mathbf{f}, \mathbf{G})$ by isomorphisms, that is, we are given a group morphism $\rho : \Gamma \rightarrow \text{Aut}(X, \mathbf{f}, \mathbf{G})$, which need not be injective. Then we can form the quotient $\tilde{X} = X/\Gamma$, a compact Kuranishi space, with projection $\pi : X \rightarrow \tilde{X}$, and $\mathbf{f}$ pushes down to $\tilde{\mathbf{f}} : \tilde{X} \rightarrow Y$ with $\mathbf{f} = \tilde{\mathbf{f}} \circ \pi$. We require that $\mathbf{G}$ should also naturally push down to extra data $\tilde{\mathbf{G}}$ for $(\tilde{X}, \tilde{\mathbf{f}})$.
- (d) If $\mathbf{G}$ is extra data for $(X, \mathbf{f})$, it has a *restriction* $\mathbf{G}|_{\partial X}$ which is extra data for $(\partial X, \mathbf{f}|_{\partial X})$.
- (e) Let $\sigma : \partial^2 X \rightarrow \partial^2 X$ be the natural involution of Definition 2.27. Suppose X has corners (not g-corners), and $\mathbf{H}$ is extra data for $(\partial X, \mathbf{f}|_{\partial X})$. Then there should exist extra data $\mathbf{G}$ for $(X, \mathbf{f})$ with $\mathbf{G}|_{\partial X} = \mathbf{H}$ if and only if $\mathbf{H}|_{\partial^2 X}$ is invariant under σ .
If also Γ is a finite subgroup of $\text{Aut}(X, \mathbf{f})$, and $\mathbf{H}$ is invariant under $\Gamma|_{\partial X}$, then we can choose $\mathbf{G}$ to be Γ -invariant.
- (f) For the case of extra data for Kuranishi homology: let Y, Z be orbifolds, and $h : Y \rightarrow Z$ a smooth map. Suppose X is a compact Kuranishi space and $\mathbf{f} : X \rightarrow Y$ is strongly smooth. If $\mathbf{G}$ is extra data for $(X, \mathbf{f})$, then there should exist extra data $h_*(\mathbf{G})$ for $(X, h \circ \mathbf{f})$. It should satisfy $(g \circ h)_*(\mathbf{G}) = g_*(h_*(\mathbf{G}))$.
- (g) For the case of extra data for Kuranishi cohomology: let Y, Z be orbifolds, and $h : Y \rightarrow Z$ a smooth, proper map. Suppose X is a compact Kuranishi space and $\mathbf{f} : X \rightarrow Z$ is a strong submersion. If $\mathbf{G}$ is extra data for $(X, \mathbf{f})$, then there should exist extra data $h^*(\mathbf{G})$ for $(Y \times_{h, Z, \mathbf{f}} X, \pi_Y)$. It should satisfy $(g \circ h)^*(\mathbf{G}) = h^*(g^*(\mathbf{G}))$.
- (h) (Mainly for Kuranishi cohomology). Let X_1, X_2, X_3 be compact Kuranishi spaces, Y an orbifold, $\mathbf{f}_e : X_e \rightarrow Y$ be strongly smooth for $e = 1, 2, 3$ with at least two $\mathbf{f}_e$ strong submersions, and $\mathbf{G}_e$ be extra data for $(X_e, \mathbf{f}_e)$ for $e = 1, 2, 3$. Then we should construct extra data $\mathbf{G}_1 \times_Y \mathbf{G}_2$ for $(X_1 \times_{\mathbf{f}_1, Y, \mathbf{f}_2} X_2, \pi_Y)$ from $\mathbf{G}_1, \mathbf{G}_2$.

This construction should be *symmetric*, in that it yields isomorphic extra data for $(X_1 \times_{\mathbf{f}_1, Y, \mathbf{f}_2} X_2, \pi_Y)$ and $(X_2 \times_{\mathbf{f}_2, Y, \mathbf{f}_1} X_1, \pi_Y)$ under the natural isomorphism $X_1 \times_{\mathbf{f}_1, Y, \mathbf{f}_2} X_2 \cong X_2 \times_{\mathbf{f}_2, Y, \mathbf{f}_1} X_1$. It should also be *associative*, in that it yields isomorphic extra data for $((X_1 \times_Y X_2) \times_Y X_3, \pi_Y)$ and $(X_1 \times_Y (X_2 \times_Y X_3), \pi_Y)$ under $(X_1 \times_Y X_2) \times_Y X_3 \cong X_1 \times_Y (X_2 \times_Y X_3)$. For Kuranishi cohomology, products should be *functorial for pullbacks* in part (g), $h^*(\mathbf{G}_1 \times_Z \mathbf{G}_2) \cong h^*(\mathbf{G}_1) \times_Y h^*(\mathbf{G}_2)$.

We shall define *gauge-fixing data* $\mathbf{G}$ (for Kuranishi homology), and *co-gauge-fixing data* $\mathbf{C}$ (for Kuranishi cohomology), which satisfy all these, and also have other technical features which will be important in Appendices B and C. The name comes from physics, where one often has an infinite-dimensional symmetry group called the gauge group, and one breaks the symmetry by ‘fixing a gauge’. It was not easy to arrange for all of Property 3.1(a)–(h) to hold at once.

3.1 Definition of (co-)gauge-fixing data

Good coordinate systems are convenient choices of finite coverings of X by Kuranishi neighbourhoods, due to Fukaya and Ono [25, Def. 6.1], [24, Lem. A1.11].

Definition 3.2. Let X be a compact Kuranishi space. A *good coordinate system* $\mathbf{I}$ for X consists of a finite indexing set I , a total order $\leq$ on I , a family $\{(V^i, E^i, s^i, \psi^i) : i \in I\}$ of compatible Kuranishi neighbourhoods on X with $V^i \neq \emptyset$ for all $i \in I$ and $X = \bigcup_{i \in I} \text{Im } \psi^i$, and for all $i, j \in I$ with $j \leq i$ a triple $(V^{ij}, \phi^{ij}, \hat{\phi}^{ij})$, where V^{ij} is an open neighbourhood of $(\psi^j)^{-1}(\text{Im } \psi^i)$ in V^j , and $(\phi^{ij}, \hat{\phi}^{ij})$ is a coordinate change from $(V^{ij}, E^j|_{V^{ij}}, s^j|_{V^{ij}}, \psi^j|_{V^{ij}})$ to (V^i, E^i, s^i, ψ^i) . When $i = j$ these should satisfy $V^{ij} = V^i$, with $(\phi^{ii}, \hat{\phi}^{ii})$ the identity map on $(V^i, \dots, \psi^i)$. When $i, j, k \in I$ with $k \leq j \leq i$ these should satisfy $\phi^{ij} \circ \phi^{jk} = \phi^{ik}$ and $\hat{\phi}^{ij} \circ \hat{\phi}^{jk} = \hat{\phi}^{ik}$ over $(\phi^{jk})^{-1}(V^{ij}) \cap V^{ik}$, and

$$\phi^{ij}(V^{ij}) \cap \phi^{ik}(V^{ik}) = \phi^{ik}((\phi^{jk})^{-1}(V^{ij}) \cap V^{ik}). \quad (32)$$

Now let Y be an orbifold and $\mathbf{f} : X \rightarrow Y$ a strongly smooth map. A *good coordinate system* $\mathbf{I}$ for $(X, \mathbf{f})$ consists of a good coordinate system as above, together with smooth $f^i : V^i \rightarrow Y$ representing $\mathbf{f}$, such that if $j \leq i$ in I then $f^j|_{V^{ij}} \equiv f^i \circ \phi^{ij}$. If $\mathbf{f}$ is a *strong submersion*, we take the f^i submersions.

The point of good coordinate systems is that one often needs to choose some geometric data (for instance, a transverse multisection, as in as [25, Th. 6.4]) on each of a system of Kuranishi neighbourhoods satisfying compatibility conditions on the overlaps. With a good coordinate system one can make these choices on each $(V^i, \dots, \psi^i)$ in the order $\leq$ on I by induction, imposing compatibility conditions on the V^{ij} . Equation (32) was added by the author; without it, in inductive arguments when making the choice on $(V^i, \dots, \psi^i)$ there may be problems at points in V^i lying in the left hand side of (32) but not the right.

We shall now define three variations on the notion of good coordinate system, which we call *very good*, *really good* and *excellent coordinate systems*. A *very good coordinate system* is a good coordinate system in which there is at most one Kuranishi neighbourhood $(V^i, \dots, \psi^i)$ of each dimension $\dim V^i$, so we use the dimension $\dim V^i$ as the index i .

Definition 3.3. Let X be a compact Kuranishi space, Y an orbifold and $\mathbf{f} : X \rightarrow Y$ a strongly smooth map. A good coordinate system $\mathbf{I}$ for $(X, \mathbf{f})$ is called *very good* if $I \subset \mathbb{N} = \{0, 1, 2, \dots\}$, and the order $\leq$ on I is the restriction of $\leq$ on $\mathbb{N}$, and $\dim V^i = i$ for all $i \in I$.

Note that it is necessary that the order $\leq$ on I is the restriction of $\leq$ on $\mathbb{N}$, since coordinate changes $(\phi^{ij}, \hat{\phi}^{ij})$ from $V^{ij} \subseteq V^j$ to V^i can exist if $\dim V^j \leq \dim V^i$, that is, if $j \leq i$, but not if $\dim V^j > \dim V^i$. A *really good coordinate system* is a very good coordinate system $\mathbf{I}$ with some extra partition of unity data $\boldsymbol{\eta}$. We will use $\boldsymbol{\eta}$ for defining really good coordinate systems on fibre products in §3.8.

Definition 3.4. Let X be a compact Kuranishi space, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ a strongly smooth map. A *really good coordinate system* $(\mathbf{I}, \boldsymbol{\eta})$ for $(X, \mathbf{f})$ is a very good coordinate system $\mathbf{I} = (I, (V^i, \dots, \psi^i), f^i : i \in I, \dots)$ for $(X, \mathbf{f})$ together with data $\boldsymbol{\eta} = (\eta_i : i \in I, \eta_i^j : i, j \in I)$ satisfying:

- (i) $\eta_i : X \rightarrow [0, 1]$ is a continuous function for each $i \in I$, with $\sum_{i \in I} \eta_i \equiv 1$, and $\eta_i > 0$ on $\text{Im } \psi^i$. If $j < i$ and $p \in X$ with $\eta_i(p) > 0$, $\eta_j(p) > 0$ and $p \in \text{Im } \psi^i$ then $p \in \text{Im } \psi^j$.
- (ii) $\eta_i^j : V^j \rightarrow [0, 1]$ is a continuous function for all $i, j \in I$, with $\sum_{i \in I} \eta_i^j \equiv 1$ on V^j , and $\eta_i^i > 0$ on V^i . If $j \leq i$ then $\phi^{ij}(V^{ij})$ is a *closed* subset of $\{v \in V^i : \eta_j^i(v) > 0\}$. If $k < j < i$ and $v \in V^k$ with $\eta_i^k(v) > 0$, $\eta_j^k(v) > 0$ and $v \in V^{ik}$ then $v \in V^{jk}$.
- (iii) $\eta_i^j|_{(s^j)^{-1}(0)} \equiv \eta_i \circ \psi^j$ for all $i, j \in I$.
- (iv) If $i, j, k \in I$ with $k \leq j$ then $\eta_i^k|_{V^{jk}} \equiv \eta_i^j \circ \phi^{jk}$.

We also impose two extra conditions on the V^i, f^i :

- (v) There should exist $M \geq 0$ such that for each $i \in I$, each $v \in V^i$ lies in at most M local boundary components, in the sense of Definition 2.4. Here if the orbifold with g-corners V^i near v is locally modelled on U/Γ near Γu as in §2.2, where Γ is a finite group acting linearly on $\mathbb{R}^n$ and U is a Γ -invariant region with g-corners in $\mathbb{R}^n$, and $u \in U$, then we define the *number of local boundary components* of V^i at v to be the number of local boundary components of U at u , even if some of these local boundary components are identified by the action of Γ .

Then the natural maps $\iota : \partial V^i \rightarrow V^i$ and $\iota : E^i|_{\partial V^i} = \partial E^i \rightarrow E^i$ satisfy $|\iota^{-1}(v)| \leq M$ and $|\iota^{-1}(e)| \leq M$ for all $v \in V^i$ and $e \in E^i$.

- (vi) There should exist a compact subset $T \subseteq Y$ with $f^i(V^i) \subseteq T$ for all $i \in I$.

If the V^i have corners (not g-corners) then (v) holds with $M = \max\{i \in I\}$.

Excellent coordinate systems are really good coordinate systems satisfying an extra condition, which we will need in §3.3 to ensure $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ is finite.

Definition 3.5. Let $(\mathbf{I}, \boldsymbol{\eta})$ be a really good coordinate system for $(X, \mathbf{f})$, and suppose $j \in I$, $l \geq 0$ and W is a connected component of $\partial^l V^j$. We call W *unnecessary* if $W \cap \partial^l V^{ij} = \emptyset$ for all $j < i \in I$ and $\text{Im } \psi^j|_W \subseteq \bigcup_{j \neq i \in I} \text{Im } \psi^i|_{\partial^l V^i} \subseteq \partial^l X$. We call $(\mathbf{I}, \boldsymbol{\eta})$ *excellent* if $\partial^l V^j$ has no unnecessary connected components for all $j \in I$ and $l \geq 0$.

Excellent coordinate systems can be split into *connected components*, a fact which will be important in §3.9.

Definition 3.6. Let Y be an orbifold, X a compact Kuranishi space, $\mathbf{f} : X \rightarrow Y$ be strongly smooth, and $(\mathbf{I}, \boldsymbol{\eta})$ an excellent coordinate system for $(X, \mathbf{f})$, with $\mathbf{I} = (I, (V^i, \dots, \psi^i) : i \in I, \dots)$. A *splitting* of the quadruple $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$ is a decomposition $X = X_1 \amalg \dots \amalg X_n$ of X as a disjoint union of compact, nonempty Kuranishi spaces $X_1, \dots, X_n$, such that for each $i \in I$ we may write $V^i = V_1^i \amalg \dots \amalg V_n^i$ for open and closed subsets $V_1^i, \dots, V_n^i$ of V^i , such that $\text{Im } \psi^i|_{V_a^i} \subseteq X_a$ for $a = 1, \dots, n$. Note that as $(\mathbf{I}, \boldsymbol{\eta})$ is excellent, every connected component of V^i intersects $\text{Im } \psi^i$, so $V_1^i, \dots, V_n^i$ are uniquely determined by $X_1, \dots, X_n$.

Write $\mathbf{f}_a = \mathbf{f}|_{X_a}$, so that $\mathbf{f}_a : X_a \rightarrow Y$ is strongly smooth. Define subsets $I_1, \dots, I_n \subseteq I$ by $I_a = \{i \in I : V_a^i \neq \emptyset\}$ for $a = 1, \dots, n$. For $i \in I_a$ let $E_a^i, s_a^i, \psi_a^i, f_a^i$ be the restrictions of E^i, s^i, ψ^i, f^i to V_a^i . For $i, j \in I_a$ set $\eta_{i,a} = \eta_i|_{X_a}$ and $\eta_{i,a}^j = \eta_i^j|_{V_a^j}$, and if $j \leq i$ write $V_a^{ij}, \phi_a^{ij}, \hat{\phi}_a^{ij}$ for the restrictions of $V^{ij}, \phi^{ij}, \hat{\phi}^{ij}$ to V_a^j, E_a^j . It is easy to show this defines an *excellent coordinate system* $(\mathbf{I}_a, \boldsymbol{\eta}_a) = (\mathbf{I}, \boldsymbol{\eta})|_{X_a}$ for $(X_a, \mathbf{f}_a)$.

Call $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$ *connected* if it does not admit a splitting $X = X_1 \amalg X_2$ as above. Since X_1, X_2 are open and closed in X and nonempty, if X is connected as a topological space then $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$ is connected, but the converse is false.

Lemma 3.7. *Let Y be an orbifold, X a compact Kuranishi space, $\mathbf{f} : X \rightarrow Y$ strongly smooth, and $(\mathbf{I}, \boldsymbol{\eta})$ an excellent coordinate system for $(X, \mathbf{f})$. Then there is a splitting $X = X_1 \amalg X_2 \amalg \dots \amalg X_n$ of $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$, unique up to the order of $X_1, \dots, X_n$, such that $(X_a, \mathbf{f}_a, \mathbf{I}_a, \boldsymbol{\eta}_a)$ is connected for all $a = 1, \dots, n$. We call $X_1, \dots, X_n$ the **connected components** of $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$.*

Proof. Write $\mathbf{I} = (I, (V^i, \dots, \psi^i) : i \in I, \dots)$. Then for each $p \in X$, there exists $i \in I$ with $p \in \text{Im } \psi^i$. Let $\tilde{V}_p^i$ be the connected component of V^i containing $(\psi^i)^{-1}(p)$. Then $U_p^i = \text{Im } \psi^i|_{\tilde{V}_p^i}$ is an open neighbourhood of p in X . Such U_p^i for all $p \in X$ form an open cover for X , so as X is compact we can choose a finite subcover $U_{p_1}^{i_1}, \dots, U_{p_m}^{i_m}$ for X .

Suppose $X = X_1 \amalg \dots \amalg X_n$ is a splitting of $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$. By Definition 3.6, if $p_b \in X_a$ with $U_{p_b}^{i_b} = \text{Im } \psi^{i_b}|_{\tilde{V}_{p_b}^{i_b}}$ then $\tilde{V}_{p_b}^{i_b} \subseteq V_a^{i_b}$, so that $U_{p_b}^{i_b} \subseteq X_a$. As $X_a \neq \emptyset$ and $X = \bigcup_{b=1}^m U_{p_b}^{i_b}$, it follows that for each $a = 1, \dots, n$ we have $p_b \in X_a$ for some b . Therefore $n \leq m$, and the number of pieces we can split $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$ into using Definition 3.6 is bounded.

Choose such a splitting $X = X_1 \amalg \dots \amalg X_n$ with n largest. Then each $(X_a, \mathbf{f}_a, \mathbf{I}_a, \boldsymbol{\eta}_a)$ is *connected*, since otherwise by splitting $(X_a, \mathbf{f}_a, \mathbf{I}_a, \boldsymbol{\eta}_a)$ into 2 pieces we could split X into $n+1$ pieces, a contradiction. If $X = X'_1 \amalg \dots \amalg X'_n$ is another such splitting, then $X_a = \coprod_{b=1, \dots, n: X_a \cap X'_b \neq \emptyset} X_a \cap X'_b$ is a splitting of $(X_a, \mathbf{f}_a, \mathbf{I}_a, \boldsymbol{\eta}_a)$. Since it is connected, we must have $X_a \cap X'_b \neq \emptyset$ for precisely one $b = 1, \dots, n$, with $X_a \cap X'_b = X_a = X'_b$. Therefore $X'_1, \dots, X'_n$ are a reordering of $X_1, \dots, X_n$, and the decomposition is unique up to order. $\square$

We can now define *gauge-fixing data* and *co-gauge-fixing data*.

Definition 3.8. Let the symmetric group S_k of permutations of $\{1, \dots, k\}$ act on $\mathbb{R}^k$ by $\sigma : (x_1, \dots, x_k) \mapsto (x_{\sigma(1)}, \dots, x_{\sigma(k)})$. Define $P = \coprod_{k=0}^{\infty} \mathbb{R}^k / S_k$. For $n = 0, 1, 2, \dots$, define $P_n \subset P$ by $P_n = \coprod_{k=0}^n \mathbb{R}^k / S_k$. Here $\mathbb{R}^0$ is a single point $(\emptyset)$, the ordered list of 0 elements of $\mathbb{R}$, and $S_0 = \{1\}$ acts trivially upon it, so $(\emptyset)$ is a special element of P . Define a *multiplication* $\mu : P \times P \rightarrow P$ by

$$\mu(S_k(x_1, \dots, x_k), S_l(y_1, \dots, y_l)) = S_{k+l}(x_1, \dots, x_k, y_1, \dots, y_l), \quad (33)$$

for $k, l \geq 0$ and $(x_1, \dots, x_k) \in \mathbb{R}^k$, $(y_1, \dots, y_l) \in \mathbb{R}^l$. Then μ is *commutative* and *associative*, with *identity* $(\emptyset)$, and also a *finite map*, since for any point $S_n(x_1, \dots, x_n)$ in P , $\mu^{-1}(S_n(x_1, \dots, x_n))$ is at most 2^n points in $P \times P$. We will use μ in defining fibre products of (co-)gauge-fixing data in §3.8.

Let X be a compact Kuranishi space, Y an orbifold and $\mathbf{f} : X \rightarrow Y$ strongly smooth. A set of *gauge-fixing data* $\mathbf{G}$ for $(X, \mathbf{f})$ consists of an excellent coordinate system $(\mathbf{I}, \boldsymbol{\eta})$ for $(X, \mathbf{f})$, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i) : i \in I, \dots)$, and maps $G^i : E^i \rightarrow P_n \subset P$ for some $n \gg 0$ and all $i \in I$, which are *globally finite*, that is, there exists $N \geq 0$ such that $|(G^i)^{-1}(p)| \leq N$ for all $p \in P$.

By calling $(X, \mathbf{f}, \mathbf{G})$ a *triple*, we will usually mean that X is a compact Kuranishi space (possibly oriented), $\mathbf{f} : X \rightarrow Y$ is strongly smooth, and $\mathbf{G}$ is gauge-fixing data for $(X, \mathbf{f})$. A *splitting* of a triple $(X, \mathbf{f}, \mathbf{G})$ is a splitting $X = X_1 \amalg \dots \amalg X_n$ of $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$ in the sense of Definition 3.6, where $\mathbf{G} = (\mathbf{I}, \boldsymbol{\eta}, G^i : i \in I)$. Then Definition 3.6 gives excellent coordinate systems $(\mathbf{I}_a, \boldsymbol{\eta}_a)$ for $(X_a, \mathbf{f}_a)$, and setting $G_a^i = G^i|_{E_a^i}$ for $i \in I_a$, it is easy to see that $\mathbf{G}|_{X_a} = \mathbf{G}_a = (\mathbf{I}_a, \boldsymbol{\eta}_a, G_a^i : i \in I_a)$ is gauge-fixing data for $(X_a, \mathbf{f}_a)$.

Call $(X, \mathbf{f}, \mathbf{G})$ *connected* if $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$ is connected in the sense of Definition 3.6. Then Lemma 3.7 implies that any triple $(X, \mathbf{f}, \mathbf{G})$ has a splitting into *connected components* $(X_1, \mathbf{f}_1, \mathbf{G}_1), \dots, (X_n, \mathbf{f}_n, \mathbf{G}_n)$, unique up to order.

Let $X, \tilde{X}$ be compact Kuranishi spaces, $\mathbf{f} : X \rightarrow Y, \tilde{\mathbf{f}} : \tilde{X} \rightarrow Y$ be strongly smooth, and $\mathbf{G}, \tilde{\mathbf{G}}$ be gauge-fixing data for $(X, \mathbf{f}), (\tilde{X}, \tilde{\mathbf{f}})$. Suppose the indexing sets $I, \tilde{I}$ of $\mathbf{I}, \tilde{\mathbf{I}}$ in $\mathbf{G}, \tilde{\mathbf{G}}$ are equal, $I = \tilde{I}$. An *isomorphism* $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \mathbf{G}) \rightarrow (\tilde{X}, \tilde{\mathbf{f}}, \tilde{\mathbf{G}})$ consists of a strong diffeomorphism $\mathbf{a} : X \rightarrow \tilde{X}$ and a collection $\mathbf{b}$ of isomorphisms $(b^i, \hat{b}^i) : (V^i, \dots, \psi^i) \rightarrow (\tilde{V}^i, \dots, \tilde{\psi}^i)$ for $i \in I = \tilde{I}$ lifting $\mathbf{a} : X \rightarrow \tilde{X}$, satisfying:

- (a) $\tilde{f}^i \circ b^i \equiv f^i$ for all $i \in I$;
- (b) $\tilde{G}^i \circ \hat{b}^i \equiv G^i$ for all $i \in I$; and
- (c) If $j \leq i$ in I then $b^j(V^{ij}) = \tilde{V}^{ij}$, and the following diagram commutes:

$$\begin{array}{ccc} (V^{ij}, \dots, \psi^j|_{V^{ij}}) & \xrightarrow{(b^j, \hat{b}^j)|_{V^{ij}}} & (\tilde{V}^{ij}, \dots, \tilde{\psi}^j|_{\tilde{V}^{ij}}) \\ \downarrow (\phi^{ij}, \hat{\phi}^{ij}) & & \downarrow (\tilde{\phi}^{ij}, \hat{\tilde{\phi}}^{ij}) \\ (V^i, \dots, \psi^i) & \xrightarrow{(b^i, \hat{b}^i)} & (\tilde{V}^i, \dots, \tilde{\psi}^i). \end{array}$$

Isomorphisms compose, and have inverses which are isomorphisms, in the obvious way. Write $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ for the *automorphism group* of $(X, \mathbf{f}, \mathbf{G})$, that is, the group of isomorphisms $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \mathbf{G}) \rightarrow (X, \mathbf{f}, \mathbf{G})$.

Definition 3.9. Let X be a compact Kuranishi space, Y an orbifold and $\mathbf{f} : X \rightarrow Y$ a strong submersion. A set of *co-gauge-fixing data* $\mathbf{C}$ for $(X, \mathbf{f})$ consists of an excellent coordinate system $(\mathbf{I}, \boldsymbol{\eta})$ for $(X, \mathbf{f})$ in the sense of §3.1, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i) : i \in I, f^i : i \in I, \dots)$, such that $f^i : V^i \rightarrow Y$ is a submersion for all $i \in I$, and maps $C^i : E^i \rightarrow P_n \subset P$ for some $n \gg 0$ and all $i \in I$, such that $C^i \times (f^i \circ \pi^i) : E^i \rightarrow P \times Y$ is a *globally finite* map for all $i \in I$, as in Definition 3.8. *Isomorphisms* $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \mathbf{C}) \rightarrow (\tilde{X}, \tilde{\mathbf{f}}, \tilde{\mathbf{C}})$ and $\text{Aut}(X, \mathbf{f}, \mathbf{C})$ are defined as in Definition 3.8, but with $\tilde{C}^i \circ \hat{b}^i \equiv C^i$ in (b).

By calling $(X, \mathbf{f}, \mathbf{C})$ a *triple*, we will usually mean that X is a compact Kuranishi space, $\mathbf{f} : X \rightarrow Y$ a strong submersion (possibly cooriented), and $\mathbf{C}$ co-gauge-fixing data for $(X, \mathbf{f})$. A *splitting* of $(X, \mathbf{f}, \mathbf{C})$ is a splitting $X = X_1 \amalg \dots \amalg X_n$ of $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$, where $\mathbf{C} = (\mathbf{I}, \boldsymbol{\eta}, C^i : i \in I)$. Then Definition 3.6 gives excellent $(\mathbf{I}_a, \boldsymbol{\eta}_a)$ for $(X_a, \mathbf{f}_a)$, and $\mathbf{C}|_{X_a} = \mathbf{C}_a = (\mathbf{I}_a, \boldsymbol{\eta}_a, C_a^i : i \in I_a)$ is co-gauge-fixing data for $(X_a, \mathbf{f}_a)$, where $C_a^i = C^i|_{E_a^i}$ for $i \in I_a$.

Call $(X, \mathbf{f}, \mathbf{C})$ *connected* if $(X, \mathbf{f}, \mathbf{I}, \boldsymbol{\eta})$ is connected in the sense of Definition 3.6. Then by Lemma 3.7, any $(X, \mathbf{f}, \mathbf{C})$ has a splitting into *connected components* $(X_1, \mathbf{f}_1, \mathbf{C}_1), \dots, (X_n, \mathbf{f}_n, \mathbf{C}_n)$, unique up to order.

Remark 3.10. Note that we do *not* require the maps $G^i : E^i \rightarrow P$, $C^i : E^i \rightarrow P$ to be continuous, or smooth, etc., they are just arbitrary maps. Nor do we impose compatibility conditions between G^i, G^j under coordinate transformations $(\phi^{ij}, \hat{\phi}^{ij})$. Their only purpose is to ensure that $\text{Aut}(X, \mathbf{f}, \mathbf{G}), \text{Aut}(X, \mathbf{f}, \mathbf{C})$ are finite, and this will follow from $G^i, C^i \times (f^i \circ \pi^i)$ globally finite.

To see why gauge-fixing data is natural for homology, and co-gauge-fixing data for cohomology, read §3.7 on pushforwards and pullbacks.

For Property 3.1(e) we define (co-)gauge-fixing data *with corners*.

Definition 3.11. Let X be a compact Kuranishi space, and $\mathbf{f} : X \rightarrow Y$ be strongly smooth. We say that a good or very good coordinate system $\mathbf{I}$, or a really good or excellent coordinate system $(\mathbf{I}, \boldsymbol{\eta})$, or gauge-fixing data $\mathbf{G} = (\mathbf{I}, \boldsymbol{\eta}, G^i : i \in I)$, or co-gauge-fixing data $\mathbf{C} = (\mathbf{I}, \boldsymbol{\eta}, C^i : i \in I)$, for $(X, \mathbf{f})$, are *without boundary*, or *with boundary*, or *with corners*, if the orbifolds V^i for all $i \in I$ are without boundary, or with boundary, or with corners, respectively. These imply that the Kuranishi space X is without boundary, or with boundary, or with corners, respectively.

Sections 3.2–3.8 prove Property 3.1(a)–(h) for (co-)gauge-fixing data. Since the two are so similar, we generally give proofs only for gauge-fixing data.

3.2 Existence of (co-)gauge-fixing data

To show that (co-)gauge-fixing data exists for every pair $(X, \mathbf{f})$, we begin by showing that $(X, \mathbf{f})$ admits good, very good, really good and excellent coordinate systems. Fukaya and Ono prove [25, Lem. 6.3], [24, Lem. A1.11]:

Proposition 3.12. *Let X be a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ a strongly smooth map, and $\{U_\alpha : \alpha \in A\}$ an open cover for X . Then there exist good coordinate systems $\mathbf{I}$ for X and $(X, \mathbf{f})$ such that for each $i \in I$ we have $\text{Im } \psi^i \subseteq U_\alpha$ for some $\alpha \in A$.*

Proposition 3.13. *Let X be a compact Kuranishi space, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ a strongly smooth map. Then there exists a very good coordinate system $\mathbf{I}$ for $(X, \mathbf{f})$.*

Proof. By Proposition 3.12 there exists a good coordinate system $\mathbf{I} = (I, \leq, (V^i, \dots, \psi^i) : i \in I, \dots)$ for $(X, \mathbf{f})$. Define $\tilde{I} = \{\dim V^i : i \in I\}$. For each $k \in \tilde{I}$, we shall define a Kuranishi neighbourhood $(\tilde{V}^k, \dots, \tilde{\psi}^k)$ on X by gluing together the $(V^i, \dots, \psi^i)$ for all $i \in I$ with $\dim V^i = k$. We first explain how to glue together two such Kuranishi neighbourhoods. Let $i, j \in I$ with $i \neq j$ and $\dim V^i = \dim V^j = k$. Either $i \leq j$ or $j \leq i$ as $\leq$ is a total order; suppose $j \leq i$. Then we have a triple $(V^{ij}, \phi^{ij}, \hat{\phi}^{ij})$ by Definition 3.2.

As $\dim V^i = \dim V^j$, it follows that the coordinate change $(\phi^{ij}, \hat{\phi}^{ij})$ is an *isomorphism* from $(V^{ij}, E^j|_{V^{ij}}, s^j|_{V^{ij}}, \psi^j|_{V^{ij}})$ to its image in $(V^i, \dots, \psi^i)$. In particular, $\phi^{ij} : V^{ij} \rightarrow V^i$ is a diffeomorphism $V^{ij} \rightarrow \text{Im } \phi^{ij}$. The obvious thing to do now is to glue V^i and V^j together using ϕ^{ij} , that is, to form the topological space $(V^i \amalg V^j)/\sim$, where $\sim$ is the equivalence relation on $V^i \amalg V^j$ generated by ϕ^{ij} , with $v \sim v'$ if $v = v'$, or $v \in V^{ij}$ and $v' = \phi^{ij}(v) \in V^i$, or vice versa. However, there is a problem: since V^{ij} and $\text{Im } \phi^{ij}$ may not be closed in V^j, V^i , this $(V^i \amalg V^j)/\sim$ may not be *Hausdorff*, and so not an orbifold.

Our solution is to make V^i, V^j a little smaller, that is, we replace them by open sets $\tilde{V}^i \subset V^i$ and $\tilde{V}^j \subset V^j$, such that $(\tilde{V}^i \amalg \tilde{V}^j)/\sim$ is Hausdorff, and so an orbifold. If $(V^i \amalg V^j)/\sim$ is not Hausdorff then there exist sequences $(v_n)_{n=1}^\infty$ in V^{ij} so that $v_n \rightarrow v^j$ in $V^j \setminus V^{ij}$ as $n \rightarrow \infty$ and $\phi^{ij}(v_n) \rightarrow v^i$ in $V^i \setminus \text{Im } \phi^{ij}$ as $n \rightarrow \infty$. Then v^i, v^j are distinct points in $(V^i \amalg V^j)/\sim$ which do not admit disjoint open neighbourhoods, contradicting Hausdorffness, since any neighbourhoods contain v_n for $n \gg 0$.

To make $(\tilde{V}^i \amalg \tilde{V}^j)/\sim$ Hausdorff, we have to ensure that for all such $(v_n)_{n=1}^\infty$ and v^i, v^j , either $v^i \notin \tilde{V}^i$ or $v^j \notin \tilde{V}^j$. Loosely speaking, the ‘boundaries’ of V^{ij} in $\tilde{V}^i$ and $\tilde{V}^j$ may not intersect. Note however that we cannot make $\tilde{V}^i, \tilde{V}^j$ too small, as we need to preserve the condition $\bigcup_{i \in I} \text{Im } \psi^i = X$, that is, the choices of $\tilde{V}^i$ for all $i \in I$ must satisfy $\bigcup_{i \in I} \text{Im } \psi^i|_{\tilde{V}^i} = X$. These conditions are consistent, since gluing $(s^i)^{-1}(0)$ and $(s^j)^{-1}(0)$ together using ϕ^{ij} yields an open subset $\text{Im } \psi^i \cup \text{Im } \psi^j$ of X , which is Hausdorff, so the points v^i, v^j which need to be excluded cannot lie in $(s^i)^{-1}(0), (s^j)^{-1}(0)$, the domains of ψ^i, ψ^j .

We need to choose open $\tilde{V}^i \subset V^i$ for all $i \in I$, which are small enough that whenever $j \leq i \in I$ with $\dim V^i = \dim V^j$ then $(\tilde{V}^i \amalg \tilde{V}^j)/\sim$ is Hausdorff, and which are large enough that $\bigcup_{i \in I} \text{Im } \psi^i|_{\tilde{V}^i} = X$. This is possible. One way to do it is to use ideas on *really good coordinate systems* below. Follow the proof of Proposition 3.15 to choose auxiliary partitions of unity $\eta_i : i \in I$ on X and $\eta_i^j : i \in I$ on V^j satisfying conditions (a)–(d) of that proof, possibly making the V^j smaller along the way, and define $\tilde{V}^i = \{v \in V^i : \eta_i^i(v) > 0\}$. As in

Proposition 3.15 the resulting $\tilde{V}^j$ have partitions of unity $\eta_i^j|_{\tilde{V}^j} : i \in I$ satisfying Definition 3.4 with $\bigcup_{i \in I} \text{Im } \psi^i|_{\tilde{V}^i} = X$, and by Remark 3.14 this ensures that $(\tilde{V}^i \amalg \tilde{V}^j)/\sim$ is Hausdorff.

This explains how to glue two Kuranishi neighbourhoods $(V^i, \dots, \psi^i)$, $(V^j, \dots, \psi^j)$ for $i \neq j \in I$ with $\dim V^i = \dim V^j = k$. To make $\tilde{V}^k$ we repeat this construction finitely many times to glue together V^i for all $i \in I$ with $\dim V^i = k$, so that $\tilde{V}^k = (\bigsqcup_{i \in I: \dim V^i = k} \tilde{V}^i)/\sim$, where $\sim$ is the equivalence relation generated by ϕ^{ij} for all $j \leq i \in I$ with $\dim V^i = \dim V^j = k$, and the $\tilde{V}^i$ are chosen small enough to make $\tilde{V}^k$ Hausdorff, and thus an orbifold.

Since the $(\phi^{ij}, \hat{\phi}^{ij})$ are isomorphisms with their images, we can also glue the restrictions of E^i, s^i, ψ^i to $\tilde{V}^i$ together to make $\tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k$ on $\tilde{V}^k$. That is, we have a Kuranishi neighbourhood $(\tilde{V}^k, \tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k)$ on X with given isomorphisms $(\tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k)|_{\tilde{V}^i} \cong (E^i, s^i, \psi^i)|_{\tilde{V}^i}$, for all $i \in I$ with $\dim V^i = k$, regarding $\tilde{V}^i$ as an open subset of both $\tilde{V}^k$ and V^i . This gives the Kuranishi neighbourhoods $(\tilde{V}^k, \tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k)$ for $k \in \tilde{I}$. Since $\bigcup_{i \in I} \text{Im } \psi^i|_{\tilde{V}^i} = X$, we have $\bigcup_{k \in \tilde{I}} \text{Im } \tilde{\psi}^k = X$.

We have smooth $f^i : V^i \rightarrow Y$ for $i \in I$ representing $\mathbf{f}$. The compatibilities $f^i \circ \phi^{ij} \equiv f^j|_{V^{ij}}$ imply that we may glue these to give smooth maps $\tilde{f}^k : \tilde{V}^k \rightarrow Y$ representing $\mathbf{f}$, with $\tilde{f}^k|_{\tilde{V}^i} = f^i|_{\tilde{V}^i}$, for all $i \in I$ with $\dim V^i = k$.

If $k, l \in \tilde{I}$ with $l < k$ then for all $i \in I$ with $\dim V^i = k$ and $j \in I$ with $\dim V^j = l$ we have $j \leq i$, so $(V^{ij}, \phi^{ij}, \hat{\phi}^{ij})$ is given. Define an open $\tilde{V}^{kl} \subseteq \tilde{V}^l$ by $\tilde{V}^{kl} = \bigcup_{i, j \in I: \dim V^i = k, \dim V^j = l} (V^{ij} \cap \tilde{V}^j \cap (\phi^{ij})^{-1}(\tilde{V}^i))$. There is then a unique coordinate transformation $(\check{\phi}^{kl}, \hat{\phi}^{kl}) : (\tilde{V}^{kl}, \tilde{E}^l|_{\tilde{V}^{kl}}, \tilde{s}^l|_{\tilde{V}^{kl}}, \tilde{\psi}^l|_{\tilde{V}^{kl}}) \rightarrow (\tilde{V}^k, \dots, \tilde{\psi}^k)$ which restricts to $(\phi^{ij}, \hat{\phi}^{ij})$ on $V^{ij} \cap \tilde{V}^j \cap (\phi^{ij})^{-1}(\tilde{V}^i)$ for all $i, j \in I$ with $\dim V^i = k$ and $\dim V^j = l$. It is now easy to check that all this data $\tilde{\mathbf{I}} = (\tilde{I}, (\tilde{V}^k, \dots, \tilde{\psi}^k), \tilde{f}^k : k \in \tilde{I}, \dots)$ satisfies the conditions for a very good coordinate system for $(X, \mathbf{f})$. This completes the proof. $\square$

Remark 3.14. In the proof of Proposition 3.13 we glued topological spaces V^j, V^i together on $V^{ij} \subseteq V^j$ and $\phi^{ij}(V^{ij}) \subseteq V^i$ using ϕ^{ij} , and we had a problem because the resulting topological space $V^i \amalg V^j/\sim$ may not be Hausdorff. The existence of functions η_i^j satisfying Definition 3.4(ii),(iv) eliminates this problem, so that $(\bigsqcup_{i \in I} V^i)/\sim$ is a Hausdorff topological space. This is useful, as it excludes some bad behaviour in constructions where we choose data over each V^i with compatibilities on the V^{ij} , as we will do many times later.

To see this, suppose $(V^i \amalg V^j)/\sim$ is not Hausdorff. Then there exists a sequence $(v_n)_{n=1}^\infty$ in V^{ij} such that $\phi^{ij}(v_n) \rightarrow v^i$ in $V^i \setminus \phi^{ij}(V^{ij})$ and $v_n \rightarrow v^j$ in $V^j \setminus V^{ij}$ as $n \rightarrow \infty$. Since η_j^i, η_j^j are continuous functions on V^i, V^j we have $\eta_j^i \circ \phi^{ij}(v_n) \rightarrow \eta_j^i(v^i)$ and $\eta_j^j(v_n) \rightarrow \eta_j^j(v^j)$ as $n \rightarrow \infty$. Definition 3.4(iv) implies that $\eta_j^i \circ \phi^{ij}(v_n) = \eta_j^j(v_n)$ for all n , so $\eta_j^i(v^i) = \eta_j^j(v^j)$. But Definition 3.4(ii) gives $\eta_j^j(v^j) > 0$, and as $\phi^{ij}(V^{ij})$ is a closed in $\{v \in V^i : \eta_j^i(v) > 0\}$ and $v^i \in \overline{\phi^{ij}(V^{ij})} \setminus \phi^{ij}(V^{ij})$ we deduce that $\eta_j^i(v^i) = 0$, a contradiction.

Proposition 3.15. *Let X be a compact Kuranishi space, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ a strongly smooth map. Then there exists a really good coordinate system $(\mathbf{I}, \boldsymbol{\eta})$ for $(X, \mathbf{f})$.*

Proof. By Proposition 3.13 we can choose a very good coordinate system $\mathbf{I}$ for $(X, \mathbf{f})$. Then $\{\text{Im } \psi^i : i \in I\}$ is an open cover for X . By general topology we may choose a partition of unity $\{\eta_i : i \in I\}$ for X subordinate to $\{\text{Im } \psi^i : i \in I\}$, that is, $\eta_i : X \rightarrow [0, 1]$ is continuous, and $\sum_{i \in I} \eta_i \equiv 1$, and the closure of the support of η_i is contained in $\text{Im } \psi^i$, that is, $\overline{\{p \in X : \eta_i(p) > 0\}} \subseteq \text{Im } \psi^i$.

Next, for each $j \in I$ we choose a partition of unity $\{\eta_i^j : i \in I\}$ on V^j , by induction in the order $\leq$ on I , satisfying:

- (a) $\eta_i^j : V^j \rightarrow [0, 1]$ is a continuous function for all $i, j \in I$, with $\sum_{i \in I} \eta_i^j \equiv 1$.
- (b) If $j \leq i$ in I then $\text{supp}(\eta_i^j) \subseteq V^{ij}$ and $\overline{\phi^{ij}(V^{ij} \cap \text{supp}(\eta_i^j))}$ is closed in V^i .
- (c) $\eta_i^j|_{(s^j)^{-1}(0)} \equiv \eta_i \circ \psi^j$ for all $i \in I$.
- (d) If $k \in I$ with $k < j$ then $\eta_i^j \circ \phi^{jk} \equiv \eta_i^k|_{V^{jk}}$.

Here $\eta_i^k|_{V^{jk}}$ makes sense in (d) since when we are choosing $\{\eta_i^j : i \in I\}$ we have already chosen $\{\eta_i^k : i \in I\}$, as $k < j$. The second part of (b) means that if $\phi^{ij}(V^{ij})$ has an ‘open edge’ in V^i , then $\eta_j^j = 0$ near this edge in V^{ij} , or equivalently by (d), $\eta_j^i|_{\phi^{ij}(V^{ij})} = 0$ near this edge in $\phi^{ij}(V^{ij})$.

Observe that conditions (c),(d) prescribe η_i^j for $i \in I$ on the subset

$$(s^j)^{-1}(0) \cup \bigcup_{k \in I: k < j} \phi^{jk}(V^{jk}). \quad (34)$$

These prescribed values are continuous on (34), and consistent on the overlaps between $(s^j)^{-1}(0)$ and $\phi^{jk}(V^{jk})$ and $\phi^{jl}(V^{jl})$ for different $k, l < j$ in I , using Definition 3.2 and conditions (a)–(d) for previous choices $\{\eta_i^k : i \in I\}$ for $k < j$. However, there is a problem in extending the η_i^j continuously from (34) to all of V^j , since $\phi^{jk}(V^{jk})$ may not be closed in V^j , and the prescribed values of η_i^j may not extend continuously to the closure of (34) in V^j .

We can avoid this problem by making all the sets V^k, V^{jk} *slightly smaller*. That is, we replace V^k and V^{jk} by open sets $\tilde{V}^j \subset V^j$ and $\tilde{V}^{jk} \subset V^{jk}$ such that the previous conditions still hold, in particular, $X = \bigcup_{i \in I} \text{Im } \psi_i|_{\tilde{V}^i}$, and in (b), we shrink the ‘open edges’ of $\phi^{jk}(V^{jk})$ in V^k by little enough that $\eta_j^j = 0$ near the new open edges, but we make $\tilde{V}^{jk}$ small enough that $\overline{\phi^{jk}(\tilde{V}^{jk})} \subseteq \phi^{jk}(V^{jk})$, taking closures in V^j , so the closure of the new set (34) lies in the old set (34). This is automatic if the closure of $\tilde{V}^{ij}$ in V^{ij} is compact. With this replacement, the η_i^j have unique continuous extensions from their prescribed values on (34) to the closure of (34).

We can then choose arbitrary continuous extensions $\eta_i^j : V^j \rightarrow [0, 1]$ from the values on this closed set to V^j , satisfying (b) and $\sum_{i \in I} \eta_i^j \equiv 1$. Therefore by induction we can choose η_i^j for all $i, j \in I$ satisfying (a)–(d), possibly with smaller sets V^k, V^{jk} . We have now constructed a very good coordinate system $\mathbf{I}$ and functions η_i, η_i^j satisfying all of Definition 3.4 except the last two parts of (i), the last three parts of (ii), and (v),(vi). But we do know that $\{p \in X : \eta_i(p) > 0\} \subseteq \text{Im } \psi^i$, and $\{v \in V^j : \eta_i^j(v) > 0\} \subseteq V^{ij}$ if $j \leq i$.

For (v), as X is compact and the map taking $p \in X$ to the number of local boundary components of X at p is upper semicontinuous, there exists $M \geq 0$ such that each $p \in X$ lies in at most M local boundary components. If $v \in (\psi^i)^{-1}(0) \subseteq V^i$ with $\psi^i(v) = p$ in X then the number of local boundary components of V^i containing v is the same as the number of local boundary components of X containing p , and so at most M . Conversely, if $v \in V^i$ and the number of local boundary components of V^i containing v is greater than M , then $v \notin (\psi^i)^{-1}(0)$. Thus, the subset of $v \in V^i$ lying in more than M local boundary components is a closed subset of V^i not intersecting $(\psi^i)^{-1}(0)$. We will delete this subset, which will make (v) hold.

For (vi), as X is compact, $f(X) \subseteq Y$ is compact. Choose an open neighbourhood U of $f(X)$ in Y whose closure $T = \bar{U}$ in Y is compact. To complete the proof we now replace V^i by the open subset $\tilde{V}^i = \{v \in V^i : \eta_i^i(v) > 0, v \text{ lies in at most } M \text{ local boundary components, } f^i(y) \in U\}$ for each $i \in I$, and replace I by $\tilde{I} = \{i \in I : \tilde{V}^i \neq \emptyset\}$, and replace V^{ij} by the open subset $\tilde{V}^{ij} = V^{ij} \cap \tilde{V}^j \cap (\phi^{ij})^{-1}(\tilde{V}^i)$ for all $i, j \in \tilde{I}$. With these replacements, we have $\text{Im } \psi^i = \{p \in X : \eta_i(p) > 0\}$ and $V^{ij} = \{v \in V^j : \eta_i^j(v) > 0\}$ for all i, j . These imply the remaining parts of (i),(ii) except $\phi^{ij}(\tilde{V}^{ij})$ closed in $\{v \in V^i : \eta_j^i(v) > 0\}$ for $j \leq i$, and this follows from (b) above. Hence $(\mathbf{I}, \boldsymbol{\eta})$ is a *really good coordinate system* for $(X, \mathbf{f})$. $\square$

Here is an algorithm for converting really good coordinate systems into excellent coordinate systems, by throwing away all unnecessary connected components of $\partial^k V^i$ for all $i \in I$ and $k \geq 0$.

Algorithm 3.16. Let $(\mathbf{I}, \boldsymbol{\eta})$ be a really good coordinate system for $(X, \mathbf{f})$. We shall construct open sets $\tilde{V}^i$ in V^i and $\tilde{V}^{ij} \subseteq V^{ij}$ for all $j \leq i \in I$, such that defining $\tilde{I} = \{i \in I : \tilde{V}^i \neq \emptyset\}$, and for $i, j \in \tilde{I}$ setting $(\tilde{V}^i, \tilde{E}^i, \tilde{s}^i, \tilde{\psi}^i) = (\tilde{V}^i, E^i|_{\tilde{V}^i}, s^i|_{\tilde{V}^i}, \psi^i|_{\tilde{V}^i})$, $\tilde{f}^i = f^i|_{\tilde{V}^i}$, $\tilde{\eta}_i = \eta_i$, $\tilde{\eta}_i^j = \eta_i^j|_{\tilde{V}^j}$, and for $j \leq i$ in $\tilde{I}$ defining $\tilde{V}^{ij} = V^{ij} \cap \tilde{V}^j \cap (\phi^{ij})^{-1}(\tilde{V}^i)$ and $(\tilde{\phi}^{ij}, \tilde{\phi}^{ij}) = (\phi^{ij}, \hat{\phi}^{ij})|_{\tilde{V}^{ij}}$, this data defines an *excellent coordinate system* $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ for $(X, \mathbf{f})$. We define $\tilde{V}^j$ by induction on decreasing $j \in I$.

In the inductive step, for fixed $j \in I$, suppose we have defined $\tilde{V}^i$ for all $i \in I$ with $j < i \in I$, and $\tilde{V}^{ik}$ for all $j < i \leq k$ in I . Define $\tilde{V}^{ij} \subseteq V^{ij} \subseteq V^j$ for all $j < i$ in I by $\tilde{V}^{ij} = (\phi^{ij})^{-1}(\tilde{V}^i)$. Then define $\tilde{V}^j$ to be the complement in V^j of the union over all $l = 0, \dots, \dim V^j$ of the images under the projection $\partial^l V^j \rightarrow V^j$ of all connected components W of $\partial^l V^j$ such that $W \cap \partial^l \tilde{V}^{ij} = \emptyset$ for all $j < i \in I$ and

$$\text{Im } \psi^j|_W \subseteq \bigcup_{i \in I: i < j} \text{Im } \psi^i|_{\partial^l V^i} \cup \bigcup_{i \in I: i > j} \text{Im } \psi^i|_{\partial^l \tilde{V}^i}. \quad (35)$$

For each l , the union of such W is an open and closed set in $\partial^l V^j$, so its image is closed in V^j . Thus $\tilde{V}^j$ is the complement of a finite union of closed sets in V^j , and so is open in V^j . Set $\tilde{V}^{jj} = \tilde{V}^j$. This completes the inductive step.

The condition (35) on W ensures that subtracting these components W does not change the fact that $X = \bigcup_{i \in I} \text{Im } \psi^i$, and therefore $X = \bigcup_{i \in I} \text{Im } \tilde{\psi}^i$. One

can prove this by induction on $i \in I$, by checking that if $X = \bigcup_{i \in I: i \leq j} \text{Im } \psi^i \cup \bigcup_{i \in I: i > j} \text{Im } \check{\psi}^i$ then $X = \bigcup_{i \in I: i < j} \text{Im } \psi^i \cup \bigcup_{i \in I: i \geq j} \text{Im } \check{\psi}^i$. Also, subtracting components W in this way does not change the fact that the last sentence of Definition 3.4(i) and the last two sentences of Definition 3.4(ii) hold. For the middle sentence of Definition 3.4(ii), note that for $j < i$ we only ever delete some component W of $\partial^l V^{ij}$ if in a previous inductive step we have also deleted the component of $\partial^l V^i$ containing $\phi^{ij}(W)$. Doing both these deletions together does not change the fact that $\phi^{ij}(V^{ij})$ is a closed subset of $\{v \in V^i : \eta_j^i(v) > 0\}$. So $(\check{I}, \check{\eta})$ is a *really good coordinate system* for $(X, \mathbf{f})$, since (I, η) is.

We claim that $(\check{I}, \check{\eta})$ is an *excellent coordinate system* for $(X, \mathbf{f})$. Suppose for a contradiction that $\check{W}$ is an unnecessary connected component of some $\partial^l \check{V}^j$. Then $\check{W}$ is an open dense subset of a unique connected component W of $\partial^l V^j$. The difference $W \setminus \check{W}$ is a union of unnecessary components of $\partial^{l'} V^j$ for $l' > l$, and does not affect the following argument. As $\check{W} \cap \check{V}^{ij} = \emptyset$ for $j < i \in I$ we have $W \cap \check{V}^{ij} = \emptyset$ for $j < i$. Also $\text{Im } \check{\psi}^j|_{\check{W}} \subseteq \bigcup_{j \neq i \in I} \text{Im } \check{\psi}^i|_{\partial^l \check{V}^i} \subseteq \partial^l X$ implies that (35) holds. So W is one of the connected components of $\partial^l V^j$ which is deleted from $\check{V}^j$, a contradiction. Thus $(\check{I}, \check{\eta})$ is excellent.

Together with Proposition 3.15 this yields:

Corollary 3.17. *Let X be a compact Kuranishi space, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ a strongly smooth map. Then there exists an excellent coordinate system (I, η) for $(X, \mathbf{f})$.*

We now prove Property 3.1(a):

Theorem 3.18. *Let X be a compact Kuranishi space, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ a strongly smooth map. Then $(X, \mathbf{f})$ admits gauge-fixing data $\mathbf{G}$.*

If Γ is a finite group of strong diffeomorphisms $\mathbf{a} : X \rightarrow X$ with $\mathbf{f} \circ \mathbf{a} = \mathbf{f}$ then we can choose $\mathbf{G}$ Γ -invariant, i.e., there is a morphism $\Gamma \rightarrow \text{Aut}(X, \mathbf{f}, \mathbf{G})$ mapping $\mathbf{a} \mapsto (\mathbf{a}, \mathbf{b}_\mathbf{a})$.

If X is without boundary, or with boundary, or with corners, then we can choose $\mathbf{G}$ without boundary, or with boundary, or with corners, respectively. All this also holds for co-gauge-fixing data $\mathbf{C}$, with $\mathbf{f}$ a strong submersion.

Proof. By Corollary 3.17 there exists an excellent coordinate system $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i), f^i : i \in I), \dots$ for $(X, \mathbf{f})$. For each $i \in I$ choose an *injective* map $G^i : E^i \rightarrow P_n$ for $n \gg 0$. This is clearly possible, and we can even choose G^i to be smooth, as a generic smooth map $G^i : E^i \rightarrow \mathbb{R}^n/S_n$ is injective provided $n > 2 \dim E^i$. As G^i is injective it is globally finite, with $N = 1$. Thus $\mathbf{G} = (\mathbf{I}, \eta, G^i : i \in I)$ is gauge-fixing data for $(X, \mathbf{f})$.

When $\mathbf{f}$ is a strong submersion, to construct co-gauge-fixing data $\mathbf{C}$ we first note that in Proposition 3.12–Corollary 3.17 we can work throughout with good, very good, really good and excellent coordinate systems in which $f^i : V^i \rightarrow Y$ are submersions. We again choose injective $C^i : E^i \rightarrow P_n$ for $n \gg 0$, and C^i injective implies $C^i \times (f^i \circ \pi^i)$ is globally finite, with $N = 1$, so $\mathbf{C} = (\mathbf{I}, \eta, C^i : i \in I)$ is co-gauge-fixing data for $(X, \mathbf{f})$.

Now let $\Gamma \subseteq \text{Aut}(X, \mathbf{f})$ be a finite subgroup. We must show we can choose $\mathbf{G}, \mathbf{C}$ to be Γ -invariant. This is similar to the problem of choosing (co-)gauge-fixing data for $(X/\Gamma, \mathbf{f} \circ \pi)$ for $\pi : X \rightarrow X/\Gamma$ the natural projection, which we have already proved is possible in the first part. We begin with the proof of Proposition 3.13, which constructs a very good coordinate system $\tilde{\mathbf{I}}$ for $(X, \mathbf{f})$ starting with a good coordinate system $\mathbf{I}$ for $(X, \mathbf{f})$, which exists by Proposition 3.12. This $\mathbf{I}$ is built from finitely many neighbourhoods $(V_p, \dots, \psi_p)$ for $p \in X$.

To choose $\mathbf{I}$ to be Γ -invariant, we take each $(V_p, \dots, \psi_p)$ to be invariant under $\{\mathbf{a} \in \Gamma : a(p) = p\}$, with V_p small enough that $\text{Im } \psi_p \cap (a \cdot \text{Im } \psi_p) = \emptyset$ whenever $\mathbf{a} \in \Gamma$ with $a(p) \neq p$. Every $p \in X$ admits such a Kuranishi neighbourhood. Given such (V_p, E_p, s_p, ψ_p) , for $\mathbf{a} \in \Gamma$ define $(V_{a(p)}, E_{a(p)}, s_{a(p)}, \psi_{a(p)})$ to be $\mathbf{a}_*(V_p, E_p, s_p, \psi_p)$, a Kuranishi neighbourhood of $a(p) \in X$.

As $(V_p, \dots, \psi_p)$ is invariant under $\{\mathbf{a} \in \Gamma : a(p) = p\}$, this $(V_{a(p)}, \dots, \psi_{a(p)})$ depends only on $a(p)$ rather than on $\mathbf{a}$, and the finite set of neighbourhoods $(V_{a(p)}, \dots, \psi_{a(p)})$ for $\mathbf{a} \in \Gamma$ is invariant under Γ , with $\text{Im } \psi_{a(p)} \cap \text{Im } \psi_{a'(p)} = \emptyset$ when $a(p) \neq a'(p)$. Thus the disjoint union $\coprod_{a(p): \mathbf{a} \in \Gamma} (V_{a(p)}, \dots, \psi_{a(p)})$ is a single Γ -invariant Kuranishi neighbourhood. We can now follow the proof of Proposition 3.12 to make $\mathbf{I}$ using Γ -invariant Kuranishi neighbourhoods $(V^i, \dots, \psi^i) = \coprod_{a(p): \mathbf{a} \in \Gamma} (V_{a(p)}, \dots, \psi_{a(p)})$.

Then in Proposition 3.13 we construct a very good coordinate system $\tilde{\mathbf{I}}$ for $(X, \mathbf{f})$ by gluing together the neighbourhoods $(V^i, \dots, \psi^i)$ of each dimension in $\mathbf{I}$. As each $(V^i, \dots, \psi^i)$ is Γ -invariant, we can do this gluing in a Γ -invariant way, to make $\tilde{\mathbf{I}}$ Γ -invariant. That is, $\mathbf{a} \in \Gamma$ lifts to $(\mathbf{a}, \mathbf{b}_{\mathbf{a}}) \in \text{Aut}(X, \mathbf{f}, \tilde{\mathbf{I}})$, where $\mathbf{b}_{\mathbf{a}} = (b_{\mathbf{a}}^i, \hat{b}_{\mathbf{a}}^i : i \in \tilde{I})$, with each $(b_{\mathbf{a}}^i, \hat{b}_{\mathbf{a}}^i) : (\tilde{V}^i, \dots, \tilde{\psi}^i) \rightarrow (\tilde{V}^i, \dots, \tilde{\psi}^i)$ an isomorphism lifting $\mathbf{a}$.

In Proposition 3.15 it is easy to extend this to a Γ -invariant really good coordinate system $(\mathbf{I}, \boldsymbol{\eta})$ for $(X, \mathbf{f})$, we just choose η_i, η_i^j to be Γ -invariant. Algorithm 3.16 then automatically yields a Γ -invariant excellent coordinate system. Thus in Corollary 3.17, we can take $(\mathbf{I}, \boldsymbol{\eta})$ to be Γ -invariant. Finally, in the first part of the theorem, rather than choosing $G^i : E^i \rightarrow P_n$, $C^i : E^i \rightarrow P_n$ injective, we take G^i, C^i to be the pullbacks to E^i of injective maps $E^i/\Gamma \rightarrow P_n$ for $n \gg 0$. Then G^i, C^i are Γ -invariant, and globally finite in the sense of Definition 3.8, with $N = |\Gamma|$. This implies $C^i \times (f^i \circ \pi^i)$ is also globally finite, with $N = |\Gamma|$. Hence we can choose $\mathbf{G}, \mathbf{C}$ to be Γ -invariant.

If X is without boundary, or with boundary, or with corners, then we can go through the whole of §3.2 working only with V^i without boundary, or with boundary, or with corners, respectively, and so construct $\mathbf{G}, \mathbf{C}$ without boundary, or with boundary, or with corners, as required. $\square$

3.3 Finiteness of automorphism groups

We now prove Property 3.1(b) for (co-)gauge-fixing data.

Proposition 3.19. *If $(\mathbf{I}, \boldsymbol{\eta})$ is an excellent coordinate system for $(X, \mathbf{f})$, with X compact, then for each $j \in I$ there are only finitely many connected components W of V^j such that $W \cap V^{ij} = \emptyset$ for all $j < i \in I$.*

Proof. Let $j \in I$ and W_a^j for $a \in A^j$ be the connected components of V^j with $W_a^j \cap V^{ij} = \emptyset$ for all $j < i \in I$. Define $U^j = V^j \setminus \bigcup_{a \in A} W_a^j$. Then V^j is the disjoint union of U^j and the W_a^j for $a \in A^j$, and U^j and each W_a^j are open and closed in V^j . Note that openness holds as V^j is an orbifold; for general topological spaces connected components are closed but need not be open. For instance, the compact subset $\{0\} \cup \{1/n : n = 1, 2, \dots\}$ in $\mathbb{R}$, has a connected component $\{0\}$ which is closed but not open.

Now $C = X \setminus (\text{Im } \psi^j|_{U^j} \cup \bigcup_{j \neq i \in I} \text{Im } \psi^i)$ is a closed subset of X , since $\text{Im } \psi^j|_{U^j}$ and each $\text{Im } \psi^i$ is open, so C is compact as X is compact and Hausdorff. Also $C \subseteq \bigcup_{a \in A^j} \text{Im } \psi^j|_{W_a^j} \subseteq X$ since $X = \bigcup_{j \in I} \text{Im } \psi^j$. And as $(\mathbf{I}, \boldsymbol{\eta})$ is excellent, Definition 3.5 implies that $C \cap \text{Im } \psi^j|_{W_a^j} \neq \emptyset$ for each $a \in A^j$, since otherwise W_a^j would be an unnecessary component of V^j . Therefore the sets $C \cap \text{Im } \psi^j|_{W_a^j}$ for $a \in A^j$ form a cover of C by nonempty, disjoint open sets. As C is compact there must be a finite subcover, but the only possible subcover is the whole thing, since the sets are nonempty and disjoint. Hence A^j is finite. $\square$

Theorem 3.20. *Let X be a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ be strongly smooth, and $\mathbf{G}$ gauge-fixing data for $(X, \mathbf{f})$. Then the automorphism group $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ is finite. The same holds for co-gauge-fixing data.*

Proof. As in Definition 3.8 we write elements of $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ as $(\mathbf{a}, \mathbf{b})$ with $\mathbf{b} = ((b^i, \hat{b}^i) : i \in I)$ for $b^i : V^i \rightarrow V^i$, $\hat{b}^i : E^i \rightarrow E^i$, and $G^i \circ \hat{b}^i \equiv G^i : E^i \rightarrow P$. As G^i is globally bounded, by Definition 3.8 there exists $N \geq 1$ such that $|(G^i)^{-1}(p)| \leq N$ for all $p \in P$.

For each such $\hat{b}^i$ and $p \in P$, $(G^i)^{-1}(p)$ is at most N points in E^i , and $\hat{b}^i$ must permute these points. But the order of a permutation of a set of at most N points must divide $N!$, so $(\hat{b}^i)^{N!}$ is the identity on $(G^i)^{-1}(p) \subset E^i$. As this is true for all $p \in P$, $(\hat{b}^i)^{N!}$ is the identity on E^i . So, every such $\hat{b}^i$ is a diffeomorphism of finite order. Thus the fixed point locus of $\hat{b}^i$ in E^i is a locally finite union of closed, embedded, connected suborbifolds of E^i , possibly of varying dimensions. Hence for each connected component F of E^i , either $\hat{b}^i$ is the identity on F , or $\hat{b}^i$ does not fix generic points of F .

Suppose for a contradiction that for some connected component F of E^i , we can find $(\mathbf{a}_1, \mathbf{b}_1), \dots, (\mathbf{a}_{N+1}, \mathbf{b}_{N+1})$ in $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ such that $\hat{b}_c^i|_F$ are distinct maps $F \rightarrow E^i$ for $c = 1, \dots, N+1$. Then for $1 \leq c < d \leq N+1$ we have finite order diffeomorphisms $\hat{b}_d^i \circ (\hat{b}_c^i)^{-1} : E^i \rightarrow E^i$ which are not the identity on $\hat{b}_c^i(F)$, so $\hat{b}_d^i \circ (\hat{b}_c^i)^{-1}$ does not fix generic points in $\hat{b}_c^i(F)$, and $\hat{b}_c^i, \hat{b}_d^i$ do not agree on generic points in F . Therefore $\hat{b}_1^i(f), \dots, \hat{b}_{N+1}^i(f)$ are distinct points in E^i for generic $f \in F$. But $G^i \circ \hat{b}_c^i(f) = G^i(f)$ for $c = 1, \dots, N+1$, so $\hat{b}_1^i(f), \dots, \hat{b}_{N+1}^i(f)$ lie in $(G^i)^{-1}(G^i(f))$, and $|(G^i)^{-1}(G^i(f))| \leq N$, a contradiction. Therefore, for each connected component F of E^i , the maps $\hat{b}^i : E^i \rightarrow E^i$ coming from $(\mathbf{a}, \mathbf{b}) \in \text{Aut}(X, \mathbf{f}, \mathbf{G})$ can realize at most N distinct maps $\hat{b}^i|_F : F \rightarrow E^i$.

Now use the notation of the proof of Proposition 3.19, so that $V^j = U^j \amalg \bigsqcup_{a \in A^j} W_a^j$ with each W_a^j a connected component of V^j , and A^j finite. Then

each $E^j|_{W_a^j}$ is a connected component of E^j , so the $\hat{b}^j|_{E^j|_{W_a^j}}$ realize at most N distinct maps. Hence, taken over all $j \in I$ and $a \in A^j$, the family of restrictions $\hat{b}^j|_{E^j|_{W_a^j}}$ can realize at most $N^{\sum_{j \in I} |A^j|}$ different possibilities.

Let $(\mathbf{a}, \mathbf{b}), (\mathbf{a}', \mathbf{b}') \in \text{Aut}(X, \mathbf{f}, \mathbf{G})$ with $\hat{b}^j|_{E^j|_{W_a^j}} = \hat{b}'^j|_{E^j|_{W_a^j}}$ for all $j \in I$ and $a \in A^j$. We will show that $(\mathbf{a}, \mathbf{b}) = (\mathbf{a}', \mathbf{b}')$. Suppose for a contradiction that $\hat{b}^j \neq \hat{b}'^j$ for some $j \in I$, and let j be largest for which this holds. From $V^j = U^j \amalg \coprod_{a \in A^j} W_a^j$ and $\hat{b}^j|_{E^j|_{W_a^j}} = \hat{b}'^j|_{E^j|_{W_a^j}}$ we see that $\hat{b}^j|_{E^j|_{U^j}} \neq \hat{b}'^j|_{E^j|_{U^j}}$. Thus, there exists some connected component W of U^j for which $\hat{b}^j|_{E^j|_W} \neq \hat{b}'^j|_{E^j|_W}$. Then $W \cap V^{ij} \neq \emptyset$ for some $j < i \in I$, by definition of W_a^j and U^j .

On $E^j|_{W \cap V^{ij}}$ we have $\hat{b}^j = \hat{b}^i \circ \hat{\phi}^{ij} = \hat{b}'^i \circ \hat{\phi}^{ij} = \hat{b}'^j$ by Definition 3.8(c), since j is largest with $\hat{b}^j \neq \hat{b}'^j$ and $j < i$, so $\hat{b}^i = \hat{b}'^i$. Hence, we know that $\hat{b}^j|_{E^j|_W} \neq \hat{b}'^j|_{E^j|_W}$, so that $\hat{b}^j, \hat{b}'^j$ differ at generic points of $E^j|_W$, but also $\hat{b}^j = \hat{b}'^j$ on $E^j|_{W \cap V^{ij}}$, which is a nonempty open set in $E^j|_W$. Choosing a generic point in $E^j|_{W \cap V^{ij}}$ gives a contradiction. Hence $\hat{b}^j = \hat{b}'^j$ for all $j \in I$. But $\hat{b}^j : E^j \rightarrow E^j$ is a lift of $b^j : V^j \rightarrow V^j$, so each $\hat{b}^j$ determines b^j . Thus $b^j = b'^j$ for $j \in I$. Also each $(b^j, \hat{b}^j)$ determines $\mathbf{a}$ over $\text{Im } \psi^j$, and $X = \bigcup_{j \in I} \text{Im } \psi^j$, so $\mathbf{b}$ determines $\mathbf{a}$. Hence $(\mathbf{a}, \mathbf{b}) = (\mathbf{a}', \mathbf{b}')$.

This proves that each $(\mathbf{a}, \mathbf{b}) \in \text{Aut}(X, \mathbf{f}, \mathbf{G})$ is determined by the restrictions $\hat{b}^j|_{E^j|_{W_a^j}}$ for all $j \in I$ and $a \in A^j$, which realize at most $N^{\sum_{j \in I} |A^j|}$ different possibilities. So $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ is finite, with $|\text{Aut}(X, \mathbf{f}, \mathbf{G})| \leq N^{\sum_{j \in I} |A^j|}$. $\square$

3.4 Quotients by finite groups

(Co-)gauge-fixing data descends to finite quotients, as in Property 3.1(c).

Definition 3.21. Let X be a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ a strongly smooth map, and $\mathbf{G}$ be gauge-fixing data for $(X, \mathbf{f})$. Suppose Γ is a finite group, and ρ an action of Γ on $(X, \mathbf{f}, \mathbf{G})$ by isomorphisms. That is, $\rho : \Gamma \rightarrow \text{Aut}(X, \mathbf{f}, \mathbf{G})$ is a group morphism. Note that we do not require ρ to be injective, so we cannot regard Γ as a subgroup of $\text{Aut}(X, \mathbf{f}, \mathbf{G})$. Write $\mathbf{G} = (\mathbf{I}, \boldsymbol{\eta}, G^i : i \in I)$, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i), f^i : i \in I, \dots)$. Then ρ induces actions of Γ on X by strong diffeomorphisms and on V^i, E^i for $i \in I$ by diffeomorphisms, where if $\gamma \in \Gamma$ with $\rho(\gamma) = (\mathbf{a}, \mathbf{b})$ and $\mathbf{b} = ((b^i, \hat{b}^i) : i \in I)$, then γ acts on X by $\mathbf{a}$, on V^i by b^i , and on E^i by $\hat{b}^i$.

Define a compact Kuranishi space $\tilde{X} = X/\Gamma$ and orbifolds $\tilde{V}^i = V^i/\Gamma$ and $\tilde{E}^i = E^i/\Gamma$ for $i \in I$. As ρ need not be injective, for example we allow the case that Γ is a nontrivial group but $\rho \equiv 1$, so that Γ acts trivially on X and V^i, E^i . In this case $\tilde{X}, \tilde{V}^i, \tilde{E}^i$ coincide with X, V^i, E^i as topological spaces, but not as Kuranishi spaces or orbifolds, since taking the quotient by Γ adds a factor of Γ to the stabilizer groups of X, V^i, E^i at every point, so that $\tilde{V}^i, \tilde{E}^i$ are *noneffective* orbifolds.

Let $\pi^i : E^i \rightarrow V^i$ be the projection. If $\gamma \in \Gamma$ and $\rho(\gamma) = (\mathbf{a}, \mathbf{b})$ then $\pi^i \circ \hat{b}^i = b^i \circ \pi^i$ by definition of $\text{Aut}(X, \mathbf{f}, \mathbf{G})$, so $\pi^i : E^i \rightarrow V^i$ is equivariant

with respect to the actions of Γ on V^i, E^i , and π^i pushes down to a submersion $\tilde{\pi}^i : \tilde{E}^i \rightarrow \tilde{V}^i$. The orbibundle structure on $\pi^i : E^i \rightarrow V^i$ descends to an orbibundle structure on $\tilde{\pi}^i : \tilde{E}^i \rightarrow \tilde{V}^i$. Since $\tilde{b}^i \circ s^i \equiv s^i \circ b^i$ for all $(\mathbf{a}, \mathbf{b}) = \rho(\gamma)$, s^i pushes down to a smooth section $\tilde{s}^i$ of $\tilde{E}^i \rightarrow \tilde{V}^i$, and $(\tilde{s}^i)^{-1}(0) = (s^i)^{-1}(0)/\Gamma$. As $\psi^i \circ b^i \equiv a \circ \psi^i$, where $a : X \rightarrow X$ is the homeomorphism underlying $\mathbf{a}$ in $(\mathbf{a}, \mathbf{b}) = \rho(\gamma)$, ψ^i pushes down to a continuous $\tilde{\psi}^i : (\tilde{s}^i)^{-1}(0) \rightarrow \tilde{X} = X/\Gamma$. Then $(\tilde{V}^i, \tilde{E}^i, \tilde{s}^i, \tilde{\psi}^i)$ is a Kuranishi neighbourhood on $\tilde{X}$.

Write $\pi_{\tilde{V}^i} : V^i \rightarrow \tilde{V}^i$ and $\pi_{\tilde{E}^i} : E^i \rightarrow \tilde{E}^i$ for the natural projections. Since $f^i : V^i \rightarrow Y$ and $G^i : E^i \rightarrow P$ are Γ -invariant they descend to $\tilde{f}^i : \tilde{V}^i \rightarrow Y$ and $\tilde{G}^i : \tilde{E}^i \rightarrow P$ with $\tilde{f}^i \circ \pi_{\tilde{V}^i} \equiv f^i$ and $\tilde{G}^i \circ \pi_{\tilde{E}^i} \equiv G^i$. Similarly $\eta_i : X \rightarrow [0, 1]$ and $\eta_i^j : V^j \rightarrow [0, 1]$ descend to $\tilde{\eta}_i : \tilde{X} \rightarrow [0, 1]$ and $\tilde{\eta}_i^j : \tilde{V}^j \rightarrow [0, 1]$. If $j \leq i$ in I then V^{ij} is a Γ -invariant subset of V^j , so $\tilde{V}^{ij} = V^{ij}/\Gamma$ is an open subset of $\tilde{V}^j = V^j/\Gamma$, and $(\phi^{ij}, \hat{\phi}^{ij})$ descends to a coordinate transformation $(\tilde{\phi}^{ij}, \hat{\tilde{\phi}}^{ij}) : (\tilde{V}^{ij}, \tilde{E}^j|_{\tilde{V}^{ij}}, \tilde{s}^j|_{\tilde{V}^{ij}}, \tilde{\psi}^j|_{\tilde{V}^{ij}}) \rightarrow (\tilde{V}^i, \dots, \tilde{\psi}^i)$. It is now easy to verify that all this comprises *gauge-fixing data* $\tilde{\mathbf{G}}$ for $(\tilde{X}, \tilde{\mathbf{f}})$. We shall also write $\tilde{\mathbf{f}} = \pi_*(\mathbf{f})$ and $\tilde{\mathbf{G}} = \pi_*(\mathbf{G})$, where $\pi : X \rightarrow \tilde{X}$ is the projection.

Exactly the same construction works for *co-gauge-fixing data* $\tilde{\mathbf{C}} = \pi_*(\mathbf{C})$.

3.5 Restriction of (co-)gauge-fixing data to boundaries

Very good, really good and excellent coordinate systems on X restrict to ∂X .

Definition 3.22. Let X be a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ be strongly smooth, and $\mathbf{I} = (I, (V^i, \dots, \psi^i), f^i : i \in I, \dots)$ be a very good coordinate system for $(X, \mathbf{f})$. Define a very good coordinate system $\tilde{\mathbf{I}} = \mathbf{I}|_{\partial X}$ for $(\partial X, \mathbf{f}|_{\partial X})$ to have indexing set $\tilde{I} = \{i : i+1 \in I, \partial V^{i+1} \neq \emptyset\}$ and Kuranishi neighbourhoods

$$(\tilde{V}^i, \tilde{E}^i, \tilde{s}^i, \tilde{\psi}^i) = (\partial V^{i+1}, E^{i+1}|_{\partial V^{i+1}}, s^{i+1}|_{\partial V^{i+1}}, \psi^{i+1}|_{\partial V^{i+1}}) \quad \text{for } i \in \tilde{I}.$$

Set $\tilde{V}^{ij} = \partial V^{(i+1)(j+1)}$ and $(\tilde{\phi}^{ij}, \hat{\tilde{\phi}}^{ij}) = (\phi^{(i+1)(j+1)}, \hat{\phi}^{(i+1)(j+1)})|_{\tilde{V}^{ij}}$ for $j \leq i$ in $\tilde{I}$. Set $\tilde{f}^i = f^{i+1}|_{\partial V^{i+1}}$ for $i \in \tilde{I}$. It is now easy to verify that $\tilde{\mathbf{I}} = (\tilde{I}, (\tilde{V}^i, \dots, \tilde{\psi}^i) : i \in \tilde{I}, \dots)$ is a *very good coordinate system* for $(\partial X, \mathbf{f}|_{\partial X})$, which we write as $\mathbf{I}|_{\partial X}$.

If $(\mathbf{I}, \boldsymbol{\eta})$ is a *really good coordinate system* for $(X, \mathbf{f})$ then we extend $\mathbf{I}|_{\partial X}$ to a really good coordinate system $(\mathbf{I}|_{\partial X}, \boldsymbol{\eta}|_{\partial X})$ in the obvious way, so that $\boldsymbol{\eta}|_{\partial X}$ consists of the functions $\tilde{\eta}_i = \eta_{i+1}|_{\partial X}$ and $\tilde{\eta}_i^j = \eta_{i+1}^{j+1}|_{\partial V^{j+1}}$ for $i, j \in \tilde{I}$. Definition 3.4(v) holds with $M-1$ in place of M , or 0 when $M=0$, as the local boundary components $\tilde{B}$ of ∂V^i at (v, B) correspond to a subset of the local boundary components B' of V^i at v with $B' \neq B$. If $(\mathbf{I}, \boldsymbol{\eta})$ is an *excellent coordinate system* for $(X, \mathbf{f})$ then $(\mathbf{I}|_{\partial X}, \boldsymbol{\eta}|_{\partial X})$ is also an excellent coordinate system, since the conditions on V^i in Definition 3.5 are stable under taking boundaries.

We can restrict (co-)gauge-fixing data to boundaries, as in Property 3.1(d).

Definition 3.23. Let X be a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ a strongly smooth map, and $\mathbf{G} = (\mathbf{I}, \boldsymbol{\eta}, G^i : i \in I)$ be gauge-fixing data for $(X, \mathbf{f})$. Definition 3.22 gives an excellent coordinate system $(\mathbf{I}|_{\partial X}, \boldsymbol{\eta}|_{\partial X})$ for $(\partial X, \mathbf{f}|_{\partial X})$. For each i in $\tilde{I} = \{i : i+1 \in I, \partial V^{i+1} \neq \emptyset\}$ we define $\tilde{G}^i = G^{i+1}|_{(E^{i+1})|_{\partial V^{i+1}}}$. Let $\iota : E^{i+1}|_{\partial V^{i+1}} = \partial E^{i+1} \rightarrow E^{i+1}$ be the natural map, M be as in Definition 3.4(v) for $(\mathbf{I}, \boldsymbol{\eta})$, and N be as in Definition 3.8 for G^{i+1} . Then $|(G^{i+1})^{-1}(p)| \leq N$ for each $p \in P$, and $|\iota^{-1}(e)| \leq M$ for each $e \in E^{i+1}|_{\partial V^{i+1}}$. Hence $\tilde{G}^i = G^{i+1} \circ \iota$ satisfies $|(\tilde{G}^i)^{-1}(p)| \leq MN$ for each $p \in P$, that is, $\tilde{G}^i$ is *globally finite*, with constant $\tilde{N} = MN$. Therefore $\mathbf{G}|_{\partial X} = (\mathbf{I}|_{\partial X}, \boldsymbol{\eta}|_{\partial X}, \tilde{G}^i : i \in \tilde{I})$ is gauge-fixing data for $(\partial X, \mathbf{f}|_{\partial X})$.

Exactly the same construction works for co-gauge-fixing data.

3.6 Extension of (co-)gauge-fixing data from boundaries

Next we prove Property 3.1(e) for (co-)gauge-fixing data.

Proposition 3.24. *Let X be a compact Kuranishi space with corners, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ be strongly smooth. Recall from Definition 2.27 that there is a natural strong diffeomorphism $\sigma : \partial^2 X \rightarrow \partial^2 X$ with $\sigma^2 = \text{id}_{\partial^2 X}$. Let $\mathbf{J}$ be a very good coordinate system for $(\partial X, \mathbf{f}|_{\partial X})$, with corners. Then there exists a very good coordinate system $\mathbf{I}$ for $(X, \mathbf{f})$ with corners with $\mathbf{I}|_{\partial X} = \mathbf{J}$ if and only if $\mathbf{J}|_{\partial^2 X}$ is invariant under σ . The analogous result holds for really good or excellent coordinate systems $(\mathbf{J}, \zeta), (\mathbf{I}, \boldsymbol{\eta})$ for $(\partial X, \mathbf{f}|_{\partial X}), (X, \mathbf{f})$.*

Proof. If $\mathbf{I}$ is a very good coordinate system with corners for $(X, \mathbf{f})$ then restrict to ∂X and then to $\partial^2 X$ gives $\tilde{\mathbf{I}} = \mathbf{I}|_{\partial^2 X}$ for $(\partial^2 X, \mathbf{f}|_{\partial^2 X})$, with indexing set $\tilde{I} = \{i : i+2 \in I, \partial^2 V^i \neq \emptyset\}$ and Kuranishi neighbourhoods $(\bar{V}^i, \dots, \bar{\psi}^i)$ with corners for $i \in \tilde{I}$ with $\bar{V}^i = \partial^2 V^{i+2}$ and $(\bar{E}^i, \bar{s}^i, \bar{\psi}^i) = (E^{i+2}, s^{i+2}, \psi^{i+2})|_{\partial^2 V^{i+2}}$. Clearly, $\tilde{\mathbf{I}}|_{\partial^2 X}$ is invariant under σ , with σ acting on $\bar{V}^i$ as the involution $\sigma : \partial^2 V^{i+2} \rightarrow \partial^2 V^{i+2}$ of Definitions 2.7 and 2.9. Thus, a necessary condition for $\mathbf{I}|_{\partial X} = \mathbf{J}$ is that $\mathbf{J}|_{\partial^2 X}$ is invariant under σ . This proves the ‘only if’ part.

For the ‘if’ part, let $\mathbf{J}$ be given and σ -invariant on $\partial^2 X$ as above. Let (W^j, F^j, t^j, ξ^j) for $j \in J$ be one of the Kuranishi neighbourhoods with corners in $\mathbf{J}$. Then $(\partial W^j, F^j|_{\partial W^j}, t^j|_{\partial W^j}, \xi^j|_{\partial W^j})$ is a Kuranishi neighbourhood with corners on $\partial^2 X$, and σ on $\partial^2 X$ lifts to an action on $(\partial W^j, \dots, \xi^j|_{\partial W^j})$. This is the condition necessary to extend $(W^j, \dots, \xi^j)$ to a Kuranishi neighbourhood $(V^{j+1}, \dots, \psi^{j+1})$ with corners on X with $(\partial V^{j+1}, \dots, \psi^{j+1}|_{\partial V^{j+1}}) = (W^j, \dots, \xi^j)$, which is ‘small’ in the sense that it extends only a little way into the interior of X , and V^{j+1} extends only a little way from $\partial V^{j+1} = W^j$.

Note that to extend the Kuranishi map s^{j+1} smoothly from its prescribed values on $\partial V^{j+1} = W^j$ to V^{j+1} requires the *Extension Principle*, Principle 2.8(c), which holds only for manifolds and orbifolds with corners, not g-corners, and this is why we restrict to Kuranishi spaces and (co-)gauge-fixing data with corners, not g-corners. The Kuranishi structure on X is defined in terms of germs, and so determines $(V^{j+1}, \dots, \psi^{j+1})$ only on an arbitrarily small open neighbourhood of $(s^{j+1})^{-1}(0)$. Away from $(s^{j+1})^{-1}(0)$ we have to choose s^{j+1} , and near ∂V^{j+1} away from $(t^j)^{-1}(0)$ we need the Extension Principle to do this.

Modifying the proof of Proposition 3.12, we can build a good (*not* very good, at this point) coordinate system $\mathbf{I}$ for $(X, \mathbf{f})$ with corners, with indexing set $I = \{j+1 : j \in J\} \amalg K$, using Kuranishi neighbourhoods $(V^{j+1}, \dots, \psi^{j+1})$ for $j \in J$ as above with V^{j+1} sufficiently small with $W^j = \partial V^{j+1}$ fixed, together with other Kuranishi neighbourhoods $(V^k, \dots, \psi^k)$ for k in some finite set K on the interior of X , with $\partial V^k = \emptyset$. Then we use Proposition 3.13 to convert $\mathbf{I}$ to a very good coordinate system $\tilde{\mathbf{I}}$ with corners for $(X, \mathbf{f})$ by gluing together the Kuranishi neighbourhoods of each dimension in $\mathbf{I}$.

In doing this, we replace the V^i by open subsets $\tilde{V}^i \subset V^i$. We choose the $\tilde{V}^{j+1}$ for $j \in J$ so that $\partial \tilde{V}^{j+1} = \partial V^{j+1} = W^j$, that is, we do not make the boundaries of the $\tilde{V}^{j+1}$ smaller. This is possible as there is at most one nonempty ∂V^i for $i \in I$ in each dimension, so the boundaries ∂V^i do not need to be glued together, and do not need to be made smaller to ensure Hausdorffness. It then follows that $\tilde{\mathbf{I}}|_{\partial X} = \mathbf{J}$, proving the ‘if’ part.

To prove the analogous ‘if’ result for *really good coordinate systems*, we first construct a very good coordinate system $\tilde{\mathbf{I}}$ with corners with $\tilde{\mathbf{I}}|_{\partial X} = \mathbf{J}$ as above, and then extend to really good $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ with $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})|_{\partial X} = (\mathbf{J}, \boldsymbol{\zeta})$. Thus we must construct continuous $\tilde{\eta}_i : X \rightarrow [0, 1]$ and $\tilde{\eta}_i^j : \tilde{V}^j \rightarrow [0, 1]$ with prescribed values ζ_{j-1} on ∂X and ζ_{i-1}^{j-1} on $\partial \tilde{V}^j$, satisfying Definition 3.4(i)–(iv). This is possible by extending Proposition 3.15.

Two new issues arise. Firstly, in constructing the η_i^j in Proposition 3.15, we made the V^j, V^{jk} smaller at each inductive step, so that η_i^j should extend continuously from its prescribed values on (34) to the closure of (34) in V^j . However, as we need $\partial \tilde{V}^j = W^{j-1}$, $\partial \tilde{V}^{jk} = W^{(j-1)(k-1)}$, we cannot make $\partial \tilde{V}^j, \partial \tilde{V}^{jk}$ smaller. This does not matter, since we have prescribed values $\tilde{\eta}_i^j|_{\partial \tilde{V}^j} \equiv \zeta_{i-1}^{j-1}$ for $\tilde{\eta}_i^j$ on $\partial \tilde{V}^j$ consistent with the prescribed values on (34), so there is no need to make $\partial \tilde{V}^j$ smaller to ensure consistency.

Secondly, Proposition 3.15 constructed a really good coordinate system satisfying the extra properties $\text{Im } \psi^i = \{p \in X : \eta_i(p) > 0\}$ and $V^{ij} = \{v \in V^j : \eta_i^j(v) > 0\}$ for all i, j , so the final sentences of both Definition 3.4(i),(ii) hold. However, the boundary data $(\mathbf{J}, \boldsymbol{\zeta})$ may not satisfy $\text{Im } \xi^j = \{p \in \partial X : \zeta_j(p) > 0\}$ and $W^{jk} = \{w \in W^k : \zeta_j^k(w) > 0\}$ for $k < j \in J$. Therefore we must relax the requirements that $\text{supp}(\eta_i) \subseteq \text{Im } \psi^i$ and $\text{supp}(\eta_i^j) \subseteq V^{ij}$ in the proof of Proposition 3.15. We do this by requiring that $\text{supp}(\eta_i)$ should be supported in the union of $\text{Im } \psi^i$ and a thin neighbourhood in X of $\text{supp}(\zeta_{i-1})$ in ∂X , and $\text{supp}(\eta_i^j)$ should be supported in the union of V^{ij} and a thin neighbourhood in V^j of $\text{supp}(\zeta_{i-1}^{j-1})$ in ∂V^j . We choose these thin neighbourhoods of $\text{supp}(\zeta_{i-1}), \text{supp}(\zeta_{i-1}^{j-1})$ small enough that the final sentences of Definition 3.4(i),(ii) for $(\mathbf{J}, \boldsymbol{\zeta})$ imply the final sentences of Definition 3.4(i),(ii) for $(\mathbf{I}, \boldsymbol{\eta})$ in the thin neighbourhoods. Then replacing V^i, V^{ij} by $\tilde{V}^i, \tilde{V}^{ij}$ as in the proof of Proposition 3.15, one can show that we get a *really good coordinate system* $(\mathbf{I}, \boldsymbol{\eta})$, with $(\mathbf{I}, \boldsymbol{\eta})|_{\partial X} = (\mathbf{J}, \boldsymbol{\zeta})$.

For *excellent coordinate systems*, if $(\mathbf{J}, \boldsymbol{\zeta})$ is excellent, we construct a really good coordinate system $(\mathbf{I}, \boldsymbol{\eta})$ for $(X, \mathbf{f})$ with corners, with $(\mathbf{I}, \boldsymbol{\eta})|_{\partial X} = (\mathbf{J}, \boldsymbol{\zeta})$, as above, and then apply Algorithm 3.16 to get an excellent coordinate system

$(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ with corners. It is automatic that $\partial\tilde{V}^i = \partial V^i = W^{i-1}$, since $(\mathbf{J}, \boldsymbol{\zeta})$ is an excellent coordinate system, so every connected component of $\partial^k W^{i-1}$ for all $k \geq 0$ is necessary, and Algorithm 3.16 does not discard any of them. Hence $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})|_{\partial X} = (\mathbf{J}, \boldsymbol{\zeta})$, as we want. $\square$

Theorem 3.25. *Suppose X is a compact Kuranishi space with corners, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ a strongly smooth map, and $\mathbf{H}$ is gauge-fixing data for $(\partial X, \mathbf{f}|_{\partial X})$ with corners. Let $\sigma : \partial^2 X \rightarrow \partial^2 X$ be the involution of Definition 2.27. Then there exists gauge-fixing data $\mathbf{G}$ for $(X, \mathbf{f})$ with corners with $\mathbf{G}|_{\partial X} = \mathbf{H}$ if and only if $\mathbf{H}|_{\partial^2 X}$ is σ -invariant.*

Suppose Γ is a finite group of strong diffeomorphisms $\mathbf{a} : X \rightarrow X$ with $\mathbf{f} \circ \mathbf{a} = \mathbf{f}$ and $\mathbf{H}$ is invariant under $\Gamma|_{\partial X}$, that is, we are given a group morphism $\Gamma \rightarrow \text{Aut}(\partial X, \mathbf{f}|_{\partial X}, \mathbf{H})$ mapping $\mathbf{a} \mapsto (\mathbf{a}|_{\partial X}, \mathbf{c}_{\mathbf{a}})$, such that the action of $(\mathbf{a}|_{\partial^2 X}, \mathbf{c}_{\mathbf{a}}|_{\partial^2 X})$ on $\mathbf{H}|_{\partial^2 X}$ commutes with the action of σ on $\mathbf{H}|_{\partial^2 X}$. Then we can choose $\mathbf{G}$ to be Γ -invariant, that is, there is a group morphism $\Gamma \rightarrow \text{Aut}(X, \mathbf{f}, \mathbf{G})$ mapping $\mathbf{a} \mapsto (\mathbf{a}, \mathbf{b}_{\mathbf{a}})$, such that $\mathbf{b}_{\mathbf{a}}|_{\partial X} = \mathbf{c}_{\mathbf{a}}$ for all $\mathbf{a} \in \Gamma$.

If $\partial Y = \emptyset$, all this also holds for co-gauge-fixing data $\mathbf{C}, \mathbf{D}$ for $(X, \mathbf{f})$, $(\partial X, \mathbf{f}|_{\partial X})$, with $\mathbf{f}$ a strong submersion.

Proof. If $\mathbf{G}$ is gauge-fixing data for $(X, \mathbf{f})$ with corners then restricting to ∂X and then to $\partial^2 X$ gives $\mathbf{G}|_{\partial^2 X}$. Clearly, $\mathbf{G}|_{\partial^2 X}$ is invariant under σ . Thus, $\mathbf{G}|_{\partial X} = \mathbf{H}$ implies that $\mathbf{H}|_{\partial^2 X}$ is σ -invariant. This proves the ‘only if’ part.

For the ‘if’ part, suppose $\mathbf{H}|_{\partial^2 X}$ is σ -invariant as above. Write $\mathbf{H} = (\mathbf{J}, \boldsymbol{\zeta}, H^j : j \in J)$, where $\mathbf{J} = (J, (W^j, F^j, t^j, \xi^j) : j \in J, \dots)$. As $(\mathbf{J}, \boldsymbol{\zeta})$ is an excellent coordinate system with corners for $(\partial X, \mathbf{f}|_{\partial X})$ and $(\mathbf{J}, \boldsymbol{\zeta})|_{\partial^2 X}$ is σ -invariant, Proposition 3.24 gives an excellent coordinate system $(\mathbf{I}, \boldsymbol{\eta})$ with corners for $(X, \mathbf{f})$ with $(\mathbf{I}, \boldsymbol{\eta})|_{\partial X} = (\mathbf{J}, \boldsymbol{\zeta})$, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i), i \in I, \dots)$.

Then $\partial E^i = F^{i-1}$ for $i \in I$ with $i-1 \in J$, and $\partial E^i = \emptyset$ for $i \in I$ with $i-1 \notin J$. When $i \in I$ with $i-1 \in J$, we have $H^{i-1} : \partial E^i \rightarrow P$ with $H^{i-1}|_{\partial^2 E^i}$ invariant under $\sigma : \partial^2 E^i \rightarrow E^i$. Therefore by Principle 2.8(a) we can choose $G^i : E^i \rightarrow P_n \subset P$ with $G^i|_{\partial E^i} \equiv H^i$. Since G^i is arbitrary, not continuous or smooth, the values of G^i on $(E^i)^\circ$ need not be related to those on ∂E^i .

Choose $G^i|_{(E^i)^\circ} : (E^i)^\circ \rightarrow P_n \subset P$ to be an arbitrary injective map, for $n \gg 0$. As H^{i-1} is globally finite with constant $N \geq 0$ in the sense of Definition 3.8, it follows that G^i is globally finite with constant $N+1$. If $i \in I$ but $i-1 \notin J$ then there are no boundary conditions on G^i , so we choose $G^i : E^i \rightarrow P_n \subset P$ to be an arbitrary injective map for $n \gg 0$, and then G^i is globally finite with $N = 1$. Thus $\mathbf{G}$ is gauge-fixing data with corners for $(X, \mathbf{f})$ with $\mathbf{G}|_{\partial X} = \mathbf{H}$, proving the ‘if’ part. For the second part on Γ actions, we combine the proof above with the second part of the proof of Theorem 3.18. The analogues for co-gauge-fixing data are immediate. $\square$

3.7 Pushforwards and pullbacks of (co-)gauge-fixing data

Our next definitions, which give Property 3.1(f),(g) for gauge-fixing data and co-gauge-fixing data, encapsulate the reasons for the differences between Definitions 3.8 and 3.9. We will use gauge-fixing data in Kuranishi homology, so it should

have a good notion of *pushforward* h_* under smooth $h : Y \rightarrow Z$. We will use co-gauge-fixing data in (compactly-supported) Kuranishi cohomology, so it should have a good notion of *pullback* h^* under smooth, proper $h : Y \rightarrow Z$.

Definition 3.26. Let Y, Z be orbifolds and $h : Y \rightarrow Z$ a smooth map. Suppose X is a compact Kuranishi space, $\mathbf{f} : X \rightarrow Y$ is strongly smooth, and $\mathbf{G} = (\mathbf{I}, \boldsymbol{\eta}, G^i : i \in I)$ is gauge-fixing data for $(X, \mathbf{f})$, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i), f^i : i \in I, \dots)$. Then $h \circ \mathbf{f} : X \rightarrow Z$ is strongly smooth. Define *pushforward gauge-fixing data* $h_*(\mathbf{G})$ for $(X, h \circ \mathbf{f})$ to be $h_*(\mathbf{G}) = (h_*(\mathbf{I}), \boldsymbol{\eta}, G^i : i \in I)$, where $h_*(\mathbf{I}) = (I, (V^i, E^i, s^i, \psi^i), h \circ f^i : i \in I, \dots)$. That is, $h_*(\mathbf{G})$ is the same as $\mathbf{G}$, except that $f^i : V^i \rightarrow Y$ for $i \in I$ are replaced by $h \circ f^i : V^i \rightarrow Z$.

Clearly $h_*(\mathbf{G})$ is gauge-fixing data for $(X, h \circ \mathbf{f})$. Note however that the same construction does *not* work for co-gauge-fixing data, as if $h : Y \rightarrow Z$ is not globally finite, then $C^i \times (f^i \circ \pi^i) : E^i \rightarrow P \times Y$ globally finite does not imply $C^i \times (h \circ f^i \circ \pi^i) : E^i \rightarrow P \times Z$ globally finite, so Definition 3.9 fails for $h_*(\mathbf{C})$.

Definition 3.27. Let Y, Z be orbifolds without boundary and $h : Y \rightarrow Z$ a smooth, proper map. Suppose X is a compact Kuranishi space, $\mathbf{f} : X \rightarrow Z$ is a strong submersion, and $\mathbf{C} = (\mathbf{I}, \boldsymbol{\eta}, C^i : i \in I)$ is co-gauge-fixing data for $(X, \mathbf{f})$, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i), f^i : i \in I, \dots)$. Then $Y \times_{h, Z, \mathbf{f}} X$ is a compact Kuranishi space, and $\pi_Y : Y \times_Z X \rightarrow Y$ is a strong submersion. We will define co-gauge-fixing data $h^*(\mathbf{C})$ for $(Y \times_Z X, \pi_Y)$, where $h^*(\mathbf{C}) = (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}}, \tilde{C}^i : i \in \tilde{I})$, with $\tilde{\mathbf{I}} = (\tilde{I}, (\tilde{V}^i, \tilde{E}^i, \tilde{s}^i, \tilde{\psi}^i), \tilde{\pi}_Y^i : i \in \tilde{I}, \dots)$.

Let $k = \dim Y - \dim Z$, and $\tilde{I} = \{i + k : i \in I\}$. Then $\dim V^i = i$ for each $i \in I$, and as $f^i : V^i \rightarrow Z$ is a submersion $i = \dim V^i \geq \dim Z$, so $i + k \geq \dim Y \geq 0$. Thus $\tilde{I} \subset \mathbb{N}$. For $i \in I$, define $(\tilde{V}^{i+k}, \tilde{E}^{i+k}, \tilde{s}^{i+k}, \tilde{\psi}^{i+k}) = (Y \times_{h, Z, f^i} V^i, \pi_{V^i}^*(E^i), s^i \circ \pi_{V^i}, h \times_Z \psi^i)$. This is a Kuranishi neighbourhood on $Y \times_{h, Z, \mathbf{f}} X$, with $\dim \tilde{V}^{i+k} = \dim V^i + \dim Y - Z = i + k$, as we want. Let $\tilde{\pi}_Y^{i+k} : \tilde{V}^{i+k} \rightarrow Y$ be the projection from the fibre product. This is a submersion, as f^i is, and represents π_Y .

For $i, j \in I$ with $j \leq i$ we have a triple $(V^{ij}, \phi^{ij}, \hat{\phi}^{ij})$. Define $\tilde{V}^{(i+k)(j+k)} = \pi_{V^j}^{-1}(V^{ij})$, where $\pi_{V^j} : \tilde{V}^{j+k} \rightarrow V^j$ is the projection from the fibre product. Define $\tilde{\phi}^{(i+k)(j+k)} : \tilde{V}^{(i+k)(j+k)} \rightarrow \tilde{V}^{i+k}$ by $\tilde{\phi}^{(i+k)(j+k)} = h^*(\phi^{ij})$, mapping $Y \times_Z V^{ij} \rightarrow Y \times_Z V^i$. Define $\hat{\phi}^{(i+k)(j+k)} : \tilde{E}^{j+k}|_{\tilde{V}^{(i+k)(j+k)}} \rightarrow (\tilde{\phi}^{(i+k)(j+k)})^*(\tilde{E}^{i+k})$ by $\hat{\phi}^{(i+k)(j+k)} = h^*(\hat{\phi}^{ij})$. Then $\tilde{\mathbf{I}} = (\tilde{I}, (\tilde{V}^i, \tilde{E}^i, \tilde{s}^i, \tilde{\psi}^i), \tilde{\pi}_Y^i : i \in \tilde{I}, (\tilde{V}^{ij}, \tilde{\phi}^{ij}, \hat{\phi}^{ij}) : j \leq i \text{ in } \tilde{I})$ is a *very good coordinate system* for $(Y \times_Z X, \pi_Y)$.

Now for $i \in I$, define $\tilde{\eta}_{i+k} : Y \times_Z X \rightarrow [0, 1]$ by $\tilde{\eta}_{i+k} = \eta_i \circ \pi_X$, and for $i, j \in I$ define $\tilde{\eta}_{i+k}^{j+k} : \tilde{V}^{j+k} \rightarrow [0, 1]$ by $\tilde{\eta}_{i+k}^{j+k} = \eta_i^j \circ \pi_{V^j}$. Write $\tilde{\boldsymbol{\eta}} = (\tilde{\eta}_i : i \in \tilde{I}, \tilde{\eta}_i^j : i, j \in \tilde{I})$. It is not difficult to check that $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ is a *really good coordinate system*, as $(\mathbf{I}, \boldsymbol{\eta})$ is. In Definition 3.4(v) we can use the same M for $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ as for $(\mathbf{I}, \boldsymbol{\eta})$, and if $T \subseteq Z$ is as in Definition 3.4(vi) for $(\mathbf{I}, \boldsymbol{\eta})$ then $\tilde{T} = h^{-1}(T) \subseteq Y$ is compact as T is compact and h is proper, and satisfies Definition 3.4(vi) for $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$.

However, although $(\mathbf{I}, \boldsymbol{\eta})$ is an *excellent coordinate system*, $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ need not be, as taking fibre products $V^i \mapsto Y \times_Z V^i = \tilde{V}^{i+k}$ may introduce unnecessary connected components of $\tilde{V}^{i+k}$ or $\partial^l \tilde{V}^{i+k}$. Let $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ be the excellent coordinate system obtained from $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ by applying Algorithm 3.16, where $\tilde{\mathbf{I}} = (\tilde{I}, (\tilde{V}^i, \tilde{E}^i, \tilde{s}^i, \tilde{\psi}^i), \tilde{\pi}_Y^i : i \in \tilde{I}, \dots)$.

Let $i \in \tilde{I}$, so that $i - k \in I$. Then $\tilde{V}^i$ is an open subset of $\tilde{V}^i = Y \times_{h, Z, f^{i-k}} V^{i-k}$, and $\tilde{E}^i \rightarrow \tilde{V}^i$ is the pullback of $E^{i-k} \rightarrow V^{i-k}$ by the projection $\pi_{V^{i-k}} : \tilde{V}^i \rightarrow V^{i-k}$. Thus there is a natural projection $\pi_{E^{i-k}} : \tilde{E}^i \rightarrow E^{i-k}$ between the total spaces of the bundles $\tilde{E}^i, E^{i-k}$. Define $\check{C}^i : \tilde{E}^i \rightarrow P$ by $\check{C}^i = C^{i-k} \circ \pi_{E^{i-k}}$. We shall show that $\check{C}^i \times (\tilde{\pi}_Y^i \circ \tilde{\pi}^i) : \tilde{E}^i \rightarrow P \times Y$ for $i \in \tilde{I}$ is *globally finite*.

Definition 3.4(vi) gives compact $T \subseteq Z$ with $f^{i-k}(V^{i-k}) \subseteq T$. The map $Z \rightarrow \mathbb{N}$ taking $z \mapsto |\text{Stab}(z)|$ is upper semicontinuous, so as T is compact there exists $L \geq 1$ such that $|\text{Stab}(z)| \leq L$ for all $z \in T$. As $C^{i-k} \times (f^{i-k} \circ \pi^{i-k}) : E^{i-k} \rightarrow P \times Z$ is globally finite, there exists $N \geq 0$ such that $|(C^{i-k} \times (f^{i-k} \circ \pi^{i-k}))^{-1}(p, z)| \leq N$ for all $(p, z) \in P \times Z$. Let $(p, y) \in P \times Y$, and $z = h(y)$. Now $\pi_{E^{i-k}} : \tilde{E}^i \rightarrow E^{i-k}$ restricts to a map

$$\pi_{E^{i-k}} : (\check{C}^i \times (\tilde{\pi}_Y^i \circ \tilde{\pi}^i))^{-1}(p, y) \longrightarrow (C^{i-k} \times (f^{i-k} \circ \pi^{i-k}))^{-1}(p, z). \quad (36)$$

Using $\tilde{E}^i \subseteq Y \times_{h, Z, f^{i-k} \circ \pi^{i-k}} E^{i-k}$ and the definition (12) of fibre products of orbifolds, we see that each e on the right hand side of (36) pulls back to a subset of $h_*(\text{Stab}(y)) \setminus \text{Stab}(z) / (f^{i-k} \circ \pi^{i-k})_*(\text{Stab}(e))$ on the left. Since $|\text{Stab}(z)| \leq L$, at most L points on the left of (36) project to each point on the right. But the right hand side of (36) has at most N elements, so the left has at most LN elements. Thus $\check{C}^i \times (\tilde{\pi}_Y^i \circ \tilde{\pi}^i)$ is globally finite, with constant $\tilde{N} = LN$. Therefore $h^*(\mathbf{C})$ is co-gauge-fixing data for $(Y \times_Z X, \pi_Y)$.

Note that the same construction does *not* work for gauge-fixing data, as if h is not globally finite then $G^i : E^i \rightarrow P$ globally finite does not imply $G^i \circ \pi_{E^i} : Y \times_Z E^i \rightarrow P$ is globally finite, so $\check{G}^i : \tilde{E}^i \rightarrow P$ may not be globally finite, and Definition 3.8 fails for $h^*(\mathbf{G})$.

Remark 3.28. We suppose Y, Z are *without boundary* in Definition 3.27 in order that $\partial(Y \times_Z X) \cong \pm Y \times_Z \partial X$. This will be needed in §4.4 to ensure that pullbacks h^* on Kuranishi cochains satisfy $d \circ h^* = h^* \circ d$. If $\partial Y \neq \emptyset$ then $\partial(Y \times_Z X)$ has an extra term $\partial Y \times_Z X$, which would invalidate the proof of $d \circ h^* = h^* \circ d$. Actually the assumption $\partial Z = \emptyset$ is unnecessary, since our definitions of (strong) submersions imply that $\mathbf{f} : X \rightarrow Z$ and $f^i : V^i \rightarrow Z$ for $i \in I$ map to Z° , so ∂Z does not affect $\partial(Y \times_Z X)$ or $\partial(Y \times_{h, Z, f^i} V^i)$.

3.8 Fibre products of (co-)gauge-fixing data

When $(\mathbf{I}, \boldsymbol{\eta})$ and $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ are really good coordinate systems for $(X, \mathbf{f})$ and $(\tilde{X}, \tilde{\mathbf{f}})$, where one of $\mathbf{f} : X \rightarrow Y$ and $\tilde{\mathbf{f}} : \tilde{X} \rightarrow Y$ is a strong submersion, we shall define a really good coordinate system $(\mathbf{I}, \boldsymbol{\eta}) \times_Y (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ for the fibre product $(X \times_Y \tilde{X}, \pi_Y)$. The construction is long and complicated.

Definition 3.29. Let $X, \tilde{X}$ be compact Kuranishi spaces, Y be an orbifold, $\mathbf{f} : X \rightarrow Y$ and $\tilde{\mathbf{f}} : \tilde{X} \rightarrow Y$ be strongly smooth and $(\mathbf{I}, \boldsymbol{\eta})$, $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ be really good coordinate systems for $(X, \mathbf{f})$ and $(\tilde{X}, \tilde{\mathbf{f}})$. Suppose one of $\mathbf{f}, \tilde{\mathbf{f}}$ is a strong submersion, and the corresponding $f^i : V^i \rightarrow Y$ or $\tilde{f}^{\tilde{i}} : \tilde{V}^{\tilde{i}} \rightarrow Y$ in $\mathbf{I}, \tilde{\mathbf{I}}$ are submersions. Then $X \times_Y \tilde{X}$ is a compact Kuranishi space, as in §2.6. As in (16), for $i \in I$ and $\tilde{i} \in \tilde{I}$ define a Kuranishi neighbourhood on $X \times_Y \tilde{X}$ by

$$\begin{aligned} (V^{(i,\tilde{i})}, E^{(i,\tilde{i})}, s^{(i,\tilde{i})}, \psi^{(i,\tilde{i})}) &= (V^i \times_{f^i, Y, \tilde{f}^{\tilde{i}}} \tilde{V}^{\tilde{i}}, \pi_{V^i}^*(E^i) \oplus \pi_{\tilde{V}^{\tilde{i}}}^*(\tilde{E}^{\tilde{i}}), \\ & (s^i \circ \pi_{V^i}) \oplus (s^{\tilde{i}} \circ \pi_{\tilde{V}^{\tilde{i}}}), (\psi^i \circ \pi_{V^i}) \times (\tilde{\psi}^{\tilde{i}} \circ \pi_{\tilde{V}^{\tilde{i}}}) \times \chi^{(i,\tilde{i})}). \end{aligned} \quad (37)$$

Let $\pi_Y^{(i,\tilde{i})} : V^{(i,\tilde{i})} \rightarrow Y$ be the projection from the fibre product. If $i, j \in I$ and $\tilde{i}, \tilde{j} \in \tilde{I}$ with $j \leq i$ and $\tilde{j} \leq \tilde{i}$ we have open sets $V^{ij} \subseteq V^j$ and $\tilde{V}^{\tilde{i}\tilde{j}} \subseteq \tilde{V}^{\tilde{j}}$ and coordinate transformations $(\phi^{ij}, \hat{\phi}^{ij})$ and $(\tilde{\phi}^{\tilde{i}\tilde{j}}, \hat{\tilde{\phi}}^{\tilde{i}\tilde{j}})$. Define $V^{(i,\tilde{i})(j,\tilde{j})} \subseteq V^{(j,\tilde{j})}$ to be $V^{ij} \times_{f^j, Y, \tilde{f}^{\tilde{j}}} \tilde{V}^{\tilde{i}\tilde{j}}$. Then $(\phi^{ij}, \hat{\phi}^{ij})$ and $(\tilde{\phi}^{\tilde{i}\tilde{j}}, \hat{\tilde{\phi}}^{\tilde{i}\tilde{j}})$ induce a coordinate transformation

$$\begin{aligned} (\phi^{(i,\tilde{i})(j,\tilde{j})}, \hat{\phi}^{(i,\tilde{i})(j,\tilde{j})}) : (V^{(i,\tilde{i})(j,\tilde{j})}, E^{(j,\tilde{j})}|_{V^{(i,\tilde{i})(j,\tilde{j})}}, s^{(j,\tilde{j})}|_{V^{(i,\tilde{i})(j,\tilde{j})}}, \\ \psi^{(j,\tilde{j})}|_{V^{(i,\tilde{i})(j,\tilde{j})}}) \longrightarrow (V^{(i,\tilde{i})}, \dots, \psi^{(i,\tilde{i})}). \end{aligned}$$

We want to build a really good coordinate system $(\dot{\mathbf{I}}, \dot{\boldsymbol{\eta}}) = (\mathbf{I}, \boldsymbol{\eta}) \times_Y (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ for $(X \times_Y \tilde{X}, \pi_Y)$ with Kuranishi neighbourhoods $(\dot{V}^k, \dots, \dot{\psi}^k)$ for $k \in \dot{I}$, using the Kuranishi neighbourhoods (37) and maps $\pi_Y^{(i,\tilde{i})}$. Since $\dim V^{(i,\tilde{i})} = \dim V^i + \dim \tilde{V}^{\tilde{i}} - \dim Y = i + \tilde{i} - \dim Y$, the obvious definition is $\dot{V}^k = \coprod_{i \in I, \tilde{i} \in \tilde{I} : i + \tilde{i} = k + \dim Y} V^{(i,\tilde{i})}$, with $(\dot{E}^k, \dot{s}^k, \dot{\psi}^k)|_{\dot{V}^{(i,\tilde{i})}} = (E^{(i,\tilde{i})}, s^{(i,\tilde{i})}, \psi^{(i,\tilde{i})})$ and $\dot{\pi}_Y^k|_{\dot{V}^{(i,\tilde{i})}} = \pi_Y^{(i,\tilde{i})}$. There are two main problems with this:

- (*) If $i \neq j \in I$ and $\tilde{i} \neq \tilde{j} \in \tilde{I}$ with $i + \tilde{i} = j + \tilde{j} = k + \dim Y$ and $\text{Im } \psi^{(i,\tilde{i})} \cap \text{Im } \psi^{(j,\tilde{j})} \neq \emptyset$ then $\dot{\psi}^k$ is not injective, so $(\dot{V}^k, \dots, \dot{\psi}^k)$ is not a Kuranishi neighbourhood; and
- (**) Suppose $i < j \in I$ and $\tilde{i} > \tilde{j} \in \tilde{I}$ with $i + \tilde{i} = k + \dim Y$ and $j + \tilde{j} = l + \dim Y$ with $l < k$, and that $\text{Im } \psi^{(i,\tilde{i})} \cap \text{Im } \psi^{(j,\tilde{j})} \neq \emptyset$. Then we expect to define an open $\dot{V}^{kl} \subset \dot{V}^l$ and a coordinate transformation $(\dot{\phi}^{kl}, \dot{\hat{\phi}}^{kl})$ from $(\dot{V}^{kl}, \dot{E}^l|_{\dot{V}^{kl}}, \dot{s}^l|_{\dot{V}^{kl}}, \dot{\psi}^l|_{\dot{V}^{kl}})$ to $(\dot{V}^k, \dots, \dot{\psi}^k)$. This should include a transformation from a nonempty subset of $V^{(j,\tilde{j})}$ to $V^{(i,\tilde{i})}$. However, since $i < j$ we have no transformation from $V^{ij} \subseteq V^j$ to V^i to build this from.

We shall deal with both these problems by making the $V^{(i,\tilde{i})}$ smaller. That is, we shall define

$$\dot{V}^k = \coprod_{i \in I, \tilde{i} \in \tilde{I} : i + \tilde{i} = k + \dim Y} \dot{V}^{(i,\tilde{i})}, \quad (38)$$

where $\dot{V}^{(i,\tilde{i})}$ is an open subset of $V^{(i,\tilde{i})}$. We shall choose these $\dot{V}^{(i,\tilde{i})}$ so that in (*) and (**) above we always have $\text{Im } \psi^{(i,\tilde{i})}|_{\dot{V}^{(i,\tilde{i})}} \cap \text{Im } \psi^{(j,\tilde{j})}|_{\dot{V}^{(j,\tilde{j})}} = \emptyset$. But we cannot make the $\dot{V}^{(i,\tilde{i})}$ too small, as we need to ensure that $X = \bigcup_{i \in I, \tilde{i} \in \tilde{I}} \text{Im } \psi^{(i,\tilde{i})}|_{\dot{V}^{(i,\tilde{i})}}$. To define the $\dot{V}^{(i,\tilde{i})}$, we will construct partitions of unity on $X \times_Y \tilde{X}$ and on the

$V^{(i,\tilde{i})}$ for all $i, \tilde{i}$ using the data $\eta, \tilde{\eta}$. We will also use these partitions of unity to construct $\dot{\eta}$ in the really good coordinate system $(\dot{I}, \dot{\eta})$ for $(X \times_Y \tilde{X}, \pi_Y)$.

For all $i, j \in I$ and $\tilde{i}, \tilde{j} \in \tilde{I}$ we will define continuous functions $\eta_{(i,\tilde{i})} : X \times_Y \tilde{X} \rightarrow [0, 1]$ and $\eta_{(i,\tilde{i})}^{(j,\tilde{j})} : V^{(j,\tilde{j})} \rightarrow [0, 1]$ satisfying the conditions:

- (a) $\sum_{i \in I, \tilde{i} \in \tilde{I}} \eta_{(i,\tilde{i})} \equiv 1$ on $X \times_Y \tilde{X}$, and $\sum_{i \in I, \tilde{i} \in \tilde{I}} \eta_{(i,\tilde{i})}^{(j,\tilde{j})} \equiv 1$ on $V^{(j,\tilde{j})}$ for all $j, \tilde{j}$.
- (b) $\eta_{(i,\tilde{i})}^{(j,\tilde{j})}|_{(s^{(j,\tilde{j})})^{-1}(0)} \equiv \eta_{(i,\tilde{i})} \circ \psi^{(j,\tilde{j})}$ for all $i, j \in I$ and $\tilde{i}, \tilde{j} \in \tilde{I}$.
- (c) Let $i, j, k \in I$ and $\tilde{i}, \tilde{j}, \tilde{k} \in \tilde{I}$ with $k \leq j$ and $\tilde{k} \leq \tilde{j}$. Then $\eta_{(i,\tilde{i})}^{(k,\tilde{k})}|_{V^{(j,\tilde{j})}(k,\tilde{k})} \equiv \eta_{(i,\tilde{i})}^{(j,\tilde{j})} \circ \phi^{(j,\tilde{j})}(k,\tilde{k})$.
- (d) If $j \leq i$ in I , $\tilde{j} \leq \tilde{i}$ in $\tilde{I}$ and $q \in X \times_Y \tilde{X}$ with $\eta_{(i,\tilde{i})}(q) > 0$, $\eta_{(j,\tilde{j})}(q) > 0$ and $q \in \text{Im } \psi^{(i,\tilde{i})}$ then $q \in \text{Im } \psi^{(j,\tilde{j})}$.
- (e) if $j \leq i$ in I and $\tilde{j} \leq \tilde{i}$ in $\tilde{I}$ then $\phi^{(i,\tilde{i})}(j,\tilde{j})(V^{(i,\tilde{i})}(j,\tilde{j}))$ is a *closed* subset of $\{v \in V^{(i,\tilde{i})} : \eta_{(j,\tilde{j})}^{(i,\tilde{i})}(v) > 0\}$. If $k \leq j \leq i$ in I and $\tilde{k} \leq \tilde{j} \leq \tilde{i}$ in $\tilde{I}$ and $v \in V^{(k,\tilde{k})}$ with $\eta_{(i,\tilde{i})}^{(k,\tilde{k})}(v) > 0$, $\eta_{(j,\tilde{j})}^{(k,\tilde{k})}(v) > 0$ and $v \in V^{(i,\tilde{i})}(k,\tilde{k})$ then $v \in V^{(j,\tilde{j})}(k,\tilde{k})$.
- (f) Suppose $i, j, k \in I$ and $\tilde{i}, \tilde{j}, \tilde{k} \in \tilde{I}$ and either $i < j$ and $\tilde{i} > \tilde{j}$, or $i > j$ and $\tilde{i} < \tilde{j}$. Then $\eta_{(i,\tilde{i})}\eta_{(j,\tilde{j})} \equiv 0$ on $X \times_Y \tilde{X}$, and $\eta_{(i,\tilde{i})}^{(k,\tilde{k})}\eta_{(j,\tilde{j})}^{(k,\tilde{k})} \equiv 0$ on $V^{(k,\tilde{k})}$. That is, the supports of $\eta_{(i,\tilde{i})}$ and $\eta_{(j,\tilde{j})}$ are disjoint, and the supports of $\eta_{(i,\tilde{i})}^{(k,\tilde{k})}$ and $\eta_{(j,\tilde{j})}^{(k,\tilde{k})}$ are disjoint.
- (g) Suppose $q \in X \times_Y X$, set $p = \pi_X(q)$ and $\tilde{p} = \pi_{\tilde{X}}(q)$, and let $i \in I$ and $\tilde{i} \in \tilde{I}$ be least with $\eta_i(p) > 0$ and $\tilde{\eta}_{\tilde{i}}(\tilde{p}) > 0$. Then $\eta_{(i,\tilde{i})}(q) > 0$ and $q \in \text{Im } \psi^{(i,\tilde{i})}$.

With the exception of (f),(g) these are analogues of Definition 3.4(i)–(iv); note that (d),(e) are partial analogues of the parts of Definition 3.4(i),(ii).

Suppose for the moment we have constructed such $\eta_{(i,\tilde{i})}, \eta_{(i,\tilde{i})}^{(j,\tilde{j})}$. Define

$$\dot{V}^{(i,\tilde{i})} = \{v \in V^{(i,\tilde{i})} : \eta_{(i,\tilde{i})}^{(i,\tilde{i})}(v) > 0\} \subseteq V^{(i,\tilde{i})}. \quad (39)$$

Then $\dot{V}^{(i,\tilde{i})}$ is open, as $\eta_{(i,\tilde{i})}^{(i,\tilde{i})}$ is continuous, and (b) with $j = i, \tilde{j} = \tilde{i}$ implies that

$$\begin{aligned} \text{Im } \psi^{(i,\tilde{i})}|_{\dot{V}^{(i,\tilde{i})}} &= \{q \in \pi_X^{-1}(\text{Im } \psi^i) \cap \pi_{\tilde{X}}^{-1}(\text{Im } \tilde{\psi}^{\tilde{i}}) : \eta_{(i,\tilde{i})}(q) > 0\} \\ &\subseteq \{q \in X \times_Y \tilde{X} : \eta_{(i,\tilde{i})}(q) > 0\}. \end{aligned} \quad (40)$$

Define $\dot{I} = \{i + \tilde{i} - \dim Y : i \in I, \tilde{i} \in \tilde{I}, \dot{V}^{(i,\tilde{i})} \neq \emptyset\}$. Since one of $\mathbf{f}, \tilde{\mathbf{f}}$ is a strong submersion, it follows that either $i \geq \dim Y$ for all $i \in I$ or $\tilde{i} \geq \dim Y$ for all $\tilde{i} \in \tilde{I}$, and so $\dot{I}$ is a finite subset of $\mathbb{N} = \{0, 1, 2, \dots\}$. For each $k \in \dot{I}$, define $\dot{V}^k$ by (38). Then $\dot{V}^k$ is an orbifold with $\dim \dot{V}^k = k$. Define an orbibundle $\dot{E}^k \rightarrow \dot{V}^k$, a smooth section $\dot{s}^k$ of $\dot{E}^k$, a continuous map $\dot{\psi}^k : (\dot{s}^k)^{-1}(0) \rightarrow X \times_Y \tilde{X}$ and a smooth map $\dot{\pi}_Y^k : \dot{V}^k \rightarrow Y$ by $(\dot{E}^k, \dot{s}^k, \dot{\psi}^k)|_{\dot{V}^{(i,\tilde{i})}} = (E^{(i,\tilde{i})}, s^{(i,\tilde{i})}, \psi^{(i,\tilde{i})})|_{\dot{V}^{(i,\tilde{i})}}$, $\dot{\pi}_Y^k|_{\dot{V}^{(i,\tilde{i})}} = \pi_Y^{(i,\tilde{i})}|_{\dot{V}^{(i,\tilde{i})}}$ for all $i, \tilde{i}$ in (38).

We claim that $(\dot{V}^k, \dots, \dot{\psi}^k)$ is a *Kuranishi neighbourhood* on $X \times_Y \tilde{X}$. Since it is a disjoint union of open subsets of Kuranishi neighbourhoods (37), the only thing to check is that $\dot{\psi}^k$ is injective, the problem identified in (*) above. It is enough to show that if $i, j \in I$ and $\tilde{i}, \tilde{j} \in \tilde{I}$ with $(i, \tilde{i}) \neq (j, \tilde{j})$ and $i + \tilde{i} = j + \tilde{j} = k + \dim Y$ then $\text{Im } \psi^{(i, \tilde{i})}|_{\dot{V}^{(i, \tilde{i})}} \cap \text{Im } \psi^{(j, \tilde{j})}|_{\dot{V}^{(j, \tilde{j})}} = \emptyset$. But these conditions imply that either $i < j$ and $\tilde{i} > \tilde{j}$, or $i > j$ and $\tilde{i} < \tilde{j}$, so (f) implies that the supports of $\eta_{(i, \tilde{i})}$ and $\eta_{(j, \tilde{j})}$ are disjoint, and thus $\text{Im } \psi^{(i, \tilde{i})}|_{\dot{V}^{(i, \tilde{i})}}, \text{Im } \psi^{(j, \tilde{j})}|_{\dot{V}^{(j, \tilde{j})}}$ are disjoint by (40). Hence $(\dot{V}^k, \dots, \dot{\psi}^k)$ is a Kuranishi neighbourhood on $X \times_Y \tilde{X}$.

Next, for $j \leq i$ in I and $\tilde{j} \leq \tilde{i}$ in $\tilde{I}$ define

$$\dot{V}^{(i, \tilde{i})(j, \tilde{j})} = V^{(i, \tilde{i})(j, \tilde{j})} \cap \dot{V}^{(j, \tilde{j})} \cap (\phi^{(i, \tilde{i})(j, \tilde{j})})^{-1}(\dot{V}^{(i, \tilde{i})}). \quad (41)$$

For $k, l \in \dot{I}$ with $l \leq k$, define

$$\dot{V}^{kl} = \bigcup_{\substack{i, j \in I, \tilde{i}, \tilde{j} \in \tilde{I}: j \leq i, \tilde{j} \leq \tilde{i}, \\ i + \tilde{i} = k + \dim Y, j + \tilde{j} = l + \dim Y}} \dot{V}^{(i, \tilde{i})(j, \tilde{j})} \subseteq \dot{V}^l. \quad (42)$$

We claim that (42) is a *disjoint union*, that is, distinct $\dot{V}^{(i, \tilde{i})(j, \tilde{j})}$ in (42) do not intersect. Since $\dot{V}^{(i, \tilde{i})(j, \tilde{j})} \subseteq \dot{V}^{(j, \tilde{j})}$ and (38) is a disjoint union, it is enough to show that if $i, i', j \in I$ and $\tilde{i}, \tilde{i}', \tilde{j} \in \tilde{I}$ with $(i, \tilde{i}) \neq (i', \tilde{i}')$, $j \leq i, j \leq i', \tilde{j} \leq \tilde{i}, \tilde{j} \leq \tilde{i}'$, $i + \tilde{i} = i' + \tilde{i}' = k + \dim Y$ and $j + \tilde{j} = l + \dim Y$ then $V^{(i, \tilde{i})(j, \tilde{j})} \cap V^{(i', \tilde{i}')(j, \tilde{j})} = \emptyset$. Now $\eta_{(i, \tilde{i})}^{(i, \tilde{i})} > 0$ on $\dot{V}^{(i, \tilde{i})}$ by (39), and $\dot{V}^{(i, \tilde{i})(j, \tilde{j})} \subseteq (\phi^{(i, \tilde{i})(j, \tilde{j})})^{-1}(\dot{V}^{(i, \tilde{i})})$ by (41), and $\eta_{(i, \tilde{i})}^{(j, \tilde{j})}|_{V^{(i, \tilde{i})(j, \tilde{j})}} \equiv \eta_{(i, \tilde{i})}^{(i, \tilde{i})} \circ \phi^{(i, \tilde{i})(j, \tilde{j})}$ by (c), so $\eta_{(i, \tilde{i})}^{(j, \tilde{j})} > 0$ on $V^{(i, \tilde{i})(j, \tilde{j})}$, and similarly $\eta_{(i', \tilde{i}')}^{(j, \tilde{j})} > 0$ on $V^{(i', \tilde{i}')(j, \tilde{j})}$. Hence $\eta_{(i, \tilde{i})}^{(j, \tilde{j})} \eta_{(i', \tilde{i}')}^{(j, \tilde{j})} > 0$ on $V^{(i, \tilde{i})(j, \tilde{j})} \cap V^{(i', \tilde{i}')(j, \tilde{j})}$. But the conditions imply that either $i < i'$ and $\tilde{i} > \tilde{i}'$, or $i > i'$ and $\tilde{i} < \tilde{i}'$, so (f) gives $\eta_{(i, \tilde{i})}^{(j, \tilde{j})} \eta_{(i', \tilde{i}')}^{(j, \tilde{j})} \equiv 0$ on $V^{(j, \tilde{j})}$. Thus $V^{(i, \tilde{i})(j, \tilde{j})} \cap V^{(i', \tilde{i}')(j, \tilde{j})} = \emptyset$, and (42) is a disjoint union.

Define a coordinate transformation

$$(\dot{\phi}^{kl}, \hat{\phi}^{kl}) : (\dot{V}^{kl}, \dot{E}^l|_{\dot{V}^{kl}}, \dot{s}^l|_{\dot{V}^{kl}}, \dot{\psi}^l|_{\dot{V}^{kl}}) \longrightarrow (\dot{V}^k, \dots, \dot{\psi}^k) \quad (43)$$

by $(\dot{\phi}^{kl}, \hat{\phi}^{kl})|_{\dot{V}^{(i, \tilde{i})(j, \tilde{j})}} = (\phi^{(i, \tilde{i})(j, \tilde{j})}, \hat{\phi}^{(i, \tilde{i})(j, \tilde{j})})|_{\dot{V}^{(i, \tilde{i})(j, \tilde{j})}}$ for all $i, j, \tilde{i}, \tilde{j}$ in (42). This is well-defined as (42) is a disjoint union. Since each $(\phi^{(i, \tilde{i})(j, \tilde{j})}, \hat{\phi}^{(i, \tilde{i})(j, \tilde{j})})$ is a coordinate transformation, to check (43) is a coordinate transformation it only remains to check that $\dot{\phi}^{kl}$ in (43) is injective. It is enough to show that if $i, j, j' \in I$ and $\tilde{i}, \tilde{j}, \tilde{j}' \in \tilde{I}$ with $(j, \tilde{j}) \neq (j', \tilde{j}')$, $j \leq i, j' \leq i, \tilde{j} \leq \tilde{i}, \tilde{j}' \leq \tilde{i}, i + \tilde{i} = k + \dim Y$ and $j + \tilde{j} = j' + \tilde{j}' = l + \dim Y$ then $\phi^{(i, \tilde{i})(j, \tilde{j})}(\dot{V}^{(i, \tilde{i})(j, \tilde{j})}) \cap \phi^{(i, \tilde{i})(j', \tilde{j}')}(\dot{V}^{(i, \tilde{i})(j', \tilde{j}')}) = \emptyset$ in $\dot{V}^{(i, \tilde{i})}$. This holds as by a similar argument $\eta_{(j, \tilde{j})}^{(i, \tilde{i})} > 0$ on $\phi^{(i, \tilde{i})(j, \tilde{j})}(\dot{V}^{(i, \tilde{i})(j, \tilde{j})})$ and $\eta_{(j', \tilde{j}')}^{(i, \tilde{i})} > 0$ on $\phi^{(i, \tilde{i})(j', \tilde{j}')}(\dot{V}^{(i, \tilde{i})(j', \tilde{j}')})$, but $\eta_{(j, \tilde{j})}^{(i, \tilde{i})} \eta_{(j', \tilde{j}')}^{(i, \tilde{i})} \equiv 0$ on $\dot{V}^{(i, \tilde{i})}$.

We claim that $\dot{\mathbf{I}} = (\dot{I}, (\dot{V}^k, \dots, \dot{\psi}^k), \dot{\pi}_Y^k : k \in \dot{I}, (\dot{V}^{kl}, \dot{\phi}^{kl}, \hat{\phi}^{kl}) : l \leq k \in \dot{I})$ is a *very good coordinate system* for $(X \times_Y \tilde{X}, \pi_Y)$. Only two nontrivial things remain to check. Firstly, (g) implies that every $q \in X \times_Y \tilde{X}$ lies in $\text{Im } \psi^{(i, \tilde{i})}|_{\dot{V}^{(i, \tilde{i})}}$ for some $i, \tilde{i}$ and hence in $\text{Im } \dot{\psi}^k$ for some k , so $X \times_Y \tilde{X} = \bigcup_{k \in \dot{I}} \text{Im } \dot{\psi}^k$, as we need for $\dot{\mathbf{I}}$ to be a good coordinate system. Secondly, we must show that $\dot{V}^{kl}$ is

an open neighbourhood of $(\psi^l)^{-1}(\text{Im } \psi^k)$ in $\dot{V}^l$ whenever $l \leq k$ in $\dot{I}$. It is enough to show that if $i, j \in I$ and $\tilde{i}, \tilde{j} \in \tilde{I}$ with $i + \tilde{i} = k + \dim Y$ and $j + \tilde{j} = l + \dim Y$ then:

- (A) if $j \leq i$ and $\tilde{j} \leq \tilde{i}$ then $\dot{V}^{(i, \tilde{i})(j, \tilde{j})}$ is an open neighbourhood of $(\psi^{(j, \tilde{j})}|_{\dot{V}^{(j, \tilde{j})}})^{-1}(\text{Im } \psi^{(i, \tilde{i})}|_{\dot{V}^{(i, \tilde{i})}})$ in $\dot{V}^{(j, \tilde{j})}$, and
- (B) if either $j > i$ and $\tilde{j} < \tilde{i}$, or $j < i$ and $\tilde{j} > \tilde{i}$, then $(\text{Im } \psi^{(i, \tilde{i})}|_{\dot{V}^{(i, \tilde{i})}}) \cap (\text{Im } \psi^{(j, \tilde{j})}|_{\dot{V}^{(j, \tilde{j})}}) = \emptyset$.

Part (A) is immediate from (41) and the fact that $V^{(i, \tilde{i})(j, \tilde{j})}$ is an open neighbourhood of $(\psi^{(j, \tilde{j})})^{-1}(\text{Im } \psi^{(i, \tilde{i})})$ in $V^{(j, \tilde{j})}$, which follows from the corresponding statements for V^{ij}, ψ^j, ψ^i in X and $\tilde{V}^{\tilde{i}\tilde{j}}, \tilde{\psi}^{\tilde{j}}, \tilde{\psi}^{\tilde{i}}$ in $\tilde{X}$. Part (B) is the problem identified in (**) above. To prove it, note that (f) gives $\eta_{(i, \tilde{i})}\eta_{(j, \tilde{j})} \equiv 0$ on $X \times_Y \tilde{X}$, but (40) yields $\eta_{(i, \tilde{i})} > 0$ on $\text{Im } \psi^{(i, \tilde{i})}|_{\dot{V}^{(i, \tilde{i})}}$ and $\eta_{(j, \tilde{j})} > 0$ on $\text{Im } \psi^{(j, \tilde{j})}|_{\dot{V}^{(j, \tilde{j})}}$. Therefore $\dot{\mathbf{I}}$ is a very good coordinate system.

Next we define the partition of unity data $\dot{\boldsymbol{\eta}}$ for $\dot{\mathbf{I}}$. For $k, l \in \dot{I}$, define continuous $\dot{\eta}_k : X \times_Y \tilde{X} \rightarrow [0, 1]$ and $\dot{\eta}_k^l : \dot{V}^l \rightarrow [0, 1]$ by

$$\dot{\eta}_k = \sum_{i \in I, \tilde{i} \in \tilde{I}: i + \tilde{i} = k + \dim Y} \eta_{(i, \tilde{i})}, \quad (44)$$

and for all $j \in I, \tilde{j} \in \tilde{I}$ with $j + \tilde{j} = l + \dim Y$,

$$\dot{\eta}_k^l|_{\dot{V}^{(j, \tilde{j})}} = \sum_{i \in I, \tilde{i} \in \tilde{I}: i + \tilde{i} = k + \dim Y} \eta_{(i, \tilde{i})}^{(j, \tilde{j})}. \quad (45)$$

All of Definition 3.4(i)–(iv) except the last sentence of (i) and the last two sentences of (ii) follow easily from (a)–(c) above, and these parts of (i), (ii) follow from (d), (e) respectively. If $M, \tilde{M}, T, \tilde{T}$ are as in Definition 3.4(v), (vi) for $(\mathbf{I}, \boldsymbol{\eta}), (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$, it is easy to see that Definition 3.4(v), (vi) hold for $(\dot{\mathbf{I}}, \dot{\boldsymbol{\eta}})$ with constant $\dot{M} = M + \tilde{M}$ and compact set $\dot{T} = T \cap \tilde{T}$. Hence $(\dot{\mathbf{I}}, \dot{\boldsymbol{\eta}})$ is a *really good coordinate system*, which we will also write as $(\mathbf{I}, \boldsymbol{\eta}) \times_Y (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$.

It remains to define the continuous functions $\eta_{(i, \tilde{i})}$ and $\eta_{(i, \tilde{i})}^{(j, \tilde{j})}$ satisfying (a)–(g) above. The obvious definitions for these are $\eta_{(i, \tilde{i})} = (\eta_i \circ \pi_X) \cdot (\tilde{\eta}_{\tilde{i}} \circ \pi_{\tilde{X}})$ and $\eta_{(i, \tilde{i})}^{(j, \tilde{j})} = (\eta_i^j \circ \pi_{V^j}) \cdot (\tilde{\eta}_{\tilde{i}}^{\tilde{j}} \circ \pi_{\tilde{V}^{\tilde{j}}})$, but these will not do as they satisfy (a)–(e) and (g) but not (f), which was crucial in solving problems (*) and (**) above.

There may be many ways to define $\eta_{(i, \tilde{i})}, \eta_{(i, \tilde{i})}^{(j, \tilde{j})}$ satisfying (a)–(g). The definition we give was chosen very carefully to ensure that Theorem 3.30 below holds, so that fibre products of really good coordinate systems are *commutative* and *associative*. It seems remarkable that such a definition is possible.

Before defining the $\eta_{(i, \tilde{i})}, \eta_{(i, \tilde{i})}^{(j, \tilde{j})}$ we set up some notation. For each $k \geq 0$, define the k -*simplex* to be

$$\Delta_k = \{(x_0, \dots, x_k) \in \mathbb{R}^{k+1} : x_i \geq 0, x_0 + \dots + x_k = 1\},$$

as in (59) below. Write $I = \{i_0, i_1, \dots, i_k\}$ with $0 \leq i_0 < i_1 < \dots < i_k$ and $\tilde{I} = \{\tilde{i}_0, \tilde{i}_1, \dots, \tilde{i}_l\}$ with $0 \leq \tilde{i}_0 < \tilde{i}_1 < \dots < \tilde{i}_l$, so that $|I| = k + 1$ and $|\tilde{I}| = l + 1$. For $a = 0, \dots, k$ and $b = 0, \dots, l$ we will define continuous functions $\zeta_{(a, b)}^{k, l} : \Delta_k \times \Delta_l \rightarrow [0, 1]$ satisfying the conditions:

- (A) $\sum_{a=0}^k \sum_{b=0}^l \zeta_{(a,b)}^{k,l} \equiv 1$ on $\Delta_k \times \Delta_l$.
- (B) $\zeta_{(a,b)}^{k,l}$ is supported on $\{(x_0, \dots, x_k), (y_0, \dots, y_l)\} \in \Delta_k \times \Delta_l : x_a > 0, y_b > 0\}$, for all a, b .
- (C) Suppose $a, a' = 0, \dots, k$ and $b, b' = 0, \dots, l$ and either $a < a'$ and $b > b'$ or $a > a'$ and $b < b'$. Then $\zeta_{(a,b)}^{k,l} \zeta_{(a',b')}^{k,l} \equiv 0$ on $\Delta_k \times \Delta_l$.
- (D) If $x_0 = \dots = x_{a-1} = 0, x_a > 0$ and $y_0 = \dots = y_{b-1} = 0, y_b > 0$ then $\zeta_{(a,b)}^{k,l}((x_0, \dots, x_k), (y_0, \dots, y_l)) > 0$.

Then we define continuous $\eta_{(i_a, \tilde{i}_b)} : X \times_Y \tilde{X} \rightarrow [0, 1]$ and $\eta_{(i_a, \tilde{i}_b)}^{(j, \tilde{j})} : V^{(j, \tilde{j})} \rightarrow [0, 1]$ by

$$\eta_{(i_a, \tilde{i}_b)}(q) = \zeta_{(a,b)}^{k,l} [(\eta_{i_0} \circ \pi_X(q), \eta_{i_1} \circ \pi_X(q), \dots, \eta_{i_k} \circ \pi_X(q)), (\tilde{\eta}_{\tilde{i}_0} \circ \pi_{\tilde{X}}(q), \tilde{\eta}_{\tilde{i}_1} \circ \pi_{\tilde{X}}(q), \dots, \tilde{\eta}_{\tilde{i}_l} \circ \pi_{\tilde{X}}(q))], \quad (46)$$

$$\eta_{(i_a, \tilde{i}_b)}^{(j, \tilde{j})}(v) = \zeta_{(a,b)}^{k,l} [(\eta_{i_0}^j \circ \pi_{V^j}(v), \eta_{i_1}^j \circ \pi_{V^j}(v), \dots, \eta_{i_k}^j \circ \pi_{V^j}(v)), (\tilde{\eta}_{\tilde{i}_0}^{\tilde{j}} \circ \pi_{\tilde{V}^{\tilde{j}}}(v), \tilde{\eta}_{\tilde{i}_1}^{\tilde{j}} \circ \pi_{\tilde{V}^{\tilde{j}}}(v), \dots, \tilde{\eta}_{\tilde{i}_l}^{\tilde{j}} \circ \pi_{\tilde{V}^{\tilde{j}}}(v))]. \quad (47)$$

Here $(\eta_{i_0} \circ \pi_X(p), \eta_{i_1} \circ \pi_X(p), \dots, \eta_{i_k} \circ \pi_X(p)) \in \Delta_k$ since $\sum_{a=0}^l \eta_{i_a} = \sum_{i \in I} \eta_i \equiv 1$ on X and $\eta_{i_a} \geq 0$ for all $a = 0, \dots, l$ as η_{i_a} maps $X \rightarrow [0, 1]$. In the same way $(\tilde{\eta}_{\tilde{i}_0} \circ \pi_{\tilde{X}}(p), \tilde{\eta}_{\tilde{i}_1} \circ \pi_{\tilde{X}}(p), \dots, \tilde{\eta}_{\tilde{i}_l} \circ \pi_{\tilde{X}}(p)) \in \Delta_l$, so (46) is well-defined, and similarly (47) is well-defined. Part (a) above follows from (A) and (46)–(47). Parts (b), (c) are immediate from (46)–(47), Definition 3.4(iii), (iv) and the identities $\pi_X \circ \psi^{(j, \tilde{j})} \equiv \psi^j \circ \pi_{V^j} : (s^{(j, \tilde{j})})^{-1}(0) \rightarrow X$, $\pi_{\tilde{X}} \circ \psi^{(j, \tilde{j})} \equiv \tilde{\psi}^{\tilde{j}} \circ \pi_{\tilde{V}^{\tilde{j}}} : (s^{(j, \tilde{j})})^{-1}(0) \rightarrow \tilde{X}$, $\pi_{V^j} \circ \phi^{(j, \tilde{j})(k, \tilde{k})} \equiv \phi^{jk} \circ \pi_{V^k} : V^{(j, \tilde{j})(k, \tilde{k})} \rightarrow V^j$, and $\pi_{\tilde{V}^{\tilde{j}}} \circ \phi^{(j, \tilde{j})(k, \tilde{k})} \equiv \tilde{\phi}^{\tilde{j}\tilde{k}} \circ \pi_{\tilde{V}^{\tilde{k}}} : V^{(j, \tilde{j})(k, \tilde{k})} \rightarrow \tilde{V}^{\tilde{j}}$.

Part (d) follows from (B), the last sentence of Definition 3.4(i) for $\text{Im } \psi^i$ and $\text{Im } \tilde{\psi}^{\tilde{i}}$, and $\text{Im } \psi^{(i, \tilde{i})} = (\pi_X \times \pi_{\tilde{X}})^{-1}(\text{Im } \psi^i \times \text{Im } \tilde{\psi}^{\tilde{i}})$. Part (e) follows from (B), the last two sentences of Definition 3.4(ii) for V^{ij} and $\tilde{V}^{\tilde{i}\tilde{j}}$ and $V^{(i, \tilde{i})(j, \tilde{j})} = V^{ij} \times_Y \tilde{V}^{\tilde{i}\tilde{j}}$. Part (f) follows from (C) and (46)–(47). For (g), suppose $q \in X \times_Y X$, set $p = \pi_X(q)$ and $\tilde{p} = \pi_{\tilde{X}}(q)$, and let $i_a \in I$ and $\tilde{i}_b \in \tilde{I}$ be least with $\eta_{i_a}(p) > 0$ and $\tilde{\eta}_{\tilde{i}_b}(\tilde{p}) > 0$. Then by (46) we have $\eta_{(i_a, \tilde{i}_b)}(q) = \zeta_{(a,b)}^{k,l}((x_0, \dots, x_k), (y_0, \dots, y_l))$ with $x_c = \eta_{i_c}(p)$ and $y_d = \tilde{\eta}_{\tilde{i}_d}(\tilde{p})$, so $x_0 = \dots = x_{a-1} = 0, x_a > 0$ and $y_0 = \dots = y_{b-1} = 0, y_b > 0$ by choice of $i_a, \tilde{i}_b$, and thus $\eta_{(i_a, \tilde{i}_b)}(q) > 0$ by (D), the first part of (g). For the second part, note that as $X = \bigcup_{j \in I} \text{Im } \psi^j$ we have $p \in \text{Im } \psi^j$ for some $j \in I$, so $\eta_j(p) > 0$ by Definition 3.4(i). Therefore $j \geq i_a$, as i_a is least with $\eta_{i_a}(p) > 0$. The final part of Definition 3.4(i) now implies that $p \in \text{Im } \psi^{i_a}$. Similarly $\tilde{p} \in \text{Im } \tilde{\psi}^{\tilde{i}_b}$. Thus $q \in \pi_X^{-1}(\text{Im } \psi^{i_a}) \cap \pi_{\tilde{X}}^{-1}(\text{Im } \tilde{\psi}^{\tilde{i}_b})$, so equation (40) implies that $q \in \text{Im } \psi^{(i, \tilde{i})}$.

Finally we construct the functions $\zeta_{(a,b)}^{k,l} : \Delta_k \times \Delta_l \rightarrow [0, 1]$. To do this we first write down a *triangulation* of $\Delta_k \times \Delta_l$ into $(k+l)$ -simplices Δ_{k+l} , and then define the $\zeta_{(a,b)}^{k,l}$ on each simplex Δ_{k+l} to be the unique affine function with prescribed values at each vertex of Δ_{k+l} . For $0 \leq a < k$ and $0 \leq b < l$, define the wall $W_{a,b} \subset \Delta_k \times \Delta_l$ to be the set of $((x_0, \dots, x_k), (y_0, \dots, y_l))$ for which

$x_0 + \cdots + x_a = y_0 + \cdots + y_b$. These walls $W_{a,b}$ for all a, b cut $\Delta_k \times \Delta_l$ into $\binom{k+l}{k}$ regions affine-isomorphic to the $(k+l)$ -simplex Δ_{k+l} .

Here is one way to describe these regions. Suppose $\{1, \dots, k+l\} = A \amalg B$ with $|A| = k$ and $|B| = l$. Write $A = \{r_0, r_1, \dots, r_{k-1}\}$ with $r_0 < \cdots < r_{k-1}$ and $B = \{s_0, s_1, \dots, s_{l-1}\}$ with $s_0 < \cdots < s_{l-1}$. Define

$$C_{A,B} = \{((x_0, \dots, x_k), (y_0, \dots, y_l)) \in \Delta_k \times \Delta_l : t_1 \leq t_2 \leq \cdots \leq t_{k+l}\},$$

where $t_c = x_0 + \cdots + x_a$ if $c = r_a$, and $t_c = y_0 + \cdots + y_a$ if $c = s_a$.

Then $C_{A,B}$ for all such A, B are the closures of the regions into which the $W_{a,b}$ divide the $\Delta_k \times \Delta_l$. Write $p_0, \dots, p_k$ for the vertices of Δ_k , where $p_a = (\delta_{0a}, \delta_{1a}, \dots, \delta_{ka})$, and $q_0, \dots, q_l$ for the vertices of Δ_l , where $q_b = (\delta_{0b}, \dots, \delta_{lb})$. Then each $C_{A,B}$ is a $(k+l)$ -simplex, with $k+l+1$ vertices, all of the form (p_c, q_d) for some c, d determined by A, B .

For $a = 0, \dots, k$ and $b = 0, \dots, l$, define $\zeta_{(a,b)}^{k,l} : \Delta_k \times \Delta_l \rightarrow \mathbb{R}$ to be affine linear on the $(k+l)$ -simplex $C_{A,B}$ for all A, B as above, and to satisfy $\zeta_{(a,b)}^{k,l}(p_c, q_d) = \delta_{ac}\delta_{bd}$ for all $c = 0, \dots, k$ and $d = 0, \dots, l$. This determines $\zeta_{(a,b)}^{k,l}$ uniquely on each $C_{A,B}$, since we have prescribed $\zeta_{(a,b)}^{k,l}$ on all the vertices (p_c, q_d) of the $(k+l)$ -simplex $C_{A,B}$, and there is a unique affine linear function taking any given values at the vertices.

Each intersection $C_{A,B} \cap C_{A',B'}$ in $\Delta_k \times \Delta_l$ is empty or a simplex Δ_m for $0 \leq m \leq k+l$, whose vertices (p_c, q_d) are the common vertices of $C_{A,B}, C_{A',B'}$. The definitions of $\zeta_{(a,b)}^{k,l}$ on $C_{A,B}$ and $C_{A',B'}$ both restrict to the unique affine linear function on $\Delta_m = C_{A,B} \cap C_{A',B'}$ with $\zeta_{(a,b)}^{k,l}(p_c, q_d) = \delta_{ac}\delta_{bd}$ on the $m+1$ vertices (p_c, q_d) of Δ_m . Thus the definitions of $\zeta_{(a,b)}^{k,l}$ on $C_{A,B}$ and $C_{A',B'}$ agree on $C_{A,B} \cap C_{A',B'}$, so $\zeta_{(a,b)}^{k,l}$ is well-defined. The maximum and minimum values of an affine linear function on a simplex are achieved at the vertices, and we have prescribed the values 0 or 1 for each vertex. Thus, $\zeta_{(a,b)}^{k,l}$ maps $\Delta_k \times \Delta_l \rightarrow [0, 1]$. Also, $\zeta_{(a,b)}^{k,l}|_{C_{A,B}}$ is continuous as it is affine linear, so since $\Delta_k \times \Delta_l = \bigcup_{A,B} C_{A,B}$ and the $C_{A,B}$ are closed, $\zeta_{(a,b)}^{k,l}$ is continuous.

We must verify (A)–(D) above. We have $\sum_{a=0}^k \sum_{b=0}^l \zeta_{(a,b)}^{k,l}(p_c, q_d) = \sum_{a=0}^k \sum_{b=0}^l \delta_{ac}\delta_{bd} = 1$. Thus, $\sum_{a=0}^k \sum_{b=0}^l \zeta_{(a,b)}^{k,l}|_{C_{A,B}}$ is the unique affine linear function which is 1 on each vertex (p_c, q_d) of $C_{A,B}$, so $\sum_{a=0}^k \sum_{b=0}^l \zeta_{(a,b)}^{k,l}|_{C_{A,B}} \equiv 1$. This holds for all $C_{A,B}$, giving (A). For (B), each $C_{A,B}$ has $k+l+1$ vertices of the form (p_c, q_d) . If (p_a, q_b) is not a vertex of $C_{A,B}$ then $\zeta_{(a,b)}^{k,l}$ is zero at each vertex (p_c, q_d) of $C_{A,B}$, so $\zeta_{(a,b)}^{k,l} \equiv 0$ on $C_{A,B}$, and (B) holds trivially on $C_{A,B}$. If (p_a, q_b) is a vertex of $C_{A,B}$ then $\zeta_{(a,b)}^{k,l}$ is 1 at this vertex and 0 at all other vertices of $C_{A,B}$. Then the support of $\zeta_{(a,b)}^{k,l}|_{C_{A,B}}$, and the subset of $C_{A,B}$ on which $x_a > 0$ and $y_b > 0$, are both the complement in $C_{A,B}$ of the unique codimension one face of $C_{A,B}$ not containing (p_a, q_b) . This proves (B).

For (C), suppose $0 \leq a < a' \leq k$ and $0 \leq b' < b \leq l$. Then (p_a, q_b) lies

strictly on one side $x_0 + \dots + x_a > y_0 + \dots + y_{b'}$ of the wall $W_{a,b'}$, and $(p_{a'}, q_{b'})$ lies strictly on the other side $x_0 + \dots + x_a < y_0 + \dots + y_{b'}$ of the wall $W_{a,b'}$. Since each $C_{A,B}$ lies on one or other side of $W_{a,b'}$, it follows that for each $C_{A,B}$, at most one of (p_a, p_b) , $(p_{a'}, p_{b'})$ can be a vertex of $C_{A,B}$, so at most one of $\zeta_{(a,b)}^{k,l}, \zeta_{(a',b')}^{k,l}$ can be nonzero on $C_{A,B}$, and $\zeta_{(a,b)}^{k,l} \zeta_{(a',b')}^{k,l} \equiv 0$ on $C_{A,B}$. Therefore $\zeta_{(a,b)}^{k,l} \zeta_{(a',b')}^{k,l} \equiv 0$ on $\Delta_k \times \Delta_l$, as this holds for all A, B . The case $0 \leq a' < a \leq k$ and $0 \leq b < b' \leq l$ follows in the same way using $W_{a',b}$. This proves (C).

For (D), let $x_0 = \dots = x_{a-1} = 0$, $x_a > 0$ and $y_0 = \dots = y_{b-1} = 0$, $y_b > 0$. For any $0 \leq a' < k$ and $0 \leq b' < l$, by considering the cases $a' < a$, $a' \geq a$ and $b' < b$, $b' \geq b$ separately it is easy to see that $((x_0, \dots, x_k), (y_0, \dots, y_l))$ and (p_a, q_b) cannot lie strictly on opposite sides of the wall $W_{a',b'}$ in $\Delta_k \times \Delta_l$. Hence there is some $(k+l)$ -simplex $C_{A,B}$ containing both $((x_0, \dots, x_k), (y_0, \dots, y_l))$ and (p_a, q_b) . Now $\zeta_{(a,b)}^{k,l}$ is affine linear on $C_{A,B}$, and is 1 on the vertex (p_a, q_b) , and 0 on the Δ_{k+l-1} face of $C_{A,B}$ opposite (p_a, q_b) , and positive otherwise. As $x_a > 0$, $y_b > 0$ we see that $((x_0, \dots, x_k), (y_0, \dots, y_l))$ cannot lie in the Δ_{k+l-1} opposite (p_a, q_b) , so $\zeta_{(a,b)}^{k,l}((x_0, \dots, x_k), (y_0, \dots, y_l)) > 0$. This proves (D), and completes the construction of $(\mathbf{I}, \boldsymbol{\eta}) \times_Y (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$, and Definition 3.29.

The construction of Definition 3.29 is *commutative* and *associative*. This will imply fibre products of (co-)gauge-fixing data are commutative and associative.

Theorem 3.30. *Let Y be an orbifold, and for $i = 1, 2, 3$ let X_i be a compact Kuranishi space, $\mathbf{f}_i : X_i \rightarrow Y$ a strong submersion, and $(\mathbf{I}_i, \boldsymbol{\eta}_i)$ a really good coordinate system for $(X_i, \mathbf{f}_i)$. Then the really good coordinate systems $(\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y (\mathbf{I}_2, \boldsymbol{\eta}_2)$ for $(X_1 \times_Y X_2, \pi_Y)$ and $(\mathbf{I}_2, \boldsymbol{\eta}_2) \times_Y (\mathbf{I}_1, \boldsymbol{\eta}_1)$ for $(X_2 \times_Y X_1, \pi_Y)$ are identified by the natural strong diffeomorphism*

$$X_1 \times_{\mathbf{f}_1, Y, \mathbf{f}_2} X_2 \cong X_2 \times_{\mathbf{f}_2, Y, \mathbf{f}_1} X_1. \quad (48)$$

Also, the really good coordinate systems $((\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y (\mathbf{I}_2, \boldsymbol{\eta}_2)) \times_Y (\mathbf{I}_3, \boldsymbol{\eta}_3)$ for $((X_1 \times_Y X_2) \times_Y X_3, \pi_Y)$ and $(\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y ((\mathbf{I}_2, \boldsymbol{\eta}_2) \times_Y (\mathbf{I}_3, \boldsymbol{\eta}_3))$ for $(X_1 \times_Y (X_2 \times_Y X_3), \pi_Y)$ are identified by the natural strong diffeomorphism

$$(X_1 \times_{\mathbf{f}_1, Y, \mathbf{f}_2} X_2) \times_{\pi_Y, Y, \mathbf{f}_3} X_3 \cong X_1 \times_{\mathbf{f}_1, Y, \pi_Y} (X_2 \times_{\mathbf{f}_2, Y, \mathbf{f}_3} X_3). \quad (49)$$

Proof. In Definition 3.29, adapting notation in the obvious way, for $i_1 \in I_1$ and $i_2 \in I_2$ we have Kuranishi neighbourhoods $(V_{12}^{(i_1, i_2)}, \dots, \psi_{12}^{(i_1, i_2)})$ on $X_1 \times_Y X_2$ and $(V_{21}^{(i_2, i_1)}, \dots, \psi_{21}^{(i_2, i_1)})$ on $X_2 \times_Y X_1$ from (37), with $V_{12}^{(i_1, i_2)} = V_1^{i_1} \times_{f_1^{i_1}, Y, f_2^{i_2}} V_2^{i_2}$ and $V_{21}^{(i_2, i_1)} = V_2^{i_2} \times_{f_2^{i_2}, Y, f_1^{i_1}} V_1^{i_1}$. Parallel to (48) we have a natural diffeomorphism $V_1^{i_1} \times_{f_1^{i_1}, Y, f_2^{i_2}} V_2^{i_2} \cong V_2^{i_2} \times_{f_2^{i_2}, Y, f_1^{i_1}} V_1^{i_1}$, that is, $V_{12}^{(i_1, i_2)} \cong V_{21}^{(i_2, i_1)}$. We also have functions $\eta_{(i_1, i_2)} : X_1 \times_Y X_2 \rightarrow [0, 1]$ and $\eta_{(i_2, i_1)} : X_2 \times_Y X_1 \rightarrow [0, 1]$, and $\eta_{(i_1, i_2)}^{(j_1, j_2)} : V_{12}^{(j_1, j_2)} \rightarrow [0, 1]$ and $\eta_{(i_2, i_1)}^{(j_2, j_1)} : V_{21}^{(j_2, j_1)} \rightarrow [0, 1]$ for $j_1 \in I_1$, $j_2 \in I_2$.

We claim that (48) identifies $\eta_{(i_1, i_2)} : X_1 \times_Y X_2 \rightarrow [0, 1]$ and $\eta_{(i_2, i_1)} : X_2 \times_Y X_1 \rightarrow [0, 1]$, and the diffeomorphism $V_{12}^{(j_1, j_2)} \cong V_{21}^{(j_2, j_1)}$ identifies $\eta_{(i_1, i_2)}^{(j_1, j_2)}$ and $\eta_{(i_2, i_1)}^{(j_2, j_1)}$. Following the definitions through, this is immediate since the $\zeta_{(a, b)}^{k, l}$ satisfy $\zeta_{(a, b)}^{k, l}((x_0, \dots, x_k), (y_0, \dots, y_l)) \equiv \zeta_{(b, a)}^{l, k}((y_0, \dots, y_l), (x_0, \dots, x_k))$.

Equation (39) now shows that the open subsets $\dot{V}_{12}^{(i_1, i_2)} \subseteq V_{12}^{(i_1, i_2)}$ and $\dot{V}_{21}^{(i_2, i_1)} \subseteq V_{21}^{(i_2, i_1)}$ are identified by the diffeomorphism $V_{12}^{(i_1, i_2)} \cong V_{21}^{(i_2, i_1)}$. So by (38), for each k we have a natural diffeomorphism between $\dot{V}_{12}^k$ and $\dot{V}_{21}^k$, which easily extends to an identification of the Kuranishi neighbourhoods $(\dot{V}_{12}^k, \dots, \dot{\psi}_{12}^k)$ in $(\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y (\mathbf{I}_2, \boldsymbol{\eta}_2)$ and $(\dot{V}_{21}^k, \dots, \dot{\psi}_{21}^k)$ in $(\mathbf{I}_2, \boldsymbol{\eta}_2) \times_Y (\mathbf{I}_1, \boldsymbol{\eta}_1)$ under (48). It is an exercise to show that all the rest of the data in $(\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y (\mathbf{I}_2, \boldsymbol{\eta}_2)$ and $(\mathbf{I}_2, \boldsymbol{\eta}_2) \times_Y (\mathbf{I}_1, \boldsymbol{\eta}_1)$ is compatible with these identifications for all k , which proves the first part of the theorem.

For the second part, we follow a similar method, except that the conditions on the $\zeta_{(a, b)}^{k, l}$ are more complicated. Let $i_e \in I_e$ for $e = 1, 2, 3$, and define

$$\begin{aligned} V_{(12)3}^{(i_1, i_2, i_3)} &= (V_1^{i_1} \times_{f_1^{i_1}, Y, f_2^{i_2}} V_2^{i_2}) \times_{\pi_Y, Y, f_3^{i_3}} V_3^{i_3} \quad \text{and} \\ V_{1(23)}^{(i_1, i_2, i_3)} &= V_1^{i_1} \times_{f_1^{i_1}, Y, \pi_Y} (V_2^{i_2} \times_{f_2^{i_2}, Y, f_3^{i_3}} V_3^{i_3}). \end{aligned}$$

These naturally extend to Kuranishi neighbourhoods $(V_{(12)3}^{(i_1, i_2, i_3)}, \dots, \psi_{(12)3}^{(i_1, i_2, i_3)})$ on $(X_1 \times_Y X_2) \times_Y X_3$ and $(V_{1(23)}^{(i_1, i_2, i_3)}, \dots, \psi_{1(23)}^{(i_1, i_2, i_3)})$ on $X_1 \times_Y (X_2 \times_Y X_3)$. As for (49), there is a natural diffeomorphism $V_{(12)3}^{(i_1, i_2, i_3)} \cong V_{1(23)}^{(i_1, i_2, i_3)}$, which extends to identifications of the Kuranishi neighbourhoods $(V_{(12)3}^{(i_1, i_2, i_3)}, \dots, \psi_{(12)3}^{(i_1, i_2, i_3)})$ and $(V_{1(23)}^{(i_1, i_2, i_3)}, \dots, \psi_{1(23)}^{(i_1, i_2, i_3)})$ under (49).

Set $I = \{i_1 + i_2 + i_3 : i_1 \in I_1, i_2 \in I_2, i_3 \in I_3\}$. Applying Definition 3.29 twice, we have functions $\eta_k : (X_1 \times_Y X_2) \times_Y X_3 \rightarrow [0, 1]$ and $\eta_k^{(i_1, i_2, i_3)} : V_{(12)3}^{(i_1, i_2, i_3)} \rightarrow [0, 1]$ for $k \in I$ satisfying analogues of Definition 3.4(i)–(iv), and open subsets $\dot{V}_{(12)3}^{(i_1, i_2, i_3)} = \{v \in V_{(12)3}^{(i_1, i_2, i_3)} : \eta_{i_1+i_2+i_3}^{(i_1, i_2, i_3)}(p) > 0\}$ in $V_{(12)3}^{(i_1, i_2, i_3)}$, such that the Kuranishi neighbourhoods in $((\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y (\mathbf{I}_2, \boldsymbol{\eta}_2)) \times_Y (\mathbf{I}_3, \boldsymbol{\eta}_3)$ are $(V_{(12)3}^k, \dots, \psi_{(12)3}^k)$ with

$$V_{(12)3}^k = \coprod_{i_1 \in I_1, i_2 \in I_2, i_3 \in I_3 : i_1 + i_2 + i_3 = k} \dot{V}_{(12)3}^{(i_1, i_2, i_3)},$$

and $(E_{(12)3}^k, s_{(12)3}^k, \psi_{(12)3}^k)|_{\dot{V}_{(12)3}^{(i_1, i_2, i_3)}} = (E_{(12)3}^{(i_1, i_2, i_3)}, s_{(12)3}^{(i_1, i_2, i_3)}, \psi_{(12)3}^{(i_1, i_2, i_3)})|_{\dot{V}_{(12)3}^{(i_1, i_2, i_3)}}$. The partition of unity data in $((\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y (\mathbf{I}_2, \boldsymbol{\eta}_2)) \times_Y (\mathbf{I}_3, \boldsymbol{\eta}_3)$ is η_k as above and η_k^l given by $\eta_k^l|_{\dot{V}_{(12)3}^{(i_1, i_2, i_3)}} \equiv \eta_k^{(i_1, i_2, i_3)}$ when $i_1 + i_2 + i_3 = l$. We also have the analogous functions, sets and Kuranishi neighbourhoods for $X_1 \times_Y (X_2 \times_Y X_3)$; for clarity, we denote the corresponding functions by $\bar{\eta}_k, \bar{\eta}_k^{(i_1, i_2, i_3)}, \bar{\eta}_k^l$.

Write $I_e = \{i_0^e, i_1^e, \dots, i_{k_e}^e\}$ with $i_0^e < i_1^e < \dots < i_{k_e}^e$ for $e = 1, 2, 3$, and write $\{i_1 + i_2 : i_1 \in I_1, i_2 \in I_2\} = \{\bar{i}_0, \bar{i}_1, \dots, \bar{i}_{\bar{k}}\}$ with $\bar{i}_0 < \bar{i}_1 < \dots < \bar{i}_{\bar{k}}$. Then

computing with Definition 3.29 shows that for $k \in I$ we have

$$\eta_k^{(i_1, i_2, i_3)}(v) = \sum_{\substack{\bar{a}=0, \dots, \bar{k}, \\ a_3=0, \dots, k_3: \bar{i}_{\bar{a}} + i_{a_3}^3 = k}} \zeta_{(\bar{a}, a_3)}^{\bar{k}, k_3}((\bar{x}_0, \dots, \bar{x}_{\bar{k}}), (\eta_{i_0^3}^{k_3} \circ \pi_{V_3^{i_3}}(v), \eta_{i_1^3}^{k_3} \circ \pi_{V_3^{i_3}}(v), \dots, \eta_{i_{k_3}^3}^{k_3} \circ \pi_{V_3^{i_3}}(v))), \quad (50)$$

where $\bar{x}_c = \sum_{\substack{\bar{a}_1=0, \dots, k_1, \\ a_2=0, \dots, k_2: i_{a_1}^1 + i_{a_2}^2 = \bar{i}_c}} \zeta_{(a_1, a_2)}^{k_1, k_2}((\eta_{i_0^1}^{k_1} \circ \pi_{V_1^{i_1}}(v), \dots, \eta_{i_{k_1}^1}^{k_1} \circ \pi_{V_1^{i_1}}(v)), (\eta_{i_0^2}^{k_2} \circ \pi_{V_2^{i_2}}(v), \dots, \eta_{i_{k_2}^2}^{k_2} \circ \pi_{V_2^{i_2}}(v))).$

There is also an analogous expression for $\bar{\eta}_k^{(i_1, i_2, i_3)} : V_{1(23)}^{(i_1, i_2, i_3)} \rightarrow [0, 1]$.

It is now easy to see that $((I_1, \eta_1) \times_Y (I_2, \eta_2)) \times_Y (I_3, \eta_3)$ and $(I_1, \eta_1) \times_Y ((I_2, \eta_2) \times_Y (I_3, \eta_3))$ are identified by (49) provided the functions $\eta_k^{(i_1, i_2, i_3)}$ on $V_{(12)3}^{(i_1, i_2, i_3)}$ and $\bar{\eta}_k^{(i_1, i_2, i_3)}$ on $V_{1(23)}^{(i_1, i_2, i_3)}$ are identified by the diffeomorphism $V_{(12)3}^{(i_1, i_2, i_3)} \cong V_{1(23)}^{(i_1, i_2, i_3)}$, for all i_1, i_2, i_3 and k . To prove this, we claim that

$$\eta_k^{(i_1, i_2, i_3)}(v) = \sum_{\substack{a_1=0, \dots, k_1, \\ a_2=0, \dots, k_2, \\ a_3=0, \dots, k_3: \\ i_{a_1}^1 + i_{a_2}^2 + i_{a_3}^3 = k}} \zeta_{(a_1, a_2, a_3)}^{k_1, k_2, k_3}((\eta_{i_0^1}^{k_1} \circ \pi_{V_1^{i_1}}(v), \dots, \eta_{i_{k_1}^1}^{k_1} \circ \pi_{V_1^{i_1}}(v)), (\eta_{i_0^2}^{k_2} \circ \pi_{V_2^{i_2}}(v), \dots, \eta_{i_{k_2}^2}^{k_2} \circ \pi_{V_2^{i_2}}(v)), (\eta_{i_0^3}^{k_3} \circ \pi_{V_3^{i_3}}(v), \dots, \eta_{i_{k_3}^3}^{k_3} \circ \pi_{V_3^{i_3}}(v))). \quad (51)$$

Here $\zeta_{(a_1, a_2, a_3)}^{k_1, k_2, k_3} : \Delta_{k_1} \times \Delta_{k_2} \times \Delta_{k_3} \rightarrow [0, 1]$ is defined by analogy with the functions $\zeta_{(a, b)}^{k, l}$ of Definition 3.29. Write points of $\Delta_{k_1} \times \Delta_{k_2} \times \Delta_{k_3}$ as $((x_0, \dots, x_{k_1}), (y_0, \dots, y_{k_2}), (z_0, \dots, z_{k_3}))$. Then we introduce three kinds of walls in $\Delta_{k_1} \times \Delta_{k_2} \times \Delta_{k_3}$, defined by the equations $x_0 + \dots + x_a = y_0 + \dots + y_b$, or $x_0 + \dots + x_a = z_0 + \dots + z_c$, or $y_0 + \dots + y_b = z_0 + \dots + z_c$, for $0 \leq a < k_1$, $0 \leq b < k_2$, $0 \leq c < k_3$. These walls triangulate $\Delta_{k_1} \times \Delta_{k_2} \times \Delta_{k_3}$ into $\frac{(k_1 + k_2 + k_3)!}{k_1! k_2! k_3!}$ $(k_1 + k_2 + k_3)$ -simplices $C_{A, B, C}$ which have vertices of the form (p_a, q_b, r_c) , where p_a, q_b, r_c are vertices of $\Delta_{k_1}, \Delta_{k_2}, \Delta_{k_3}$ respectively. Then we define $\zeta_{(a_1, a_2, a_3)}^{k_1, k_2, k_3}$ to be the unique function which is affine linear on each $C_{A, B, C}$ and satisfies $\zeta_{(a_1, a_2, a_3)}^{k_1, k_2, k_3}(p_a, q_b, r_c) = \delta_{a_1 a} \delta_{a_2 b} \delta_{a_3 c}$ for all a, b, c .

To prove (51) from (50), we first show that on each simplex $C_{A, B, C}$ on which the $\zeta_{(a_1, a_2, a_3)}^{k_1, k_2, k_3}$ in (51) are affine linear maps in (50) to some simplices $C_{A', B'}$ and $C_{A'', B''}$ in the domains of $\zeta_{(\bar{a}, a_3)}^{\bar{k}, k_3}$ and $\zeta_{(a_1, a_2)}^{k_1, k_2}$ in which $\zeta_{(\bar{a}, a_3)}^{\bar{k}, k_3}, \zeta_{(a_1, a_2)}^{k_1, k_2}$ are affine linear. Since compositions of affine linear functions are affine linear, it follows that on each simplex $C_{A, B, C}$, the functions in (50) and (51) are affine linear. But it also quickly follows that (50) and (51) coincide on the vertices (p_a, q_b, r_c) of $\Delta_{k_1} \times \Delta_{k_2} \times \Delta_{k_3}$, since both are 1 if $i_a^1 + i_b^2 + i_c^3 = k$ and 0 otherwise. Therefore (50) and (51) coincide on each $C_{A, B, C}$, and so on the whole of $\Delta_{k_1} \times \Delta_{k_2} \times \Delta_{k_3}$.

This proves (51). In the same way we obtain an analogous expression for $\bar{\eta}_k^{(i_1, i_2, i_3)}(v)$. Identifying $V_{(12)3}^{(j_1, j_2, j_3)}$ and $V_{1(23)}^{(j_1, j_2, j_3)}$ using the natural diffeomorphism, these expressions coincide, so $\eta_k^{(i_1, i_2, i_3)}$ and $\bar{\eta}_k^{(i_1, i_2, i_3)}$ are identified. This completes the proof of Theorem 3.30. $\square$

Note that associativity in Theorem 3.30 worked because of special properties of the $\zeta_{(a,b)}^{k,l}$ in Definition 3.29, implying that (50),(51) are equivalent. We define fibre products of (co-)gauge-fixing data, and prove Property 3.1(h). As we want products on cohomology, not homology, we explain the co-gauge-fixing case.

Definition 3.31. Let $X, \tilde{X}$ be compact Kuranishi spaces, Y an orbifold, and $f : X \rightarrow Y, \tilde{f} : \tilde{X} \rightarrow Y$ be strong submersions. Suppose $\mathbf{C} = (\mathbf{I}, \boldsymbol{\eta}, C^i : i \in I)$ and $\tilde{\mathbf{C}} = (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}}, \tilde{C}^{\tilde{i}} : \tilde{i} \in \tilde{I})$ are co-gauge-fixing data for (X, f) and $(\tilde{X}, \tilde{f})$. Let $(\mathbf{I}, \boldsymbol{\eta}) \times_Y (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ be the really good coordinate system for $(X \times_Y \tilde{X}, \pi_Y)$ given in Definition 3.29. Write $((\mathbf{I}, \boldsymbol{\eta}) \times_Y (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}}))^\sim$ for the excellent coordinate system for $(X \times_Y \tilde{X}, \pi_Y)$ constructed from $(\mathbf{I}, \boldsymbol{\eta}) \times_Y (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ by Algorithm 3.16.

In the notation of Definition 3.29 we have $(\mathbf{I}, \boldsymbol{\eta}) \times_Y (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}}) = (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$, where $\tilde{\mathbf{I}} = (\tilde{I}, (\tilde{V}^k, \tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k), \tilde{\pi}_Y^k : k \in \tilde{I}, \dots)$, and $\tilde{V}^k = \coprod_{i \in I, \tilde{i} \in \tilde{I} : i + \tilde{i} = k + \dim Y} \tilde{V}^{(i, \tilde{i})}$ by (38), with $\tilde{V}^{(i, \tilde{i})}$ an open subset in $V^{(i, \tilde{i})} = V^i \times_{f^i, Y, \tilde{f}^{\tilde{i}}} \tilde{V}^{\tilde{i}}$. Also $\tilde{E}^k|_{\tilde{V}^{(i, \tilde{i})}} = E^{(i, \tilde{i})}|_{\tilde{V}^{(i, \tilde{i})}}$, where $E^{(i, \tilde{i})} = \pi_{V^i}^*(E^i) \oplus \pi_{\tilde{V}^{\tilde{i}}}^*(\tilde{E}^{\tilde{i}})$ by (37). Thus by Algorithm 3.16 we may write $((\mathbf{I}, \boldsymbol{\eta}) \times_Y (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}}))^\sim = (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$, where $\tilde{\mathbf{I}} = (\tilde{I}, (\tilde{V}^k, \tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k), \tilde{\pi}_Y^k : k \in \tilde{I}, \dots)$, with $\tilde{V}^k$ an open subset of $\tilde{V}^k$ for each $k \in \tilde{I}$, and $\tilde{I} = \{k \in \tilde{I} : \tilde{V}^k \neq \emptyset\}$, and $(\tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k) = (\tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k)|_{\tilde{V}^k}, \tilde{\pi}_Y^k = \tilde{\pi}_Y^k|_{\tilde{V}^k}$ for all $k \in \tilde{I}$.

For $k \in \tilde{I}$, define $\tilde{C}^k : \tilde{E}^k \rightarrow P$ by $\tilde{C}^k(\tilde{e}) = \mu(C^i \circ \pi_{E^i}(\tilde{e}), \tilde{C}^{\tilde{i}} \circ \pi_{\tilde{E}^{\tilde{i}}}(\tilde{e}))$ whenever $\tilde{e}$ lies in $\tilde{E}^k \cap (\tilde{E}^k|_{\tilde{V}^k \cap \tilde{V}^{(i, \tilde{i})}}) \subseteq E^{(i, \tilde{i})}$, for all $i \in I$ and $\tilde{i} \in \tilde{I}$ with $i + \tilde{i} = k + \dim Y$, where $\pi_{E^i} : E^{(i, \tilde{i})} \rightarrow E^i$ and $\pi_{\tilde{E}^{\tilde{i}}} : E^{(i, \tilde{i})} \rightarrow \tilde{E}^{\tilde{i}}$ are the projections from $E^{(i, \tilde{i})} = \pi_{V^i}^*(E^i) \oplus \pi_{\tilde{V}^{\tilde{i}}}^*(\tilde{E}^{\tilde{i}}) \cong E^i \times_Y \tilde{E}^{\tilde{i}}$, and $\mu : P \times P \rightarrow P$ is as in (33). We shall show $\tilde{C}^k \times (\tilde{f}^k \circ \tilde{\pi}^k) : \tilde{E}^k \rightarrow P \times Y$ is *globally finite*.

Let $i \in I$ and $\tilde{i} \in \tilde{I}$ with $i + \tilde{i} = k + \dim Y$. Then $C^i : E^i \rightarrow P_{n^i} \subset P$ and $\tilde{C}^{\tilde{i}} : \tilde{E}^{\tilde{i}} \rightarrow P_{\tilde{n}^{\tilde{i}}} \subset P$ for some $n^i, \tilde{n}^{\tilde{i}} \gg 0$. Definition 3.4(vi) gives compact $T, \tilde{T} \subseteq Y$ with $f^i(V^i) \subseteq T$ and $\tilde{f}^{\tilde{i}}(\tilde{V}^{\tilde{i}}) \subseteq \tilde{T}$. The map $Y \rightarrow \mathbb{N}$ taking $y \mapsto |\text{Stab}(y)|$ is upper semicontinuous, so as $T \cap \tilde{T}$ is compact there exists $L \geq 1$ such that $|\text{Stab}(y)| \leq L$ for all $y \in T \cap \tilde{T}$. As $C^i \times (f^i \circ \pi^i) : E^i \rightarrow P \times Y$ and $\tilde{C}^{\tilde{i}} \times (\tilde{f}^{\tilde{i}} \circ \tilde{\pi}^{\tilde{i}}) : \tilde{E}^{\tilde{i}} \rightarrow P \times Y$ are globally finite, there exist $N^i, \tilde{N}^{\tilde{i}} \geq 0$ such that $|(C^i \times (f^i \circ \pi^i))^{-1}(p, y)| \leq N^i$ and $|(\tilde{C}^{\tilde{i}} \times (\tilde{f}^{\tilde{i}} \circ \tilde{\pi}^{\tilde{i}}))^{-1}(\tilde{p}, y)| \leq \tilde{N}^{\tilde{i}}$ for all $p, \tilde{p} \in P$ and $y \in Y$. For $p' \in P$ and $y \in Y$, consider the projection

$$\pi_{E^i} \times \pi_{\tilde{E}^{\tilde{i}}} : ((\tilde{C}^k \times (\tilde{f}^k \circ \tilde{\pi}^k))^{-1}(p', y)) \cap E^{(i, \tilde{i})} \longrightarrow \prod_{\substack{(p, \tilde{p}) \in P \times P : \\ \mu(p, \tilde{p}) = p'}} (C^i \times (f^i \circ \pi^i))^{-1}(p, y) \times (\tilde{C}^{\tilde{i}} \times (\tilde{f}^{\tilde{i}} \circ \tilde{\pi}^{\tilde{i}}))^{-1}(\tilde{p}, y). \quad (52)$$

Since C^i maps $E^i \rightarrow P_{n^i}$, $\tilde{C}^{\tilde{i}}$ maps $\tilde{E}^{\tilde{i}} \rightarrow P_{\tilde{n}^{\tilde{i}}}$, and μ maps $P_{n^i} \times P_{\tilde{n}^{\tilde{i}}} \rightarrow P_{n^i + \tilde{n}^{\tilde{i}}}$, the only nonempty terms in (52) occur when $p \in P_{n^i}, \tilde{p} \in P_{\tilde{n}^{\tilde{i}}}$ and $p' \in P_{n^i + \tilde{n}^{\tilde{i}}}$. From (33) we see that each $p' \in P_{n^i + \tilde{n}^{\tilde{i}}}$ may be written as $\mu(p, \tilde{p})$ for at most $\binom{n^i + \tilde{n}^{\tilde{i}}}{n^i}$ pairs $(p, \tilde{p})$. Thus on the second line of (52) there are at most $\binom{n^i + \tilde{n}^{\tilde{i}}}{n^i}$ nonempty sets, each of which has at most $N^i \tilde{N}^{\tilde{i}}$ elements, so the second line of (52) is a finite set of size at most $\binom{n^i + \tilde{n}^{\tilde{i}}}{n^i} N^i \tilde{N}^{\tilde{i}}$.

But $\tilde{E}^k \cap E^{(i, \tilde{i})}$ is an open subset of $E^i \times_Y \tilde{E}^{\tilde{i}}$, so the definition (12) of fibre products of orbifolds implies that if $e \in E^i$ and $\tilde{e} \in \tilde{E}^{\tilde{i}}$ with $f^i \circ \pi^i(e) = y =$

$\tilde{f}^{\tilde{i}} \circ \tilde{\pi}^{\tilde{i}}(\tilde{e})$ then the set of $\tilde{e} \in \tilde{E}^k \cap E^{(i,\tilde{i})}$ with $(\pi_{E^i} \times \pi_{\tilde{E}^{\tilde{i}}})(\tilde{e}) = (e, \tilde{e})$ is a subset of $(f^i \circ \pi^i)_*(\text{Stab}(e)) \setminus \text{Stab}(y) / (\tilde{f}^{\tilde{i}} \circ \tilde{\pi}^{\tilde{i}})_*(\text{Stab}(\tilde{e}))$, and so consists of at most L points, as $|\text{Stab}(y)| \leq L$. Hence each point on the second line of (52) pulls back to at most L points on the first line. Therefore for all $(p', y) \in P \times Y$ we have

$$|((\check{C}^k \times (\check{f}^k \circ \check{\pi}^k))^{-1}(p', y)) \cap E^{(i,\tilde{i})}| \leq L \binom{n^i + \tilde{n}^{\tilde{i}}}{n^i} N^i \tilde{N}^{\tilde{i}}.$$

Summing this over all $i, \tilde{i}$ implies that $\check{C}^k \times (\check{f}^k \circ \check{\pi}^k)$ is *globally finite*, with constant $\tilde{N}^k = \sum_{i,\tilde{i}: i+\tilde{i}=k+\dim Y} L \binom{n^i + \tilde{n}^{\tilde{i}}}{n^i} N^i \tilde{N}^{\tilde{i}}$. Hence $\check{\mathbf{C}} = (\check{\mathbf{I}}, \check{\boldsymbol{\eta}}, \check{C}^k : k \in \check{I})$ is *co-gauge-fixing data* for $(X \times_Y \tilde{X}, \boldsymbol{\pi}_Y)$, which we write as $\mathbf{C} \times_Y \tilde{\mathbf{C}}$.

Now suppose that $\mathbf{f} : X \rightarrow Y$ is strongly smooth and $\mathbf{G}$ is gauge-fixing data for $(X, \mathbf{f})$, and $\tilde{\mathbf{f}} : \tilde{X} \rightarrow Y$ is a strong submersion and $\tilde{\mathbf{C}}$ is co-gauge-fixing data for $(\tilde{X}, \tilde{\mathbf{f}})$. Then the construction above, replacing $C^i, \check{C}^k$ by $G^i, \check{G}^k$, yields *gauge-fixing data* $\check{\mathbf{G}} = \mathbf{G} \times_Y \tilde{\mathbf{C}}$ for $(X \times_Y \tilde{X}, \boldsymbol{\pi}_Y)$. The proof that the $\check{G}^k$ are globally finite is slightly different, but the definitions are unchanged.

Proposition 3.32. *Let Y be an orbifold without boundary, and for $a = 1, 2, 3$ let X_a be a compact Kuranishi space, $\mathbf{f}_a : X_a \rightarrow Y$ a strong submersion, and $\mathbf{C}_a$ co-gauge-fixing data for $(X_a, \mathbf{f}_a)$. Then:*

- (a) *The natural strong diffeomorphism $\mathbf{a} : X_1 \times_Y X_2 \rightarrow X_2 \times_Y X_1$ extends to an isomorphism in the sense of Definition 3.9*

$$(\mathbf{a}, \mathbf{b}) : (X_1 \times_Y X_2, \boldsymbol{\pi}_Y, \mathbf{C}_1 \times_Y \mathbf{C}_2) \longrightarrow (X_2 \times_Y X_1, \boldsymbol{\pi}_Y, \mathbf{C}_2 \times_Y \mathbf{C}_1). \quad (53)$$

- (b) *The natural strong diffeomorphism $\mathbf{a}' : \partial(X_1 \times_Y X_2) \rightarrow \partial X_1 \times_Y X_2 \amalg X_1 \times_Y \partial X_2$ extends to an isomorphism*

$$(\mathbf{a}', \mathbf{b}') : (\partial(X_1 \times_Y X_2), \boldsymbol{\pi}_Y, (\mathbf{C}_1 \times_Y \mathbf{C}_2)|_{\partial(X_1 \times_Y X_2)}) \longrightarrow \quad (54)$$

$$(\partial X_1 \times_Y X_2, \boldsymbol{\pi}_Y, (\mathbf{C}_1|_{\partial X_1}) \times_Y \mathbf{C}_2) \amalg (X_1 \times_Y \partial X_2, \boldsymbol{\pi}_Y, \mathbf{C}_1 \times_Y (\mathbf{C}_2|_{\partial X_2})).$$

- (c) *The natural strong diffeomorphism $\mathbf{a}'' : (X_1 \times_Y X_2) \times_Y X_3 \rightarrow X_1 \times_Y (X_2 \times_Y X_3)$ extends to an isomorphism*

$$(\mathbf{a}'', \mathbf{b}'') : ((X_1 \times_Y X_2) \times_Y X_3, \boldsymbol{\pi}_Y, (\mathbf{C}_1 \times_Y \mathbf{C}_2) \times_Y \mathbf{C}_3) \longrightarrow \quad (55)$$

$$(X_1 \times_Y (X_2 \times_Y X_3), \boldsymbol{\pi}_Y, \mathbf{C}_1 \times_Y (\mathbf{C}_2 \times_Y \mathbf{C}_3)).$$

- (d) *If instead $\mathbf{f}_1 : X_1 \rightarrow Y$ is strongly smooth and $\mathbf{G}_1$ is gauge-fixing data for $(X_1, \mathbf{f}_1)$, then in (b),(c) above $\mathbf{a}', \mathbf{a}''$ extend to isomorphisms*

$$(\mathbf{a}', \mathbf{b}') : (\partial(X_1 \times_Y X_2), \boldsymbol{\pi}_Y, (\mathbf{G}_1 \times_Y \mathbf{C}_2)|_{\partial(X_1 \times_Y X_2)}) \longrightarrow \quad (56)$$

$$(\partial X_1 \times_Y X_2, \boldsymbol{\pi}_Y, (\mathbf{G}_1|_{\partial X_1}) \times_Y \mathbf{C}_2) \amalg (X_1 \times_Y \partial X_2, \boldsymbol{\pi}_Y, \mathbf{G}_1 \times_Y (\mathbf{C}_2|_{\partial X_2})),$$

$$(\mathbf{a}'', \mathbf{b}'') : ((X_1 \times_Y X_2) \times_Y X_3, \boldsymbol{\pi}_Y, (\mathbf{G}_1 \times_Y \mathbf{C}_2) \times_Y \mathbf{C}_3) \longrightarrow \quad (57)$$

$$(X_1 \times_Y (X_2 \times_Y X_3), \boldsymbol{\pi}_Y, \mathbf{G}_1 \times_Y (\mathbf{C}_2 \times_Y \mathbf{C}_3)),$$

Proof. By Theorem 3.30, the strong diffeomorphism $\mathbf{a}$ identifies the really good coordinate systems $(\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y (\mathbf{I}_2, \boldsymbol{\eta}_2)$ and $(\mathbf{I}_2, \boldsymbol{\eta}_2) \times_Y (\mathbf{I}_1, \boldsymbol{\eta}_1)$. So applying Algorithm 3.16, $\mathbf{a}$ also identifies $((\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y (\mathbf{I}_2, \boldsymbol{\eta}_2))^\sim$ and $((\mathbf{I}_2, \boldsymbol{\eta}_2) \times_Y (\mathbf{I}_1, \boldsymbol{\eta}_1))^\sim$. That is, there is a natural isomorphism $\mathbf{b} : ((\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y (\mathbf{I}_2, \boldsymbol{\eta}_2))^\sim \rightarrow ((\mathbf{I}_2, \boldsymbol{\eta}_2) \times_Y (\mathbf{I}_1, \boldsymbol{\eta}_1))^\sim$ compatible with $\mathbf{a}$. Write the really good coordinate system in $((\mathbf{I}_1, \boldsymbol{\eta}_1) \times_Y (\mathbf{I}_2, \boldsymbol{\eta}_2))^\sim$ as $(\check{I}_{12}, (\check{V}_{12}^k, \dots, \check{\psi}_{12}^k) : k \in \check{I}_{12}, \dots)$ with functions $\check{C}_{12}^k : \check{E}_{12}^k \rightarrow P$, and similarly for $((\mathbf{I}_2, \boldsymbol{\eta}_2) \times_Y (\mathbf{I}_1, \boldsymbol{\eta}_1))^\sim$ with $\check{I}_{21}, \dots$.

To show that $(\mathbf{a}, \mathbf{b})$ is an isomorphism in (53), it only remains to check that $\check{C}_{21}^k \circ \hat{b}^k \equiv \check{C}_{12}^k$ for all $k \in \check{I}_{12} = \check{I}_{21}$. But this is obvious since $C_{12}^k|_{\check{E}^k \cap E_{12}^{(i_1, i_2)}} \equiv \mu \circ ((C_1^{i_1} \circ \pi_{E_1^{i_1}}) \times (C_2^{i_2} \circ \pi_{E_2^{i_2}}))$ for $i_1 \in I_1, i_2 \in I_2$ with $i_1 + i_2 = k + \dim Y$, and $C_{21}^k|_{\check{E}^k \cap E_{21}^{(i_2, i_1)}} \equiv \mu \circ ((C_2^{i_2} \circ \pi_{E_2^{i_2}}) \times (C_1^{i_1} \circ \pi_{E_1^{i_1}}))$, and we have $(\pi_{E_1^{i_1}} \times \pi_{E_2^{i_2}}) \circ \hat{b}^k \equiv \pi_{E_1^{i_1}} \times \pi_{E_2^{i_2}}$ as maps $\check{E}^k \cap E_{12}^{(i_1, i_2)} \rightarrow E_1^{i_1} \times E_2^{i_2}$, and μ in (33) is *commutative*. This proves part (a).

For (b)–(d) we follow a similar method, using the fact that μ in (33) is *associative* for (55) and (57), but there is one complication. The argument above proves that the excellent coordinate systems $((\mathbf{I}_1 \times_Y \mathbf{I}_2) \times_Y \mathbf{I}_3)^\sim$ and $(\mathbf{I}_1 \times_Y (\mathbf{I}_2 \times_Y \mathbf{I}_3))^\sim$ are isomorphic under $\mathbf{a}''$. But we need to know that $((\mathbf{I}_1 \times_Y \mathbf{I}_2) \times_Y \mathbf{I}_3)^\sim$ and $(\mathbf{I}_1 \times_Y (\mathbf{I}_2 \times_Y \mathbf{I}_3))^\sim$ are isomorphic. One can show that $((\mathbf{I}_1 \times_Y \mathbf{I}_2) \times_Y \mathbf{I}_3)^\sim = ((\mathbf{I}_1 \times_Y \mathbf{I}_2) \times_Y \mathbf{I}_3)^\sim$ and $(\mathbf{I}_1 \times_Y (\mathbf{I}_2 \times_Y \mathbf{I}_3))^\sim = (\mathbf{I}_1 \times_Y (\mathbf{I}_2 \times_Y \mathbf{I}_3))^\sim$, that is, applying Algorithm 3.16 twice gives the same result as applying it once. Equations (55) and (57) follow. $\square$

Products are also functorial for proper pullbacks, and pushforwards. The proof is similar to that of Proposition 3.32, so we leave it as an exercise.

Proposition 3.33. *Suppose $h : Y \rightarrow Z$ is a smooth, proper map of orbifolds.*

- (a) *Let X_i be a compact Kuranishi space, $\mathbf{f}_i : X_i \rightarrow Z$ a strong submersion, and $\mathbf{C}_i$ co-gauge-fixing data for $(X_i, \mathbf{f}_i)$ for $i = 1, 2$. Then the natural strong diffeomorphism $\mathbf{a} : Y \times_Z (X_1 \times_Z X_2) \rightarrow (Y \times_Z X_1) \times_Y (Y \times_Z X_2)$ extends to an isomorphism*

$$(\mathbf{a}, \mathbf{b}) : (Y \times_Z (X_1 \times_Z X_2), \pi_Y, h^*(\mathbf{C}_1 \times_Z \mathbf{C}_2)) \longrightarrow ((Y \times_Z X_1) \times_Y (Y \times_Z X_2), \pi_Y, h^*(\mathbf{C}_1) \times_Y h^*(\mathbf{C}_2)).$$

- (b) *Let X_1, X_2 be compact Kuranishi spaces, $\mathbf{f}_1 : X_1 \rightarrow Y$ strongly smooth, $\mathbf{G}_1$ gauge-fixing data for $(X_1, \mathbf{f}_1)$, $\mathbf{f}_2 : X_2 \rightarrow Z$ a strong submersion, and $\mathbf{C}_2$ co-gauge-fixing data for $(X_2, \mathbf{f}_2)$. Then the natural strong diffeomorphism $\mathbf{a}' : X_1 \times_{h \circ \mathbf{f}_1, Z, \mathbf{f}_2} X_2 \rightarrow X_1 \times_{\mathbf{f}_1, Y, \pi_Y} (Y \times_{h, Z, \mathbf{f}_2} X_2)$ extends to an isomorphism*

$$(\mathbf{a}', \mathbf{b}') : (X_1 \times_Z X_2, \pi_Z, h_*(\mathbf{G}_1) \times_Z \mathbf{C}_2) \longrightarrow (X_1 \times_Y (Y \times_Z X_2), h \circ \pi_Y, h_*(\mathbf{G}_1 \times_Y h^*(\mathbf{C}_2))).$$

Remark 3.34. (a) The construction of fibre products $\mathbf{C} \times_Y \tilde{\mathbf{C}}$ in Definition 3.31 can actually be split into two stages. Given compact Kuranishi spaces $X, \tilde{X}$, orbifolds $Y, \tilde{Y}$, strongly smooth maps $\mathbf{f} : X \rightarrow Y$ and $\tilde{\mathbf{f}} : \tilde{X} \rightarrow \tilde{Y}$, and co-gauge-fixing data $\mathbf{C}, \tilde{\mathbf{C}}$ for $(X, \mathbf{f})$ and $(\tilde{X}, \tilde{\mathbf{f}})$, we can define *product co-gauge-fixing data* $\mathbf{C} \times \tilde{\mathbf{C}}$ for $(X \times \tilde{X}, \mathbf{f} \times \tilde{\mathbf{f}})$, that is, for the strongly smooth map $\mathbf{f} \times \tilde{\mathbf{f}} : X \times \tilde{X} \rightarrow Y \times \tilde{Y}$. To do this we simply replace fibre products over Y by ordinary products (fibre products over a point) throughout the construction of $\mathbf{C} \times_Y \tilde{\mathbf{C}}$. Note that the construction of $\mathbf{C} \times \tilde{\mathbf{C}}$ is still extremely complex. Products of gauge-fixing data $\mathbf{C} \times \tilde{\mathbf{C}}$ are commutative and associative, by an easy modification of Proposition 3.32.

Then to construct $\mathbf{C} \times_Y \tilde{\mathbf{C}}$, take $\tilde{Y} = Y$, and define $\Delta : Y \rightarrow Y \times Y$ to be the diagonal map, $\Delta : y \mapsto (y, y)$. Then $\mathbf{C} \times_Y \tilde{\mathbf{C}} = \Delta^*(\mathbf{C} \times \tilde{\mathbf{C}})$, that is, $\mathbf{C} \times_Y \tilde{\mathbf{C}}$ is the pullback of $\mathbf{C} \times \tilde{\mathbf{C}}$ under Δ , as in §3.7.

If $\mathbf{f} : X \rightarrow Y$, $\tilde{\mathbf{f}} : \tilde{X} \rightarrow \tilde{Y}$ are strongly smooth and $\mathbf{G}, \tilde{\mathbf{G}}$ are gauge-fixing data for $(X, \mathbf{f})$, $(\tilde{X}, \tilde{\mathbf{f}})$, then the definition of $\mathbf{C} \times \tilde{\mathbf{C}}$ generalizes immediately to give *product gauge-fixing data* $\mathbf{G} \times \tilde{\mathbf{G}}$ for $(X \times \tilde{X}, \mathbf{f} \times \tilde{\mathbf{f}})$, where $\mathbf{f} \times \tilde{\mathbf{f}} : X \times \tilde{X} \rightarrow Y \times \tilde{Y}$ is strongly smooth.

(b) We can use products $\mathbf{C} \times \tilde{\mathbf{C}}$ in (a) and pullbacks to define generalizations of fibre products of co-gauge-fixing data to more than one orbifold Y , which we leave as an exercise. For example, if X_1, X_2 are compact Kuranishi spaces, Y_3, Y_4, Y_5 are orbifolds, $\mathbf{f}_{13} \times \mathbf{f}_{14} : X_1 \rightarrow Y_3 \times Y_4$, $\mathbf{f}_{24} \times \mathbf{f}_{25} : X_2 \rightarrow Y_4 \times Y_5$ strong submersions, and $\mathbf{C}_1, \mathbf{C}_2$ are co-gauge-fixing data for $(X_1, \mathbf{f}_{13} \times \mathbf{f}_{14})$ and $(X_2, \mathbf{f}_{24} \times \mathbf{f}_{25})$, then we can define co-gauge-fixing data $\mathbf{C}_1 \times_{Y_4} \mathbf{C}_2$ for $(X_1 \times_{\mathbf{f}_{14}, Y_4, \mathbf{f}_{24}} X_2, (\mathbf{f}_{13} \circ \pi_{X_1}) \times \pi_{Y_4} \times (\mathbf{f}_{25} \circ \pi_{X_2}))$ over $Y_3 \times Y_4 \times Y_5$.

We will generalize these ideas on products of (co-)gauge-fixing data in our discussion of Kuranishi (co)homology as a *bivariant theory* in §4.8.

3.9 Effective gauge-fixing data and co-gauge-fixing data

For reasons that will become clearer in Chapter 4, because we allow triples $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \mathbf{C})$ to have finite automorphism groups, as in §3.3, we can only use them to define Kuranishi (co)homology theories $KH_*, KH^*(Y; R)$ in which the coefficient ring R is a $\mathbb{Q}$ -algebra, with the homology theory isomorphic to singular homology $KH_*(Y; R) \cong H_*^{\text{si}}(Y; R)$. Essentially this is because R must contain $|\text{Aut}(X, \mathbf{f}, \mathbf{G})|^{-1}$, to enable us to average over $\text{Aut}(X, \mathbf{f}, \mathbf{G})$.

We will also define *effective Kuranishi (co)homology* $KH_*^{\text{ef}}, KH_{\text{ec}}^*(Y; R)$, which work for R any commutative ring, with $KH_*^{\text{ef}}(Y; R) \cong H_*^{\text{si}}(Y; R)$ and (for Y a manifold) $KH_{\text{ec}}^*(Y; R) \cong H_{\text{cs}}^*(Y; R)$. To make this work, we need to ensure that (connected) triples $(X, \mathbf{f}, \underline{\mathbf{G}})$ or $(X, \mathbf{f}, \underline{\mathbf{C}})$ have only *trivial automorphism groups* $\{1\}$. Our goal now is to define notions of *effective* (co-)gauge-fixing data $\underline{\mathbf{G}}, \underline{\mathbf{C}}$, and partially extend §3.2–§3.8 to them.

The basic idea is that we shall require the maps $G^i, C^i \times (f^i \circ \pi^i)$ of Definitions 3.8 and 3.9 to be *injective on* $(E^i)^\circ$ as well as globally finite, which easily implies that $(X, \mathbf{f}, \underline{\mathbf{G}})$ and $(X, \mathbf{f}, \underline{\mathbf{C}})$ have only trivial automorphisms. However, there are significant problems in carrying this out. Here are some of them:

- (i) Let $(X, \mathbf{f}, \underline{\mathbf{G}})$ be a triple, with $\underline{\mathbf{G}}$ effective gauge-fixing data. Gauge-fixing data must restrict to boundaries, so we can consider $(\partial^2 X, \mathbf{f}|_{\partial^2 X}, \underline{\mathbf{G}}|_{\partial^2 X})$. Let $\sigma : \partial^2 X \rightarrow \partial^2 X$ be the natural involution of Definition 2.27. Then $\sigma \in \text{Aut}(\partial^2 X, \mathbf{f}|_{\partial^2 X}, \underline{\mathbf{G}}|_{\partial^2 X})$. This is incompatible with $\underline{\mathbf{G}}|_{\partial^2 X}$ being *effective* gauge-fixing data. Thus, restricting effective (co-)gauge-fixing data to boundaries will not yield effective (co-)gauge-fixing data.
- (ii) Even if we did define a homology theory $KH_*^{\text{ef}}(Y; R)$ using some notion of effective gauge-fixing data, we could not prove $KH_*^{\text{ef}}(Y; R) \cong H_*^{\text{si}}(Y; R)$ for rings $R \not\supseteq \mathbb{Q}$ unless we make some restriction on the *stabilizer groups* of Kuranishi spaces X in triples $(X, \mathbf{f}, \underline{\mathbf{G}})$, because otherwise we cannot perturb X to a manifold or orbifold, and triangulate it by simplices.
- (iii) If $f^i \circ \pi^i : E^i \rightarrow Y$, $\tilde{f}^i \circ \tilde{\pi}^i : \tilde{E}^i \rightarrow Y$ are submersions of orbifolds, the definition (12) of $E^i \times_Y \tilde{E}^i$ involving $f_*(\text{Stab}(p)) \backslash \text{Stab}(f(p)) / f'_*(\text{Stab}(p'))$ means that the projection $\pi_{E^i} \times \pi_{\tilde{E}^i} : E^i \times_Y \tilde{E}^i \rightarrow E^i \times \tilde{E}^i$ need not be injective, only finite, if Y is an orbifold, not a manifold. This can cause $\tilde{C}^k \times (\tilde{f}^k \circ \tilde{\pi}^k) : (\tilde{E}^k)^\circ \rightarrow P \times Y$ in fibre products of effective co-gauge-fixing data $\underline{\mathbf{C}}, \tilde{\underline{\mathbf{C}}}$ in Definition 3.31 not to be injective, so that $\underline{\mathbf{C}} \times_Y \tilde{\underline{\mathbf{C}}}$ is not effective co-gauge-fixing data.
- (iv) Here is a second problem with taking fibre products of effective co-gauge-fixing data. Let $(X, \mathbf{f}, \underline{\mathbf{C}})$ be a triple, with $\underline{\mathbf{C}}$ effective co-gauge-fixing data and $\mathbf{f} : X \rightarrow Y$ a strong submersion. Consider the triple $(X \times_Y X, \pi_Y, \underline{\mathbf{C}} \times_Y \underline{\mathbf{C}})$, with $\underline{\mathbf{C}} \times_Y \underline{\mathbf{C}}$ as in §3.8. This has an obvious $\mathbb{Z}_2$ -symmetry exchanging the two factors of $X, \underline{\mathbf{C}}$. So $\text{Aut}(X \times_Y X, \pi_Y, \underline{\mathbf{C}} \times_Y \underline{\mathbf{C}})$ may be nontrivial, and $\underline{\mathbf{C}} \times_Y \underline{\mathbf{C}}$ is not effective.

To overcome each of these problems, we have to sacrifice one of the good properties of our theory.

Problems (i)–(iii) force us to restrict the Kuranishi spaces X and morphisms $\mathbf{f}$ allowed in triples $(X, \mathbf{f}, \underline{\mathbf{G}})$ or $(X, \mathbf{f}, \underline{\mathbf{C}})$ in the theory. To solve (i) we require the injectivity conditions to hold only for each *connected component* of $(\partial^k X, \mathbf{f}|_{\partial^k X}, \underline{\mathbf{G}}|_{\partial^k X})$. Then we can ensure that $\text{Aut}(X, \mathbf{f}, \underline{\mathbf{G}}) = \{1\}$ for *connected* triples $(X, \mathbf{f}, \underline{\mathbf{G}})$, which is enough to make the theory work. If $\sigma : \partial^2 X \rightarrow \partial^2 X$ acts freely on the connected components of $(\partial^2 X, \mathbf{f}|_{\partial^2 X}, \underline{\mathbf{G}}|_{\partial^2 X})$, there is then no contradiction in (i). This constrains the codimension k corners $\partial^k X$ for $k = 2, 3, \dots$. To deal with (ii) we restrict to *effective* Kuranishi spaces X , defined below, which can always be perturbed to effective orbifolds.

For (iii) we require the smooth maps $f^i \circ \pi^i : E^i \rightarrow Y$ in triples $(X, \mathbf{f}, \underline{\mathbf{C}})$ to be surjective on stabilizer groups, which makes $\pi_{E^i} \times \pi_{\tilde{E}^i} : E^i \times_Y \tilde{E}^i \rightarrow E^i \times \tilde{E}^i$ injective. Thus, Property 3.1(a) will not hold: only some, not all, pairs $(X, \mathbf{f})$ admit effective (co-)gauge-fixing data. This will limit the applications of the theory — for instance, in §6.2 we cannot define Gromov–Witten type invariants in $KH_*^{\text{ef}}(Y; \mathbb{Z})$ by taking X to be a moduli space $\bar{\mathcal{M}}_{g,m}(J, \beta)$ of J -holomorphic curves in a symplectic manifold (M, ω) , since such $\bar{\mathcal{M}}_{g,m}(J, \beta)$ may not be effective.

Problem (iv) means that products $\underline{\mathbf{C}}_1 \times_Y \underline{\mathbf{C}}_2$ of effective co-gauge-fixing data

cannot be defined to be commutative, and therefore the cup product $\cup$ on effective Kuranishi cochains $KC_{\text{ec}}^*(Y; R)$ will not be supercommutative. Thus, symmetry in Property 3.1(h) fails. We will deal with this by replacing $P = \coprod_{k=0}^{\infty} \mathbb{R}^k / S_k$ in Definition 3.8 by $\underline{P} = \coprod_{k=0}^{\infty} \mathbb{R}^k$, and taking effective (co-)gauge-fixing data $\underline{G}, \underline{C}$ to include maps $\underline{G}^i, \underline{C}^i : E^i \rightarrow \underline{P}$. The product $\underline{\mu} : \underline{P} \times \underline{P} \rightarrow \underline{P}$ corresponding to μ in (33) is associative but not commutative, and $\underline{\mu} : \mathbb{R}^k \times \mathbb{R}^l \rightarrow \mathbb{R}^{k+l}$ is injective, which will be enough to preserve the injectivity conditions for effective co-gauge-fixing data. In Remark 4.33 we explain a reason from algebraic topology why $\cup$ cannot be supercommutative for cohomology theories over $R = \mathbb{Z}$.

We define *effective* Kuranishi neighbourhoods and Kuranishi spaces.

Definition 3.35. An orbifold Y is *effective* if for all orbifold charts (U, Γ, ϕ) on Y in Definition 2.9 with $U \subset \mathbb{R}^n$, the representation of Γ on $\mathbb{R}^n$ is *effective*. Here a representation is *effective* if every non-identity element of Γ acts nontrivially on $\mathbb{R}^n$. Equivalently, an orbifold is effective if generic $y \in Y$ have $\text{Stab}_Y(y) = \{1\}$.

A Kuranishi neighbourhood (V, E, s, ψ) on X is called *effective* if V is an effective orbifold, and for each $v \in V$, the stabilizer group $\text{Stab}_V(v)$ acts trivially on the fibre $E|_v$ of E over v . That is, in the notation of Definition 2.13, if (U, Γ, ϕ) is an orbifold chart on V and $(E_U, \Gamma, \hat{\phi})$ the associated orbifold chart on E , then for all $u \in U$, the action of $\{\gamma \in \Gamma : \gamma \cdot u = u\}$ on $E_U|_u$ is trivial.

If $(V_p, \dots, \psi_p), (V_q, \dots, \psi_q)$ are Kuranishi neighbourhoods on X , $(\phi_{pq}, \hat{\phi}_{pq}) : (V_q, \dots, \psi_q) \rightarrow (V_p, \dots, \psi_p)$ is a coordinate change, and $(V_p, \dots, \psi_p)$ is effective, it is easy to show that $(V_q, \dots, \psi_q)$ is also effective.

A Kuranishi space X is called *effective* if for all $p \in X$, all sufficiently small Kuranishi neighbourhoods $(V_p, \dots, \psi_p)$ in the germ at p are effective.

Here are the reasons for this definition:

- For proving isomorphism of effective Kuranishi homology $KH_*^{\text{ef}}(Y; R)$ with singular homology $H_*^{\text{si}}(Y; R)$, chains involving *effective* orbifolds are as good as chains involving manifolds. An effective orbifold can be triangulated by simplices to define a singular chain.
- However, if Y is a *non-effective* orbifold, then for generic $y \in Y$, $\text{Stab}(y)$ is some nontrivial finite group Γ . When Y is triangulated by simplices to define a singular chain, they should be weighted by $|\Gamma|^{-1}$. This only makes sense if $|\Gamma|^{-1}$ lies in the coefficient ring R , that is, if R is a $\mathbb{Q}$ -algebra.
- If (V, E, s, ψ) is an *effective* Kuranishi neighbourhood, and $\tilde{s}$ is a small generic perturbation of s , then $\tilde{s}$ is transverse and $\tilde{s}^{-1}(0)$ is an effective orbifold. This is because the obstruction to deforming s to transverse is nontrivial actions of the stabilizer groups $\text{Stab}(v)$ on the fibres $E|_v$ of E , and $\tilde{s}^{-1}(0)$ is effective as V is effective.
- If X is a compact, *effective* Kuranishi space, then by choosing a finite cover of X by effective Kuranishi neighbourhoods (V^i, E^i, s^i, ψ^i) , and perturbing the s^i , we can deform X to a compact, effective orbifold $\tilde{X}$, which we can then triangulate by simplices to get a singular chain.

We can now define *effective* gauge-fixing data.

Definition 3.36. Modifying P, μ in Definition 3.8, define $\underline{P} = \coprod_{k=0}^{\infty} \mathbb{R}^k$ and $\underline{\mu} : \underline{P} \times \underline{P} \rightarrow \underline{P}$ by

$$\underline{\mu}((x_1, \dots, x_k), (y_1, \dots, y_l)) = (x_1, \dots, x_k, y_1, \dots, y_l). \quad (58)$$

Then $\underline{\mu}$ is *associative*, with *identity* $(\emptyset)$, but is *not* commutative. The obvious projection $\Pi : \underline{P} \rightarrow P$ mapping $\Pi : (x_1, \dots, x_k) \mapsto S_k(x_1, \dots, x_k)$ satisfies $\mu(\Pi(p), \Pi(q)) = \Pi(\underline{\mu}(p, q))$ for $p, q \in \underline{P}$.

Let X be a compact, effective Kuranishi space, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ strongly smooth. A set of *effective gauge-fixing data* for $(X, \mathbf{f})$ is $\underline{\mathbf{G}} = (\mathbf{I}, \boldsymbol{\eta}, \underline{\mathbf{G}}^i : i \in I)$, where $(\mathbf{I}, \boldsymbol{\eta})$ is an excellent coordinate system for $(X, \mathbf{f})$ with $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i) : i \in I, \dots)$, and $\underline{\mathbf{G}}^i : E^i \rightarrow \underline{P}$ for $i \in I$, satisfying:

- (a) (V^i, E^i, s^i, ψ^i) is an effective Kuranishi neighbourhood for all $i \in I$; and
- (b) Write $\partial^l X = X_1^l \amalg \dots \amalg X_{n^l}^l$ for the splitting of $(\partial^l X, \mathbf{f}|_{\partial^l X}, \mathbf{I}|_{\partial^l X}, \boldsymbol{\eta}|_{\partial^l X})$ into connected components in Lemma 3.7 for $l \geq 0$, and $(\mathbf{I}_a^l, \boldsymbol{\eta}_a^l) = (\mathbf{I}|_{X_a^l}, \boldsymbol{\eta}|_{X_a^l})$, with $\mathbf{I}_a^l = (I_a^l, (V_a^{l,i}, \dots, \psi_a^{l,i}) : i \in I_a^l, \dots)$. Then $(E_a^{l,i})^\circ \subseteq E_a^{l,i} \subseteq \partial^l E^{i+l}$, where $(E_a^{l,i})^\circ$ is the *interior* of $E_a^{l,i}$, so we can consider $\underline{\mathbf{G}}^{i+l}|_{(E_a^{l,i})^\circ} : (E_a^{l,i})^\circ \rightarrow \underline{P}$. We require that $\underline{\mathbf{G}}^{i+l}|_{(E_a^{l,i})^\circ}$ should map $(E_a^{l,i})^\circ$ to $\mathbb{R}^{k_a^l} \subset \underline{P}$ for some $k_a^l \geq 0$ which depends on l, a but *not* on $i \in I_a^l$, for all $l \geq 0$, $a = 1, \dots, n^l$ and $i \in I_a^l$. We also require that $\coprod_{i \in I_a^l} \underline{\mathbf{G}}^{i+l}|_{(E_a^{l,i})^\circ} : \coprod_{i \in I_a^l} (E_a^{l,i})^\circ \rightarrow \mathbb{R}^{k_a^l} \subset \underline{P}$ should be *injective* for all $l \geq 0$ and $a = 1, \dots, n^l$.

Here (a) implies X is effective, as for sufficiently small $(V_p, \dots, \psi_p)$ in the germ at $p \in X$ we have $(\phi_p^i, \hat{\phi}_p^i) : (V_p, \dots, \psi_p) \rightarrow (V^i, \dots, \psi^i)$ for some $i \in I$, and $(V^i, \dots, \psi^i)$ effective implies $(V_p, \dots, \psi_p)$ effective, as in Definition 3.35.

Define *triples* $(X, \mathbf{f}, \underline{\mathbf{G}})$, *splittings* $X = X_1 \amalg \dots \amalg X_n$ of triples $(X, \mathbf{f}, \underline{\mathbf{G}})$, *connected triples* $(X, \mathbf{f}, \underline{\mathbf{G}})$, *isomorphisms* $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \underline{\mathbf{G}}) \rightarrow (\tilde{X}, \tilde{\mathbf{f}}, \tilde{\underline{\mathbf{G}}})$, and *automorphism groups* $\text{Aut}(X, \mathbf{f}, \underline{\mathbf{G}})$ as in Definition 3.8.

If $\underline{\mathbf{G}}$ is effective gauge-fixing data for $(X, \mathbf{f})$, we define the *restriction* $\underline{\mathbf{G}}|_{\partial X}$ as in Definition 3.9. Since the conditions in (a),(b) above are stable under taking boundaries, $V^i, E^i \mapsto \partial V^i, \partial E^i$, $\underline{\mathbf{G}}|_{\partial X}$ is effective gauge-fixing data for $(\partial X, \mathbf{f}|_{\partial X})$.

Let $\underline{\mathbf{G}} = (\mathbf{I}, \boldsymbol{\eta}, \underline{\mathbf{G}}^i : i \in I)$ be effective gauge-fixing data for $(X, \mathbf{f})$. We shall show that $\Pi \circ \underline{\mathbf{G}}^i : E^i \rightarrow P$ is a *globally finite* map, where $\Pi : \underline{P} \rightarrow P$ is as above. To see this, note that $E^i = \bigcup_{l \geq 0, a=1, \dots, n^l : E_a^{l,i-l} \neq \emptyset} \iota((E_a^{l,i-l})^\circ)$, where $\iota : \partial^l E_a^i \rightarrow E_a^i$ is the natural map, and $\underline{\mathbf{G}}^i$ maps $(E_a^{l,i-l})^\circ \rightarrow \mathbb{R}^{k_a^l} \subset \underline{P}$, and $\Pi : \mathbb{R}^{k_a^l} \rightarrow \mathbb{R}^{k_a^l}/S_{k_a^l}$ pulls back one point to at most $k_a^l!$ points. Hence if $p \in P$ then $|\Pi \circ \underline{\mathbf{G}}^i|^{-1}(p)|$ is at most N points, where $N = \sum_{l \geq 0, a=1, \dots, n^l : E_a^{l,i-l} \neq \emptyset} k_a^l!$. As this is a finite sum, $\Pi \circ \underline{\mathbf{G}}^i$ is globally finite, and so $\Pi(\underline{\mathbf{G}}) = \mathbf{G} = (\mathbf{I}, \boldsymbol{\eta}, \Pi \circ \underline{\mathbf{G}}^i : i \in I)$ is *gauge-fixing data* for $(X, \mathbf{f})$. This gives a natural map from effective gauge-fixing data to gauge-fixing data.

Here are analogues of Definitions 3.35 and 3.36 for co-gauge-fixing data.

Definition 3.37. Let X be a topological space, (V, E, s, ψ) a Kuranishi neighbourhood on X , Y an orbifold, and $f : V \rightarrow Y$ a submersion. We call the pair $(V, E, s, \psi), f$ *coeffective* if for all v in V with $y = f(v)$ in Y , the morphism $f_* : \text{Stab}_V(v) \rightarrow \text{Stab}_Y(y)$ is surjective, and the subgroup $\text{Ker } f_* \subseteq \text{Stab}_V(v)$ acts effectively on $T_v V$, and $\text{Stab}_V(v)$ acts trivially on the fibre $E|_v$ of E at v .

If $(V, E, s, \psi), f$ is coeffective and $e \in E$ with $\pi(e) = v$ in V and $f(v) = y$ in Y then $\pi_* : \text{Stab}_E(e) \rightarrow \text{Stab}_Y(y)$ is an isomorphism as $\text{Stab}_V(v)$ acts trivially on $E|_v$, so $(f \circ \pi)_* = f_* \circ \pi_* : \text{Stab}_E(e) \rightarrow \text{Stab}_Y(y)$ is surjective. This will help us deal with problem (iii) above.

Suppose Y is an *effective* orbifold. Then $(V, E, s, \psi), f$ is coeffective if and only if (V, E, s, ψ) is effective and $f : V \rightarrow Y$ is surjective on stabilizer groups. The only nontrivial thing to prove is that $(V, E, s, \psi), f$ coeffective implies V is effective. If v is generic in V then $y = f(v)$ is generic in Y as f is a submersion, so $\text{Stab}_Y(y) = \{1\}$ as Y is effective, giving $\text{Ker } f_* = \text{Stab}_V(v)$. Thus $\text{Stab}_V(v)$ acts effectively on $T_v V$, which implies that generic v' close to v in V have $\text{Stab}_V(v') = \{1\}$. Hence generic points in V have trivial stabilizers, and V is effective. When Y is a *manifold*, $f : V \rightarrow Y$ is trivially surjective on stabilizer groups, so $(V, E, s, \psi), f$ is coeffective if and only if (V, E, s, ψ) is effective.

If $(V_p, \dots, \psi_p), (V_q, \dots, \psi_q)$ are Kuranishi neighbourhoods on X , $(\phi_{pq}, \hat{\phi}_{pq}) : (V_q, \dots, \psi_q) \rightarrow (V_p, \dots, \psi_p)$ is a coordinate change, $f_p : V_p \rightarrow Y$ and $f_q : V_q \rightarrow Y$ are submersions with $f_q \equiv f_p \circ \phi_{pq}$ and $(V_p, \dots, \psi_p), f_p$ is coeffective, it is easy to show that $(V_q, \dots, \psi_q), f_q$ is also coeffective.

Let X be a Kuranishi space, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ a strong submersion. We call $(X, \mathbf{f})$ *coeffective* if for all $p \in X$ and all sufficiently small Kuranishi neighbourhoods $(V_p, \dots, \psi_p)$ in the germ at p with $f_p : V_p \rightarrow Y$ representing $\mathbf{f}$, the pair $(V_p, \dots, \psi_p), f_p$ is coeffective.

Definition 3.38. Now let X be a compact Kuranishi space, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ a strong submersion with $(X, \mathbf{f})$ coeffective. A set of *effective co-gauge-fixing data* for $(X, \mathbf{f})$ is $\underline{\mathbf{C}} = (\mathbf{I}, \boldsymbol{\eta}, \underline{\mathbf{C}}^i : i \in I)$, where $(\mathbf{I}, \boldsymbol{\eta})$ is an excellent coordinate system for $(X, \mathbf{f})$ with $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i), f^i : i \in I, \dots)$, and $\underline{\mathbf{C}}^i : E^i \rightarrow \underline{P}$ are maps for $i \in I$, satisfying

- (a) the pair $(V^i, E^i, s^i, \psi^i), f^i$ is coeffective for all $i \in I$; and
- (b) Write $\partial^l X = X_1^l \amalg \dots \amalg X_{n^l}^l$ for the splitting of $(\partial^l X, \mathbf{f}|_{\partial^l X}, \mathbf{I}|_{\partial^l X}, \boldsymbol{\eta}|_{\partial^l X})$ into connected components in Lemma 3.7 for $l \geq 0$, and $(\mathbf{I}_a^l, \boldsymbol{\eta}_a^l) = (\mathbf{I}|_{X_a^l}, \boldsymbol{\eta}|_{X_a^l})$, with $\mathbf{I}_a^l = (I_a^l, (V_a^{l,i}, \dots, \psi_a^{l,i}) : i \in I_a^l, \dots)$. Then $(E_a^{l,i})^\circ \subseteq E_a^{l,i} \subseteq \partial^l E^{i+l}$, so we can consider $\underline{\mathbf{C}}^{i+l}|_{(E_a^{l,i})^\circ} : (E_a^{l,i})^\circ \rightarrow \underline{P}$. We require that $\underline{\mathbf{C}}^{i+l}|_{(E_a^{l,i})^\circ}$ should map $(E_a^{l,i})^\circ$ to $\mathbb{R}^{k_a^l} \subset \underline{P}$ for some $k_a^l \geq 0$ which depends on l, a but *not* on $i \in I_a^l$, for all $l \geq 0, a = 1, \dots, n^l$ and $i \in I_a^l$. We also require that $\coprod_{i \in I_a^l} (\underline{\mathbf{C}}^{i+l} \times (f^{i+l} \circ \pi^{i+l}))|_{(E_a^{l,i})^\circ} : \coprod_{i \in I_a^l} (E_a^{l,i})^\circ \rightarrow \underline{P} \times Y$ should be *injective* for all $l \geq 0$ and $a = 1, \dots, n^l$.

Here (a) implies $(X, \mathbf{f})$ is coeffective, as for sufficiently small $(V_p, \dots, \psi_p), f_p$ in the germ at $p \in X$ we have $(\phi_p^i, \hat{\phi}_p^i) : (V_p, \dots, \psi_p) \rightarrow (V^i, \dots, \psi^i)$ for some $i \in I$ with $f_p \equiv f^i \circ \phi_p^i$, and $(V^i, \dots, \psi^i), f^i$ coeffective implies $(V_p, \dots, \psi_p), f_p$ coeffective as in Definition 3.37.

As in Definition 3.36 we define *triples* $(X, \mathbf{f}, \underline{\mathbf{C}})$, *splittings* $X = X_1 \amalg \dots \amalg X_n$ of $(X, \mathbf{f}, \underline{\mathbf{C}})$, *connected* $(X, \mathbf{f}, \underline{\mathbf{C}})$, *isomorphisms* $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \underline{\mathbf{C}}) \rightarrow (\tilde{X}, \tilde{\mathbf{f}}, \tilde{\underline{\mathbf{C}}})$, *automorphism groups* $\text{Aut}(X, \mathbf{f}, \underline{\mathbf{C}})$, *restrictions* $\underline{\mathbf{C}}|_{\partial X}$, and a natural map $\underline{\mathbf{C}} \mapsto \Pi(\underline{\mathbf{C}})$ from effective co-gauge-fixing data to co-gauge-fixing data.

Here are partial analogues of Theorems 3.18, 3.20 and 3.25, which show that versions of Property 3.1(a),(b) and (e) hold for effective (co-)gauge-fixing data.

Theorem 3.39. (a) *Let X be a compact, effective Kuranishi space with $\partial^2 X = \emptyset$, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ a strongly smooth map. Then $(X, \mathbf{f})$ admits effective gauge-fixing data $\underline{\mathbf{G}}$.*

(b) *If $\underline{\mathbf{G}}$ is effective gauge-fixing data for $(X, \mathbf{f})$ and $(X, \mathbf{f}, \underline{\mathbf{G}})$ is connected then $\text{Aut}(X, \mathbf{f}, \underline{\mathbf{G}}) = \{1\}$.*

(c) *Suppose X is a compact, effective Kuranishi space with corners, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ a strongly smooth map, and $\underline{\mathbf{H}}$ is effective gauge-fixing data for $(\partial X, \mathbf{f}|_{\partial X})$ with corners. Let $\sigma : \partial^2 X \rightarrow \partial^2 X$ be the involution of Definition 2.27. Then there exists effective gauge-fixing data $\underline{\mathbf{G}}$ for $(X, \mathbf{f})$ with corners with $\underline{\mathbf{G}}|_{\partial X} = \underline{\mathbf{H}}$ if and only if $\underline{\mathbf{H}}|_{\partial^2 X}$ is σ -invariant.*

Parts (a)–(c) also hold for effective co-gauge-fixing data $\underline{\mathbf{C}}, \underline{\mathbf{D}}$, taking $\mathbf{f}$ to be a strong submersion and $(X, \mathbf{f})$ to be coeffective rather than X effective.

Proof. For (a), we modify the proof of Theorem 3.18. In the first part, since X is effective and $\partial^2 X = \emptyset$, in Proposition 3.13 we can choose the very good coordinate system $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i), f^i : i \in I), \dots)$ for $(X, \mathbf{f})$ with $(V^i, \dots, \psi^i)$ effective and $\partial^2 V^i = \emptyset$ for all $i \in I$. Then Definition 3.36(a) holds. In Theorem 3.18 we choose the $\underline{\mathbf{G}}^i : E^i \rightarrow \underline{\mathbf{P}}$ such that $\coprod_{i \in I} \underline{\mathbf{G}}^i : \coprod_{i \in I} E^i \rightarrow \mathbb{R}^k \subset \underline{\mathbf{P}}$ is injective, for some $k \gg 0$. As $\partial^2 V^i = \emptyset$ we have $\partial^2 E^i = \emptyset$, so the immersion $\iota : \partial E^i \rightarrow E^i$ is injective, rather than merely finite. Hence $\coprod_{i \in I} \underline{\mathbf{G}}^i|_{\partial E^i} : \coprod_{i \in I} \partial E^i \rightarrow \underline{\mathbf{P}}$ is injective. As $\partial^l E^i = \emptyset$ for $l \geq 2$, Definition 3.36(b) follows.

Part (b) is a simplification of the proof of Theorem 3.20: for $(\mathbf{a}, \mathbf{b})$ in $\text{Aut}(X, \mathbf{f}, \underline{\mathbf{G}})$, as $\underline{\mathbf{G}}^i|_{(E^i)^\circ} : (E^i)^\circ \rightarrow \underline{\mathbf{P}}$ is injective, the diffeomorphisms $\hat{b}^i : E^i \rightarrow E^i$ with $\underline{\mathbf{G}}^i \circ \hat{b}^i \equiv \underline{\mathbf{G}}^i : E^i \rightarrow \underline{\mathbf{P}}$ must be the identity on $(E^i)^\circ$, so $\hat{b}^i$ is the identity by continuity. Therefore $b^i : V^i \rightarrow V^i$ is the identity, as $\hat{b}^i$ lifts b^i . Since $(b^i, \hat{b}^i)$ determines $\mathbf{a}$ on $\text{Im } \psi^i$ and $X = \bigcup_{i \in I} \text{Im } \psi^i$ this gives $\mathbf{a} = \text{id}_X$, so $\text{Aut}(X, \mathbf{f}, \underline{\mathbf{G}}) = \{1\}$. Part (c) is also a simple modification of Theorem 3.25: for the ‘if’ part, if $\underline{\mathbf{H}}|_{\partial^2 X}$ is σ -invariant, then $\underline{\mathbf{H}}$ prescribes consistent values for $\underline{\mathbf{G}}^i : E^i \rightarrow \underline{\mathbf{P}}$ on $E^i \setminus (E^i)^\circ$ for all $i \in I$, satisfying Definition 3.36(b) for $l \geq 1$. We then choose $\underline{\mathbf{G}}^i|_{(E^i)^\circ} : (E^i)^\circ \rightarrow \underline{\mathbf{P}}$ for $i \in I$ such that $\coprod_{i \in I} \underline{\mathbf{G}}^i|_{(E^i)^\circ} : \coprod_{i \in I} (E^i)^\circ \rightarrow \mathbb{R}^k \subset \underline{\mathbf{P}}$ is injective for some $k \gg 0$. $\square$

Remark 3.40. In Theorem 3.39(a) we omit the parts on finite groups Γ in Property 3.1(a), since Theorem 3.39(b) shows that we cannot have nontrivial symmetry groups Γ of $(X, \mathbf{f}, \underline{\mathbf{G}})$ for connected X . We also suppose $\partial^2 X = \emptyset$.

This can be weakened, but some assumption on $\partial^2 X$ is necessary for effective gauge-fixing data to exist on $(X, \mathbf{f})$, as the following argument shows.

Suppose that $\underline{\mathbf{G}}$ is effective gauge-fixing data for $(X, \mathbf{f})$ with X oriented, and that $\partial^2 X$ is nonempty, and connected as a topological space. Restrict $\underline{\mathbf{G}}$ to ∂X and $\partial^2 X$ to get effective gauge-fixing data $\underline{\mathbf{G}}|_{\partial^2 X}$ for $(\partial^2 X, \mathbf{f}|_{\partial^2 X})$. Then $(\partial^2 X, \mathbf{f}|_{\partial^2 X}, \underline{\mathbf{G}}|_{\partial^2 X})$ is a connected triple, as $\partial^2 X$ is a connected topological space, so $\text{Aut}(\partial^2 X, \mathbf{f}|_{\partial^2 X}, \underline{\mathbf{G}}|_{\partial^2 X}) = \{1\}$ by Theorem 3.39(b). However, the natural involution $\sigma : \partial^2 X \rightarrow \partial^2 X$ clearly preserves $\underline{\mathbf{G}}|_{\partial^2 X}$ and lies in $\text{Aut}(\partial^2 X, \mathbf{f}|_{\partial^2 X}, \underline{\mathbf{G}}|_{\partial^2 X}) = \{1\}$, a contradiction, as σ is orientation-reversing.

Following Definitions 3.26 and 3.27, we can define *pushforwards* $h_*(\underline{\mathbf{G}})$ and *pullbacks* $h^*(\underline{\mathbf{C}})$ of effective (co-)gauge-fixing data $\underline{\mathbf{G}}, \underline{\mathbf{C}}$.

Definition 3.41. Let Y, Z be orbifolds and $h : Y \rightarrow Z$ a smooth map. Suppose X is a compact, effective Kuranishi space, $\mathbf{f} : X \rightarrow Y$ is strongly smooth, and $\underline{\mathbf{G}} = (\mathbf{I}, \boldsymbol{\eta}, \underline{\mathbf{G}}^i : i \in I)$ is effective gauge-fixing data for $(X, \mathbf{f})$, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i), f^i : i \in I, \dots)$. Then $h \circ \mathbf{f} : X \rightarrow Z$ is strongly smooth. Define $h_*(\underline{\mathbf{G}})$ to be $h_*(\underline{\mathbf{G}}) = (h_*(\mathbf{I}), \boldsymbol{\eta}, \underline{\mathbf{G}}^i : i \in I)$, where $h_*(\mathbf{I}) = (I, (V^i, E^i, s^i, \psi^i), h \circ f^i : i \in I, \dots)$. Then $h_*(\underline{\mathbf{G}})$ is effective gauge-fixing data for $(X, h \circ \mathbf{f})$.

Definition 3.42. Let Y, Z be orbifolds without boundary and $h : Y \rightarrow Z$ a smooth, proper map. Suppose X is a compact Kuranishi space, $\mathbf{f} : X \rightarrow Y$ a strong submersion with $(X, \mathbf{f})$ coeffective, and $\underline{\mathbf{C}} = (\mathbf{I}, \boldsymbol{\eta}, \underline{\mathbf{C}}^i : i \in I)$ is effective co-gauge-fixing data for $(X, \mathbf{f})$, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i), f^i : i \in I, \dots)$. Then $Y \times_{h, Z, \mathbf{f}} X$ is a compact Kuranishi space, and $\pi_Y : Y \times_Z X \rightarrow Y$ is a strong submersion.

Define an excellent coordinate system $(\check{\mathbf{I}}, \check{\boldsymbol{\eta}})$ for $(Y \times_Z X, \pi_Y)$ as in Definition 3.27. Then $\check{I} \subseteq \{i + k : i \in I\}$, where $k = \dim Y - \dim Z$, and for each $i \in \check{I}$, $\check{V}^i$ is an open subset of $Y \times_{h, Z, f^{i-k}} V^{i-k}$, and $\check{E}^i$ an open subset of $Y \times_{h, Z, f^{i-k} \circ \pi^{i-k}} E^{i-k}$. Define $\check{\underline{\mathbf{C}}}^i : \check{E}^i \rightarrow \underline{\mathbf{P}}$ by $\check{\underline{\mathbf{C}}}^i = \underline{\mathbf{C}}^{i-k} \circ \pi_{E^{i-k}}$. We claim that $h^*(\underline{\mathbf{C}}) = (\check{\mathbf{I}}, \check{\boldsymbol{\eta}}, \check{\underline{\mathbf{C}}}^i : i \in \check{I})$ is *effective co-gauge-fixing data* for $(Y \times_Z X, \pi_Y)$.

We must verify Definition 3.38(a),(b) hold. For (a), as the pair $(V^{i-k}, \dots, \psi^{i-k}), f^{i-k}$ is coeffective and $\check{V}^i \subseteq Y \times_{h, Z, f^{i-k}} V^{i-k}$, $\check{E}^i \subseteq Y \times_{h, Z, f^{i-k} \circ \pi^{i-k}} E^{i-k}$, one can show directly that the pair $(\check{V}^i, \dots, \check{\psi}^i), \check{\pi}_Y^i$ is coeffective. This then implies that $(Y \times_Z X, \pi_Y)$ is coeffective, as in Definition 3.38.

For (b), if $\partial^l X = \coprod_{a=1}^{m^l} X_a^l$ and $\partial^l(Y \times_Z X) = \coprod_{b=1}^{n^l} (Y \times_Z X)_b^l$ are the splittings of $(\partial^l X, \mathbf{f}|_{\partial^l X}, \mathbf{I}|_{\partial^l X}, \boldsymbol{\eta}|_{\partial^l X})$ and $(\partial^l(Y \times_Z X), \pi_Y|_{\partial^l(Y \times_Z X)}, \check{\mathbf{I}}|_{\partial^l(Y \times_Z X)}, \check{\boldsymbol{\eta}}|_{\partial^l(Y \times_Z X)})$ into connected components in Lemma 3.7, then for each $b = 1, \dots, n^l$, $(Y \times_Z X)_b^l$ is a connected component of $(Y \times_Z (X_a^l), \pi_Y|_{Y \times_Z X_a^l}, \check{\mathbf{I}}|_{Y \times_Z X_a^l}, \check{\boldsymbol{\eta}}|_{Y \times_Z X_a^l})$ for some $a = 1, \dots, m^l$. If $(\check{V}_b^{l,i}, \check{E}_b^{l,i}, \check{s}_b^{l,i}, \check{\psi}_b^{l,i})$ is a Kuranishi neighbourhood on $(Y \times_Z X)_b^l$ in $\check{\mathbf{I}}_b^l$, then $(\check{E}_b^{l,i})^\circ \subseteq Y \times_Z (E_a^{l,i-k})^\circ$, and $\check{\underline{\mathbf{C}}}^{i+l}|_{(\check{E}_b^{l,i})^\circ} \equiv \underline{\mathbf{C}}^{i-k+l} \circ \pi_{(E_a^{l,i-k})^\circ}$. As $\underline{\mathbf{C}}^{i-k+l}$ maps $(E_a^{l,i-k})^\circ \rightarrow \mathbb{R}^{k_a^l} \subset \underline{\mathbf{P}}$, it follows that $\check{\underline{\mathbf{C}}}^{i+l}$ maps $(\check{E}_b^{l,i})^\circ \rightarrow \mathbb{R}^{k_a^l} \subset \underline{\mathbf{P}}$, giving the first part of Definition 3.38(b). For the second, since $\coprod_{i-k \in I_a^l} (\underline{\mathbf{C}}^{i-k+l} \times (f^{i-k+l} \circ \pi^{i-k+l}))|_{(E_a^{l,i-k})^\circ} :$

$\coprod_{i-k \in I_a^l} (E_a^{l,i-k})^\circ \rightarrow \underline{P} \times Z$ is injective by Definition 3.38(b), and $\coprod_{i \in \tilde{I}_b^l} (\tilde{E}_b^{l,i})^\circ \subset Y \times_Z (\coprod_{i-k \in I_a^l} (E_a^{l,i-k})^\circ)$, it follows that $\coprod_{i \in \tilde{I}_b^l} (\tilde{\mathcal{C}}^{i+l} \times (\tilde{f}^{i+l} \circ \tilde{\pi}^{i+l}))|_{(\tilde{E}_b^{l,i})^\circ} : \coprod_{i \in \tilde{I}_b^l} (\tilde{E}_b^{l,i})^\circ \rightarrow \underline{P} \times Y$ is injective, as we want.

This last part is a little more subtle than it appears. The important point is this: suppose (V, E, s, ψ) is a Kuranishi neighbourhood on X , $f : V \rightarrow Z$ a submersion with $(V, E, s, \psi), f$ coeffective, and $\underline{C} : E \rightarrow \underline{P}$ is a map with $\underline{C} \times (f \circ \pi) : E \rightarrow \underline{P} \times Z$ injective. As $(V, E, s, \psi), f$ is coeffective, $f \circ \pi : E \rightarrow Z$ is surjective on stabilizer groups. Therefore the biquotient terms $f_*(\text{Stab}(p) \backslash \text{Stab}(f(p)) / f'_*(\text{Stab}(p')))$ in the definition (12) of the orbifold fibre product $Y \times_{h,Z,f \circ \pi} E$ are all points, so that $\pi_E \times \pi_Y : Y \times_Z E \rightarrow E \times Y$ is injective. Combining this with $\underline{C} \times (f \circ \pi)$ injective implies that $(\underline{C} \circ \pi_E) \times \pi_Y : Y \times_Z E \rightarrow \underline{P} \times Y$ is injective. Thus, we need the coeffective assumptions for $\underline{C}$ in Definition 3.38(a) to prove the injectivity for $h^*(\underline{C})$ in Definition 3.38(b).

Pushforwards and pullbacks are *functorial*, that is, $(g \circ h)_*(\underline{G}) = g_* \circ h_*(\underline{G})$, and $(g \circ h)^*(\underline{C}) = h^* \circ g^*(\underline{C})$, as for (co-)gauge-fixing data. We modify Definition 3.31 to define *fibre products* $\underline{C} \times_Y \tilde{\underline{C}}, \underline{G} \times_Y \tilde{\underline{C}}$ of effective (co-)gauge-fixing data.

Definition 3.43. Let $X, \tilde{X}$ be compact Kuranishi spaces, Y an orbifold, and $\mathbf{f} : X \rightarrow Y, \tilde{\mathbf{f}} : \tilde{X} \rightarrow Y$ be strong submersions with $(X, \mathbf{f}), (\tilde{X}, \tilde{\mathbf{f}})$ coeffective. Suppose $\underline{C} = (I, \eta, \underline{C}^i : i \in I)$ and $\tilde{\underline{C}} = (\tilde{I}, \tilde{\eta}, \tilde{\underline{C}}^{\tilde{i}} : \tilde{i} \in \tilde{I})$ are effective co-gauge-fixing data for $(X, \mathbf{f})$ and $(\tilde{X}, \tilde{\mathbf{f}})$. Let $(I, \eta) \times_Y (\tilde{I}, \tilde{\eta})$ be the really good coordinate system for $(X \times_Y \tilde{X}, \pi_Y)$ given in Definition 3.29, and $(\tilde{I}, \tilde{\eta}) = ((I, \eta) \times_Y (\tilde{I}, \tilde{\eta}))^\sim$ the excellent coordinate system for $(X \times_Y \tilde{X}, \pi_Y)$ constructed from $(I, \eta) \times_Y (\tilde{I}, \tilde{\eta})$ by Algorithm 3.16. In the notation of Definitions 3.29 and 3.31 we have $\tilde{I} = (\tilde{I}, (\tilde{V}^k, \tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k), \tilde{\pi}_Y^k : k \in \tilde{I}, \dots)$, with $\tilde{V}^k, \tilde{E}^k$ open sets in $\coprod_{i \in I, \tilde{i} \in \tilde{I} : i + \tilde{i} = k + \dim Y} V^{(i, \tilde{i})}, \coprod_{i \in I, \tilde{i} \in \tilde{I} : i + \tilde{i} = k + \dim Y} E^{(i, \tilde{i})}$, where $V^{(i, \tilde{i})} = V^i \times_{f^i, Y, \tilde{f}^{\tilde{i}}} \tilde{V}^{\tilde{i}}$ and $E^{(i, \tilde{i})} = E^i \times_{f^i \circ \pi^i, Y, \tilde{f}^{\tilde{i}} \circ \tilde{\pi}^{\tilde{i}}} \tilde{E}^{\tilde{i}}$.

For $k \in \tilde{I}$, define $\tilde{\underline{C}}^k : \tilde{E}^k \rightarrow \underline{P}$ by $\tilde{\underline{C}}^k(\tilde{e}) = \underline{\mu}(\underline{C}^i \circ \pi_{E^i}(\tilde{e}), \tilde{\underline{C}}^{\tilde{i}} \circ \pi_{\tilde{E}^{\tilde{i}}}(\tilde{e}))$ whenever $\tilde{e}$ lies in $\tilde{E}^k \cap E^{(i, \tilde{i})}$, for all $i \in I$ and $\tilde{i} \in \tilde{I}$ with $i + \tilde{i} = k + \dim Y$, where $\pi_{E^i} : E^{(i, \tilde{i})} \rightarrow E^i$ and $\pi_{\tilde{E}^{\tilde{i}}} : E^{(i, \tilde{i})} \rightarrow \tilde{E}^{\tilde{i}}$ are the projections from $E^{(i, \tilde{i})} = E^i \times_Y \tilde{E}^{\tilde{i}}$, and $\underline{\mu} : \underline{P} \times \underline{P} \rightarrow \underline{P}$ is as in (58). We shall show that $\underline{C} \times_Y \tilde{\underline{C}} = \tilde{\underline{C}} = (\tilde{I}, \tilde{\eta}, \tilde{\underline{C}}^{\tilde{i}} : \tilde{i} \in \tilde{I})$ is *effective co-gauge-fixing data* for $(X \times_Y \tilde{X}, \pi_Y)$.

We must verify Definition 3.38(a),(b) hold. The proof is similar to that for $h^*(\underline{C})$ in Definition 3.42. For (a), as $(V^i, \dots, \psi^i), f^i$ and $(\tilde{V}^{\tilde{i}}, \dots, \tilde{\psi}^{\tilde{i}}), \tilde{f}^{\tilde{i}}$ are coeffective one can show directly that $(V^{(i, \tilde{i})}, \dots, \psi^{(i, \tilde{i})}), \pi_Y^{(i, \tilde{i})}$ in (37) is coeffective. Therefore $(\tilde{V}^k, \dots, \tilde{\psi}^k), \tilde{\pi}_Y^k$ is coeffective over the open set $\tilde{V}^k \cap V^{(i, \tilde{i})}$ in $V^{(i, \tilde{i})}$, and as this holds for all $i, \tilde{i}$ with $i + \tilde{i} = k + \dim Y$, $(\tilde{V}^k, \dots, \tilde{\psi}^k), \tilde{\pi}_Y^k$ is coeffective. This proves (a).

For Definition 3.38(b), first consider the case $l = 0$, and suppose $(X, \mathbf{f}, \underline{C}), (\tilde{X}, \tilde{\mathbf{f}}, \tilde{\underline{C}})$ are connected. Then Definition 3.38(b) implies that $\underline{C}^i$ maps $(E^i)^\circ \rightarrow \mathbb{R}_1^{k_1^0} \subset \underline{P}$ for some $k_1^0 \geq 0$ independent of i and all $i \in I$, and $\tilde{\underline{C}}^{\tilde{i}}$ maps $(\tilde{E}^{\tilde{i}})^\circ \rightarrow \mathbb{R}_1^{\tilde{k}_1^0} \subset \underline{P}$ for some $\tilde{k}_1^0 \geq 0$ independent of $\tilde{i}$ and all $\tilde{i} \in \tilde{I}$. As $(\tilde{E}^k)^\circ \subseteq \coprod_{i \in I, \tilde{i} \in \tilde{I} : i + \tilde{i} = k + \dim Y} (E^i)^\circ \times_Y (\tilde{E}^{\tilde{i}})^\circ$, it follows from (58) that $\tilde{\underline{C}}^k$ maps

$(\check{E}^k)^\circ \rightarrow \mathbb{R}^{k_1^0 + \tilde{k}_1^0} \subset \underline{P}$ for all $k \in \check{I}$, where $k_1^0 + \tilde{k}_1^0$ is independent of $k \in \check{I}$. This proves the first part of Definition 3.38(b) in this case.

For the second part, observe that $\coprod_{k \in \check{I}} (\check{\underline{C}}^k \times (\check{f}^k \circ \check{\pi}^k))$ is the composition

$$\coprod_{k \in \check{I}} (\check{E}^k)^\circ \hookrightarrow \left(\coprod_{i \in I} (E^i)^\circ \right) \times_Y \frac{\left(\coprod_{i \in I} \underline{C}^i \circ \pi_{E^i} \right) \times \left(\coprod_{i \in \check{I}} \check{\underline{C}}^i \circ \pi_{\check{E}^i} \right) \times \pi_Y}{\left(\coprod_{i \in \check{I}} (\check{E}^i)^\circ \right)} \xrightarrow{\quad} \mathbb{R}^{k_1^0} \times \mathbb{R}^{\tilde{k}_1^0} \times Y \xrightarrow{\underline{\mu} \times \text{id}_Y} \mathbb{R}^{k_1^0 + \tilde{k}_1^0} \times Y.$$

Here the first arrow ‘ $\hookrightarrow$ ’ is an inclusion of open sets, and so injective. The second arrow is injective as $\coprod_{i \in I} (\underline{C}^i \times (f^i \circ \pi^i)) : \coprod_{i \in I} (E^i)^\circ \rightarrow \mathbb{R}^{k_1^0} \times Y \subset \underline{P} \times Y$ is injective, and $\coprod_{i \in \check{I}} (\check{\underline{C}}^i \times (\check{f}^i \circ \check{\pi}^i)) : \coprod_{i \in \check{I}} (\check{E}^i)^\circ \rightarrow \mathbb{R}^{\tilde{k}_1^0} \times Y \subset \underline{P} \times Y$ is injective, and $\pi_{E^i} \times \pi_{\check{E}^i} : E^i \times_Y \check{E}^i \rightarrow E^i \times \check{E}^i$ is injective as $f^i \circ \pi^i : E^i \rightarrow Y$, $\check{f}^i \circ \check{\pi}^i : \check{E}^i \rightarrow Y$ are surjective on stabilizer groups. The third arrow is injective as $\underline{\mu} : \mathbb{R}^{k_1^0} \times \mathbb{R}^{\tilde{k}_1^0} \rightarrow \mathbb{R}^{k_1^0 + \tilde{k}_1^0}$ in (58) is injective.

Therefore the composition $\coprod_{k \in \check{I}} (\check{\underline{C}}^k \times (\check{f}^k \circ \check{\pi}^k))$ is injective, and Definition 3.38(b) holds in the case when $l = 0$ and $(X, \underline{f}, \underline{C}), (\tilde{X}, \tilde{\underline{f}}, \tilde{\underline{C}})$ are connected. For the general case, we apply the argument above with connected components $(X_a^j, \underline{f}|_{X_a^j}, \underline{C}|_{X_a^j})$ of $(\partial^j X, \underline{f}|_{\partial^j X}, \underline{C}|_{\partial^j X})$ and $(\tilde{X}_a^{\tilde{j}}, \tilde{\underline{f}}|_{\tilde{X}_a^{\tilde{j}}}, \tilde{\underline{C}}|_{\tilde{X}_a^{\tilde{j}}})$ of $(\partial^{\tilde{j}} \tilde{X}, \tilde{\underline{f}}|_{\partial^{\tilde{j}} \tilde{X}}, \tilde{\underline{C}}|_{\partial^{\tilde{j}} \tilde{X}})$ in place of $(X, \underline{f}, \underline{C}), (\tilde{X}, \tilde{\underline{f}}, \tilde{\underline{C}})$, with $j + \tilde{j} = l$. This proves Definition 3.38(b), so $\underline{C} \times_Y \tilde{\underline{C}}$ is *effective co-gauge-fixing data*.

Now let X be a compact, effective Kuranishi space, Y an orbifold, $\underline{f} : X \rightarrow Y$ strongly smooth, $\underline{G}$ effective gauge-fixing data for $(X, \underline{f})$, $\tilde{X}$ a compact Kuranishi space, $\tilde{\underline{f}} : \tilde{X} \rightarrow Y$ a strong submersion with $(\tilde{X}, \tilde{\underline{f}})$ coefficient, and $\tilde{\underline{C}}$ be effective co-gauge-fixing data for $(\tilde{X}, \tilde{\underline{f}})$. Then we can modify Definition 3.31 in the same way to define effective gauge-fixing data $\underline{G} \times_Y \tilde{\underline{C}}$ for $(X \times_Y \tilde{X}, \pi_Y)$, defining $\check{\underline{G}}^k : \check{E}^k \rightarrow \underline{P}$ by $\check{\underline{G}}^k(\check{e}) = \underline{\mu}(\underline{G}^i \circ \pi_{E^i}(\check{e}), \tilde{\underline{C}}^i \circ \pi_{\check{E}^i}(\check{e}))$. The proof that $\underline{G} \times_Y \tilde{\underline{C}}$ satisfies Definition 3.36 is similar to the above.

The maps $\underline{G} \mapsto \Pi(\underline{G})$, $\underline{C} \mapsto \Pi(\underline{C})$ from effective (co-)gauge-fixing data to (co-)gauge-fixing data in Definitions 3.36 and 3.38 are compatible with fibre products, that is, $\Pi(\underline{C} \times_Y \tilde{\underline{C}}) = \Pi(\underline{C}) \times_Y \Pi(\tilde{\underline{C}})$ and $\Pi(\underline{G} \times_Y \tilde{\underline{C}}) = \Pi(\underline{G}) \times_Y \Pi(\tilde{\underline{C}})$, because $\mu(\Pi(p), \Pi(q)) = \Pi(\mu(p, q))$ for $p, q \in \underline{P}$.

From Propositions 3.32 and 3.33, with proofs almost unchanged, we deduce:

Proposition 3.44. *Effective (co-)gauge-fixing data satisfies analogues of Propositions 3.32(b),(c),(d) and 3.33, taking X_a effective or $(X_a, \underline{f}_a)$ coefficient where appropriate. However, the analogue of Proposition 3.32(a) does not apply, that is, fibre products of effective co-gauge-fixing data are not commutative, since $\underline{\mu}$ in (58) is not commutative.*

Problem (iv) at the beginning of §3.9 implies that it is necessary for fibre products $\underline{C} \times_Y \tilde{\underline{C}}$ of effective co-gauge-fixing data not to be commutative. We give another reason, involving *Steenrod squares*, in Remark 4.33.

4 Kuranishi homology and cohomology

We now define two slightly different homology theories of an orbifold Y : *Kuranishi homology* $KH_*(Y; R)$, which is defined when R is a $\mathbb{Q}$ -algebra, and *effective Kuranishi homology* $KH_*^{\text{ef}}(Y; R)$, which is defined when R is a commutative ring. Both of them are isomorphic to *singular homology* $H_*^{\text{si}}(Y; R)$. They are variations of the same idea, which is to define a homology theory using as chains strongly smooth maps $\mathbf{f} : X \rightarrow Y$ from a compact Kuranishi space X .

However, we show in §4.9 that just using isomorphism classes $[X, \mathbf{f}]$ of pairs $(X, \mathbf{f})$ as chains would make $KH_*(Y; R) = 0$ for all Y, R . The problems are caused by pairs $(X, \mathbf{f})$ whose automorphism groups are infinite. To avoid them, we include (*effective*) *gauge-fixing data* $\mathbf{G}$ on X , as in Chapter 3, and take chains to be generated by isomorphism classes $[X, \mathbf{f}, \mathbf{G}]$ of triples $(X, \mathbf{f}, \mathbf{G})$.

We also define Poincaré dual cohomology theories, *Kuranishi cohomology* $KH^*(Y; R)$ for R a $\mathbb{Q}$ -algebra, and *effective Kuranishi cohomology* $KH_{\text{ec}}^*(Y; R)$ for R a commutative ring. Here $KH^*(Y; R)$ is always isomorphic to *compactly supported cohomology* $H_{\text{cs}}^*(Y; R)$, and $KH_{\text{ec}}^*(Y; R) \cong H_{\text{cs}}^*(Y; R)$ if Y is a manifold. The main point of introducing them is as a tool to use in areas where Kuranishi spaces naturally arise, such as closed or open Gromov–Witten theory, or Lagrangian Floer cohomology.

Kuranishi (co)homology $KH_*, KH^*(Y; R)$ is better behaved at the (co)chain level than effective Kuranishi (co)homology $KH_*^{\text{ef}}, KH_{\text{ec}}^*(Y; R)$. For example, the cup product $\cup$ on Kuranishi cochains $KC^*(Y; R)$ is supercommutative, but the cup product $\cup$ on effective Kuranishi cochains $KC_{\text{ec}}^*(Y; R)$ is not. Also, we can form chains $[X, \mathbf{f}, \mathbf{G}]$ in $KC_*(Y; R)$ for X an arbitrary compact oriented Kuranishi space, but for $KC_*^{\text{ef}}(Y; R)$, X must satisfy restrictions on its stabilizer groups and corners.

The advantage of effective Kuranishi (co)homology $KH_*^{\text{ef}}, KH_{\text{ec}}^*(Y; R)$ is that they work over any commutative ring R , such as $\mathbb{Z}$, not just over $\mathbb{Q}$ -algebras. This will be important in the author’s approach to the integrality conjecture for Gopakumar–Vafa invariants, to be discussed in Chapter 6. We will also use effective Kuranishi homology in showing Kuranishi homology is isomorphic to singular homology, as we will first prove that $KH_*^{\text{ef}}(Y; R) \cong H_*^{\text{si}}(Y; R)$ for R a commutative ring, and then that $KH_*(Y; R) \cong KH_*^{\text{ef}}(Y; R)$ for R a $\mathbb{Q}$ -algebra.

4.1 Classical homology and cohomology

Let Y be a topological space and R a commutative ring. Then there are many ways of defining *homology groups* $H_k(Y; R)$ of Y with coefficients in R , for $k = 0, 1, 2, \dots$, for example, singular homology, cellular homology (of a CW-complex), and Borel–Moore homology. A general feature of all these theories is that if Y is sufficiently well-behaved, for instance, if Y is a manifold, then the homology groups obtained from any two homology theories are canonically isomorphic. (Theorems 4.8 and 4.9 below are results of this kind.) So for many purposes, it does not matter which homology theory one uses.

There are also many ways of defining *cohomology groups* $H^k(Y; R)$ and

compactly-supported cohomology groups $H_{\text{cs}}^k(Y; R)$, for example, singular cohomology, Čech cohomology, and de Rham cohomology (of a smooth manifold, over $\mathbb{R}$). Again, if Y is sufficiently well-behaved, then the (compactly-supported) cohomology groups obtained from any two cohomology theories are isomorphic.

We briefly recall the definition of *singular homology*, from Bredon [10, §IV].

Definition 4.1. For $k = 0, 1, \dots$, the k -simplex Δ_k is

$$\Delta_k = \{(x_0, \dots, x_k) \in \mathbb{R}^{k+1} : x_i \geq 0, x_0 + \dots + x_k = 1\}. \quad (59)$$

It is a compact, oriented k -manifold, with boundary and corners.

Let Y be a topological space, and R a commutative ring. Define $C_k^{\text{si}}(Y; R)$ to be the R -module spanned by *singular simplices*, which are continuous maps $\sigma : \Delta_k \rightarrow Y$. Elements of $C_k^{\text{si}}(Y; R)$, which are called *singular chains*, are finite sums $\sum_{a \in A} \rho_a \sigma_a$, where A is a finite indexing set, $\rho_a \in R$, and $\sigma_a : \Delta_k \rightarrow Y$ is continuous for $a \in A$.

The *boundary operator* $\partial : C_k^{\text{si}}(Y; R) \rightarrow C_{k-1}^{\text{si}}(Y; R)$ is [10, §IV.1]:

$$\partial : \sum_{a \in A} \rho_a \sigma_a \mapsto \sum_{a \in A} \sum_{j=0}^k (-1)^j \rho_a (\sigma_a \circ F_j^k), \quad (60)$$

where for $j = 0, \dots, k$ the map $F_j^k : \Delta_{k-1} \rightarrow \Delta_k$ is given by $F_j^k(x_0, \dots, x_{k-1}) = (x_0, \dots, x_{j-1}, 0, x_j, \dots, x_{k-1})$. As a manifold with boundary, we have $\partial\Delta_k = \coprod_{j=0}^k \Delta_j^k$, where Δ_j^k is the connected component of $\partial\Delta_k$ on which $x_j \equiv 0$, so that $F_j^k : \Delta_{k-1} \rightarrow \Delta_j^k$ is a diffeomorphism. The orientation on Δ_k induces one on $\partial\Delta_k$, and so induces orientations on Δ_j^k for $j = 0, \dots, k$. It is easy to show that under $F_j^k : \Delta_{k-1} \rightarrow \Delta_j^k$, the orientations of Δ_{k-1} and Δ_j^k differ by a factor $(-1)^j$, which is why this appears in (60). So ∂ in (60) basically restricts from Δ_k to $\partial\Delta_k$, as oriented manifolds with boundary and corners.

From (60) we find that $\partial \circ \partial = 0$, since each codimension 2 face Δ_{k-2} of Δ_k contributes twice to $\partial \circ \partial$, once with sign 1 and once with sign -1 . Thus we may define the *singular homology group*

$$H_k^{\text{si}}(Y; R) = \frac{\text{Ker}(\partial : C_k^{\text{si}}(Y; R) \rightarrow C_{k-1}^{\text{si}}(Y; R))}{\text{Im}(\partial : C_{k+1}^{\text{si}}(Y; R) \rightarrow C_k^{\text{si}}(Y; R))}.$$

If Y is a smooth manifold or orbifold, we can instead define $C_k^{\text{si}}(Y; R)$, $H_k^{\text{si}}(Y; R)$ using *smooth* maps $\sigma : \Delta_k \rightarrow Y$, as in [10, §V.5], which we call *smooth singular simplices*, and this gives the same homology groups. We shall always take $C_k^{\text{si}}(Y; R)$ and $H_k^{\text{si}}(Y; R)$ to be defined using smooth singular simplices.

We will not choose a particular cohomology theory. We will only be interested in compactly-supported cohomology of manifolds or orbifolds, and this can be characterized in terms of (singular) homology by Poincaré duality isomorphisms, as in (61) below. So when we prove that (effective) Kuranishi cohomology is isomorphic to compactly-supported cohomology, we will go via singular homology using Poincaré duality.

Here are some general properties of homology and cohomology, which can mostly be found in Bredon [10].

- Let Y, Z be topological spaces, $h : Y \rightarrow Z$ a continuous map, and R a commutative ring. Then there is a *pushforward map* $h_* : H_*(Y; R) \rightarrow H_*(Z; R)$. On singular homology, this is induced by $h_* : C_*^{\text{si}}(Y; R) \rightarrow C_*^{\text{si}}(Z; R)$ mapping $h_* : \sum_{a \in A} \rho_a \sigma_a \mapsto \sum_{a \in A} \rho_a (h \circ \sigma_a)$. Pushforwards are functorial, $(g \circ h)_* = g_* \circ h_*$.
- Let Y, Z be topological spaces, $h : Y \rightarrow Z$ a continuous map, and R a commutative ring. Then there is a *pullback map* $h^* : H^*(Z; R) \rightarrow H^*(Y; R)$ on cohomology. Pullbacks are functorial, $(g \circ h)^* = h^* \circ g^*$.

For *compactly-supported* cohomology, we can only define pullbacks $h^* : H_{\text{cs}}^*(Z; R) \rightarrow H_{\text{cs}}^*(Y; R)$ if $h : Y \rightarrow Z$ is *proper*, that is, if $h^{-1}(S) \subseteq Y$ is compact in Y whenever $S \subseteq Z$ is compact in Z . This is because h^* must pull compactly-supported cochains in Z back to compactly-supported cochains in Y .

- There is a natural morphism $H_{\text{cs}}^*(Y; R) \rightarrow H^*(Y; R)$. If Y is compact this is an isomorphism, $H_{\text{cs}}^*(Y; R) \cong H^*(Y; R)$.
- There are associative, supercommutative, graded multiplications on both cohomology and compactly-supported cohomology, the *cup product* $\cup : H^k(Y; R) \times H^l(Y; R) \rightarrow H^{k+l}(Y; R)$. There is an *identity* $1 \in H^0(Y; R)$, and if Y is compact there is an identity in $H_{\text{cs}}^0(Y; R)$.

There are also *cap products* $\cap : H_k(Y; R) \times H^l(Y; R) \rightarrow H_{k-l}(Y; R)$ and $\cap : H_k(Y; R) \times H_{\text{cs}}^l(Y; R) \rightarrow H_{k-l}(Y; R)$, which make $H_*(Y; R)$ into a *module* over $H^*(Y; R)$ and $H_{\text{cs}}^*(Y; R)$.

- Suppose Y is an oriented manifold, of dimension n , without boundary, and not necessarily compact. Then there are *Poincaré duality isomorphisms*

$$\text{Pd} : H_{\text{cs}}^k(Y; R) \longrightarrow H_{n-k}(Y; R) \quad (61)$$

between compactly-supported cohomology, and homology.

If Y is also *compact* then it has a *fundamental class* $[Y] \in H_n(Y; R)$, and we can write the Poincaré duality map Pd of (61) in terms of the cap product by $\text{Pd}(\alpha) = [Y] \cap \alpha$ for $\alpha \in H_{\text{cs}}^k(Y; R)$. If Y is noncompact and not too badly behaved, a similar interpretation is possible, but the fundamental class $[Y]$ exists not in $H_n(Y; R)$ but in the ‘non-compactly-supported homology group’ $H_n^{\text{nc}}(Y; R)$ which is $H_n(\bar{Y}, \{\infty\}; R)$ in relative homology, where $\bar{Y} = Y \cup \{\infty\}$ is the one-point compactification of Y .

Poincaré duality for manifolds Y with boundary and (g-)corners will be discussed in §4.5.

- If Y is an oriented n -manifold without boundary then Poincaré duality and the cup product on $H_{\text{cs}}^*(Y; R)$ induce an associative, supercommutative *intersection product* $\bullet : H_k(Y; R) \times H_l(Y; R) \rightarrow H_{k+l-n}(Y; R)$ on homology $H_*(Y; R)$. This can be described geometrically at the chain level in singular homology, as in Bredon [10, VI.11] and Lefschetz [46, §IV].

- Fulton and MacPherson [27] define the formalism of *bivariant theories*, which simultaneously generalize homology and cohomology. In §4.8 we will explain this, and show how (effective) Kuranishi (co)homology can be extended to a bivariant theory, *at the level of (co)chains*.
- Here is an important point: *by the homology or cohomology of an orbifold Y , we will always mean the (co)homology of the underlying topological space*. However, there are other ways to define (co)homology of an orbifold, which also use the orbifold structure of Y , and even additional data.

Behrend [7] defines versions of de Rham cohomology $H_{\text{dR}}^*(Y; \mathbb{R})$, singular homology $H_*^{\text{Be}}(Y; R)$ and singular cohomology $H_{\text{Be}}^*(Y; R)$ of an orbifold, regarded as a stack. In Behrend's notation, the underlying topological space $\bar{Y}$ of an orbifold Y is called the *coarse moduli space* of Y . Behrend shows [7, Prop. 36] that $H_*^{\text{Be}}(Y; R) \cong H_*^{\text{si}}(\bar{Y}; R)$ and $H_{\text{Be}}^*(Y; R) \cong H^*(\bar{Y}; R)$ when R is a $\mathbb{Q}$ -algebra, but for general R such as $R = \mathbb{Z}$ the (co)homology of Y and $\bar{Y}$ may differ.

Also, if Y is an orbifold with an almost complex structure, Chen and Ruan [13] define the *orbifold cohomology* $H_{\text{CR}}^*(Y; R)$, which appears to be natural in Gromov–Witten theory and String Theory of orbifolds. In general $H_{\text{CR}}^*(Y; R)$ is different from both $H^*(\bar{Y}; R)$ and $H_{\text{Be}}^*(Y; R)$.

We shall be interested in how Poincaré duality for manifolds extends to *orbifolds* Y . Satake [66, Th. 3] showed that Poincaré duality isomorphisms (61) exist when Y is an oriented orbifold and R is a $\mathbb{Q}$ -algebra. However, *Poincaré duality need not hold for orbifolds for general R* . (Here, as in the last remark above, we are discussing Poincaré duality for the (co)homology of the underlying topological space.) Poincaré duality can fail at two levels.

Let Y be an oriented n -orbifold, and suppose for simplicity that Y is compact. Firstly, if Y is a *non-effective* orbifold, then one can in general define the fundamental class $[Y] \in H_n(Y; R)$ only if R is a $\mathbb{Q}$ -algebra. This is because generic points in Y have non-trivial stabilizer group Γ , so working say in singular homology, when we triangulate Y by simplices to define a cycle representing $[Y]$ we must weight each simplex by $\pm|\Gamma|^{-1}$, so we need $|\Gamma|^{-1} \in R$, and $\mathbb{Q} \subseteq R$. Since Pd in (61) is defined using $[Y]$, for non-effective Y and general R the Poincaré duality map Pd is undefined.

Secondly, if Y is an *effective* orbifold, we can define $[Y]$ and Pd in (61), but Pd may not be an isomorphism. Here is an example of a compact, oriented, effective orbifold for which Pd is not an isomorphism over $\mathbb{Z}$.

Example 4.2. Let $Y = (\mathbb{CP}^1 \times \mathbb{CP}^1)/\mathbb{Z}_2$, where the $\mathbb{Z}_2$ -action is generated by $\sigma : \mathbb{CP}^1 \times \mathbb{CP}^1 \rightarrow \mathbb{CP}^1 \times \mathbb{CP}^1$ mapping $\sigma : ([x_0, x_1], [y_0, y_1]) \mapsto ([x_0, -x_1], [y_0, -y_1])$. It is a compact 4-orbifold, with an orientation induced by the complex structure, that has four orbifold points $\{[1, 0], [0, 1]\} \times \{[1, 0], [0, 1]\}$ modelled on $\mathbb{R}^4/\{\pm 1\}$.

Let α, β in $H_2(Y; \mathbb{Z}) \subset H_2(Y; \mathbb{Q})$ be the homology classes of the suborbifolds $(\mathbb{CP}^1 \times \{[1, 0]\})/\mathbb{Z}_2$ and $(\{[1, 0]\} \times \mathbb{CP}^1)/\mathbb{Z}_2$. Since Poincaré duality holds over $\mathbb{Q}$, there is an intersection product $\bullet : H_2(Y; \mathbb{Q}) \times H_2(Y; \mathbb{Q}) \rightarrow H_0(Y; \mathbb{Q}) \cong \mathbb{Q}$.

As $(\mathbb{CP}^1 \times \{[1, 0]\})/\mathbb{Z}_2$, $(\{[1, 0]\} \times \mathbb{CP}^1)/\mathbb{Z}_2$ intersect transversely in one point $([1, 0], [1, 0])$, which has stabilizer group $\mathbb{Z}_2$, it follows that $\alpha \bullet \beta = \frac{1}{2} \in \mathbb{Q}$.

If Poincaré duality (61) were an isomorphism for $R = \mathbb{Z}$ then there would exist an intersection product $\bullet : H_2(Y; \mathbb{Z}) \times H_2(Y; \mathbb{Z}) \rightarrow H_0(Y; \mathbb{Z}) \cong \mathbb{Z}$ agreeing with that on $H_*(Y; \mathbb{Q})$ under the morphism $H_*(Y; \mathbb{Z}) \rightarrow H_*(Y; \mathbb{Q})$ induced by $\mathbb{Z} \hookrightarrow \mathbb{Q}$. However, this contradicts $\alpha, \beta \in H_2(Y; \mathbb{Z})$ but $\alpha \bullet \beta = \frac{1}{2} \notin H_0(Y; \mathbb{Z}) \subset H_0(Y; \mathbb{Q})$. This means that α, β do not lie in the image of Pd (though $2\alpha, 2\beta$ do). Thus Pd in (61) is not surjective, and so not an isomorphism.

4.2 (Effective) Kuranishi homology

We now define the (effective) Kuranishi homology groups of an orbifold.

Definition 4.3. Let Y be an orbifold. Consider triples $(X, \mathbf{f}, \mathbf{G})$, where X is a compact oriented Kuranishi space, $\mathbf{f} : X \rightarrow Y$ is strongly smooth, and $\mathbf{G}$ is gauge-fixing data for $(X, \mathbf{f})$. Write $[X, \mathbf{f}, \mathbf{G}]$ for the isomorphism class of $(X, \mathbf{f}, \mathbf{G})$ under isomorphisms $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \mathbf{G}) \rightarrow (\tilde{X}, \tilde{\mathbf{f}}, \tilde{\mathbf{G}})$ as in Definition 3.8, where in addition we require $\mathbf{a}$ to identify the orientations of $X, \tilde{X}$.

Let R be a $\mathbb{Q}$ -algebra, for instance $\mathbb{Q}, \mathbb{R}$ or $\mathbb{C}$. For each $k \in \mathbb{Z}$, define $KC_k(Y; R)$ to be the R -module of finite R -linear combinations of isomorphism classes $[X, \mathbf{f}, \mathbf{G}]$ for which $\text{vdim } X = k$, with the relations:

- (i) Let $[X, \mathbf{f}, \mathbf{G}]$ be an isomorphism class, and write $-X$ for X with the opposite orientation. Then in $KC_k(Y; R)$ we have

$$[X, \mathbf{f}, \mathbf{G}] + [-X, \mathbf{f}, \mathbf{G}] = 0. \quad (62)$$

- (ii) Let $[X, \mathbf{f}, \mathbf{G}]$ be an isomorphism class, and suppose there exists an isomorphism $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \mathbf{G}) \rightarrow (X, \mathbf{f}, \mathbf{G})$ in the sense of Definition 3.8, such that $\mathbf{a}$ reverses the orientation of X . Then

$$[X, \mathbf{f}, \mathbf{G}] = 0 \quad \text{in } KC_k(Y; R). \quad (63)$$

- (iii) Let $[X, \mathbf{f}, \mathbf{G}]$ be an isomorphism class, and let $X = X_1 \amalg X_2 \amalg \cdots \amalg X_n$ be a *splitting* of $(X, \mathbf{f}, \mathbf{G})$, in the sense of Definition 3.8. Then

$$[X, \mathbf{f}, \mathbf{G}] = \sum_{a=1}^n [X_a, \mathbf{f}|_{X_a}, \mathbf{G}|_{X_a}] \quad \text{in } KC_k(Y; R). \quad (64)$$

- (iv) Let $[X, \mathbf{f}, \mathbf{G}]$ be an isomorphism class, Γ a finite group, and ρ an action of Γ on $(X, \mathbf{f}, \mathbf{G})$ by orientation-preserving automorphisms. That is, $\rho : \Gamma \rightarrow \text{Aut}(X, \mathbf{f}, \mathbf{G})$ is a group morphism, and if $\rho(\gamma) = (\mathbf{a}, \mathbf{b})$ for $\gamma \in \Gamma$ then $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \mathbf{G}) \rightarrow (X, \mathbf{f}, \mathbf{G})$ is an isomorphism with $\mathbf{a} : X \rightarrow X$ orientation-preserving. Note that we do not require ρ to be injective, so we cannot regard Γ as a subgroup of $\text{Aut}(X, \mathbf{f}, \mathbf{G})$. Then Γ acts on X , and $\tilde{X} = X/\Gamma$ is a compact oriented Kuranishi space, with a projection $\pi : X \rightarrow \tilde{X}$. As in Property 3.1(c), Definition 3.21 gives strongly smooth

$\pi_*(\mathbf{f}) = \tilde{\mathbf{f}} : \tilde{X} \rightarrow Y$ and gauge-fixing data $\pi_*(\mathbf{G}) = \tilde{\mathbf{G}}$ for $(\tilde{X}, \tilde{\mathbf{f}})$. Thus $[X/\Gamma, \pi_*(\mathbf{f}), \pi_*(\mathbf{G})]$ is an isomorphism class. We impose the relation

$$[X/\Gamma, \pi_*(\mathbf{f}), \pi_*(\mathbf{G})] = \frac{1}{|\Gamma|} [X, \mathbf{f}, \mathbf{G}] \quad \text{in } KC_k(Y; R). \quad (65)$$

Elements of $KC_k(Y; R)$ will be called *Kuranishi chains*. We require R to be a $\mathbb{Q}$ -algebra so that the factor $\frac{1}{|\Gamma|}$ in (65) makes sense.

Define the *boundary operator* $\partial : KC_k(Y; R) \rightarrow KC_{k-1}(Y; R)$ by

$$\partial : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a] \mapsto \sum_{a \in A} \rho_a [\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a}], \quad (66)$$

where A is a finite indexing set and $\rho_a \in R$ for $a \in A$. This is a morphism of R -modules. Clearly, ∂ takes each relation (62)–(65) in $KC_k(Y; R)$ to the corresponding relation in $KC_{k-1}(Y; R)$, and so ∂ is well-defined.

Recall from Definition 2.27 that if X is an oriented Kuranishi space then there is a natural orientation-reversing strong diffeomorphism $\sigma : \partial^2 X \rightarrow \partial^2 X$, with $\sigma^2 = \text{id}_{\partial^2 X}$. If $[X, \mathbf{f}, \mathbf{G}]$ is an isomorphism class then this σ extends to an isomorphism (σ, τ) of $(\partial^2 X, \mathbf{f}|_{\partial^2 X}, \mathbf{G}|_{\partial^2 X})$, where J is the indexing set of $\mathbf{I}|_{\partial^2 X}$, and for each $j \in J$, $(\tau^j, \hat{\tau}^j)$ is the automorphism of $(\partial^2 V^{j+2}, \dots, \psi^{j+2}|_{\partial^2 V^{j+2}})$ induced by the orientation-reversing involution $\sigma : \partial^2 V^{j+2} \rightarrow \partial^2 V^{j+2}$ described in Definitions 2.7 and 2.9. So part (ii) in $KC_{k-2}(Y; R)$ yields

$$[\partial^2 X, \mathbf{f}|_{\partial^2 X}, \mathbf{G}|_{\partial^2 X}] = 0 \quad \text{in } KC_{k-2}(Y; R).$$

Clearly this implies that $\partial \circ \partial = 0$ as a map $KC_k(Y; R) \rightarrow KC_{k-2}(Y; R)$.

Define the *Kuranishi homology group* $KH_k(Y; R)$ of Y for $k \in \mathbb{Z}$ to be

$$KH_k(Y; R) = \frac{\text{Ker}(\partial : KC_k(Y; R) \rightarrow KC_{k-1}(Y; R))}{\text{Im}(\partial : KC_{k+1}(Y; R) \rightarrow KC_k(Y; R))}.$$

This is a well-defined R -module, as $\partial \circ \partial = 0$.

Note that if 2 is invertible in R , as it is when R is a $\mathbb{Q}$ -algebra, then Definition 4.3(ii) follows from Definition 4.3(i), since $(\mathbf{a}, \mathbf{b})$ in (ii) implies the isomorphism classes $[X, \mathbf{f}, \mathbf{G}]$ and $[-X, \mathbf{f}, \mathbf{G}]$ are equal, so (62) gives $2[X, \mathbf{f}, \mathbf{G}] = 0$, and multiplying by $\frac{1}{2}$ gives (63). But we state it as a separate axiom as we shall re-use it in our next definition, in which 2 need not be invertible in R .

Definition 4.4. Let Y be an orbifold. Consider triples $(X, \mathbf{f}, \underline{\mathbf{G}})$, where X is a compact, oriented, *effective* Kuranishi space, $\mathbf{f} : X \rightarrow Y$ is strongly smooth, and $\underline{\mathbf{G}}$ is *effective* gauge-fixing data for $(X, \mathbf{f})$. Write $[X, \mathbf{f}, \underline{\mathbf{G}}]$ for the isomorphism class of $(X, \mathbf{f}, \underline{\mathbf{G}})$ under isomorphisms $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \underline{\mathbf{G}}) \rightarrow (\tilde{X}, \tilde{\mathbf{f}}, \tilde{\underline{\mathbf{G}}})$, where $\mathbf{a}$ identifies the orientations of $X, \tilde{X}$.

Let R be a commutative ring, for instance $R = \mathbb{Z}, \mathbb{Z}_p, \mathbb{Q}, \mathbb{R}$ or $\mathbb{C}$. For each $k \in \mathbb{Z}$, define $KC_k^{\text{ef}}(Y; R)$ to be the R -module of finite R -linear combinations of isomorphism classes $[X, \mathbf{f}, \underline{\mathbf{G}}]$ as above for which $\text{vdim } X = k$, with the

analogues of relations Definition 4.3(i)–(iii), but *not* Definition 4.3(iv). Elements of $KC_k^{\text{ef}}(Y; R)$ will be called *effective Kuranishi chains*.

Define the *boundary operator* $\partial : KC_k^{\text{ef}}(Y; R) \rightarrow KC_{k-1}^{\text{ef}}(Y; R)$ by

$$\partial : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a] \mapsto \sum_{a \in A} \rho_a [\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a}],$$

as in (66). Then $\partial \circ \partial = 0$. Define the *effective Kuranishi homology group* $KH_k^{\text{ef}}(Y; R)$ of Y for $k \in \mathbb{Z}$ to be

$$KH_k^{\text{ef}}(Y; R) = \frac{\text{Ker}(\partial : KC_k^{\text{ef}}(Y; R) \rightarrow KC_{k-1}^{\text{ef}}(Y; R))}{\text{Im}(\partial : KC_{k+1}^{\text{ef}}(Y; R) \rightarrow KC_k^{\text{ef}}(Y; R))}.$$

Remark 4.5. (a) We define $KC_k, KH_k, KC_k^{\text{ef}}, KH_k^{\text{ef}}(Y; R)$ for all $k \in \mathbb{Z}$, not just for $k \geq 0$. This is because nontrivial Kuranishi spaces X can have $\text{vdim } X = k$ for any $k \in \mathbb{Z}$. Theorem 4.8 and Corollary 4.10 will show that $KH_k(Y; R) = KH_k^{\text{ef}}(Y; R) = 0$ when $k < 0$, so we may as well restrict to $k \geq 0$ when considering homology. However, at the chain level we need to allow $k < 0$, as $KC_k, KC_k^{\text{ef}}(Y; R)$ can be nonzero for any $k \in \mathbb{Z}$. To define $KH_0, KH_0^{\text{ef}}(Y; R)$ we use $\partial : KC_0(Y; R) \rightarrow KC_{-1}(Y; R)$ and $\partial : KC_0^{\text{ef}}(Y; R) \rightarrow KC_{-1}^{\text{ef}}(Y; R)$, so we need to consider -1 -chains. Also, the inductive proofs in Appendices B and C can implicitly involve Kuranishi chains of arbitrarily negative degree.

(b) The relation Definition 4.3(iv) in $KC_*(Y; R)$ is not necessary to get a well-behaved homology theory, and we omit it in the definition of $KC_*^{\text{ef}}(Y; R)$. We include it because it will be useful in some of our applications, in particular, the open Gromov–Witten invariants to be discussed in §6.7 and [39], for which we need it to prove equation (170) of Corollary 6.30. The author believes that Theorem 4.9 below would still hold if we defined $KC_*(Y; R)$ omitting Definition 4.3(iv), and in Remark C.1(a) we sketch how to modify the proof in this case.

Note that if we omit Definition 4.3(iv) then the definitions of $KC_*(Y; R)$ and $KH_*(Y; R)$ make sense for R an arbitrary commutative ring, rather than a $\mathbb{Q}$ -algebra, since we no longer need R to contain the factors $|\Gamma|^{-1}$ in (65). However, as discussed in Remark C.1(b), we would still need R a $\mathbb{Q}$ -algebra to prove Theorem 4.9. In fact the author has an outline proof that if we defined $KC_*(Y; R), KH_*(Y; R)$ omitting Definition 4.3(iv) for R a general commutative ring, then $KH_*(Y; R) \cong H_*^{\text{si}}(Y; R \otimes_{\mathbb{Z}} \mathbb{Q})$.

(c) One could probably weaken the condition that X is *effective* in effective Kuranishi chains $[X, \mathbf{f}, \mathbf{G}]$, and still get a homology theory with $KH_*^{\text{ef}}(Y; R) \cong H_*^{\text{si}}(Y; R)$ for all commutative rings R , including $R = \mathbb{Z}$. In §5.6 we will define the *orbifold strata* $X^{\Gamma, \rho}$ of a Kuranishi space X , and in Remark 5.20 we will show that if X is an oriented, *effective* Kuranishi space then $\text{vdim } X^{\Gamma, \rho} \leq \text{vdim } X - 2$ for all nonempty orbifold strata $X^{\Gamma, \rho}$ with $\Gamma \neq \{1\}$.

For a compact, oriented Kuranishi space X without boundary, the condition that $\text{vdim } X^{\Gamma, \rho} \leq \text{vdim } X - 2$ for all $X^{\Gamma, \rho} \neq \emptyset$ with $\Gamma \neq \{1\}$ is enough to ensure that the virtual class of X can be defined in homology over $\mathbb{Z}$, not just over $\mathbb{Q}$. This holds because if we take a generic (single-valued) perturbation $\tilde{X}$ of

such a Kuranishi space X , although $\tilde{X}$ may not be a manifold or orbifold, the singularities of $\tilde{X}$ due to the orbifold strata $X^{\Gamma, \rho}$ occur in codimension at least 2, so $\tilde{X}$ is a *pseudo-manifold*, and thus has an integral fundamental class. This is the important fact underlying the definition of Gromov–Witten invariants in *integral* homology for *semi-positive* symplectic manifolds, as discussed briefly in §6.3, and proved in McDuff and Salamon [56, §7].

It seems likely that one could define some modification of $KH_*^{\text{ef}}(Y; R)$ isomorphic to $H_*^{\text{si}}(Y; R)$ for all commutative rings R , with chains $[X, \mathbf{f}, \underline{\mathbf{G}}]$ in which we require $\text{vdim } X^{\Gamma, \rho} \leq \text{vdim } X - 2$ for all $X^{\Gamma, \rho} \neq \emptyset$ with $\Gamma \neq \{1\}$, instead of requiring X to be effective. This modification might then have interesting applications, for instance in defining Lagrangian Floer cohomology over $\mathbb{Z}$ for semi-positive symplectic manifolds. But we will not attempt this here.

4.3 Morphisms between singular and Kuranishi homology

There is a natural projection $\Pi_{\text{ef}}^{\text{Kh}}$ from effective Kuranishi chains and homology to Kuranishi chains and homology. Also, (effective) Kuranishi chains and homology have functorial pushforwards h_* by smooth maps $h : Y \rightarrow Z$.

Definition 4.6. Let Y be an orbifold, and R a commutative ring. Define

$$\begin{aligned} \Pi_{\text{ef}}^{\text{Kh}} : KC_k^{\text{ef}}(Y; R) &\longrightarrow KC_k(Y; R \otimes_{\mathbb{Z}} \mathbb{Q}) \quad \text{by} \\ \Pi_{\text{ef}}^{\text{Kh}} : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a] &\longmapsto \sum_{a \in A} \pi(\rho_a) [X_a, \mathbf{f}_a, \Pi(\underline{\mathbf{G}}_a)], \end{aligned} \quad (67)$$

for $k \in \mathbb{Z}$, where $\pi : R \rightarrow R \otimes_{\mathbb{Z}} \mathbb{Q}$ is the natural ring morphism, and $\Pi(\underline{\mathbf{G}}_a)$ is as in Definition 3.36. If R is a $\mathbb{Q}$ -algebra then $R \otimes_{\mathbb{Z}} \mathbb{Q} = R$ and π is the identity. The $\Pi_{\text{ef}}^{\text{Kh}}$ map relations in $KC_k^{\text{ef}}(Y; R)$ to relations in $KC_k(Y; R \otimes_{\mathbb{Z}} \mathbb{Q})$, and so are well-defined. They satisfy $\Pi_{\text{ef}}^{\text{Kh}} \circ \partial = \partial \circ \Pi_{\text{ef}}^{\text{Kh}}$, so they induce morphisms

$$\Pi_{\text{ef}}^{\text{Kh}} : KH_k^{\text{ef}}(Y; R) \longrightarrow KH_k(Y; R \otimes_{\mathbb{Z}} \mathbb{Q}). \quad (68)$$

Let Y, Z be orbifolds, and $h : Y \rightarrow Z$ a smooth map. Using the notation of Definitions 3.26 and 3.41, define the *pushforwards*

$$\begin{aligned} h_* : KC_k(Y; R) &\longrightarrow KC_k(Z; R), \quad h_* : KC_k^{\text{ef}}(Y; R) \longrightarrow KC_k^{\text{ef}}(Z; R) \\ \text{by } h_* : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a] &\longmapsto \sum_{a \in A} \rho_a [X_a, h \circ \mathbf{f}_a, h_*(\underline{\mathbf{G}}_a)] \\ \text{and } h_* : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a] &\longmapsto \sum_{a \in A} \rho_a [X_a, h \circ \mathbf{f}_a, h_*(\underline{\mathbf{G}}_a)]. \end{aligned} \quad (69)$$

These take relations (i)–(iv) or (i)–(iii) in $KC_k, KC_k^{\text{ef}}(Y; R)$ to relations (i)–(iv) or (i)–(iii) in $KC_k, KC_k^{\text{ef}}(Z; R)$, and so are well-defined. They satisfy $h_* \circ \partial = \partial \circ h_*$, so they induce morphisms of homology groups

$$h_* : KH_k(Y; R) \longrightarrow KH_k(Z; R) \text{ and } h_* : KH_k^{\text{ef}}(Y; R) \longrightarrow KH_k^{\text{ef}}(Z; R).$$

Pushforward is functorial, that is, $(g \circ h)_* = g_* \circ h_*$, on chains and homology.

Definition 4.7. Let Y be an orbifold, Δ_k be the k -simplex of (59), and $\sigma : \Delta_k \rightarrow Y$ be a smooth map. Then Δ_k is a compact oriented manifold with corners, so we may regard it as an oriented Kuranishi space, and σ as a strongly smooth map.

Define an excellent coordinate system $(\mathbf{I}_{\Delta_k}, \boldsymbol{\eta}_{\Delta_k})$ for (Δ_k, σ) to have indexing set $I_{\Delta_k} = \{k\}$, Kuranishi neighbourhood $(V_{\Delta_k}^k, E_{\Delta_k}^k, s_{\Delta_k}^k, \psi_{\Delta_k}^k) = (\Delta_k, \Delta_k, 0, \text{id}_{\Delta_k})$, and map $\sigma^k : V_{\Delta_k}^k \rightarrow Y$ representing σ given by $\sigma^k = \sigma$. Here $E_{\Delta_k}^k \rightarrow V_{\Delta_k}^k$ is the zero vector bundle, so that as manifolds $E_{\Delta_k}^k = V_{\Delta_k}^k = \Delta_k$. Set $\boldsymbol{\eta}_{\Delta_k} = (\eta_{k, \Delta_k}, \eta_{k, \Delta_k}^k)$, where $\eta_{k, \Delta_k} \equiv 1$ on Δ_k and $\eta_{k, \Delta_k}^k \equiv 1$ on $V_{\Delta_k}^k$. Define $G_{\Delta_k}^k : E_{\Delta_k}^k \rightarrow P$ and $\underline{G}_{\Delta_k}^k : E_{\Delta_k}^k \rightarrow \underline{P}$ by

$$\begin{aligned} G_{\Delta_k}^k((x_0, \dots, x_k)) &= S_{l+1} \cdot (y_0, y_0 + y_1, \dots, y_0 + y_1 + \dots + y_l), \\ \underline{G}_{\Delta_k}^k((x_0, \dots, x_k)) &= (y_0, y_0 + y_1, \dots, y_0 + y_1 + \dots + y_l), \end{aligned} \quad (70)$$

where $y_0, y_1, \dots, y_l$ are those $x_0, x_1, \dots, x_k$ which are nonzero, in the same order, so that $(x_0, \dots, x_k)$ is $(y_0, y_1, \dots, y_l)$ with $k-l$ zeros inserted in the list. Then

$$G_{\Delta_k}^k \circ F_j^k \equiv G_{\Delta_{k-1}}^{k-1}, \quad \underline{G}_{\Delta_k}^k \circ F_j^k \equiv \underline{G}_{\Delta_{k-1}}^{k-1} \text{ for } j = 0, \dots, k, \quad (71)$$

with F_j^k as in §4.1. Also, for $(x_0, \dots, x_k) \in \Delta_k^\circ$ we have $x_j > 0$ for all j , so that $(y_0, \dots, y_l) = (x_0, \dots, x_k)$, and $G_{\Delta_k}^k : (x_0, \dots, x_k) \mapsto S_{k+1} \cdot (x_0, \dots, x_0 + \dots + x_k)$, and $\underline{G}_{\Delta_k}^k : (x_0, \dots, x_k) \mapsto (x_0, \dots, x_0 + \dots + x_k)$. Since $x_0, x_0 + x_1, \dots, x_0 + \dots + x_k$ are arranged in strictly increasing order, $S_{k+1} \cdot (x_0, \dots, x_0 + \dots + x_k)$ determines $x_0, x_0 + x_1, \dots, x_0 + \dots + x_k$ and hence $x_0, \dots, x_k$. Thus $G_{\Delta_k}^k|_{\Delta_k^\circ}$ is injective. Similarly, $\underline{G}_{\Delta_k}^k|_{\Delta_k^\circ}$ maps $\Delta_k^\circ \rightarrow \mathbb{R}^{k+1} \subset \underline{P}$, and is injective. It is now easy to check that Definitions 3.8 and 3.36 hold, so that $\mathbf{G}_{\Delta_k}$ is *gauge-fixing data*, and $\underline{\mathbf{G}}_{\Delta_k}$ is *effective gauge-fixing data*, for (Δ_k, σ) .

Taking R to be a $\mathbb{Q}$ -algebra in (72), define maps

$$\Pi_{\text{si}}^{\text{Kh}} : C_k^{\text{si}}(Y; R) \rightarrow KC_k(Y; R), \quad \Pi_{\text{si}}^{\text{Kh}} : \sum_{a \in A} \rho_a \sigma_a \mapsto \sum_{a \in A} \rho_a [\Delta_k, \sigma_a, \mathbf{G}_{\Delta_k}], \quad (72)$$

$$\Pi_{\text{si}}^{\text{ef}} : C_k^{\text{si}}(Y; R) \rightarrow KC_k^{\text{ef}}(Y; R), \quad \Pi_{\text{si}}^{\text{ef}} : \sum_{a \in A} \rho_a \sigma_a \mapsto \sum_{a \in A} \rho_a [\Delta_k, \sigma_a, \underline{\mathbf{G}}_{\Delta_k}]. \quad (73)$$

As $\partial \Delta_k = \coprod_{j=1}^k (-1)^j F_j^k(\Delta_{k-1})$ in oriented $(k-1)$ -manifolds, (71) gives

$$\partial[\Delta_k, \sigma_a, \mathbf{G}_{\Delta_k}] = \sum_{j=0}^k (-1)^j [\Delta_{k-1}, (\sigma_a \circ F_j^k), \mathbf{G}_{\Delta_{k-1}}] \quad (74)$$

in $KC_{k-1}(Y; R)$. Comparing (60), (66) and (74) shows that $\partial \circ \Pi_{\text{si}}^{\text{Kh}} = \Pi_{\text{si}}^{\text{Kh}} \circ \partial$. Similarly $\partial \circ \Pi_{\text{si}}^{\text{ef}} = \Pi_{\text{si}}^{\text{ef}} \circ \partial$. Therefore (72)–(73) induce projections

$$\Pi_{\text{si}}^{\text{Kh}} : H_k^{\text{si}}(Y; R) \longrightarrow KH_k(Y; R), \quad (75)$$

$$\Pi_{\text{si}}^{\text{ef}} : H_k^{\text{si}}(Y; R) \longrightarrow KH_k^{\text{ef}}(Y; R), \quad (76)$$

requiring R to be a $\mathbb{Q}$ -algebra in (75). The morphisms (67), (72), (73) on chains, and (68), (75), (76) on homology, satisfy

$$\Pi_{\text{ef}}^{\text{Kh}} \circ \Pi_{\text{si}}^{\text{ef}} = \Pi_{\text{si}}^{\text{Kh}}. \quad (77)$$

Our next two theorems, the most important of the book, will be proved in Appendices B and C respectively, after preparatory work in Appendix A.

Theorem 4.8. *Let Y be an orbifold and R a commutative ring. Then the morphism $\Pi_{\text{si}}^{\text{ef}} : H_k^{\text{si}}(Y; R) \rightarrow KH_k^{\text{ef}}(Y; R)$ in (76) is an isomorphism for $k \geq 0$, with $KH_k^{\text{ef}}(Y; R) = 0$ when $k < 0$.*

Theorem 4.9. *Let Y be an orbifold and R a $\mathbb{Q}$ -algebra. Then the morphism $\Pi_{\text{ef}}^{\text{Kh}} : KH_k^{\text{ef}}(Y; R) \rightarrow KH_k(Y; R)$ in (68) is an isomorphism for $k \in \mathbb{Z}$.*

These and equation (77) imply:

Corollary 4.10. *Let Y be an orbifold and R a $\mathbb{Q}$ -algebra. Then the morphism $\Pi_{\text{si}}^{\text{Kh}} : H_k^{\text{si}}(Y; R) \rightarrow KH_k(Y; R)$ in (75) is an isomorphism for $k \geq 0$, with $KH_k(Y; R) = 0$ when $k < 0$.*

Remark 4.11. (a) Theorems 4.8 and 4.9 are our substitute for results of Fukaya and Ono on *multisections* and *virtual cycles* [25, §3, §6], [24, §A1]. If X is a compact Kuranishi space, Y an orbifold, and $\mathbf{f} : X \rightarrow Y$ is strongly smooth, then by choosing an abstract, multivalued perturbation of X called a multisection which locally turns X into a (non-Hausdorff) smooth manifold X^{per} , with weights in $\mathbb{Q}$, and then triangulating X^{per} by simplices, Fukaya and Ono produce a singular chain $C(X, \mathbf{f}) \in C_{\text{vdim } X}^{\text{si}}(Y; \mathbb{Q})$ called the *virtual chain*, depending on many choices. If $\partial X = \emptyset$ then $\partial C(X, \mathbf{f}) = 0$, and $[C(X, \mathbf{f})] \in H_{\text{vdim } X}^{\text{si}}(Y; \mathbb{Q})$ is independent of choices.

The relation of this to Theorems 4.8 and 4.9 and Corollary 4.10 is that if $\partial X = \emptyset$ then choosing any gauge-fixing data $\mathbf{G}$ for $(X, \mathbf{f})$, we have a class $[X, \mathbf{f}, \mathbf{G}] \in KH_{\text{vdim } X}(Y; \mathbb{Q})$, and $\Pi_{\text{si}}^{\text{Kh}} : [C(X, \mathbf{f})] \mapsto [X, \mathbf{f}, \mathbf{G}]$. That is, in proving $\Pi_{\text{si}}^{\text{Kh}}$ is invertible we construct an inverse $(\Pi_{\text{si}}^{\text{Kh}})^{-1} : KH_*(Y; \mathbb{Q}) \rightarrow H_*^{\text{si}}(Y; \mathbb{Q})$. Very roughly speaking, at the chain level $(\Pi_{\text{si}}^{\text{Kh}})^{-1}$ takes $[X, \mathbf{f}, \mathbf{G}]$ to $C(X, \mathbf{f})$, Fukaya and Ono's virtual chain for $(X, \mathbf{f})$.

In proving Theorems 4.8 and 4.9 in Appendices B and C, we encounter most of the problems Fukaya and Ono were tackling with their virtual cycle technology, and we freely borrow their ideas for our proof, in particular, good coordinate systems, and inductive choices of small perturbations. But we do not use multisections: instead we use Definition 4.3(iv) to represent each Kuranishi homology class by a sum of chains $[X, \mathbf{f}, \mathbf{G}]$ in which X has trivial stabilizer groups, so that we can lift to effective Kuranishi homology, and then we can perturb each X to a manifold using single-valued perturbations.

(b) The proofs of Theorems 4.8 and 4.9 use almost no properties of the target space Y . In the proofs, given some $[X, \mathbf{f}, \mathbf{G}]$ or $[X, \mathbf{f}, \underline{\mathbf{G}}]$, we have a cover of X by Kuranishi neighbourhoods (V^i, E^i, s^i, ψ^i) with maps $f^i : V^i \rightarrow Y$ and $G^i : E^i \rightarrow P$ or $\underline{G}^i : E^i \rightarrow \underline{P}$. All we require of these maps f^i is that they satisfy $f^j|_{V^{ij}} \equiv f^i \circ \phi^{ij}$ when $j \leq i$ in I . The proofs involve cutting V^i into pieces, changing $G^i, \underline{G}^i$, and deforming s^i , which changes X as $X \cong (s^i)^{-1}(0)$, but we never change the maps f^i .

Because of this, Theorems 4.8 and 4.9 and their proofs would still hold, essentially unchanged, if we took Y to be an *arbitrary topological space*, rather

than an orbifold, and maps $\mathbf{f} : X \rightarrow Y$ in chains $[X, \mathbf{f}, \mathbf{G}]$ or $[X, \mathbf{f}, \mathbf{\underline{G}}]$ to be *strongly continuous*, rather than strongly smooth. We will need Y an orbifold and $\mathbf{f}$ strongly smooth or a strong submersion in §4.7 to define *products* $\cup, \cap, \bullet$ on Kuranishi (co)homology, but to establish isomorphism with singular homology, this is unnecessary.

In the same way, Theorems 4.8 and 4.9 and their proofs will still hold, essentially unchanged, if we take Y to be an *infinite-dimensional* manifold or orbifold, such as a *loop space* $\mathcal{LM}$ of smooth maps $\lambda : S^1 \rightarrow M$ for M a finite-dimensional manifold, and $\mathbf{f} : X \rightarrow Y$ to be strongly smooth in the appropriate sense. (That is, we prove that $KH_*, KH_*^{\text{ef}}(Y; R)$ are isomorphic to singular homology $H_*^{\text{si}}(Y; R)$ defined using smooth singular simplices in Y .) This will be important in [40], when we apply Kuranishi (co)homology to the *String Topology* programme of Chas and Sullivan [11].

We will define a homology theory which computes $H_*^{\text{si}}(\mathcal{LM}, M; \mathbb{Q})$, the singular homology of $\mathcal{LM}$ relative to constant loops $M \subset \mathcal{LM}$, and on which Chas and Sullivan's String Topology operations are defined *at the chain level*, without any transversality assumptions, and satisfy the desired identities *exactly* at the chain level, not just up to homotopy. We will use this to prove results sketched by Fukaya [23, §6], which use moduli spaces of J -holomorphic discs in a symplectic manifold (M, ω) with boundary in a Lagrangian L to define chains in our homology theory of the loop space of L satisfying identities involving String Topology operations.

4.4 Kuranishi cohomology and Poincaré duality for $\partial Y = \emptyset$

Here is our dual notion of *Kuranishi cohomology* of an orbifold Y . The differences with Kuranishi homology are that we replace strongly smooth maps by strong submersions, and orientations by coorientations, and gauge-fixing data by co-gauge-fixing data, and we grade $[X, \mathbf{f}, \mathbf{C}]$ by codimension $\dim Y - \text{vdim } X$ rather than by dimension $\text{vdim } X$. In this section we restrict to Y *without boundary*. The extension to $\partial Y \neq \emptyset$ will be explained in §4.5.

Definition 4.12. Let Y be an orbifold without boundary. Consider triples $(X, \mathbf{f}, \mathbf{C})$, where X is a compact Kuranishi space, $\mathbf{f} : X \rightarrow Y$ is a strong submersion with $(X, \mathbf{f})$ cooriented, and $\mathbf{C}$ is co-gauge-fixing data for $(X, \mathbf{f})$. Write $[X, \mathbf{f}, \mathbf{C}]$ for the isomorphism class of $(X, \mathbf{f}, \mathbf{C})$ under isomorphisms $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \mathbf{C}) \rightarrow (\tilde{X}, \tilde{\mathbf{f}}, \tilde{\mathbf{C}})$ as in Definition 3.9, where $\mathbf{a}$ must identify the coorientations of $(X, \mathbf{f}), (\tilde{X}, \tilde{\mathbf{f}})$.

Let R be a $\mathbb{Q}$ -algebra. For $k \in \mathbb{Z}$, define $KC^k(Y; R)$ to be the R -module of finite R -linear combinations of isomorphism classes $[X, \mathbf{f}, \mathbf{C}]$ for which $\text{vdim } X = \dim Y - k$, with the analogues of relations Definition 4.3(i)–(iv), replacing gauge-fixing data $\mathbf{G}$ by co-gauge-fixing data $\mathbf{C}$. Elements of $KC^k(Y; R)$ are called *Kuranishi cochains*. Define $d : KC^k(Y; R) \rightarrow KC^{k+1}(Y; R)$ by

$$d : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{C}_a] \longmapsto \sum_{a \in A} \rho_a [\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{C}_a|_{\partial X_a}]. \quad (78)$$

As in Definition 4.3 we have $d \circ d = 0$. Define the *Kuranishi cohomology groups*

$KH^k(Y; R)$ of Y for $k \in \mathbb{Z}$ to be

$$KH^k(Y; R) = \frac{\text{Ker}(d : KC^k(Y; R) \rightarrow KC^{k+1}(Y; R))}{\text{Im}(d : KC^{k-1}(Y; R) \rightarrow KC^k(Y; R))}.$$

Let Y, Z be orbifolds without boundary, and $h : Y \rightarrow Z$ be a smooth, proper map. Using the notation of Definition 3.27, define the *pullback* $h^* : KC^k(Z; R) \rightarrow KC^k(Y; R)$ by

$$h^* : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{C}_a] \mapsto \sum_{a \in A} \rho_a [Y \times_{h, Z, \mathbf{f}_a} X_a, \pi_Y, h^*(\mathbf{C}_a)]. \quad (79)$$

Here the coorientation for $(X_a, \mathbf{f}_a)$ pulls back to a coorientation for $(Y \times_{h, Z, \mathbf{f}_a} X_a, \pi_Y)$, as follows. Let $p \in X_a$ and (V_p, E_p, s_p, ψ_p) be a sufficiently small Kuranishi neighbourhood in the germ at p , with submersion $f_p : V_p \rightarrow Z$ representing $\mathbf{f}$. Then the coorientation for $(X, \mathbf{f})$ gives orientations on the fibres of the orbibundle $(\text{Ker } df_p \oplus E_p) \rightarrow V_p$. As in (16), we lift $(V_p, \dots, \psi_p)$ to a Kuranishi neighbourhood

$$(V_p^Y, E_p^Y, s_p^Y, \psi_p^Y) = (Y \times_{h, Z, f_p} V_p, \pi_{V_p}^*(E_p), s_p \circ \pi_{V_p}, \pi_Y \times (\psi_p \circ \pi_{V_p}) \times \chi)$$

on $Y \times_Z X_a$. Then the orbibundle $(\text{Ker } d\pi_Y \oplus E_p^Y) \rightarrow V_p^Y$ is naturally isomorphic to $\pi_Y^*(\text{Ker } df_p \oplus E_p) \rightarrow V_p$, so the orientations on the fibres of $(\text{Ker } df_p \oplus E_p) \rightarrow V_p$ lift to orientations on the fibres of $(\text{Ker } d\pi_Y \oplus E_p^Y) \rightarrow V_p^Y$, which define a *coorientation* for $(Y \times_Z X_a, \pi_Y)$.

These $h^* : KC^k(Z; R) \rightarrow KC^k(Y; R)$ satisfy $h^* \circ d = d \circ h^*$, so they induce morphisms of cohomology groups $h^* : KH^k(Z; R) \rightarrow KH^k(Y; R)$. Pullbacks are functorial, that is, $(g \circ h)^* = h^* \circ g^*$, on both cochains and cohomology.

Remark 4.13. In Kuranishi cochains $[X, \mathbf{f}, \mathbf{C}]$ we assume X is compact, so that $f(X) \subseteq Y$ is also compact. Thus, our cochains are *compactly-supported*, and $KH^*(Y; R)$ is a form of *compactly-supported cohomology*.

One could also try to define versions of Kuranishi cohomology analogous to ordinary cohomology. The chains should be $[X, \mathbf{f}, \mathbf{C}]$ for X a *not necessarily compact* Kuranishi space, and $\mathbf{f} : X \rightarrow Y$ a *proper* strong submersion. The notion of co-gauge-fixing data $\mathbf{C}$ for $(X, \mathbf{f})$ when X is noncompact would certainly need revision. Working with noncompact X raises important issues to do with finiteness — finiteness of indexing sets for good coordinate systems, finiteness of $\text{Aut}(X, \mathbf{f}, \mathbf{G})$ in Theorem 3.20 (this relies on Proposition 3.19, which uses the compactness of X), and so on. As we do not need noncompact Kuranishi spaces in any of our applications [4, 39–41], we will not develop this.

We shall prove *Poincaré duality* for Kuranishi (co)homology. We do not assume Y is compact.

Definition 4.14. Let Y be an orbifold which is oriented, without boundary, and of dimension n , and R a $\mathbb{Q}$ -algebra. Then for $\mathbf{f} : X \rightarrow Y$ a strong submersion, as in Definition 2.32 there is a 1-1 correspondence between coorientations for $(X, \mathbf{f})$ and orientations for X . Choose an injective map $G_Y : Y \rightarrow P_m \subset P$,

which is possible for $m \gg 0$. If $\mathbf{C} = (\mathbf{I}, \boldsymbol{\eta}, C^i : i \in I)$ is co-gauge-fixing data for $(X, \mathbf{f})$ with $\mathbf{I} = (I, (V^i, \dots, \psi^i), f^i : i \in I, \dots)$, define $\mathbf{G}_{\mathbf{C}} = (\mathbf{I}, \boldsymbol{\eta}, G^i : i \in I)$, where $G^i : E^i \rightarrow P$ is given by $G^i = \mu \circ ((G_Y \circ f^i \circ \pi^i) \times C^i)$, for μ in (33). Definition 3.8 for $\mathbf{C}$ says that C^i maps $E^i \rightarrow P_n \subset P$ for some $n \gg 0$, and $C^i \times (f^i \circ \pi^i) : E^i \rightarrow P \times Y$ is globally finite. Together with $G_Y : Y \rightarrow P_m$ injective and $\mu : P_m \times P_n \rightarrow P_{m+n}$ globally finite this implies that G^i maps $E^i \rightarrow P_{m+n}$ and is globally finite, so that $\mathbf{G}_{\mathbf{C}}$ is *gauge-fixing data* for $(X, \mathbf{f})$.

Define R -module morphisms $\Pi_{\text{Kch}}^{\text{Kh}} : KC^k(Y; R) \rightarrow KC_{n-k}(Y; R)$ for $k \in \mathbb{Z}$ by $\Pi_{\text{Kch}}^{\text{Kh}} : [X, \mathbf{f}, \mathbf{C}] \mapsto [X, \mathbf{f}, \mathbf{G}_{\mathbf{C}}]$, using the coorientation for $(X, \mathbf{f})$ from $[X, \mathbf{f}, \mathbf{C}] \in KC^k(Y; R)$ to determine the orientation on X for $[X, \mathbf{f}, \mathbf{G}_{\mathbf{C}}] \in KC_{n-k}(Y; R)$. Then $\Pi_{\text{Kch}}^{\text{Kh}}$ takes Definitions 4.3(i)–(iv) to the same relations, so they are well-defined, and $\partial \circ \Pi_{\text{Kch}}^{\text{Kh}} = \Pi_{\text{Kch}}^{\text{Kh}} \circ d$, so they induce morphisms of (co)homology groups

$$\Pi_{\text{Kch}}^{\text{Kh}} : KH^k(Y; R) \longrightarrow KH_{n-k}(Y; R). \quad (80)$$

We will now construct an inverse $\Pi_{\text{Kch}}^{\text{Kh}}$ for (80). On the (co)chain level, we cannot just define $\Pi_{\text{Kch}}^{\text{Kh}} : [X, \mathbf{f}, \mathbf{G}] \mapsto [X, \mathbf{f}, \mathbf{C}_{\mathbf{G}}]$, turning gauge-fixing data $\mathbf{G}$ into co-gauge-fixing data $\mathbf{C}_{\mathbf{G}}$, since $\mathbf{f}$ may not be a strong submersion. Instead, we *modify the Kuranishi structure of X* to make $\mathbf{f}$ a strong submersion.

Write $\pi : TY \rightarrow Y$ for the tangent orbibundle of Y , and $z : Y \rightarrow TY$ for the zero section. Choose an orbifold Riemannian metric g on Y . Then we can define the *exponential map* $\exp : V \rightarrow Y$, a smooth submersion from an open neighbourhood V of $z(Y)$ in TY , such that for $v \in V$ with $\pi(v) = y$ we have $\exp(v) = \gamma(1)$, where $\gamma : [0, 1] \rightarrow Y$ is the unique geodesic interval with $\gamma(0) = y$, $\dot{\gamma}(0) = v$ and $\nabla_{\dot{\gamma}(t)} \dot{\gamma}(t) = 0$ for $t \in [0, 1]$. Choose V such that $\exp$ extends to the closure $\bar{V}$ of V in TY , and $\pi : \bar{V} \rightarrow Y$ is a *proper* map. This is possible provided V is small enough.

Define a new Kuranishi structure κ_Y on Y by the single Kuranishi neighbourhood $(V, E, s, \psi) = (V, \pi^*(TY), \text{id}_{TY}, z^{-1})$. Here $s = \text{id}_{TY}$ is just the identity map on TY , restricted to V and regarded as a section of the bundle $E = \pi^*(TY) \rightarrow V$. Thus $s^{-1}(0)$ is $z(Y)$, so $\psi = z^{-1} : z(Y) \rightarrow Y$ makes sense. This induces κ_Y by taking the germ of κ_Y at each $p \in Y$ to be the equivalence class of (V, E, s, ψ) , regarded as a Kuranishi neighbourhood at p , and the coordinate changes between such neighbourhoods to be the identity.

Define strong submersions $\boldsymbol{\pi} : (Y, \kappa_Y) \rightarrow Y$ and $\mathbf{exp} : (Y, \kappa_Y) \rightarrow Y$ by $\pi : V \rightarrow Y$ and $\exp : V \rightarrow Y$ on the Kuranishi neighbourhood (V, E, s, ψ) . The continuous maps induced by $\boldsymbol{\pi}$ and $\mathbf{exp}$ are both the identity map $\text{id} : Y \rightarrow Y$, but nonetheless $\boldsymbol{\pi}$ and $\mathbf{exp}$ are different as strongly smooth maps, provided that $\dim Y > 0$. Also, $z : Y \rightarrow V$ induces an embedding $\mathbf{z} : Y \rightarrow (Y, \kappa_Y)$ inducing the identity $\text{id} : Y \rightarrow Y$. Choose an injective map $C_Y : E \rightarrow P_{m'} \subset P$ for some m' , which is possible for $m' \gg 0$.

Suppose X is a compact oriented Kuranishi space, $\mathbf{f} : X \rightarrow Y$ is strongly smooth, and $\mathbf{G}$ is gauge-fixing data for $(X, \mathbf{f})$. We will define modifications $X^Y, \mathbf{f}^Y, \mathbf{C}_{\mathbf{G}}^Y$ of $X, \mathbf{f}, \mathbf{G}$ such that X^Y is a compact Kuranishi space, $\mathbf{f}^Y : X^Y \rightarrow Y$ is a strong submersion with $(X^Y, \mathbf{f}^Y)$ cooriented, and $\mathbf{C}_{\mathbf{G}}^Y$ is co-gauge-fixing

data for $(X^Y, \mathbf{f}^Y)$. Define X^Y to be the fibre product $(Y, \kappa_Y) \times_{\pi, Y, \mathbf{f}} X$. The underlying topological space of X^Y is X , so X^Y is really X with a new Kuranishi structure. Define a strong submersion $\mathbf{f}^Y : X^Y \rightarrow Y$ by $\mathbf{f}^Y = \mathbf{exp} \circ \pi_{(Y, \kappa_Y)}$. There is a natural coorientation on $((Y, \kappa_Y), \pi)$, and combining this with the orientation on X gives an orientation on X^Y , as in Convention 2.33(c). As Y, X^Y are oriented and $\mathbf{f}^Y : X^Y \rightarrow Y$ is a strong submersion, as in Definition 2.32 we obtain a *coorientation* for $(X^Y, \mathbf{f}^Y)$.

Write $\mathbf{G} = (\mathbf{I}, \boldsymbol{\eta}, G^i : i \in I)$ with $\mathbf{I} = (I, (V^i, \dots, \psi^i), f^i : i \in I, \dots)$. Define $I^Y = \{i + n : i \in I\}$. Following (16), for each $i \in I$ define a Kuranishi neighbourhood on X^Y to be

$$(V^{i+n, Y}, E^{i+n, Y}, s^{i+n, Y}, \psi^{i+n, Y}) = (V \times_{\pi, Y, f^i} V^i, \pi_V^*(E) \oplus \pi_{V^i}^*(E^i), (s \circ \pi_V) \oplus (s^i \circ \pi_{V^i}), (\psi \circ \pi_V) \times (\psi^i \circ \pi_{V^i}) \times \chi^i). \quad (81)$$

Define $f^{i+n, Y} : V^{i+n, Y} \rightarrow Y$ by $f^{i+n, Y} = \mathbf{exp} \circ \pi_V$. Define $\eta_{i+n}^Y : X^Y \rightarrow [0, 1]$ by $\eta_{i+n}^Y \equiv \eta_i$, identifying $X^Y \cong X$. For $i, j \in I$, define $\eta_{i+n}^{j+n, Y} : V^{j+n, Y} \rightarrow [0, 1]$ by $\eta_{i+n}^{j+n, Y} \equiv \eta_i^j \circ \pi_{V^j}$. When $j \leq i$ in I , define $V^{(i+n)(j+n), Y} \subseteq V^{j+n, Y}$ to be $V \times_{\pi, Y, f^j} V^{ij}$, and let $(\phi^{(i+n)(j+n), Y}, \hat{\phi}^{(i+n)(j+n), Y}) : (V^{(i+n)(j+n), Y}, \dots, \psi^{j+n, Y}|_{V^{(i+n)(j+n), Y}}) \rightarrow (V^{i+n, Y}, \dots, \psi^{i+n, Y})$ be induced by the identity on (V, E, s, ψ) , and $(\phi^{ij}, \hat{\phi}^{ij})$.

As $(\mathbf{I}, \boldsymbol{\eta})$ is a really good coordinate system for $(X, \mathbf{f})$, it is easy to show $((I^Y, (V^{k, Y}, \dots, \psi^{k, Y}), f^{k, Y} : k \in I^Y, \dots), \eta_k^Y, \eta_k^{l, Y} : k, l \in I^Y)$ is a really good coordinate system for $(X^Y, \mathbf{f}^Y)$. To verify Definition 3.4(vi), let $T \subseteq Y$ be the compact subset for $(\mathbf{I}, \boldsymbol{\eta})$ in Definition 3.4(vi). Then $\pi^{-1}(T) \cap \bar{V}$ is compact as $\pi : \bar{V} \rightarrow Y$ is proper, so $T^Y = \mathbf{exp}(\pi^{-1}(T) \cap \bar{V}) \subseteq Y$ is well-defined as $\mathbf{exp}$ extends to $\bar{V}$, and compact as $\pi^{-1}(T) \cap \bar{V}$ is compact. Since $f^{k-n}(V^{k-n}) \subseteq T$ we find that $f^{k, Y}(V^{k, Y}) \subseteq T^Y$ for all $k \in I^Y$, proving Definition 3.4(vi).

Let $(\tilde{\mathbf{I}}^Y, \tilde{\boldsymbol{\eta}}^Y)$ be the excellent coordinate system for $(X^Y, \mathbf{f}^Y)$ constructed from this by Algorithm 3.16, with $\tilde{\mathbf{I}}^Y = (\tilde{I}^Y, (\tilde{V}^{k, Y}, \dots, \tilde{\psi}^{k, Y}) : k \in \tilde{I}^Y, \dots)$. For each $k \in \tilde{I}^Y$, define $\check{C}^{k, Y} : \check{E}^{k, Y} \rightarrow P$ by $\check{C}^{k, Y} = \mu \circ ((C_Y \circ \pi_E) \times (G^{k-n} \circ \pi_{E^{k-n}}))$. Since C_Y maps $E \rightarrow P_{m'} \subset P$, and G^{k-n} maps $E^{k-n} \rightarrow P_{n'} \subset P$ for some $n' \geq 0$, and μ maps $P_{m'} \times P_{n'} \rightarrow P_{m'+n'}$, it follows that $\check{C}^{k, Y}$ maps $\check{E}^{k, Y} \rightarrow P_{m'+n'}$, one of the conditions of Definition 3.9.

We will prove $\check{C}^{k, Y}$ is *globally finite*. Observe that $\check{C}^{k, Y}$ is the composition

$$\check{E}^{k, Y} \hookrightarrow E \times_Y E^{k-n} \xrightarrow{\pi_E \times \pi_{E^{k-n}}} E \times E^{k-n} \xrightarrow{C_Y \times G^{k-n}} P_{m'} \times P_{n'} \xrightarrow{\mu} P_{m'+n'}. \quad (82)$$

We shall show that each morphism in (82) is globally finite, in the sense of Definition 3.8. The first map in (82) is an inclusion of open sets, and so globally finite with $N = 1$. For the second, let $T \subseteq Y$ be as above. The map $Y \rightarrow \mathbb{N}$ taking $y \mapsto |\text{Stab}(y)|$ is upper semicontinuous, so as T is compact there exists $L \geq 1$ such that $|\text{Stab}(y)| \leq L$ for all $y \in T$. As $f^{k-n}(V^{k-n}) \subseteq T$, in the formula (12) for the orbifold fibre product $E \times_{\pi, Y, f^{k-n} \circ \pi^{k-n}} E^{k-n}$ the biquotient term $f_*(\text{Stab}(p)) \backslash \text{Stab}(f(p)) / f'_*(\text{Stab}(p'))$ is at most L points, so $\pi_E \times \pi_{E^{k-n}}$ is globally finite with $N = L$.

The third morphism $C_Y \times G^{k-n}$ in (82) is globally finite as C_Y is injective and G^{k-n} is globally finite. For the last, (33) implies that $\mu : P_{m'} \times P_{n'} \rightarrow P_{m'+n'}$ is globally finite with $N = \binom{m'+n'}{m'}$. Thus the composition $\tilde{C}^{k,Y}$ is globally finite, and $\mathbf{C}_G^Y = (\tilde{I}^Y, \tilde{\eta}^Y, \tilde{C}^{k,Y} : k \in \tilde{I}^Y)$ is *co-gauge-fixing data* for $(X^Y, \mathbf{f}^Y)$.

Define R -module morphisms $\Pi_{\text{Kh}}^{\text{Kch}} : KC_{n-k}(Y; R) \rightarrow KC^k(Y; R)$ by $\Pi_{\text{Kh}}^{\text{Kch}} : [X, \mathbf{f}, \mathbf{G}] \mapsto [X^Y, \mathbf{f}^Y, \mathbf{C}_G^Y]$. These are well-defined, and $d \circ \Pi_{\text{Kh}}^{\text{Kch}} = \Pi_{\text{Kh}}^{\text{Kch}} \circ \partial$, so they induce morphisms of (co)homology groups

$$\Pi_{\text{Kh}}^{\text{Kch}} : KH_{n-k}(Y; R) \longrightarrow KH^k(Y; R). \quad (83)$$

Theorem 4.15. *Let Y be an oriented n -orbifold without boundary. Then $\Pi_{\text{Kh}}^{\text{Kch}}$ in (83) is the inverse of $\Pi_{\text{Kch}}^{\text{Kh}}$ in (80), so they are both isomorphisms. Also (80), (83) are independent of choices in their constructions.*

Proof. If $\dim Y = 0$ then taking $G_Y \equiv (\emptyset)$, $C_Y \equiv (\emptyset)$ we see that $\Pi_{\text{Kh}}^{\text{Kch}}$ is inverse to $\Pi_{\text{Kch}}^{\text{Kh}}$ at the chain level, and the theorem is trivial. So suppose $n = \dim Y > 0$. Define a Kuranishi structure κ' on $[0, 1] \times Y$ as follows. On $[0, \frac{1}{2}] \times Y$ this comes from the orbifold structure on $[0, 1] \times Y$. On $[\frac{1}{2}, 1] \times Y$ it comes from the product Kuranishi structure on $[0, 1] \times (Y, \kappa_Y)$. Near $\{\frac{1}{2}\} \times Y$ we glue these two Kuranishi structures together using $\mathbf{z} : Y \rightarrow (Y, \kappa_Y)$ in Definition 4.14. Define strong submersions $\pi, \mathbf{exp} : ([0, 1] \times Y, \kappa') \rightarrow Y$ to be induced by $\text{id}_Y : Y \rightarrow Y$ on $[0, \frac{1}{2}] \times Y$ and by $\pi, \mathbf{exp} : (Y, \kappa_Y) \rightarrow Y$ on $[\frac{1}{2}, 1] \times Y$. Using the orientation on $[0, 1]$ we can construct a natural *coorientation* for $(([0, 1] \times Y, \kappa'), \pi)$, independent of the orientation on Y .

We shall define an excellent coordinate system $(\mathbf{J}_Y, \boldsymbol{\zeta}_Y)$ for $(([0, 1] \times Y, \kappa'), \pi)$. The indexing set is $J_Y = \{n+1, 2n+1\}$, with Kuranishi neighbourhoods

$$\begin{aligned} (V_Y^{n+1}, \dots, \psi_Y^{n+1}) &= ([0, \frac{1}{2}] \times Y, [0, \frac{1}{2}] \times Y, 0, \text{id}_{[0, \frac{1}{2}] \times Y}), \\ (V_Y^{2n+1}, \dots, \psi_Y^{2n+1}) &= ((\frac{1}{3}, 1] \times V, (\frac{1}{3}, 1] \times E, s \circ \pi_E, \text{id}_{(\frac{1}{3}, 1] \times \psi}). \end{aligned} \quad (84)$$

Here $E_Y^{n+1} \rightarrow V_Y^{n+1}$ is the zero vector bundle, so $E_Y^{n+1} \cong V_Y^{n+1}$ as manifolds. Define $V_Y^{(2n+1)(n+1)} = (\frac{1}{3}, \frac{1}{2}] \times Y \subset V_Y^{n+1}$, with coordinate change $(\phi_Y^{(2n+1)(n+1)}, \hat{\phi}_Y^{(2n+1)(n+1)}) : (V_Y^{(2n+1)(n+1)}, \dots, \psi_Y^{n+1}|_{V^{(2n+1)(n+1)}}) \rightarrow (V_Y^{2n+1}, \dots, \psi_Y^{2n+1})$ given by $\phi_Y^{(2n+1)(n+1)} = \text{id}_{(\frac{1}{3}, \frac{1}{2}] \times z}$, where $z : Y \rightarrow V$ is the zero section $z : Y \rightarrow TY$, recalling that V is a neighbourhood of $z(Y)$ in TY , and $\hat{\phi}_Y^{(2n+1)(n+1)} = \text{id}_{(\frac{1}{3}, \frac{1}{2}] \times (z_E \circ z)}$, where $z_E : V \rightarrow E$ is the zero section of E .

These Kuranishi neighbourhoods (84) are compatible with the Kuranishi structure κ' on $[0, 1] \times Y$, since for each point p in $\text{Im } \psi_Y^{n+1}$ or $\text{Im } \psi_Y^{2n+1}$, there is a sufficiently small $(V_p, \dots, \psi_p)$ in the germ of κ' at p and a coordinate change from $(V_p, \dots, \psi_p)$ to $(V_Y^{n+1}, \dots, \psi_Y^{n+1})$ or $(V_Y^{2n+1}, \dots, \psi_Y^{2n+1})$. Also $\pi, \mathbf{exp}$ are represented by $\pi_Y^{n+1} = \exp_Y^{n+1} = \pi_Y : [0, \frac{1}{2}] \times Y \rightarrow Y$ on $(V_Y^{n+1}, \dots, \psi_Y^{n+1})$ and by $\pi_Y^{2n+1} = \pi \circ \pi_V, \exp_Y^{2n+1} = \exp \circ \pi_V : (\frac{1}{3}, 1] \times V \rightarrow Y$ on $(V_Y^{2n+1}, \dots, \psi_Y^{2n+1})$.

Let $\zeta : [0, 1] \rightarrow [0, 1]$ be continuous with $\zeta \equiv 0$ on $[0, \frac{1}{3}]$ and $\zeta \equiv 1$ on $[\frac{1}{2}, 1]$. Define $\zeta_{j,Y} : [0, 1] \times Y \rightarrow [0, 1]$ for $j \in J_Y$ by $\zeta_{n+1,Y}(t, y) = 1 - \zeta(t)$

and $\zeta_{2n+1,Y}(t,y) = \zeta(t)$. Define $\zeta_{i,Y}^j : V^j \rightarrow [0,1]$ for $i,j \in J_Y$ by $\zeta_{n+1,Y}^j \equiv 1 - \zeta \circ \pi_{[0,1]}$ and $\zeta_{2n+1,Y}^j \equiv \zeta \circ \pi_{[0,1]}$. This data defines an *excellent coordinate system* $(\mathbf{J}_Y, \boldsymbol{\zeta}_Y)$ for $(([0,1] \times Y, \kappa'), \boldsymbol{\pi})$.

Choose maps $C_Y^j : E_Y^j \rightarrow P_m \subset P$ for $j \in J_Y$ and $m \gg 0$ such that $C_Y^{m+1}|_{\{0\} \times Y} \equiv (\emptyset)$ and $C_Y^{2n+1}|_{\{1\} \times E} \equiv \mu \circ (C_Y \times (G_Y \circ \pi))$, identifying $\{1\} \times E \cong E$, and $C_Y^j|_{(E_Y^j)^\circ} : (E_Y^j)^\circ \rightarrow P$ is injective. Then $C_Y^j \times \pi_Y$ is globally finite, and $\mathbf{C}_Y = (\mathbf{J}_Y, \boldsymbol{\zeta}_Y, C_Y^j : j \in J_Y)$ is *co-gauge-fixing data* for $(([0,1] \times Y, \kappa'), \boldsymbol{\pi})$.

Define R -module morphisms $\Xi^k : KC^k(Y; R) \rightarrow KC^{k-1}(Y; R)$ by

$$\Xi^k : [X, \mathbf{f}, \mathbf{C}] \mapsto [([0,1] \times Y, \kappa') \times_{\pi_Y, \mathbf{f}} X, \mathbf{exp} \circ \boldsymbol{\pi}_{([0,1] \times Y, \kappa')}, \mathbf{D}_\mathbf{C}], \quad (85)$$

where $(([0,1] \times Y, \kappa') \times_{\pi_Y, \mathbf{f}} X, \mathbf{exp} \circ \boldsymbol{\pi}_{([0,1] \times Y, \kappa')})$ has a coorientation constructed from that of $(X, \mathbf{f})$, and the co-gauge-fixing data $\mathbf{D}_\mathbf{C}$ is defined as follows. We form co-gauge-fixing data $\mathbf{C}_Y \times_Y \mathbf{C}$ for $(([0,1] \times Y, \kappa') \times_{\pi_Y, \mathbf{f}} X, \boldsymbol{\pi}_Y)$ as in Definition 3.31. Then we take $\mathbf{D}_\mathbf{C}$ to be the same as $\mathbf{C}_Y \times_Y \mathbf{C}$, except that the maps $\tilde{\pi}_Y^k : \tilde{V}^k \rightarrow Y$ for $k \in \tilde{I}$, representing $\boldsymbol{\pi}_Y$, are replaced by $\text{exp}^k = \text{exp}^j \circ \pi_{V_Y^j}$ on $\tilde{V}^k \cap (V_Y^j \times_Y V^i)$, representing $\mathbf{exp} \circ \boldsymbol{\pi}_{([0,1] \times Y, \kappa')}$.

To verify $\mathbf{D}_\mathbf{C}$ is co-gauge-fixing data, there is one nontrivial thing to check. The proof that $\mathbf{C}_Y \times_Y \mathbf{C}$ is co-gauge-fixing data shows that $\check{C}^k \times (\tilde{\pi}_Y^k \circ \tilde{\pi}^k) : \check{E}^k \rightarrow P \times Y$ is a globally finite map, but we have replaced $\tilde{\pi}_Y^k$ by exp^k , so we must prove that $\check{C}^k \times (\text{exp}^k \circ \tilde{\pi}^k) : \check{E}^k \rightarrow P \times Y$ is globally finite. Write $\check{E}^k = \check{E}^{k'} \amalg \check{E}^{k''}$, where $\check{E}^{k'}$ lies over V_Y^{n+1} and $\check{E}^{k''}$ lies over V_Y^{2n+1} . As $\pi_Y^{n+1} = \text{exp}_Y^{n+1}$ on V_Y^{n+1} , $\tilde{\pi}_Y^k$ and exp^k coincide on $\check{E}^{k'}$, so $\check{C}^k \times (\text{exp}^k \circ \tilde{\pi}^k)$ is globally finite on $\check{E}^{k'}$ as $\check{C}^k \times (\tilde{\pi}_Y^k \circ \tilde{\pi}^k)$ is. On $\check{E}^{k''}$ we use the fact that $C_Y^{2n+1} : E_Y^{2n+1} \rightarrow P_m \subset P$ is globally finite to deduce that $\check{C}^k : \check{E}^{k''} \rightarrow P$ is globally finite, following the proof that $\mathbf{G} \times_Y \tilde{\mathbf{C}}$ is gauge-fixing data, and temporarily treating $\mathbf{C}_Y$ as gauge-fixing data. But $\check{C}^k : \check{E}^{k''} \rightarrow P$ globally finite implies $\check{C}^k \times (\text{exp}^k \circ \tilde{\pi}^k) : \check{E}^{k''} \rightarrow P \times Y$ is globally finite, as we want.

From (23) we find that in cooriented Kuranishi spaces we have

$$\begin{aligned} \partial([0,1] \times Y, \kappa') \times_{\pi_Y, \mathbf{f}} X, \mathbf{exp} \circ \boldsymbol{\pi}_{[0,1] \times Y} &\cong -(\{0\} \times X, \mathbf{f}) \amalg \\ \amalg (\{1\} \times X^Y, \mathbf{f}^Y) &- ([0,1] \times Y, \kappa') \times_{\pi_Y, \mathbf{f}|_{\partial X}} \partial X, \mathbf{exp} \circ \boldsymbol{\pi}_{[0,1] \times Y}. \end{aligned}$$

Adding co-gauge-fixing data to this, in $KC^k(Y; R)$ we have

$$\begin{aligned} d([0,1] \times Y, \kappa') \times_{\pi_Y, \mathbf{f}} X, \mathbf{exp} \circ \boldsymbol{\pi}_{[0,1] \times Y}, \mathbf{D}_\mathbf{C} &= -[X, \mathbf{f}, \mathbf{C}] \\ [X^Y, \mathbf{f}^Y, \mathbf{C}_{\mathbf{G}_\mathbf{C}}^Y] &- ([0,1] \times Y, \kappa') \times_{\pi_Y, \mathbf{f}|_{\partial X}} \partial X, \mathbf{exp} \circ \boldsymbol{\pi}_{[0,1] \times Y}, \mathbf{D}_{\mathbf{C}|_{\partial X}}, \end{aligned} \quad (86)$$

where $\mathbf{G}_\mathbf{C}$ is the gauge-fixing data for $(X, \mathbf{f})$ constructed from $\mathbf{C}$ in Definition 4.14, and $\mathbf{C}_{\mathbf{G}_\mathbf{C}}^Y$ is the co-gauge-fixing data for $(X^Y, \mathbf{f}^Y)$ constructed from $\mathbf{G}_\mathbf{C}$ in Definition 4.14. The first term on the r.h.s. of (86) holds because C_Y^{m+1} restricts to $(\emptyset)$ on $\{0\} \times Y$, where $(\emptyset)$ is the identity for μ . The second term holds because C_Y^{2n+1} restricts on $\{1\} \times E$ to $\mu \circ (C_Y \times (G_Y \circ \pi))$, which combines the extra factor $G_Y \circ \pi$ in going from $\mathbf{C}$ to $\mathbf{G}_\mathbf{C}$ with the extra factor C_Y in going from $\mathbf{G}_\mathbf{C}$ to $\mathbf{C}_{\mathbf{G}_\mathbf{C}}^Y$, and using associativity of μ .

Combining (85), (86) and Definition 4.14 shows that

$$d \circ \Xi^k = -\text{id} + \Pi_{\text{Kch}}^{\text{Kh}} \circ \Pi_{\text{Kch}}^{\text{Kh}} - \Xi^{k+1} \circ d : KC^k(Y; R) \longrightarrow KC^k(Y; R). \quad (87)$$

Passing to cohomology, this implies $\Pi_{\text{Kch}}^{\text{Kh}} \circ \Pi_{\text{Kch}}^{\text{Kh}} : KH^k(Y; R) \rightarrow KH^k(Y; R)$ is the identity. An almost identical proof yields $\Upsilon^{n-k} : KC_{n-k}(Y; R) \rightarrow KC_{n-k+1}(Y; R)$ satisfying

$$\partial \circ \Upsilon^{n-k} = -\text{id} + \Pi_{\text{Kch}}^{\text{Kh}} \circ \Pi_{\text{Kch}}^{\text{Kh}} - \Upsilon^{n-k-1} \circ \partial : KC_{n-k}(Y; R) \rightarrow KC_{n-k}(Y; R).$$

In defining Υ^{n-k} we replace the condition on C_Y^{2n+1} above by $C_Y^{2n+1}|_{\{1\} \times E} \equiv \mu \circ ((G_Y \circ \exp \circ \pi_V) \times C_Y)$, identifying $\{1\} \times E \cong E$. Passing to homology shows that $\Pi_{\text{Kch}}^{\text{Kh}} \circ \Pi_{\text{Kch}}^{\text{Kh}} : KH_{n-k}(Y; R) \rightarrow KH_{n-k}(Y; R)$ is the identity. Therefore (83) is the inverse of (80).

For the final part of the theorem, by the first part it is enough to show that (80) is independent of choices, and the only choice in its construction is that of $G_Y : Y \rightarrow P_m \subset P$. If G_Y, G'_Y are possible choices inducing operators $\Pi_{\text{Kch}}^{\text{Kh}}, \Pi_{\text{Kch}}^{\text{Kh}'}$ on (co)chains, then choose globally finite $G_{[0,1] \times Y} : [0, 1] \times Y \rightarrow P_{m''} \subset P$ for $m'' \gg 0$ with $G_{[0,1] \times Y}|_{\{0\} \times Y} = G_Y$, $G_{[0,1] \times Y}|_{\{1\} \times Y} = G'_Y$.

Let $\mathbf{f} : X \rightarrow Y$ be a cooriented strong submersion and $\mathbf{C}$ be co-gauge-fixing data for $(X, \mathbf{f})$, where $\mathbf{C} = (\mathbf{I}, \boldsymbol{\eta}, C^i : i \in I)$ with $\mathbf{I} = (I, (V^i, \dots, \psi^i), f^i : i \in I, \dots)$. Set $\tilde{I} = \{i + 1 : i \in I\}$. For $i \in I$, define $(\tilde{V}^{i+1}, \tilde{E}^{i+1}, \tilde{s}^{i+1}, \tilde{\psi}^{i+1}) = ([0, 1] \times V^i, [0, 1] \times E^i, s^i \circ \pi_{V^i}, \text{id}_{[0,1]} \times \psi^i)$, $\tilde{f}^{i+1} = f^i \circ \pi_{V^i} : \tilde{V}^{i+1} \rightarrow Y$, and $\tilde{\eta}_{i+1} : [0, 1] \times X \rightarrow [0, 1]$ by $\tilde{\eta}_{i+1} = \eta_i \circ \pi_X$, and $H^{i+1} : \tilde{E}^{i+1} \rightarrow P$ by $H^{i+1}(u, e) = \mu(G_{[0,1] \times Y}(u, f^i \circ \pi_{V^i}(e)), C^i(e))$.

For $i, j \in I$, define $\tilde{V}^{(i+1)(j+1)} = [0, 1] \times V^{ij}$, $\tilde{\phi}^{(i+1)(j+1)} = \text{id}_{[0,1]} \times \phi^{ij}$, $\tilde{\phi}^{(i+1)(j+1)} = \text{id}_{[0,1]} \times \phi^{ij}$ and $\tilde{\eta}_{i+1}^{j+1} : \tilde{V}^{j+1} \rightarrow [0, 1]$ by $\tilde{\eta}_{i+1}^{j+1} = \eta_i^j \circ \pi_{V^j}$. Write $\tilde{\mathbf{I}} = (\tilde{I}, (\tilde{V}^i, \dots, \tilde{\psi}^i), \tilde{f}^i : i \in \tilde{I}, \dots)$, $\tilde{\boldsymbol{\eta}} = (\tilde{\eta}_i : i \in \tilde{I}, \tilde{\eta}_i^j : i, j \in \tilde{I})$, and $\mathbf{H}_{\mathbf{C}} = (\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}}, H^i : i \in \tilde{I})$. Then $\mathbf{H}_{\mathbf{C}}$ is *gauge-fixing data* for $([0, 1] \times X, \mathbf{f} \circ \pi_X)$.

Define R -module morphisms $\Theta^k : KC^k(Y; R) \rightarrow KC_{n-k-1}(Y; R)$ by

$$\Theta^k : [X, \mathbf{f}, \mathbf{C}] \longmapsto [[0, 1] \times X, \mathbf{f} \circ \pi_X, \mathbf{H}_{\mathbf{C}}].$$

A similar proof to (87) using $G_{[0,1] \times Y}|_{\{0\} \times Y} = G_Y$, $G_{[0,1] \times Y}|_{\{1\} \times Y} = G'_Y$ gives

$$\partial \circ \Theta^k = -\Pi_{\text{Kch}}^{\text{Kh}} + \Pi_{\text{Kch}}^{\text{Kh}'} - \Theta^{k+1} \circ d : KC^k(Y; R) \rightarrow KC_{n-k}(Y; R).$$

Passing to (co)homology yields $\Pi_{\text{Kch}}^{\text{Kh}} = \Pi_{\text{Kch}}^{\text{Kh}'} : KH^k(Y; R) \rightarrow KH_{n-k}(Y; R)$. This completes the proof. $\square$

We show Kuranishi and compactly-supported cohomology are isomorphic.

Definition 4.16. Let Y be an n -orbifold without boundary, and R a $\mathbb{Q}$ -algebra. We will define R -module morphisms $\Pi_{\text{cs}}^{\text{Kh}} : H_{\text{cs}}^k(Y; R) \rightarrow KH^k(Y; R)$ for all $k \geq 0$. First suppose Y is orientable, and choose an orientation. Define $\Pi_{\text{cs}}^{\text{Kh}}$ to be the composition

$$H_{\text{cs}}^k(Y; R) \xrightarrow{\text{Pd}} H_{n-k}^{\text{si}}(Y; R) \xrightarrow{\Pi_{\text{si}}^{\text{Kh}}} KH_{n-k}(Y; R) \xrightarrow{\Pi_{\text{Kh}}^{\text{Kh}}} KH^k(Y; R),$$

where $\text{Pd}, \Pi_{\text{si}}^{\text{Kh}}, \Pi_{\text{Kh}}^{\text{Kh}}$ are as in (61), (75), (83) respectively. Changing the orientation of Y changes the signs of $\text{Pd}, \Pi_{\text{Kh}}^{\text{Kh}}$. Thus $\Pi_{\text{cs}}^{\text{Kh}}$ is independent of the choice of orientation on Y .

Even if Y is not orientable, it has an *orientation bundle*, a principal $\mathbb{Z}_2$ -bundle $O \rightarrow Y$. The argument above still works if we replace $H_*^{\text{si}}(Y; R)$ and $KH_*(Y; R)$ by the *twisted homology groups* $H_*^{\text{si}}, KH_*(Y; O \times_{\{\pm 1\}} R)$. Geometrically, $H_k^{\text{si}}(Y; O \times_{\{\pm 1\}} R)$ is defined using chains (σ, o) where $\sigma : \Delta_k \rightarrow Y$ is a smooth map and o is an orientation of the fibres of $f^*(TY) \rightarrow \Delta_k$, with $(\sigma, -o) = -(\sigma, o)$, and $KH_*(Y; O \times_{\{\pm 1\}} R)$ is defined using chains $[X, \mathbf{f}, \mathbf{G}]$ with $(X, \mathbf{f})$ cooriented, rather than X oriented. (One can still make sense of coorientations for $(X, \mathbf{f})$ even if $\mathbf{f}$ is only strongly smooth).

Poincaré duality for orbifolds Y without boundary when R is a $\mathbb{Q}$ -algebra as discussed in §4.1, Corollary 4.10, and Theorem 4.15 imply:

Corollary 4.17. *Let Y be an orbifold without boundary, and R a $\mathbb{Q}$ -algebra. Then $\Pi_{\text{cs}}^{\text{Kh}} : H_{\text{cs}}^k(Y; R) \rightarrow KH^k(Y; R)$ is an isomorphism for $k \geq 0$, with $KH^k(Y; R) = 0$ when $k < 0$.*

4.5 The case $\partial Y \neq \emptyset$, and relative Kuranishi (co)homology

We now generalize §4.4 to define Kuranishi cohomology $KH^*(Y; R)$ for orbifolds Y with $\partial Y \neq \emptyset$. To show that $KH^*(Y; R) \cong H_{\text{cs}}^*(Y; R)$ using Poincaré duality involves *relative* Kuranishi homology and cohomology, so we define these as well. We will be brief, and give proofs only in the case $\partial^2 Y = \emptyset$.

We first explain the parts of classical (relative) (co)homology theory we will need. In §4.1 we discussed *Poincaré duality isomorphisms* (61) for manifolds and orbifolds Y without boundary. Here is how to extend this to manifolds with boundary and g-corners. Suppose Y is an oriented manifold with g-corners, of dimension n , not necessarily compact. Write $\iota : \partial Y \rightarrow Y$ for the natural immersion, and $\iota(\partial Y)$ for its image as a subset of Y . Then equation (61) extends to two families of Poincaré duality isomorphisms:

$$\text{Pd}' : H_{\text{cs}}^k(Y, \iota(\partial Y); R) \longrightarrow H_{n-k}(Y; R), \quad (88)$$

$$\text{Pd}'' : H_{\text{cs}}^k(Y; R) \longrightarrow H_{n-k}(Y, \iota(\partial Y); R), \quad (89)$$

where $H_{\text{cs}}^*(Y, \iota(\partial Y); R)$ is the *compactly-supported relative cohomology* and $H_*(Y, \iota(\partial Y); R)$ the *relative homology* of $(Y, \iota(Y))$. The usual form in which Poincaré duality is stated, as in Bredon [10, §VI.9] for instance, is (89) for *compact* Y , with $H^*(Y; R)$ in place of $H_{\text{cs}}^*(Y; R)$.

The interior Y° is a manifold without boundary, and there are natural isomorphisms $H_{\text{cs}}^*(Y, \iota(\partial Y); R) \cong H_{\text{cs}}^*(Y^\circ; R)$, $H_*(Y; R) \cong H_*(Y^\circ; R)$ which identify Pd' in (88) with Pd in (61) for Y° . Thus, (88) follows trivially from (61).

If $\partial^2 Y \neq \emptyset$ then $\iota : \partial Y \rightarrow Y$ is not injective, and its image $\iota(\partial Y)$ is not a manifold. However, if $\partial^2 Y = \emptyset$ then ι is injective and $\iota(\partial Y)$ is a submanifold of Y . *Relative singular homology* when $\partial^2 Y = \emptyset$ is the homology of the chain

complex $(C_*^{\text{si}}(Y, \iota(\partial Y); R), \partial)$, where

$$C_k^{\text{si}}(Y, \iota(\partial Y); R) = C_k^{\text{si}}(Y; R) / \iota_*(C_k^{\text{si}}(\partial Y; R)),$$

and $\iota_* : C_k^{\text{si}}(\partial Y; R) \rightarrow C_k^{\text{si}}(Y; R)$ is as in §4.1. This gives a short exact sequence of chain complexes:

$$0 \longrightarrow C_*^{\text{si}}(\partial Y; R) \xrightarrow{\iota_*} C_*^{\text{si}}(Y; R) \xrightarrow{\pi} C_*^{\text{si}}(Y, \iota(\partial Y); R) \longrightarrow 0, \quad (90)$$

where π is the projection $C_*^{\text{si}}(Y; R) \rightarrow C_*^{\text{si}}(Y; R) / \iota_*(C_*^{\text{si}}(\partial Y; R))$. Equation (90) induces a long exact sequence of (relative) homology groups:

$$\cdots \rightarrow H_k^{\text{si}}(\partial Y; R) \xrightarrow{\iota_*} H_k^{\text{si}}(Y; R) \xrightarrow{\pi_*} H_k^{\text{si}}(Y, \iota(\partial Y); R) \xrightarrow{\partial_*} H_{k-1}^{\text{si}}(\partial Y; R) \rightarrow \cdots \quad (91)$$

Similarly, when $\partial^2 Y = \emptyset$ we have a long exact sequence of cohomology groups:

$$\cdots \rightarrow H_{\text{cs}}^{k-1}(\partial Y; R) \xrightarrow{\partial^*} H_{\text{cs}}^k(Y, \iota(\partial Y); R) \xrightarrow{\pi^*} H_{\text{cs}}^k(Y; R) \xrightarrow{\iota^*} H_{\text{cs}}^k(\partial Y; R) \rightarrow \cdots \quad (92)$$

Now ∂Y is an oriented $(n-1)$ -manifold without boundary as $\partial^2 Y = \emptyset$, so Poincaré duality isomorphisms (61) hold for ∂Y . These combine with the isomorphisms (88)–(89) and sequences (91)–(92) in a commutative diagram

$$\begin{array}{ccccccc} \cdots & \rightarrow & H_{\text{cs}}^{k-1}(\partial Y; R) & \xrightarrow{\partial^*} & H_{\text{cs}}^k(Y, \iota(\partial Y); R) & \xrightarrow{\pi^*} & H_{\text{cs}}^k(Y; R) \xrightarrow{\iota^*} \cdots \\ & & \cong \downarrow \text{Pd} & & \cong \downarrow \text{Pd}' & & \cong \downarrow \text{Pd}'' \\ \cdots & \rightarrow & H_{n-k}^{\text{si}}(\partial Y; R) & \xrightarrow{\iota_*} & H_{n-k}^{\text{si}}(Y; R) & \xrightarrow{\pi_*} & H_{n-k}^{\text{si}}(Y, \iota(\partial Y); R) \xrightarrow{\partial_*} \cdots \end{array}$$

As in §4.1, the Poincaré duality isomorphisms (88)–(89) also hold for orbifolds Y provided R is a $\mathbb{Q}$ -algebra. We now define *relative Kuranishi homology*.

Definition 4.18. Let Y, Z be orbifolds, $h : Y \rightarrow Z$ a smooth map, and R a $\mathbb{Q}$ -algebra. Then Definition 4.6 gives $h_* : KC_k(Y; R) \rightarrow KC_k(Z; R)$ with $\partial \circ h_* = h_* \circ \partial$. Define the *relative Kuranishi chains* $KC_k(Z, h(Y); R)$ to be

$$KC_k(Z, h(Y); R) = KC_k(Z; R) / h_*(KC_k(Y; R)).$$

Then $\partial : KC_k(Z; R) \rightarrow KC_{k-1}(Z; R)$ descends to $\partial : KC_k(Z, h(Y); R) \rightarrow KC_{k-1}(Z, h(Y); R)$ with $\partial^2 = 0$. Define *relative Kuranishi homology* $KH_*(Z, h(Y); R)$ to be the homology of $(KC_*(Z, h(Y); R), \partial)$.

Suppose now that $h : Z \rightarrow Y$ is an *embedding*. Then $h_* : KC_*(Y; R) \rightarrow KC_*(Z; R)$ and $h_* : C_*^{\text{si}}(Y; R) \rightarrow C_*^{\text{si}}(Z; R)$ are injective, so we have a commutative diagram of chain complexes with exact rows:

$$\begin{array}{ccccccc} 0 & \rightarrow & C_*^{\text{si}}(Y; R) & \xrightarrow{h_*} & C_*^{\text{si}}(Z; R) & \xrightarrow{\pi} & C_*^{\text{si}}(Z, h(Y); R) \longrightarrow 0 \\ & & \downarrow \Pi_{\text{si}}^{\text{Kh}} & & \downarrow \Pi_{\text{si}}^{\text{Kh}} & & \downarrow \Pi_{\text{rsi}}^{\text{Kh}} \\ 0 & \rightarrow & KC_*(Y; R) & \xrightarrow{h_*} & KC_*(Z; R) & \xrightarrow{\pi} & KC_*(Z, h(Y); R) \longrightarrow 0, \end{array} \quad (93)$$

where $\Pi_{\text{rsi}}^{\text{rKh}}$ is the projection from relative singular chains to relative Kuranishi chains defined as for $\Pi_{\text{si}}^{\text{Kh}}$ in Definition 4.6.

Equation (93) induces a commutative diagram of homology groups

$$\begin{array}{ccccccc}
\cdots \longrightarrow & H_k^{\text{si}}(Y; R) & \xrightarrow{h_*} & H_k^{\text{si}}(Z; R) & \xrightarrow{\pi_*} & H_k^{\text{si}}(Z, h(Y); R) & \xrightarrow{\partial_*} & H_{k-1}^{\text{si}}(Y; R) \longrightarrow \cdots \\
& \cong \downarrow \Pi_{\text{si}}^{\text{Kh}} & & \cong \downarrow \Pi_{\text{si}}^{\text{Kh}} & & \downarrow \Pi_{\text{rsi}}^{\text{rKh}} & & \cong \downarrow \Pi_{\text{si}}^{\text{Kh}} \\
\cdots \longrightarrow & KH_k(Y; R) & \xrightarrow{h_*} & KH_k(Z; R) & \xrightarrow{\pi_*} & KH_k(Z, h(Y); R) & \xrightarrow{\partial_*} & KH_{k-1}(Y; R) \longrightarrow \cdots,
\end{array} \tag{94}$$

with long exact rows. Corollary 4.10 implies that the columns $\Pi_{\text{si}}^{\text{Kh}}$ in (94) are isomorphisms. Since the rows are exact, properties of long exact sequences then imply that the remaining column $\Pi_{\text{rsi}}^{\text{rKh}}$ is an isomorphism. This implies:

Corollary 4.19. *Suppose $h : Y \rightarrow Z$ is a smooth embedding of orbifolds, and R is a $\mathbb{Q}$ -algebra. Then $\Pi_{\text{rsi}}^{\text{rKh}} : H_k^{\text{si}}(Z, h(Y); R) \rightarrow KH_k(Z, h(Y); R)$ is an isomorphism for all $k \geq 0$, with $KH_k(Z, h(Y); R) = 0$ for $k < 0$.*

The author expects that the assumption h is an embedding is unnecessary in Corollary 4.19. Note that if Y is an orbifold with $\partial^2 Y = \emptyset$ then $\iota : \partial Y \rightarrow Y$ is an embedding, so $\Pi_{\text{rsi}}^{\text{rKh}} : H_k^{\text{si}}(Y, \iota(\partial Y); R) \rightarrow KH_k(Y, \iota(\partial Y); R)$ is an isomorphism. Next we define $KH^*(Y; R)$ and $KH^*(Y, \iota(\partial Y); R)$ when Y is an orbifold with g-corners, allowing $\partial Y \neq \emptyset$.

Definition 4.20. Let X be a Kuranishi space, Y an orbifold with g-corners, and $\mathbf{f} : X \rightarrow Y$ a strong submersion. Then if $\partial Y \neq \emptyset$, as for submersions of manifolds in Definition 2.5 we have a decomposition $\partial X = \partial_+^{\mathbf{f}} X \amalg \partial_-^{\mathbf{f}} X$ and strong submersions $\mathbf{f}_+ : \partial_+^{\mathbf{f}} X \rightarrow Y$ and $\mathbf{f}_- : \partial_-^{\mathbf{f}} X \rightarrow \partial Y$, where $\mathbf{f}_+ = \mathbf{f}|_{\partial_+^{\mathbf{f}} X}$. In particular, if $\partial_-^{\mathbf{f}} X \neq \emptyset$ then $\mathbf{f}|_{\partial X}$ is *not* a strong submersion, but only strongly smooth.

Let Y be an orbifold and R a $\mathbb{Q}$ -algebra. When $\partial Y = \emptyset$, Definition 4.12 defined the Kuranishi cochains $KC^*(Y; R)$, with $d : KC^k(Y; R) \rightarrow KC^{k+1}(Y; R)$ given by (78). If $\partial Y \neq \emptyset$, equation (78) is no longer valid, since $\mathbf{f}_a|_{\partial X_a}$ may not be a strong submersion. Instead, when $\partial Y \neq \emptyset$ we define $KC^*(Y; R)$ exactly as in Definition 4.12, but we define $d : KC^k(Y; R) \rightarrow KC^{k+1}(Y; R)$ by

$$d : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{C}_a] \mapsto \sum_{a \in A} \rho_a [\partial_+^{\mathbf{f}_a} X_a, \mathbf{f}_{a,+}, \mathbf{C}_a|_{\partial_+^{\mathbf{f}_a} X_a}].$$

Since the natural orientation-reversing involution $\sigma : \partial^2 X_a \rightarrow \partial^2 X_a$ restricts to an involution of $\partial_+^{\mathbf{f}_a, +}(\partial_+^{\mathbf{f}_a} X_a)$, relation Definition 4.3(ii) still implies that $d^2 = 0$. We then define *Kuranishi cohomology* $KH^*(Y; R)$ to be the cohomology of $(KC^*(Y; R), d)$.

Define a morphism $\iota^* : KC^k(Y; R) \rightarrow KC^k(\partial Y; R)$ by

$$\iota^* : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{C}_a] \mapsto (-1)^k \sum_{a \in A} \rho_a [\partial_-^{\mathbf{f}_a} X_a, \mathbf{f}_{a,-}, \mathbf{C}_a|_{\partial_-^{\mathbf{f}_a} X_a}], \tag{95}$$

where $\mathbf{f}_{a,-} : \partial_-^{\mathbf{f}_a} X_a \rightarrow \partial Y$ is a strong submersion as above. Then ι^* takes relations Definition 4.3(i)–(iv) in $KC^k(Y; R)$ to the same relations in $KC^k(\partial Y; R)$, and so is well-defined. We think of ι^* as the *pullback* under $\iota : \partial Y \rightarrow Y$.

If X is an oriented Kuranishi space and $\mathbf{f} : X \rightarrow Y$ a strong submersion, then the natural orientation-reversing involution $\sigma : \partial^2 X \rightarrow \partial^2 X$ restricts to an orientation-reversing isomorphism $\sigma : \partial_+^{\mathbf{f}}(\partial_+^{\mathbf{f}} X) \rightarrow \partial_+^{\mathbf{f}}(\partial_+^{\mathbf{f}} X)$. Using this we see that $d \circ \iota^* = \iota^* \circ d$ as maps $KC^k(Y; R) \rightarrow KC^{k+1}(\partial Y; R)$. The purpose of the factor $(-1)^k$ in (95) is to make $d \circ \iota^* = \iota^* \circ d$, so that ι^* is a chain map; without it we would get $d \circ \iota^* = -\iota^* \circ d$, since σ is orientation-reversing. Since ι^* is a chain map it induces morphisms $\iota^* : KH^k(Y; R) \rightarrow KH^k(\partial Y; R)$.

Define the *relative Kuranishi cochains* $KC^k(Y, \iota(Y); R)$ to be the kernel of $\iota^* : KC^k(Y; R) \rightarrow KC^k(\partial Y; R)$. Then $d : KC^k(Y; R) \rightarrow KC^{k+1}(Y; R)$ restricts to $d : KC^k(Y, \iota(Y); R) \rightarrow KC^{k+1}(Y, \iota(Y); R)$, since $d \circ \iota^* = \iota^* \circ d$. Define *relative Kuranishi cohomology* $KH^*(Y, \iota(Y); R)$ to be the cohomology of $(KC^k(Y, \iota(Y); R), d)$.

For the rest of this section we restrict to the case that $\partial^2 Y = \emptyset$, so that $\iota : \partial Y \rightarrow Y$ is an embedding. Then one can show that $\iota^* : KC^k(Y; R) \rightarrow KC^k(\partial Y; R)$ is *surjective*; one way to do this is to identify $Y \cong [0, \epsilon] \times \partial Y$ near ∂Y for small $\epsilon > 0$, and then for each $[X, \mathbf{f}, \mathbf{C}] \in KC^k(\partial Y; R)$, to construct explicit $[[0, \epsilon] \times X, \mathbf{f}', \mathbf{C}'] \in KC^k(Y; R)$ with $\iota^* : [[0, \epsilon] \times X, \mathbf{f}', \mathbf{C}'] \mapsto [X, \mathbf{f}, \mathbf{C}]$. Thus we have a short exact sequence of cochain complexes

$$0 \longrightarrow KC^*(Y, \iota(Y); R) \xrightarrow{\text{inc}} KC^*(Y; R) \xrightarrow{\iota^*} KC^*(\partial Y; R) \longrightarrow 0,$$

where ‘inc’ is the inclusion. This induces a long exact sequence in cohomology, the analogue of (92):

$$\cdots \rightarrow KH^{k-1}(\partial Y; R) \xrightarrow{\partial^*} KH^k(Y, \iota(\partial Y); R) \xrightarrow{\text{inc}^*} KH^k(Y; R) \xrightarrow{\iota^*} KH^k(\partial Y; R) \rightarrow \cdots.$$

Our goal is to prove that $KH^*(Y; R) \cong H_{\text{cs}}^*(Y; R)$ and $KH^*(Y, \iota(Y); R) \cong H_{\text{cs}}^*(Y, \iota(Y); R)$, generalizing Corollary 4.17. To do this we need analogues of the Poincaré duality morphisms $\Pi_{\text{Kch}}^{\text{Kh}}, \Pi_{\text{Kh}}^{\text{Kch}}$ of Definition 4.14 for $KH^*(Y; R)$ and $KH^*(Y, \iota(Y); R)$. That is, we wish to define chain maps

$$\Pi_{\text{rKch}}^{\text{Kh}} : KC^k(Y, \iota(\partial Y); R) \longrightarrow KC_{n-k}(Y; R), \quad (96)$$

$$\Pi_{\text{Kh}}^{\text{rKch}} : KC_{n-k}(Y; R) \longrightarrow KC^k(Y, \iota(\partial Y); R), \quad (97)$$

$$\Pi_{\text{Kch}}^{\text{rKh}} : KC^k(Y; R) \longrightarrow KC_{n-k}(Y, \iota(\partial Y); R), \quad (98)$$

$$\Pi_{\text{rKh}}^{\text{Kch}} : KC_{n-k}(Y, \iota(\partial Y); R) \dashrightarrow KC^k(Y; R), \quad (99)$$

which should induce the analogues of Pd', Pd'' and their inverses.

We define $\Pi_{\text{rKch}}^{\text{Kh}}, \Pi_{\text{Kch}}^{\text{rKh}}$ in (96), (98) to map $[X, \mathbf{f}, \mathbf{C}] \mapsto [X, \mathbf{f}, \mathbf{G}_{\mathbf{C}}]$ exactly as for $\Pi_{\text{rKch}}^{\text{Kh}}$ in Definition 4.14. The proof that $\partial \circ \Pi_{\text{Kch}}^{\text{rKh}} = \Pi_{\text{Kch}}^{\text{rKh}} \circ d$ then becomes

nontrivial: we have

$$\begin{aligned}\partial \circ \Pi_{\text{Kch}}^{\text{rKh}} : [X, \mathbf{f}, \mathbf{C}] &\longmapsto [\partial X, \mathbf{f}|_{\partial X}, \mathbf{G}_{\mathbf{C}}|_{\partial X}], \quad \text{and} \\ \Pi_{\text{Kch}}^{\text{rKh}} \circ d : [X, \mathbf{f}, \mathbf{C}] &\longmapsto [\partial_+^{\mathbf{f}} X, \mathbf{f}_+, \mathbf{G}_{\mathbf{C}}|_{\partial_+^{\mathbf{f}} X}].\end{aligned}$$

The difference between these two is $[\partial_-^{\mathbf{f}} X, \mathbf{f}|_{\partial_-^{\mathbf{f}} X}, \mathbf{G}_{\mathbf{C}}|_{\partial_-^{\mathbf{f}} X}]$, but $\mathbf{f}|_{\partial_-^{\mathbf{f}} X} = \iota \circ \mathbf{f}_-$, so this difference lies in $\iota_*(KC_k(\partial Y; R))$, and is zero in $KC_{n-k-1}(Y, \iota(\partial Y); R)$, giving $\partial \circ \Pi_{\text{Kch}}^{\text{rKh}} = \Pi_{\text{Kch}}^{\text{rKh}} \circ d$. To prove that $\partial \circ \Pi_{\text{rKch}}^{\text{Kh}} = \Pi_{\text{rKch}}^{\text{Kh}} \circ d$, we show that $\partial \circ \Pi_{\text{rKch}}^{\text{Kh}} - \Pi_{\text{rKch}}^{\text{Kh}} \circ d = (-1)^k \iota_* \circ \iota^*$, and then note that $KC^k(Y, \iota(\partial Y); R)$ is the kernel of ι^* . Hence $\Pi_{\text{rKch}}^{\text{Kh}}, \Pi_{\text{Kch}}^{\text{rKh}}$ are well-defined chain maps, and induce morphisms $\Pi_{\text{rKch}}^{\text{Kh}} : KH^k(Y, \iota(\partial Y); R) \rightarrow KH_{n-k}(Y; R)$ and $\Pi_{\text{Kch}}^{\text{rKh}} : KH^k(Y; R) \rightarrow KH_{n-k}(Y, \iota(\partial Y); R)$.

We can also define $\Pi_{\text{Kh}}^{\text{Kch}}$ in (97) generalizing $\Pi_{\text{Kh}}^{\text{Kch}}$ in Definition 4.14, though this is a little more complicated, since one must also add in a correction term over ∂Y to ensure that $\Pi_{\text{Kh}}^{\text{Kch}}$ maps to $\text{Ker } \iota^*$ in $KC^k(Y; R)$. It is a chain map, and induces $\Pi_{\text{Kh}}^{\text{Kch}} : KH_{n-k}(Y; R) \rightarrow KH^k(Y, \iota(\partial Y); R)$. We can then prove the analogue of Theorem 4.15, that $\Pi_{\text{rKch}}^{\text{Kh}} : KH^k(Y, \iota(\partial Y); R) \rightarrow KH_{n-k}(Y; R)$ and $\Pi_{\text{Kh}}^{\text{Kch}} : KH_{n-k}(Y; R) \rightarrow KH^k(Y, \iota(\partial Y); R)$ are inverse, and so are both isomorphisms. This is not surprising, since we expect $KH^*(Y, \iota(\partial Y); R) \cong KH^*(Y^\circ; R)$ and $KH_*(Y; R) \cong KH_*(Y^\circ; R)$ as for (88) above, and this analogue would follow immediately from Theorem 4.15.

However, the author does not have a good definition for $\Pi_{\text{rKch}}^{\text{Kch}}$ in (99), on the (co)chain level. Instead, we consider the commutative diagram

$$\begin{array}{ccccccc} \cdots & \longrightarrow & KH^{k-1}(\partial Y; R) & \longrightarrow & KH^k(Y, \partial Y; R) & \longrightarrow & KH^k(Y; R) \longrightarrow \cdots \\ & & \cong \downarrow \Pi_{\text{Kch}}^{\text{Kh}} & & \downarrow \Pi_{\text{rKch}}^{\text{Kh}} & & \cong \downarrow \Pi_{\text{rKch}}^{\text{Kh}} \\ \cdots & \longrightarrow & KH_{n-k}(\partial Y; R) & \longrightarrow & KH_{n-k}(Y; R) & \longrightarrow & KH_{n-k}(Y, \partial Y; R) \longrightarrow \cdots, \end{array} \quad (100)$$

with exact rows. Here the left hand column is an isomorphism by Theorem 4.15, as ∂Y is an oriented $(n-1)$ -manifold without boundary, and the right hand column is an isomorphism by the analogue of Theorem 4.15 discussed above. Thus the central column is also an isomorphism, by properties of long exact sequences. Therefore we have proved:

Theorem 4.21. *Suppose Y is an oriented n -orbifold with $\partial^2 Y = \emptyset$, and R is a $\mathbb{Q}$ -algebra. Then we have isomorphisms $\Pi_{\text{rKch}}^{\text{Kh}} : KH^k(Y, \iota(\partial Y); R) \rightarrow KH_{n-k}(Y; R)$ and $\Pi_{\text{Kch}}^{\text{rKh}} : KH^k(Y; R) \rightarrow KH_{n-k}(Y, \iota(\partial Y); R)$.*

The author expects the theorem holds without assuming $\partial^2 Y = \emptyset$. The proof for $\Pi_{\text{rKch}}^{\text{Kh}}$ looks straightforward, but for $\Pi_{\text{Kch}}^{\text{rKh}}$ the long exact sequence argument in (100) will need modification. Here is the analogue of Definition 4.16:

Definition 4.22. Let Y be an n -orbifold, and R a $\mathbb{Q}$ -algebra. We will define R -module morphisms $\Pi_{\text{rcs}}^{\text{rKh}} : H_{\text{cs}}^k(Y, \iota(\partial Y); R) \rightarrow KH^k(Y, \iota(\partial Y); R)$ and $\Pi_{\text{cs}}^{\text{Kch}} : H_{\text{cs}}^k(Y; R) \rightarrow KH^k(Y; R)$ for all $k \geq 0$. First suppose Y is orientable, and

choose an orientation. Define $\Pi_{\text{rcs}}^{\text{rKch}}$ and $\Pi_{\text{cs}}^{\text{rKch}}$ to be the compositions

$$\begin{aligned} H_{\text{cs}}^k(Y, \iota(\partial Y); R) &\xrightarrow{\text{Pd}'} H_{n-k}^{\text{si}}(Y; R) \xrightarrow{\Pi_{\text{si}}^{\text{Kh}}} KH_{n-k}(Y; R) \xrightarrow{\Pi_{\text{Kh}}^{\text{rKch}}} KH^k(Y, \iota(\partial Y); R), \\ H_{\text{cs}}^k(Y; R) &\xrightarrow{\text{Pd}''} H_{n-k}^{\text{si}}(Y, \iota(\partial Y); R) \xrightarrow{\Pi_{\text{rsi}}^{\text{Kh}}} KH_{n-k}(Y, \iota(\partial Y); R) \xrightarrow{\Pi_{\text{rKh}}^{\text{rKch}}} KH^k(Y; R). \end{aligned}$$

Changing the orientation of Y changes the signs of Pd' , Pd'' , $\Pi_{\text{Kh}}^{\text{rKch}}$, $\Pi_{\text{rKh}}^{\text{rKch}}$. Thus $\Pi_{\text{rcs}}^{\text{rKch}}$ and $\Pi_{\text{cs}}^{\text{rKch}}$ are independent of the choice of orientation on Y . If Y is not orientable, we define $\Pi_{\text{rcs}}^{\text{rKch}}$, $\Pi_{\text{cs}}^{\text{rKch}}$ using *twisted* (relative) homology groups, by the method of Definition 4.16.

Poincaré duality, Corollaries 4.10 and 4.19, and Theorem 4.21 imply:

Corollary 4.23. *Let Y be an orbifold with $\partial^2 Y = \emptyset$, and R a $\mathbb{Q}$ -algebra. Then $\Pi_{\text{rcs}}^{\text{rKch}} : H_{\text{cs}}^k(Y, \iota(\partial Y); R) \rightarrow KH^k(Y, \iota(\partial Y); R)$ and $\Pi_{\text{cs}}^{\text{rKch}} : H_{\text{cs}}^*(Y; R) \rightarrow KH^*(Y; R)$ are isomorphisms.*

The author expects the corollary holds without assuming $\partial^2 Y = \emptyset$.

The other material we give on Kuranishi cochains $KC^*(Y; R)$ and Kuranishi cohomology $KH^*(Y; R)$ when $\partial Y = \emptyset$, such as pullbacks h^* in §4.4, effective Kuranishi cohomology in §4.6, and cup and cap products in §4.7, all has generalizations to the case $\partial Y \neq \emptyset$ using the ideas of this section. We leave this as an exercise for the reader.

4.6 Effective Kuranishi cohomology

We extend §4.4 to *effective* Kuranishi cohomology.

Definition 4.24. Let Y be an orbifold without boundary. Consider triples $(X, \mathbf{f}, \underline{\mathbf{C}})$, where X is a compact Kuranishi space, $\mathbf{f} : X \rightarrow Y$ is a strong submersion with $(X, \mathbf{f})$ coeffective and cooriented, and $\underline{\mathbf{C}}$ is effective co-gauge-fixing data for $(X, \mathbf{f})$. Write $[X, \mathbf{f}, \underline{\mathbf{C}}]$ for the isomorphism class of $(X, \mathbf{f}, \underline{\mathbf{C}})$ under coorientation-preserving isomorphisms $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \underline{\mathbf{C}}) \rightarrow (\tilde{X}, \tilde{\mathbf{f}}, \tilde{\underline{\mathbf{C}}})$.

Let R be a commutative ring. For each $k \in \mathbb{Z}$, define $KC_{\text{ec}}^k(Y; R)$ to be the R -module of finite R -linear combinations of isomorphism classes $[X, \mathbf{f}, \underline{\mathbf{C}}]$ as above for which $\text{vdim } X = \dim Y - k$, with the analogues of relations Definition 4.3(i)–(iii), replacing gauge-fixing data $\mathbf{G}$ by effective co-gauge-fixing data $\underline{\mathbf{C}}$. Elements of $KC_{\text{ec}}^k(Y; R)$ are called *effective Kuranishi cochains*. Define $d : KC_{\text{ec}}^k(Y; R) \rightarrow KC_{\text{ec}}^{k+1}(Y; R)$ by (78), replacing $\mathbf{C}_a$ by $\underline{\mathbf{C}}_a$. Then $d \circ d = 0$. Define the *effective Kuranishi cohomology groups* $KH_{\text{ec}}^k(Y; R)$ of Y to be

$$KH_{\text{ec}}^k(Y; R) = \frac{\text{Ker}(d : KC_{\text{ec}}^k(Y; R) \rightarrow KC_{\text{ec}}^{k+1}(Y; R))}{\text{Im}(d : KC_{\text{ec}}^{k-1}(Y; R) \rightarrow KC_{\text{ec}}^k(Y; R))}$$

for $k \in \mathbb{Z}$. Arguably one should call this ‘coeffective Kuranishi cohomology’, as it is defined using $[X, \mathbf{f}, \underline{\mathbf{C}}]$ with $(X, \mathbf{f})$ coeffective, but we choose not to.

As for $\Pi_{\text{ef}}^{\text{Kh}}$ in Definition 4.6, define

$$\begin{aligned} \Pi_{\text{ec}}^{\text{Kch}} : KC_{\text{ec}}^k(Y; R) &\longrightarrow KC^k(Y; R \otimes_{\mathbb{Z}} \mathbb{Q}) \quad \text{by} \\ \Pi_{\text{ec}}^{\text{Kch}} : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{\mathbf{C}}_a] &\longmapsto \sum_{a \in A} \pi(\rho_a) [X_a, \mathbf{f}_a, \Pi(\underline{\mathbf{C}}_a)]. \end{aligned} \quad (101)$$

Then $\Pi_{\text{ec}}^{\text{Kch}} \circ d = d \circ \Pi_{\text{ec}}^{\text{Kch}}$, so they induce morphisms of cohomology groups

$$\Pi_{\text{ec}}^{\text{Kch}} : KH_{\text{ec}}^k(Y; R) \longrightarrow KH^k(Y; R \otimes_{\mathbb{Z}} \mathbb{Q}). \quad (102)$$

Let Y, Z be orbifolds without boundary, and $h : Y \rightarrow Z$ be a smooth, proper map. As in Definition 4.12, define the *pullback* $h^* : KC_{\text{ec}}^k(Z; R) \rightarrow KC_{\text{ec}}^k(Y; R)$ by (79), with $\underline{\mathbf{C}}_a$ in place of $\mathbf{C}_a$. These satisfy $h^* \circ d = d \circ h^*$, and so induce morphisms $h^* : KH_{\text{ec}}^k(Z; R) \rightarrow KH_{\text{ec}}^k(Y; R)$. We have $(g \circ h)^* = h^* \circ g^*$ and $\Pi_{\text{ec}}^{\text{Kch}} \circ h^* = h^* \circ \Pi_{\text{ec}}^{\text{Kch}}$, on both cochains and cohomology.

We can also try to extend Definition 4.14 and Theorem 4.15 to effective Kuranishi (co)homology. However, as in §4.1 Poincaré duality fails for general orbifolds Y and commutative rings R , so something must go wrong. The problem is in generalizing the morphisms $\Pi_{\text{Kh}}^{\text{Kh}} : KC^k(Y; R) \rightarrow KC_{n-k}(Y; R)$ and $\Pi_{\text{Kh}}^{\text{Kch}} : KC_{n-k}(Y; R) \rightarrow KC^k(Y; R)$ to the effective case. We can define $\Pi_{\text{ec}}^{\text{ef}} : KC_{\text{ec}}^k(Y; R) \rightarrow KC_{n-k}^{\text{ef}}(Y; R)$ only if Y is an *effective* orbifold, as otherwise $(V^i, \dots, \psi^i), f^i$ coeffective does not imply $(V^i, \dots, \psi^i)$ effective. This corresponds to the observation in §4.1 that if Y is not effective then we cannot define the Poincaré duality map Pd in (61), which is the analogue of $\Pi_{\text{ec}}^{\text{ef}}$.

Worse still, we can define $\Pi_{\text{ef}}^{\text{ec}} : KC_{n-k}^{\text{ef}}(Y; R) \rightarrow KC_{\text{ec}}^k(Y; R)$ only if Y is a *manifold*, since otherwise the Kuranishi neighbourhoods (81) may not be coeffective, as the stabilizer groups of V can act nontrivially on the fibres of E . This corresponds to the observation in §4.1 that if Y is an effective orbifold, but not a manifold, then Pd in (61) is defined but may not be invertible, so we cannot define its inverse Pd^{-1} , which is the analogue of $\Pi_{\text{ef}}^{\text{ec}}$.

Definition 4.25. Let Y be an *effective* orbifold without boundary, which is oriented of dimension n , and R a commutative ring. Choose an injective map $\underline{\mathbf{G}}_Y : Y \rightarrow \mathbb{R}^m \subset \underline{\mathbf{P}}$ for some $m \gg 0$. Suppose X is a compact, oriented Kuranishi space, $\mathbf{f} : X \rightarrow Y$ is a strong submersion, $(X, \mathbf{f})$ is coeffective and cooriented, and $\underline{\mathbf{C}}$ is effective co-gauge-fixing data for $(X, \mathbf{f})$. Define effective gauge-fixing data $\underline{\mathbf{G}}_{\underline{\mathbf{C}}}$ for $(X, \mathbf{f})$ as for $\mathbf{G}_{\mathbf{C}}$ in Definition 4.14, but with $\underline{\mathbf{G}}^i = \underline{\mu} \circ ((\underline{\mathbf{G}}_Y \circ \mathbf{f}^i \circ \pi^i) \times \underline{\mathbf{C}}^i)$, for $\underline{\mu}$ as in (58). A similar proof to those for $h^*(\underline{\mathbf{C}})$ and $\underline{\mathbf{C}} \times_Y \tilde{\mathbf{C}}$ in Definitions 3.42 and 3.43 shows that $\underline{\mathbf{G}}_{\underline{\mathbf{C}}}$ is *effective gauge-fixing data*. To verify Definition 3.36(a), note that as $(V^i, \dots, \psi^i), f^i$ in $\underline{\mathbf{C}}$ is coeffective and Y is effective, $(V^i, \dots, \psi^i)$ is effective as in Definition 3.37.

Define R -module morphisms $\Pi_{\text{ec}}^{\text{ef}} : KC_{\text{ec}}^k(Y; R) \rightarrow KC_{n-k}^{\text{ef}}(Y; R)$ for $k \in \mathbb{Z}$ by $\Pi_{\text{ec}}^{\text{ef}} : [X, \mathbf{f}, \underline{\mathbf{C}}] \mapsto [X, \mathbf{f}, \underline{\mathbf{G}}_{\underline{\mathbf{C}}}]$, using the coorientation for $(X, \mathbf{f})$ and orientation on Y to determine the orientation on X . Then $\partial \circ \Pi_{\text{ec}}^{\text{ef}} = \Pi_{\text{ec}}^{\text{ef}} \circ d$, so they induce morphisms of (co)homology groups

$$\Pi_{\text{ec}}^{\text{ef}} : KH_{\text{ec}}^k(Y; R) \longrightarrow KH_{n-k}^{\text{ef}}(Y; R). \quad (103)$$

Now suppose Y is a *manifold*, without boundary. Let X be a compact, effective, oriented Kuranishi space, $\mathbf{f} : X \rightarrow Y$ a strongly smooth map, and $\underline{\mathbf{G}}$ be effective gauge-fixing data for $(X, \mathbf{f})$. Define $X^Y, \mathbf{f}^Y$ as in Definition 4.14. As Y is a manifold, one can show from (81) that $(X^Y, \mathbf{f}^Y)$ is coeffective. Define $\underline{\mathbf{C}}_{\underline{\mathbf{G}}}^Y$ for $(X^Y, \mathbf{f}^Y)$ as for $\mathbf{C}_{\mathbf{G}}^Y$ in Definition 4.14, except that we set $\check{\mathbf{C}}^{k,Y} = \underline{\mu} \circ ((\underline{\mathbf{C}}_Y \circ \pi_E) \times (\underline{\mathbf{G}}^{k-n} \circ \pi_{E^{k-n}}))$, where $\underline{\mathbf{C}}_Y : E \rightarrow \mathbb{R}^{m'} \subset \underline{P}$ is injective for some $m' \gg 0$. Again, we can check that $\underline{\mathbf{C}}_{\underline{\mathbf{G}}}^Y$ is *effective co-gauge-fixing data*.

Define R -module morphisms $\Pi_{\text{ef}}^{\text{ec}} : KC_{n-k}^{\text{ef}}(Y; R) \rightarrow KC_{\text{ec}}^k(Y; R)$ by $\Pi_{\text{ef}}^{\text{ec}} : [X, \mathbf{f}, \underline{\mathbf{G}}] \mapsto [X^Y, \mathbf{f}^Y, \underline{\mathbf{C}}_{\underline{\mathbf{G}}}^Y]$. Then $d \circ \Pi_{\text{ef}}^{\text{ec}} = \Pi_{\text{ef}}^{\text{ec}} \circ \partial$, so they induce morphisms of (co)homology groups

$$\Pi_{\text{ef}}^{\text{ec}} : KH_{n-k}^{\text{ef}}(Y; R) \longrightarrow KH_{\text{ec}}^k(Y; R). \quad (104)$$

Theorem 4.15, Definition 4.16 and Corollary 4.17 then generalize as follows.

Theorem 4.26. *Let Y be an oriented n -manifold without boundary, and R a commutative ring. Then $\Pi_{\text{ef}}^{\text{ec}}$ in (104) is the inverse of $\Pi_{\text{ec}}^{\text{ef}}$ in (103), so they are both isomorphisms. Also (103)–(104) are independent of choices.*

Definition 4.27. Let Y be an n -manifold without boundary, and R a commutative ring. We will define R -module morphisms $\Pi_{\text{cs}}^{\text{ec}} : H_{\text{cs}}^k(Y; R) \rightarrow KH_{\text{ec}}^k(Y; R)$ for all $k \geq 0$. First suppose Y is oriented. Define $\Pi_{\text{cs}}^{\text{ec}}$ to be the composition

$$H_{\text{cs}}^k(Y; R) \xrightarrow{\text{Pd}} H_{n-k}^{\text{si}}(Y; R) \xrightarrow{\Pi_{\text{si}}^{\text{ef}}} KH_{n-k}^{\text{ef}}(Y; R) \xrightarrow{\Pi_{\text{ef}}^{\text{ec}}} KH_{\text{ec}}^k(Y; R),$$

where $\text{Pd}, \Pi_{\text{si}}^{\text{ef}}, \Pi_{\text{ef}}^{\text{ec}}$ are as in (61), (76), (104). Independence of the orientation of Y , and extension to non-orientable Y , are as in Definition 4.16.

Corollary 4.28. *Let Y be a manifold without boundary, and R a commutative ring. Then $\Pi_{\text{cs}}^{\text{ec}} : H_{\text{cs}}^*(Y; R) \rightarrow KH_{\text{ec}}^*(Y; R)$ is an isomorphism.*

This still leaves the question of whether $H_{\text{cs}}^*(Y; R) \cong KH_{\text{ec}}^*(Y; R)$ for Y a general orbifold without boundary, and R a commutative ring. The author expects this is true, even though our proof via homology and Poincaré duality fails in this case. When R is a $\mathbb{Q}$ -algebra, one approach might be to generalize the proof of Theorem 4.9 in Appendix C to show that $\Pi_{\text{ec}}^{\text{Kch}} : KH_{\text{ec}}^k(Y; R) \rightarrow KH^k(Y; R)$ in (102) is an isomorphism, and then use Corollary 4.17.

4.7 Products on (effective) Kuranishi (co)homology

There are *cup products* $\cup$, *cap products* $\cap$ and *intersection products* $\bullet$ on (effective) Kuranishi (co)chains and (co)homology.

Definition 4.29. Let Y be an orbifold without boundary, and R a $\mathbb{Q}$ -algebra. Define the *cup product* $\cup : KC^k(Y; R) \times KC^l(Y; R) \rightarrow KC^{k+l}(Y; R)$ by

$$[X, \mathbf{f}, \mathbf{C}] \cup [\tilde{X}, \tilde{\mathbf{f}}, \tilde{\mathbf{C}}] = [X \times_{\mathbf{f}, Y, \tilde{\mathbf{f}}} \tilde{X}, \pi_Y, \mathbf{C} \times_Y \tilde{\mathbf{C}}], \quad (105)$$

extended R -bilinearly. Here $\pi_Y : X \times_{f,Y,\tilde{f}} \tilde{X} \rightarrow Y$ is the projection from the fibre product, which is a strong submersion as both $f, \tilde{f}$ are strong submersions, and $C \times_Y \tilde{C}$ is as in Definition 3.31. As for d , it is easy to show $\cup$ takes relations (i)–(iv) in both $KC^k(Y; R)$ and $KC^l(Y; R)$ to the same relations in $KC^{k+l}(Y; R)$. Thus $\cup$ is well-defined.

Proposition 3.32(a)–(c) and (23)–(25) imply that for $\gamma \in KC^k(Y; R)$, $\delta \in KC^l(Y; R)$ and $\epsilon \in KC^m(Y; R)$ we have

$$\gamma \cup \delta = (-1)^{kl} \delta \cup \gamma, \quad (106)$$

$$d(\gamma \cup \delta) = (d\gamma) \cup \delta + (-1)^k \gamma \cup (d\delta) \text{ and } (\gamma \cup \delta) \cup \epsilon = \gamma \cup (\delta \cup \epsilon). \quad (107)$$

Therefore in the usual way $\cup$ induces an associative, supercommutative product $\cup : KH^k(Y; R) \times KH^l(Y; R) \rightarrow KH^{k+l}(Y; R)$ given for $\gamma \in KC^k(Y; R)$ and $\delta \in KC^l(Y; R)$ with $d\gamma = d\delta = 0$ by

$$(\gamma + \text{Im } d_{k-1}) \cup (\delta + \text{Im } d_{l-1}) = (\gamma \cup \delta) + \text{Im } d_{k+l-1}.$$

Now suppose Y is compact, and of dimension n . Then $\text{id}_Y : Y \rightarrow Y$ is a (strong) submersion, with a trivial coorientation giving the positive sign to the zero vector bundle over Y . Define co-gauge-fixing data C_Y for (Y, id_Y) by $I = \{n\}$, $(V^n, E^n, s^n, \psi^n) = (Y, Y, 0, \text{id}_Y)$, $\text{id}_Y^n = \text{id}_Y : V^n \rightarrow Y$, and $\eta_n : Y \rightarrow [0, 1]$, $\eta_n^n : V^n \rightarrow [0, 1]$ and $C^n : E^n \rightarrow P$ given by $\eta_n \equiv 1$, $\eta_n^n \equiv 1$ and $C^n \equiv (\emptyset)$, where $(\emptyset) \in P$ is the identity for μ , as in Definition 3.8. Then $[Y, \text{id}_Y, C_Y] \in KC^0(Y; R)$, with $d[Y, \text{id}_Y, C_Y] = 0$. Following the definitions through shows that for $[X, f, C] \in KC^k(Y; R)$ we have

$$[Y, \text{id}_Y, C_Y] \cup [X, f, C] = [X, f, C] \cup [Y, \text{id}_Y, C_Y] = [X, f, C].$$

Thus $[Y, \text{id}_Y, C_Y]$ is the identity for $\cup$, at the cochain level. Passing to cohomology, $[[Y, \text{id}_Y, C_Y]] \in KH^0(Y; R)$ is the identity for $\cup$ in $KH^*(Y; R)$. We call $[[Y, \text{id}_Y, C_Y]]$ the *fundamental class* of Y .

Define the *cap product* $\cap : KC_k(Y; R) \times KC^l(Y; R) \rightarrow KC_{k-l}(Y; R)$ by

$$[X, f, G] \cap [\tilde{X}, \tilde{f}, \tilde{C}] = [X \times_{f,Y,\tilde{f}} \tilde{X}, \pi_Y, G \times_Y \tilde{C}], \quad (108)$$

extended R -bilinearly, with $G \times_Y \tilde{C}$ as in Definition 3.31. For $\gamma \in KC_k(Y; R)$ and $\delta, \epsilon \in KC^*(Y; R)$, the analogue of (107), using Proposition 3.32(d), is

$$\partial(\gamma \cap \delta) = (\partial\gamma) \cap \delta + (-1)^{\dim Y - k} \gamma \cap (d\delta), \quad (\gamma \cap \delta) \cap \epsilon = \gamma \cap (\delta \cup \epsilon). \quad (109)$$

If also Y is compact then $\gamma \cap [Y, \text{id}_Y, C_Y] = \gamma$. Thus $\cap$ induces a *cap product* $\cap : KH_k(Y; R) \times KH^l(Y; R) \rightarrow KH_{k-l}(Y; R)$. These products $\cap$ make Kuranishi chains and homology into *modules* over Kuranishi cochains and cohomology.

Let Y, Z be orbifolds without boundary, and $h : Y \rightarrow Z$ a smooth, proper map. Then Proposition 3.33 implies that pullbacks h^* and pushforwards h_* are compatible with $\cup, \cap$ on (co)chains, in the sense that if $\alpha \in KC_*(Y; R)$ and $\beta, \gamma \in KC^*(Z; R)$ then

$$h^*(\beta \cup \gamma) = h^*(\beta) \cup h^*(\gamma) \quad \text{and} \quad h_*(\alpha \cap h^*(\beta)) = h_*(\alpha) \cap \beta. \quad (110)$$

Since $\cup, \cap, h^*, h_*$ are compatible with d, ∂ , passing to (co)homology shows that (110) also holds for $\alpha \in KH_*(Y; R)$ and $\beta, \gamma \in KH^*(Z; R)$. If Z is compact then Y is, with $h^*([Z, \text{id}_Z, \mathbf{C}_Z]) = [Y, \text{id}_Y, \mathbf{C}_Y]$ in $KC^0(Y; R)$ and $h^*([Z, \text{id}_Z, \mathbf{C}_Z]) = [[Y, \text{id}_Y, \mathbf{C}_Y]]$ in $KH^0(Y; R)$.

To summarize: Kuranishi cochains $KC^*(Y; R)$ form a *supercommutative, associative, differential graded R -algebra*, and Kuranishi cohomology $KH^*(Y; R)$ is a *supercommutative, associative, graded R -algebra*. These algebras are *with identity* if Y is compact without boundary, and *without identity* otherwise. Pullbacks h^* induce *algebra morphisms* on both cochains and cohomology. Kuranishi chains $KC_*(Y; R)$ are a *graded module* over $KC^*(Y; R)$, and Kuranishi homology $KH_*(Y; R)$ is a *graded module* over $KH^*(Y; R)$.

Let Y be oriented, of dimension n . Then Poincaré duality, Theorem 4.15, and the cup product $\cup$ on Kuranishi cohomology induce an *intersection product* $\bullet : KH_k(Y; R) \times KH_l(Y; R) \rightarrow KH_{k+l-n}(Y; R)$ on Kuranishi homology. It is supercommutative (with degrees shifted by n) and associative, with identity in $KH_n(Y; R)$ if Y is compact. Intersection products are *not* preserved by pushforwards $h_* : KH_*(Y; R) \rightarrow KH_*(Z; R)$, the degrees of $h_*(\alpha) \bullet h_*(\beta)$ and $h_*(\alpha \bullet \beta)$ differ by $\dim Y - \dim Z$.

On chains we can define $\bullet : KC_k(Y; R) \times KC_l(Y; R) \rightarrow KC_{k+l-n}(Y; R)$ by

$$\alpha \bullet \beta = \Pi_{Kch}^{Kh}(\Pi_{Kh}^{Kch}(\alpha) \cup \Pi_{Kh}^{Kch}(\beta)), \quad (111)$$

where $\Pi_{Kch}^{Kh}, \Pi_{Kh}^{Kch}$ are as in Definition 4.14. Then $\bullet$ on chains induces $\bullet$ on homology, and $\bullet$ on chains is supercommutative, but *not* associative, and does not have an identity. Thus, the intersection product $\bullet$ is not as well-behaved on Kuranishi chains as the cup and cap products $\cup, \cap$.

Remark 4.30. It is an attractive and unusual feature of our theory of Kuranishi (co)homology that the products $\cup, \cap$ are everywhere defined, and supercommutative, and associative, *at the (co)chain level*, as well as on (co)homology. Furthermore, pushforwards and pullbacks are functorial and compatible with $\cup, \cap$ at the (co)chain level, and for compact Y there is an identity $[Y, \text{id}_Y, \mathbf{C}_Y]$ for $\cup, \cap$ at the cochain level. There are few homology or cohomology theories for which this holds — it works for de Rham cohomology, for instance, but fails for singular and simplicial (co)homology.

The fact that products $\cup, \cap$ are *everywhere defined* on (co)chains, rather than only under some transversality assumptions such as transverse intersection of chains in Y , is a feature of working with Kuranishi spaces rather than manifolds or orbifolds. If X is a manifold or orbifold then submersions $f : X \rightarrow Y$ only exist if $\dim X \geq \dim Y$. So for defining (co)homology using pairs (X, f) as (co)chains with X a manifold or orbifold (for example, singular homology, with $X = \Delta_k$), making f a submersion is too strong an assumption, and this leads to transversality problems when defining products. But for Kuranishi spaces X we can have strong submersions $\mathbf{f} : X \rightarrow Y$ with $\text{vdim } X < \dim Y$, since the obstruction bundles on X decrease $\text{vdim } X$. Thus we can use pairs $(X, \mathbf{f})$ with $\mathbf{f}$ a strong submersion as cochains, and $\cup, \cap$ are everywhere defined.

A lot of work in Chapter 3 went into defining (co-)gauge-fixing data with associative, supercommutative products. This good behaviour of Kuranishi cochains will be very important, and lead to huge simplifications, in the sequels [4, 39, 40], where we have to work at the cochain level.

Definition 4.31. Let Y be an orbifold without boundary, and R a commutative ring. For cup and cap products on *effective* Kuranishi (co)chains and (co)homology, we follow Definition 4.29, except that we use the fibre products $\underline{\mathcal{C}} \times_Y \tilde{\mathcal{C}}$, $\underline{\mathcal{G}} \times_Y \tilde{\mathcal{C}}$ of effective (co-)gauge-fixing data from Definition 3.43. Thus we define $\cup : KC_{ec}^k(Y; R) \times KC_{ec}^l(Y; R) \rightarrow KC_{ec}^{k+l}(Y; R)$ by

$$[X, \underline{f}, \underline{\mathcal{C}}] \cup [\tilde{X}, \tilde{f}, \tilde{\mathcal{C}}] = [X \times_{\underline{f}, Y, \tilde{f}} \tilde{X}, \pi_Y, \underline{\mathcal{C}} \times_Y \tilde{\mathcal{C}}].$$

Then $\cup$ satisfies (107) by Propositions 3.32(b),(c) and 3.44, but it does not satisfy (106), since fibre products of effective co-gauge-fixing data are not commutative. Hence $\cup$ induces an associative cup product

$$\cup : KH_{ec}^k(Y; R) \times KH_{ec}^l(Y; R) \longrightarrow KH_{ec}^{k+l}(Y; R). \quad (112)$$

Suppose Y is compact of dimension n . As for $\underline{\mathcal{C}}_Y$ in Definition 4.29, define effective co-gauge-fixing data $\underline{\mathcal{C}}_Y$ for (Y, id_Y) by $I = \{n\}$, $(V^n, E^n, s^n, \psi^n) = (Y, Y, 0, \text{id}_Y)$, $\text{id}_Y^n = \text{id}_Y : V^n \rightarrow Y$, and $\eta_n : Y \rightarrow [0, 1]$, $\eta_n^n : V^n \rightarrow [0, 1]$ and $\underline{\mathcal{C}}^n : E^n \rightarrow \underline{P}$ given by $\eta_n \equiv 1$, $\eta_n^n \equiv 1$ and $\underline{\mathcal{C}}^n \equiv (\emptyset)$, where $(\emptyset) \in \underline{P}$ is the identity for $\underline{P}$, as in Definition 3.36. Then $[Y, \text{id}_Y, \underline{\mathcal{C}}_Y] \in KC_{ec}^0(Y; R)$, with $d[Y, \text{id}_Y, \underline{\mathcal{C}}_Y] = 0$, and $[Y, \text{id}_Y, \underline{\mathcal{C}}_Y]$ is the identity for $\cup$, at the cochain level. Passing to cohomology, $[[Y, \text{id}_Y, \underline{\mathcal{C}}_Y]] \in KH_{ec}^0(Y; R)$ is the identity for $\cup$ in $KH_{ec}^*(Y; R)$. We call $[[Y, \text{id}_Y, \underline{\mathcal{C}}_Y]]$ the *fundamental class* of Y .

Define the *cap product* $\cap : KC_k^{\text{ef}}(Y; R) \times KC_{ec}^l(Y; R) \rightarrow KC_{k-l}^{\text{ef}}(Y; R)$ by

$$[X, \underline{f}, \underline{\mathcal{G}}] \cap [\tilde{X}, \tilde{f}, \tilde{\mathcal{C}}] = [X \times_{\underline{f}, Y, \tilde{f}} \tilde{X}, \pi_Y, \underline{\mathcal{G}} \times_Y \tilde{\mathcal{C}}].$$

It satisfies (109), by Propositions 3.32(d) and 3.44. Thus $\cap$ induces a *cap product*

$$\cap : KH_k^{\text{ef}}(Y; R) \times KH_{ec}^l(Y; R) \longrightarrow KH_{k-l}^{\text{ef}}(Y; R). \quad (113)$$

If Y is compact then $\alpha \cap [Y, \text{id}_Y, \underline{\mathcal{C}}_Y] = \alpha$ for all α in $KC_*^{\text{ef}}(Y; R)$, and $\beta \cap [[Y, \text{id}_Y, \underline{\mathcal{C}}_Y]] = \beta$ for all β in $KH_*^{\text{ef}}(Y; R)$.

Let Y, Z be orbifolds without boundary, and $h : Y \rightarrow Z$ a smooth, proper map. Then Propositions 3.33 and 3.44 imply (110) holds for $\alpha \in KC_*^{\text{ef}}(Y; R)$ and $\beta, \gamma \in KC_{ec}^*(Z; R)$, and hence for $\alpha \in KH_*^{\text{ef}}(Y; R)$ and $\beta, \gamma \in KH_{ec}^*(Z; R)$. If Z is compact then Y is, with $h^*([Z, \text{id}_Z, \underline{\mathcal{C}}_Z]) = [Y, \text{id}_Y, \underline{\mathcal{C}}_Y]$ in $KC_{ec}^0(Y; R)$ and $h^*([Z, \text{id}_Z, \underline{\mathcal{C}}_Z]) = [[Y, \text{id}_Y, \underline{\mathcal{C}}_Y]]$ in $KH_{ec}^0(Y; R)$.

Let Y be an oriented n -manifold without boundary. Then Poincaré duality, Theorem 4.26, and $\cup$ in (112) induce an associative *intersection product* $\bullet : KH_k^{\text{ef}}(Y; R) \times KH_l^{\text{ef}}(Y; R) \rightarrow KH_{k+l-n}^{\text{ef}}(Y; R)$ on effective Kuranishi homology. On chains we can define $\bullet : KC_k^{\text{ef}}(Y; R) \times KC_l^{\text{ef}}(Y; R) \rightarrow KC_{k+l-n}^{\text{ef}}(Y; R)$ by

$$\alpha \bullet \beta = \Pi_{ec}^{\text{ef}}(\Pi_{ec}^{\text{ec}}(\alpha) \cup \Pi_{ec}^{\text{ec}}(\beta)). \quad (114)$$

This is neither associative nor supercommutative, at the chain level.

Since $\mu : P \times P \rightarrow P$, $\underline{\mu} : \underline{P} \times \underline{P} \rightarrow \underline{P}$ and $\Pi : \underline{P} \rightarrow P$ in Definitions 3.8 and 3.36 satisfy $\Pi(\underline{\mu}(p, q)) = \mu(\Pi(p), \Pi(q))$ for all $p, q \in \underline{P}$, it easily follows that the morphisms $\Pi_{\text{ef}}^{\text{Kh}}, \Pi_{\text{ec}}^{\text{Kh}}$ from effective Kuranishi (co)chains to Kuranishi (co)chains in (67) and (101) satisfy

$$\begin{aligned} \Pi_{\text{ec}}^{\text{Kh}}(\gamma) \cup \Pi_{\text{ec}}^{\text{Kh}}(\delta) &= \Pi_{\text{ec}}^{\text{Kh}}(\gamma \cup \delta), & \Pi_{\text{ef}}^{\text{Kh}}(\alpha) \cap \Pi_{\text{ec}}^{\text{Kh}}(\gamma) &= \Pi_{\text{ef}}^{\text{Kh}}(\alpha \cap \gamma) \\ \text{and} \quad \Pi_{\text{ef}}^{\text{Kh}}(\alpha) \bullet \Pi_{\text{ef}}^{\text{Kh}}(\beta) &= \Pi_{\text{ef}}^{\text{Kh}}(\alpha \bullet \beta), \end{aligned} \quad (115)$$

for all $\alpha, \beta \in KC_{\text{ec}}^{\text{ef}}(Y; R)$ and $\gamma, \delta \in KC_{\text{ec}}^*(Y; R)$, supposing Y oriented for the third equation. Passing to (co)homology, the morphisms $\Pi_{\text{ef}}^{\text{Kh}}, \Pi_{\text{ec}}^{\text{Kh}}$ in (68) and (102) also satisfy (115) for all $\alpha, \beta \in KH_{\text{ec}}^{\text{ef}}(Y; R)$ and $\gamma, \delta \in KH_{\text{ec}}^*(Y; R)$.

Although $\cup$ on $KC_{\text{ec}}^*(Y; R)$ is not supercommutative, it is up to homotopy, and so $\cup$ on $KH_{\text{ec}}^*(Y; R)$ is supercommutative.

Proposition 4.32. *Let Y be an orbifold without boundary and R a commutative ring. Then $\cup$ on $KH_{\text{ec}}^*(Y; R)$ in (112) satisfies (106), that is, $\cup$ is supercommutative. If Y is an oriented manifold without boundary then $\bullet$ is supercommutative on $KH_{\text{ec}}^{\text{ef}}(Y; R)$.*

Proof. Let $X, \tilde{X}$ be compact Kuranishi spaces, $\mathbf{f} : X \rightarrow Y$, $\tilde{\mathbf{f}} : \tilde{X} \rightarrow Y$ coefficientive, cooriented strong submersions, and $\underline{\mathbf{C}}, \underline{\tilde{\mathbf{C}}}$ effective co-gauge-fixing data for $(X, \mathbf{f}), (\tilde{X}, \tilde{\mathbf{f}})$. Then Definition 3.43 defines effective co-gauge-fixing data $\underline{\mathbf{C}} \times_Y \underline{\tilde{\mathbf{C}}}, \underline{\tilde{\mathbf{C}}} \times_Y \underline{\mathbf{C}}$ for $(X \times_Y \tilde{X}, \pi_Y)$ and $(\tilde{X} \times_Y X, \pi_Y)$. Our problem is that the natural isomorphism $X \times_Y \tilde{X} \cong \tilde{X} \times_Y X$ does not identify $\underline{\mathbf{C}} \times_Y \underline{\tilde{\mathbf{C}}}$ with $\underline{\tilde{\mathbf{C}}} \times_Y \underline{\mathbf{C}}$, because in the definition $\underline{\tilde{\mathbf{C}}}^k(\tilde{e}) = \underline{\mu}(\underline{\mathbf{C}}^i \circ \pi_{E^i}(\tilde{e}), \tilde{\mathbf{C}}^i \circ \pi_{\tilde{E}^i}(\tilde{e}))$ of $\underline{\tilde{\mathbf{C}}}^k : \tilde{E}^k \rightarrow \underline{P}$ in $\underline{\mathbf{C}} \times_Y \underline{\tilde{\mathbf{C}}}$, the function $\underline{\mu}$ in (58) is not commutative.

As in Definition 3.31, write $(\tilde{\mathbf{I}}, \tilde{\boldsymbol{\eta}})$ for the excellent coordinate system for $(X \times_Y \tilde{X}, \pi_Y)$ in $\underline{\mathbf{C}} \times_Y \underline{\tilde{\mathbf{C}}}$, where $\tilde{\mathbf{I}} = (\tilde{I}, (\tilde{V}^k, \tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k), \tilde{\pi}_Y^k : k \in \tilde{I}, \dots)$. Define an excellent coordinate system $(\mathbf{J}, \boldsymbol{\zeta})$ for $([0, 1] \times (X \times_Y \tilde{X}), \pi_Y)$ to have indexing set $J = \{k + 1 : k \in \tilde{I}\}$, Kuranishi neighbourhoods

$$(\tilde{V}^{k+1}, \tilde{E}^{k+1}, \tilde{s}^{k+1}, \tilde{\psi}^{k+1}) = ([0, 1] \times \tilde{V}^k, [0, 1] \times \tilde{E}^k, \text{id}_{[0,1]} \times \tilde{s}^k, \text{id}_{[0,1]} \times \tilde{\psi}^k)$$

for $k \in \tilde{I}$, maps $\tilde{\pi}_Y^{k+1} : \tilde{V}^{k+1} \rightarrow Y$, $\zeta_{k+1} : [0, 1] \times (X \times_Y \tilde{X}) \rightarrow [0, 1]$ and $\zeta_{k+1}^{l+1} : \tilde{V}^{l+1} \rightarrow [0, 1]$ given by $\tilde{\pi}_Y^{k+1} = \tilde{\pi}_Y^k \circ \pi_{\tilde{V}^k}$, $\zeta_{k+1} = \tilde{\eta}_k \circ \pi_{X \times_Y \tilde{X}}$ and $\zeta_{k+1}^{l+1} = \tilde{\eta}_k^l \circ \pi_{\tilde{V}^l}$, and triples $(\tilde{V}^{(k+1)(l+1)}, \tilde{\phi}^{(k+1)(l+1)}, \hat{\phi}^{(k+1)(l+1)})$ given by $\tilde{V}^{(k+1)(l+1)} = [0, 1] \times \tilde{V}^{kl}$, $\tilde{\phi}^{(k+1)(l+1)} = \text{id}_{[0,1]} \times \tilde{\phi}^{kl}$ and $\hat{\phi}^{(k+1)(l+1)} = \text{id}_{[0,1]} \times \hat{\phi}^{kl}$.

Define $\underline{\nu} : [0, 1] \times \underline{P} \times \underline{P} \rightarrow \underline{P}$ by

$$\underline{\nu}(t, (x_1, \dots, x_k), (y_1, \dots, y_l)) = \begin{cases} (x_1, \dots, x_k, y_1, \dots, y_l), & t = 0, \\ (t, x_1, \dots, x_k, y_1, \dots, y_l), & t \in (0, 1), \\ (y_1, \dots, y_l, x_1, \dots, x_k), & t = 1. \end{cases} \quad (116)$$

Then $\underline{\nu}(0, p, q) = \underline{\mu}(p, q)$ and $\underline{\nu}(1, p, q) = \underline{\mu}(q, p)$. Define $\underline{D}^{k+1} : \tilde{E}^{k+1} \rightarrow P$ for $k \in \tilde{I}$ and $u \in [0, 1]$, $\check{e} \in \tilde{E}^k \cap E^{(i, \tilde{i})}$ by

$$\underline{D}^{k+1}(u, \check{e}) = \underline{\nu}(u, \underline{C}^i \circ \pi_{E^i}(\check{e}), \tilde{\underline{C}}^{\tilde{i}} \circ \pi_{\tilde{E}^{\tilde{i}}}(\check{e})). \quad (117)$$

A similar proof to those for $h^*(\underline{C})$ and $\underline{C} \times_Y \tilde{\underline{C}}$ in Definitions 3.42 and 3.43 shows that $\underline{D}_{\underline{C}, \tilde{\underline{C}}} = ((\underline{J}, \underline{\zeta}), \underline{D}^j : j \in J)$ is *effective co-gauge-fixing data* for $([0, 1] \times (X \times_Y \tilde{X}), \pi_Y)$.

Using (23) we see that in cooriented Kuranishi spaces we have

$$\begin{aligned} \partial([0, 1] \times (X \times_Y \tilde{X}), \pi_Y) &\cong -(\{0\} \times (X \times_Y \tilde{X}), \pi_Y) \\ &\amalg (\{1\} \times (X \times_Y \tilde{X}), \pi_Y) \amalg -([0, 1] \times (\partial X \times_Y \tilde{X}), \pi_Y) \\ &\amalg -(-1)^{\text{vdim } X - \dim Y}([0, 1] \times (X \times_Y \partial \tilde{X}), \pi_Y). \end{aligned} \quad (118)$$

Let us add co-gauge-fixing data to (118). The restriction of $\underline{D}_{\underline{C}, \tilde{\underline{C}}}$ to the first term on the r.h.s. is isomorphic to $\underline{C} \times_Y \tilde{\underline{C}}$ under $\{0\} \times (X \times_Y \tilde{X}) \cong X \times_Y \tilde{X}$, since $\underline{D}^{k+1} : [0, 1] \times \tilde{E}^k \rightarrow P$ in (117) restricts on $\{0\} \times \tilde{E}^k \cong \tilde{E}^k$ to $\tilde{\underline{C}}^k : \tilde{E}^k \rightarrow P$ in the definition of $\underline{C} \times_Y \tilde{\underline{C}}$, by the case $t = 0$ in (116). The restriction of $\underline{D}_{\underline{C}, \tilde{\underline{C}}}$ to the second term is isomorphic to $\tilde{\underline{C}} \times_Y \underline{C}$ under $\{1\} \times (X \times_Y \tilde{X}) \cong \tilde{X} \times_Y X$, since $\underline{D}^{k+1}$ restricts on $\{1\} \times \tilde{E}^k \cong \tilde{E}^k$ to $\tilde{\underline{C}}^k$ with the rôles of $\underline{C}, \tilde{\underline{C}}$ reversed, by the case $t = 1$ in (116). In cooriented Kuranishi spaces we have $(\{1\} \times (X \times_Y \tilde{X}), \pi_Y) \cong (-1)^{(\text{vdim } X - \dim Y)(\text{vdim } \tilde{X} - \dim Y)}(\tilde{X} \times_Y X, \pi_Y)$ by (24). The restrictions of $\underline{D}_{\underline{C}, \tilde{\underline{C}}}$ to the third and fourth terms are $\underline{D}_{\underline{C}|_{\partial X}, \tilde{\underline{C}}}$ and $\underline{D}_{\underline{C}, \tilde{\underline{C}}|_{\partial \tilde{X}}}$. Putting all this together, in $KC_{\text{ec}}^*(Y; R)$ we have

$$\begin{aligned} d[[0, 1] \times (X \times_Y \tilde{X}), \pi_Y, \underline{D}_{\underline{C}, \tilde{\underline{C}}}] &= -[X \times_Y \tilde{X}, \pi_Y, \underline{C} \times_Y \tilde{\underline{C}}] \\ &\quad + (-1)^{(\text{vdim } X - \dim Y)(\text{vdim } \tilde{X} - \dim Y)}[\tilde{X} \times_Y X, \pi_Y, \tilde{\underline{C}} \times_Y \underline{C}] \\ &\quad - [[0, 1] \times (\partial X \times_Y \tilde{X}), \pi_Y, \underline{D}_{\underline{C}|_{\partial X}, \tilde{\underline{C}}}] \\ &\quad - (-1)^{\text{vdim } X - \dim Y}[[0, 1] \times (X \times_Y \partial \tilde{X}), \pi_Y, \underline{D}_{\underline{C}, \tilde{\underline{C}}|_{\partial \tilde{X}}}]. \end{aligned} \quad (119)$$

Define R -bilinear $\Xi^{k, l} : KC_{\text{ec}}^k(Y; R) \times KC_{\text{ec}}^l(Y; R) \rightarrow KC_{\text{ec}}^{k+l-1}(Y; R)$ by

$$\Xi^{k, l} : ([X, \underline{f}, \underline{C}], [\tilde{X}, \tilde{\underline{f}}, \tilde{\underline{C}}]) \mapsto [[0, 1] \times (X \times_Y \tilde{X}), \pi_Y, \underline{D}_{\underline{C}, \tilde{\underline{C}}}].$$

Using (119) and putting $\text{vdim } X = \dim Y - k$, $\text{vdim } \tilde{X} = \dim Y - l$ yields

$$d \circ \Xi^{k, l}(\alpha, \beta) = -\alpha \cup \beta + (-1)^{kl} \beta \cup \alpha - \Xi^{k+1, l}(d\alpha, \beta) - (-1)^{k} \Xi^{k, l+1}(d\alpha, d\beta)$$

for $\alpha \in KC_{\text{ec}}^k(Y; R)$ and $\beta \in KC_{\text{ec}}^l(Y; R)$. Passing to cohomology, this gives $\gamma \cup \delta = (-1)^{kl} \delta \cup \gamma$ for $\gamma \in KH_{\text{ec}}^k(Y; R)$, $\delta \in KH_{\text{ec}}^l(Y; R)$, that is, $\cup$ is supercommutative on $KH_{\text{ec}}^*(Y; R)$. The last part is immediate as $\cup, \bullet$ are identified by $\Pi_{\text{ec}}^{\text{ef}}$ in (103). $\square$

Remark 4.33. We have now defined Kuranishi cohomology $KH^*(Y; R)$, which has the advantage of a cup product $\cup$ which is supercommutative and associative on cochains, but the disadvantage that it works only for R a $\mathbb{Q}$ -algebra. We have also defined effective Kuranishi cohomology $KH_{ec}^*(Y; R)$, which has the advantage that it works for R any commutative ring, but the disadvantage that the cup product $\cup$ is not supercommutative on cochains.

In fact this dichotomy is necessary: it should not be possible to define a cohomology theory which both works when $R = \mathbb{Z}$, and has a cup product $\cup$ on cochains that is everywhere defined and supercommutative. This because of *Steenrod squares* [10, §VI.15–§VI.16], which are cohomology operations defined using the failure of $\cup$ to be supercommutative at the cochain level. Since Steenrod squares are known to be nontrivial when $R = \mathbb{Z}$ (though they are trivial for $R = \mathbb{Q}$), they are an obstruction to defining such a cohomology theory.

To make our theories of effective Kuranishi (co)homology work over $R = \mathbb{Z}$ and compute $H_*^{si}(Y; R)$ and $H_{cs}^*(Y; R)$, we had to sacrifice some good properties of Kuranishi (co)homology, including supercommutativity of $\cup$ on cochains. The argument above suggests these sacrifices were unavoidable.

The next theorem, proved in Appendix D, shows that (at least for manifolds Y) the isomorphisms $\Pi_{cs}^{Kch}, \Pi_{cs}^{ec}, \Pi_{si}^{Kh}, \Pi_{si}^{ef}$ identify the products $\cup, \cap, \bullet$ on $H_{cs}^*, H_*^{si}(Y; R)$ with those on $KH^*, KH_*(Y; R)$ and $KH_{ec}^*, KH_{ec}^*(Y; R)$.

Theorem 4.34. *Let Y be a manifold without boundary, and R a commutative ring (for $KH_{ec}^*, KH_{ec}^*(Y; R)$) or $\mathbb{Q}$ -algebra (for $KH_*, KH^*(Y; R)$). Then*

- (a) *The isomorphisms $\Pi_{cs}^{Kch} : H_{cs}^*(Y; R) \rightarrow KH^*(Y; R)$ and $\Pi_{cs}^{ec} : H_{cs}^*(Y; R) \rightarrow KH_{ec}^*(Y; R)$ identify the cup product $\cup$ on $H_{cs}^*(Y; R)$ with those on $KH^*, KH_{ec}^*(Y; R)$. That is, for all $\alpha, \beta \in H_{cs}^*(Y; R)$ we have*

$$\Pi_{cs}^{Kch}(\alpha \cup \beta) = (\Pi_{cs}^{Kch}(\alpha)) \cup (\Pi_{cs}^{Kch}(\beta)), \quad \Pi_{cs}^{ec}(\alpha \cup \beta) = (\Pi_{cs}^{ec}(\alpha)) \cup (\Pi_{cs}^{ec}(\beta)).$$

- (b) *$\Pi_{si}^{Kh}, \Pi_{si}^{ef}, \Pi_{cs}^{Kch}, \Pi_{cs}^{ec}$ identify the cap product $\cap : H_*^{si}(Y; R) \times H_{cs}^*(Y; R) \rightarrow H_*^{si}(Y; R)$ with those on (effective) Kuranishi (co)homology.*
- (c) *If Y is oriented then $\Pi_{si}^{Kh}, \Pi_{si}^{ef}$ identify the intersection product $\bullet$ on $H_*^{si}(Y; R)$ with those on $KH_*, KH_{ec}^*(Y; R)$.*
- (d) *If Y is compact then $\Pi_{cs}^{Kch}, \Pi_{cs}^{ec}$ take the identity 1 in $H_{cs}^0(Y; R)$ to the identities $[Y, \text{id}_Y, \underline{C}_Y], [Y, \text{id}_Y, \underline{C}_Y]$ in $KH^0, KH_{ec}^0(Y; R)$.*

4.8 Kuranishi (co)homology as a bivariant theory, at the (co)chain level

For manifolds, because of Poincaré duality, homology and cohomology can be identified into a single theory with *products* (cup, cap and intersection products), *pushforwards* (covariant functoriality), and *pullbacks* (contravariant functoriality). Fulton and MacPherson [27] defined *bivariant theories*, a way of naturally

integrating pairs of a homology theory and a cohomology theory into a single larger theory, which works for singular spaces.

We now summarize the notion of a bivariate theory, following [27, §1.1]. Let $\mathcal{C}$ be a category. A bivariate theory from $\mathcal{C}$ to abelian groups (or graded vector spaces, etc.) associates a group $T(f : X \rightarrow Y)$ to each morphism $f : X \rightarrow Y$ in $\mathcal{C}$. It also has three operations:

- **Products.** If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are morphisms in $\mathcal{C}$, and $\alpha \in T(f : X \rightarrow Y)$, $\beta \in T(Y \rightarrow Z)$, we can form the product $\beta \cdot \alpha \in T(g \circ f : X \rightarrow Z)$.
- **Pushforwards.** For a certain class of morphisms $f : X \rightarrow Y$ in $\mathcal{C}$ called *confined morphisms*, if $g : Y \rightarrow Z$ is a morphism in $\mathcal{C}$ and $\alpha \in T(g \circ f : X \rightarrow Z)$ we can form the pushforward $f_*(\alpha) \in T(g : Y \rightarrow Z)$.
- **Pullbacks.** For a certain class of commutative squares in $\mathcal{C}$ called *independent squares*, if

$$\begin{array}{ccc} X' & \xrightarrow{\quad} & X \\ f' \downarrow & g' & \downarrow f \\ Y' & \xrightarrow{\quad g \quad} & Y, \end{array} \quad (120)$$

is an independent square then for each $\alpha \in T(f : X \rightarrow Y)$, we can form the pullback $g^*(\alpha) \in T(f' : X' \rightarrow Y')$.

All this data must satisfy a number of compatibility axioms.

The following example, taken from [27, §3.1] modified as in [27, §3.3.2], combines homology and compactly-supported cohomology into a bivariate theory.

Example 4.35. Let **Top** be the category of topological spaces X which admit an embedding as a closed subspace of $\mathbb{R}^n$ for some $n \gg 0$ (in particular, X must be Hausdorff and paracompact), and morphisms are continuous maps. Let R be a commutative ring. Suppose $f : X \rightarrow Y$ is a morphism in **Top** with X *compact*. Choose an embedding $\phi : X \rightarrow \mathbb{R}^n$ for some $n \gg 0$, and define $H^k(f : X \rightarrow Y; R) = H^{k+n}(Y \times \mathbb{R}^n, Y \times \mathbb{R}^n \setminus (f \times \phi)(X); R)$, using relative cohomology. Then $H^*(f : X \rightarrow Y; R)$ is a graded R -module, which turns out to be independent of the choice of ϕ up to canonical isomorphism. For general morphisms in **Top**, we define $H^*(f : X \rightarrow Y; R)$ to be the direct limit $\varinjlim H^*(f \circ p : K \rightarrow Y; R)$ over all $p : K \rightarrow X$ in **Top** with K compact.

If $\overline{X}$ is a topological space in **Top**, we can associate two natural morphisms in **Top** to X : the projection to a point $\pi : X \rightarrow \{0\}$, and the identity map $\text{id}_X : X \rightarrow X$. By standard algebraic topology one can show that $H^k(\pi : X \rightarrow \{0\}; R) \cong H_{-k}(X; R)$ and $H^k(\text{id}_X : X \rightarrow X; R) \cong H_{\text{cs}}^k(X; R)$. Thus, the bivariate theory H^* specializes to homology and compactly-supported cohomology over R .

Products are defined as in [27, §3.1.7]; the product $H^k(\text{id}_X : X \rightarrow X; R) \times H^l(\text{id}_X : X \rightarrow X; R) \rightarrow H^{k+l}(\text{id}_X : X \rightarrow X; R)$ is $\cup : H_{\text{cs}}^k(X; R) \times H_{\text{cs}}^l(X; R) \rightarrow H_{\text{cs}}^{k+l}(X; R)$, and the product $H^k(\pi : X \rightarrow \{0\}; R) \times H^l(\text{id}_X : X \rightarrow X; R) \rightarrow H^{k+l}(\pi : X \rightarrow \{0\}; R)$ is $\cap : H_{-k}(X; R) \times H_{\text{cs}}^l(X; R) \rightarrow H_{-k-l}(X; R)$.

All morphisms in **Top** are confined, and *pushforwards* are defined as in [27, §3.1.8]. For $f : X \rightarrow Y$ a morphism in **Top**, the pushforward $f_* : H^k(\pi : X \rightarrow \{0\}; R) \rightarrow H^k(\pi : Y \rightarrow \{0\}; R)$ is $f_* : H_{-k}(X; R) \rightarrow H_{-k}(Y; R)$ on homology.

We define *independent squares* to be squares (120) which are Cartesian, so that $X' \cong X \times_{f, Y', g} Y$, and with g proper, so that g' is also proper. *Pullbacks* are defined as in [27, §3.1.6]. If $g : Y' \rightarrow Y$ is a proper morphism in **Top** then $g^* : H^k(\text{id}_Y : Y \rightarrow Y; R) \rightarrow H^k(\text{id}_{Y'} : Y' \rightarrow Y; R)$ is $g^* : H_{\text{cs}}^k(Y; R) \rightarrow H_{\text{cs}}^k(Y'; R)$ on compactly-supported cohomology.

Let **Man** be the category of manifolds, with morphisms smooth maps. Applying the natural functor **Man** $\rightarrow$ **Top** taking a manifold to its underlying topological space, the bivariant theory H^* of Example 4.35 induces a bivariant theory on **Man**. However, because of Poincaré duality the picture simplifies, and we can write H^* on **Man** more directly. The following is an equivalent way of defining the bivariant theory of Example 4.35 on the category **Man**.

Example 4.36. Let **Man** be the category of manifolds, with morphisms smooth maps, and R be a commutative ring. Suppose $f : X \rightarrow Y$ is a morphism in **Man**, with $\dim X = m$ and $\dim Y = n$. Define

$$\begin{aligned} H^k(f : X \rightarrow Y; R) &= H_{\text{cs}}^{k+m-n}(X; O_X \otimes f^*(O_Y) \times_{\{\pm 1\}} R) \\ &\cong H_{n-k}(X; f^*(O_Y) \times_{\{\pm 1\}} R), \end{aligned} \quad (121)$$

where as in Definition 4.16, O_X, O_Y are the *orientation bundles* of X, Y , which are principal $\mathbb{Z}_2$ -bundles over X, Y , so that $O_X \otimes f^*(O_Y)$ is a principal $\mathbb{Z}_2$ -bundle over X , and $H_{\text{cs}}^k(X; O_X \otimes f^*(O_Y) \times_{\{\pm 1\}} R)$ is compactly-supported cohomology of X with coefficients in R twisted by $O_X \otimes f^*(O_Y)$, and $H_*(X; f^*(O_Y) \times_{\{\pm 1\}} R)$ is homology of X with coefficients in R twisted by $f^*(O_Y)$, and the groups on the r.h.s. of (121) are isomorphic by Poincaré duality.

If $f : X \rightarrow Y$ is $\pi : X \rightarrow \{0\}$ then O_Y and $f^*(O_Y)$ are trivial and $n = 0$, so $H^k(\pi : X \rightarrow \{0\}; R) \cong H_{-k}(X; R)$. If $f : X \rightarrow Y$ is $\text{id}_X : X \rightarrow X$ then $f^*(O_Y) \cong O_X$ and $O_X \otimes f^*(O_Y)$ is the trivial $\mathbb{Z}_2$ -bundle over X , so that $H^k(\text{id}_X : X \rightarrow X; R) = H_{\text{cs}}^k(X; R)$. Thus, (121) specializes to homology and compactly-supported cohomology.

To define products, note that the cup product generalizes to twisted cohomology as follows: if $P \rightarrow X$ and $Q \rightarrow X$ are principal $\mathbb{Z}_2$ -bundles then there is a cup product $\cup : H_{\text{cs}}^k(X; P \times_{\{\pm 1\}} R) \times H_{\text{cs}}^l(X; Q \times_{\{\pm 1\}} R) \rightarrow H_{\text{cs}}^{k+l}(X; (P \otimes Q) \times_{\{\pm 1\}} R)$. Suppose $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are smooth with $\dim X = m$, $\dim Y = n$ and $\dim Z = o$. For $\alpha \in H^k(f : X \rightarrow Y; R)$ and $\beta \in H^l(g : Y \rightarrow Z; R)$, define $\beta \cdot \alpha = f^*(\beta) \cup \alpha$. Here $\alpha \in H_{\text{cs}}^{k+m-n}(X; O_X \otimes f^*(O_Y) \times_{\{\pm 1\}} R)$ and $f^*(\beta) \in H_{\text{cs}}^{l+n-o}(X; f^*(O_Y \otimes g^*(O_Z)) \times_{\{\pm 1\}} R)$, and $(O_X \otimes f^*(O_Y)) \otimes f^*(O_Y \otimes g^*(O_Z)) \cong O_X \otimes (g \circ f)^*(O_Z)$ as principal $\mathbb{Z}_2$ -bundles over X , so that $f^*(\beta) \cup \alpha$ lies in $H_{\text{cs}}^{k+l+m-o}(X; O_X \otimes (g \circ f)^*(O_Z) \times_{\{\pm 1\}} R) = H^{k+l}(g \circ f : X \rightarrow Z; R)$.

All morphisms in **Man** are confined. *Pushforwards* $f_* : H^k(g \circ f : X \rightarrow Z; R) \rightarrow H^k(g : Y \rightarrow Z; R)$ are given by pushforwards $f_* : H_{o-k}(X; (g \circ f)^*(O_Z) \times_{\{\pm 1\}} R) \rightarrow H_{o-k}(Y; g^*(O_Z) \times_{\{\pm 1\}} R)$ in twisted homology. *Independent squares* are Cartesian squares (120) in **Man** with g proper, and hence g'

proper. Then with $\dim X = m$, $\dim Y = n$, $\dim X' = m'$, $\dim Y' = n'$, the Cartesian property gives $(g')^*(O_X \otimes f^*(O_Y)) \cong O_{X'} \otimes (f')^*(O_{Y'})$ and $m - n = m' - n'$. So we define $g^* : H^k(f : X \rightarrow Y; R) \rightarrow H^k(f' : X' \rightarrow Y'; R)$ to be

$$\begin{aligned} (g')^* : H_{\text{cs}}^{k+m-n}(X; O_X \otimes f^*(O_Y) \times_{\{\pm 1\}} R) &\longrightarrow \\ H_{\text{cs}}^{k+m-n}(X'; (g')^*(O_X \otimes f^*(O_Y)) \times_{\{\pm 1\}} R) &\cong \\ H_{\text{cs}}^{k+m'-n'}(X'; O_{X'} \otimes (f')^*(O_{Y'}) \times_{\{\pm 1\}} R), & \end{aligned}$$

using proper pullbacks on twisted compactly-supported cohomology.

For *oriented* manifolds X, Y , with O_X, O_Y trivial, equation (121) becomes

$$H^k(f : X \rightarrow Y; R) = H_{\text{cs}}^{k+m-n}(X; R) \cong H_{n-k}(X; R),$$

and we lose the dependence on f, Y entirely. Thus, for oriented manifolds and smooth maps, the bivariant theories of Examples 4.35 and 4.36 reduce to homology, compactly-supported cohomology, and Poincaré duality. That is, *bivariant (co)homology theories really give us nothing new for manifolds*, they are of interest *only for singular spaces*.

As our Kuranishi (co)homology theories are defined only for manifolds and orbifolds, and bivariant (co)homology theories give nothing new for manifolds, one might expect that there is nothing to be gained from recasting Kuranishi (co)homology as a bivariant theory. However, as we have already emphasized, Kuranishi (co)homology is particularly well behaved at the (co)chain level, with pushforwards, pullbacks, and cup and cap products on (co)chains all well-defined and with all the properties one would wish.

We will now explain how to formulate (effective) Kuranishi (co)homology as a bivariant theory *at the (co)chain level*. The author does not know of any other (co)homology theories for which this can be done. This bivariant framework is likely to be valuable in situations in which one is forced to work with (co)chains rather than just with (co)homology, for example, in defining A_∞ -algebras in Lagrangian Floer cohomology, as in [4] and sketched in §6.6.

Definition 4.37. Let **Orb** be the category of orbifolds without boundary, with morphisms smooth maps, and R be a commutative ring. Roughly speaking, we will define a bivariant theory from **Orb** to the category of differential graded R -modules, generalizing Kuranishi (co)chains. However, we will define $KC^*(g : Y \rightarrow Z; R)$ only when $g : Y \rightarrow Z$ is a *submersion* of orbifolds Y, Z .

Let Y, Z be orbifolds without boundary and $g : Y \rightarrow Z$ be a submersion. Consider triples $(X, \mathbf{f}, \mathbf{B})$, where X is a compact Kuranishi space, $\mathbf{f} : X \rightarrow Y$ is a strongly smooth map such that $g \circ \mathbf{f} : X \rightarrow Z$ is a strong submersion, and we are given a coorientation for $(X, g \circ \mathbf{f})$, and $\mathbf{B} = (\mathbf{I}, \boldsymbol{\eta}, B^i : i \in I)$, where $(\mathbf{I}, \boldsymbol{\eta})$ is an excellent coordinate system for $(X, \mathbf{f})$ with $\mathbf{I} = (I, (V^i, \dots, \psi^i), f^i : i \in I, \dots)$, and $g \circ f^i : V^i \rightarrow Z$ are submersions representing $g \circ \mathbf{f}$ for all $i \in I$, and $B^i : E^i \rightarrow P$ for $i \in I$ are maps such that each B^i maps $E^i \rightarrow P_l \subset P$ for $l \gg 0$, and $B^i \times (g \circ f^i \circ \pi^i) : E^i \rightarrow P \times Y$ is a globally finite map for all $i \in I$. Such $\mathbf{B}$ generalize the notions of (co-)gauge-fixing data in §3.1, since

if $g : Y \rightarrow Z$ is $\pi : Y \rightarrow \{0\}$ then $\mathbf{B}$ is gauge-fixing data for $(X, \mathbf{f})$, and if $g : Y \rightarrow Z$ is $\text{id}_Y : Y \rightarrow Y$ then $\mathbf{B}$ is co-gauge-fixing data for $(X, \mathbf{f})$. We call $\mathbf{B}$ *bi-gauge-fixing data*.

Define *isomorphisms* $(\mathbf{a}, \mathbf{b})$ of such triples $(X, \mathbf{f}, \mathbf{B})$ as for (co-)gauge-fixing data in §3.1. Write $[X, \mathbf{f}, \mathbf{B}]$ for the isomorphism class of $(X, \mathbf{f}, \mathbf{B})$ under isomorphisms $(\mathbf{a}, \mathbf{b}) : (X, \mathbf{f}, \mathbf{B}) \rightarrow (\tilde{X}, \tilde{\mathbf{f}}, \tilde{\mathbf{B}})$ which identify the coorientations of $(X, g \circ \mathbf{f})$ and $(\tilde{X}, g \circ \tilde{\mathbf{f}})$.

For each $k \in \mathbb{Z}$, define $KC^k(g : Y \rightarrow Z; R)$ to be the R -module of finite R -linear combinations of isomorphism classes $[X, \mathbf{f}, \mathbf{G}]$ for which $\text{vdim } X = \dim Z - k$, with the analogues of relations Definition 4.3(i)–(iv). Then $KC^k(\pi : Y \rightarrow \{0\}; R) = KC_{-k}(Y; R)$ and $KC^k(\text{id}_Y : Y \rightarrow Y; R) = KC^k(Y; R)$, so this generalizes the definitions of Kuranishi (co)chains in §4.2 and §4.4. Elements of $KC^*(g : Y \rightarrow Z; R)$ will be called *Kuranishi bichains*.

Define $d : KC^k(g : Y \rightarrow Z; R) \rightarrow KC^{k+1}(g : Y \rightarrow Z; R)$ by

$$d : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{B}_a] \mapsto \sum_{a \in A} \rho_a [\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{B}_a|_{\partial X_a}],$$

as in (66) and (78). Then $d \circ d = 0$. Define $KH^*(g : Y \rightarrow Z; R)$ to be the cohomology of $(KC^*(g : Y \rightarrow Z; R), d)$. Then $KH^k(\pi : Y \rightarrow \{0\}; R) = KH_{-k}(Y; R)$ and $KH^k(\text{id}_Y : Y \rightarrow Y; R) = KH^k(Y; R)$. Combining the proofs of Corollary 4.10 and Theorem 4.15 shows that $KH^k(g : Y \rightarrow Z; R) \cong H_{\dim Z - k}^{\text{si}}(Y; g^*(O_Z) \times_{\{\pm 1\}} R)$, as in (121).

We define the three bivariant operations on Kuranishi bichains:

- **Products.** Let X, Y, Z be orbifolds without boundary, and $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be submersions. Define a product $KC^k(g : Y \rightarrow Z; R) \times KC^l(f : X \rightarrow Y; R) \rightarrow KC^{k+l}(g \circ f : X \rightarrow Z; R)$ by

$$[W, \mathbf{e}, \mathbf{B}] \cdot [\tilde{W}, \tilde{\mathbf{e}}, \tilde{\mathbf{B}}] = [W \times_{\mathbf{e}, Y, f \circ \tilde{\mathbf{e}}} \tilde{W}, \tilde{\mathbf{e}} \circ \pi_{\tilde{W}}, \mathbf{B} \times_Y \tilde{\mathbf{B}}], \quad (122)$$

extended R -bilinearly. Here $[W, \mathbf{e}, \mathbf{B}] \in KC^k(g : Y \rightarrow Z; R)$, so that $\mathbf{e} : W \rightarrow Y$ is strongly smooth, and $[\tilde{W}, \tilde{\mathbf{e}}, \tilde{\mathbf{B}}] \in KC^l(f : X \rightarrow Y; R)$ so that $f \circ \tilde{\mathbf{e}} : \tilde{W} \rightarrow Y$ is a strong submersion. Thus $W \times_{\mathbf{e}, Y, f \circ \tilde{\mathbf{e}}} \tilde{W}$ is a compact Kuranishi space, and $\tilde{\mathbf{e}} \circ \pi_{\tilde{W}} : W \times_{\mathbf{e}, Y, f \circ \tilde{\mathbf{e}}} \tilde{W}$ is strongly smooth.

We have $f \circ (\tilde{\mathbf{e}} \circ \pi_{\tilde{W}}) = (f \circ \tilde{\mathbf{e}}) \circ \pi_{\tilde{W}} = \pi_Y = \mathbf{e} \circ \pi_W$. Composing with g then gives $(g \circ f) \circ (\tilde{\mathbf{e}} \circ \pi_{\tilde{W}}) = (g \circ \mathbf{e}) \circ \pi_W$. But $g \circ \mathbf{e}$ is a strong submersion as $[W, \mathbf{e}, \mathbf{B}] \in KC^k(g : Y \rightarrow Z; R)$, and π_W is a strong submersion as $f \circ \tilde{\mathbf{e}}$ is, by properties of fibre products. Hence $(g \circ f) \circ (\tilde{\mathbf{e}} \circ \pi_{\tilde{W}})$ is a strong submersion, as we need for the r.h.s. of (122) to lie in $KC^{k+l}(g \circ f : X \rightarrow Z; R)$. The coorientations for $(W, g \circ \mathbf{e})$ and $(\tilde{W}, f \circ \tilde{\mathbf{e}})$ combine to give a coorientation for $(W \times_{\mathbf{e}, Y, f \circ \tilde{\mathbf{e}}} \tilde{W}, (g \circ f) \circ (\tilde{\mathbf{e}} \circ \pi_{\tilde{W}}))$, using $(g \circ f) \circ (\tilde{\mathbf{e}} \circ \pi_{\tilde{W}}) = (g \circ \mathbf{e}) \circ \pi_W$. The fibre product of bi-gauge-fixing data $\mathbf{B} \times_Y \tilde{\mathbf{B}}$ is defined as for $\mathbf{C} \times_Y \tilde{\mathbf{C}}$ in §3.8.

When $X = Y = Z$ and $f = g = \text{id}_Y$, equation (122) agrees with $\cup$ in (105), and when $X = Y$, $f = \text{id}_Y$, $Z = \{0\}$ and $g = \pi : Y \rightarrow \{0\}$, equation (122) agrees with $\cap$ in (108). Thus, the product (122) generalizes $\cup, \cap$ on Kuranishi (co)chains.

- **Pushforwards.** Let X, Y, Z be orbifolds without boundary, and $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be smooth with $g, g \circ f$ submersions. Define $f_* : KC^k(g \circ f : X \rightarrow Z; R) \rightarrow KC^k(g : Y \rightarrow Z; R)$ by

$$f_* : \sum_{a \in A} \rho_a [W_a, e_a, B_a] \mapsto \sum_{a \in A} \rho_a [W_a, f \circ e_a, f_*(B_a)],$$

as in (69), where $f_*(B_a)$ is defined as for $h_*(G)$ in §3.7. Note that as $(g \circ f) \circ e_a$ is a strong submersion with $(W_a, (g \circ f) \circ e_a)$ cooriented, trivially $g \circ (f \circ e_a)$ is a strong submersion with $(W_a, g \circ (f \circ e_a))$ cooriented.

- **Pullbacks.** Let (120) be a Cartesian square of orbifolds without boundary, with f (and hence f') a submersion, and g (and hence g') proper. Define $g^* : KC^k(f : X \rightarrow Y; R) \rightarrow KC^k(f' : X' \rightarrow Y'; R)$ by

$$g^* : \sum_{a \in A} \rho_a [W_a, e_a, B_a] \mapsto \sum_{a \in A} \rho_a [X' \times_{g', X, e_a} W_a, \pi_{X'}, (g')^*(B_a)],$$

as in (79), where $(g')^*(B_a)$ is defined as for $h^*(C)$ in §4.8. Since $f \circ e_a : W_a \rightarrow Y$ is a strong submersion and (120) is Cartesian, $f' \circ \pi_{X'} : X' \times_{g', X, e_a} W_a \rightarrow Y'$ is a strong submersion. Also, using the coorientation for $(W_a, f \circ e_a)$ we construct a coorientation for $(X' \times_{g', X, e_a} W_a, f' \circ \pi_{X'})$.

It is easy to show that these operations satisfy the axioms for a bivariant theory in [27, §2.2]. Also, the three operations are compatible with differentials d on $KC^*(g : Y \rightarrow Z; R)$, and so induce operations on $KH^*(g : Y \rightarrow Z; R)$.

It is easy to generalize Definition 4.37 to *effective* Kuranishi (co)homology. The differences are that we restrict to submersions $g : Y \rightarrow Z$ for which $g_* : \text{Stab}_Y(y) \rightarrow \text{Stab}_Z(g(y))$ is *surjective* for all $y \in Y$, and in defining $KC_{\text{ef}}^k(g : Y \rightarrow Z; R)$, we use triples $(X, \mathbf{f}, \underline{B})$ in which $(X, g \circ \mathbf{f})$ is also *coeffective*, and $\underline{B} = (\mathbf{I}, \boldsymbol{\eta}, \underline{B}^i : i \in I)$, where $(V^i, \dots, \psi^i), g \circ f^i$ is *coeffective* for all $i \in I$, and $\underline{B}^i : E^i \rightarrow \underline{P}$ are maps such that $\underline{B}^i \times (g \circ f^i \circ \pi^i) : E^i \rightarrow \underline{P} \times Z$ satisfy the conditions of Definition 3.38.

4.9 Why we need (co-)gauge-fixing data

To show that we really do need some form of (co-)gauge-fixing data to define sensible (co)homology theories using Kuranishi spaces as chains, we will now prove that if we omit (co-)gauge-fixing data from the definitions above, then Kuranishi (co)homology is always zero.

Let Y be an orbifold without boundary, and R a commutative ring. Define *naïve Kuranishi chains* $KC_k^{\text{na}}(Y; R)$ to be the R -module spanned by isomorphism classes $[X, \mathbf{f}]$ of pairs $(X, \mathbf{f})$ with X a compact, oriented Kuranishi space with $\text{vdim } X = k$ and $\mathbf{f} : X \rightarrow Y$ strongly smooth, with the analogues of relations Definition 4.3(i)–(iii). The boundary operator is $\partial : [X, \mathbf{f}] \mapsto [\partial X, \mathbf{f}|_{\partial X}]$. Let *naïve Kuranishi homology* $KH_*^{\text{na}}(Y; R)$ be the homology of $(KC_*^{\text{na}}(Y; R), \partial)$.

Similarly, define *naïve Kuranishi cochains* $KC_{\text{na}}^k(Y; R)$ to be spanned by isomorphism classes $[X, \mathbf{f}]$ with X a compact Kuranishi space with $\text{vdim } X =$

$\dim Y - k$ and $\mathbf{f} : X \rightarrow Y$ a cooriented strong submersion, and the same relations. Define $d : [X, \mathbf{f}] \mapsto [\partial X, \mathbf{f}|_{\partial X}]$, and let *naïve Kuranishi cohomology* $KH_{\text{na}}^*(Y; R)$ be the cohomology of $(KC_{\text{na}}^*(Y; R), d)$. One can then define cup and cap products $\cup, \cap$ on $KC_{\text{na}}^*, KC_{\text{na}}^{\text{na}}(Y; R)$ and $KH_{\text{na}}^*, KH_{\text{na}}^{\text{na}}(Y; R)$, as in §4.7, but omitting all (co-)gauge-fixing data.

Take Y to be a point $\{0\}$, and for any Kuranishi space X write $\pi : X \rightarrow \{0\}$ for the trivial projection. Let R be a commutative ring. In the next example we show that the identity $[[\{0\}, \text{id}_{\{0\}}]]$ in $KH_{\text{na}}^*(\{0\}; R)$ is zero. From this we will deduce that $KH_{\text{na}}^*(Y; R) = KH_{\text{na}}^{\text{na}}(Y; R) = 0$ for all Y .

Example 4.38. Let $L \rightarrow \mathbb{CP}^1$ be the complex line bundle $\mathcal{O}(1)$. Define a Kuranishi space X_k for $k \in \mathbb{Z}$ to be the topological space $\mathbb{CP}^1$ with the Kuranishi structure induced by the Kuranishi neighbourhood $(\mathbb{CP}^1, L^k, 0, \text{id}_{\mathbb{CP}^1})$, with obstruction bundle $L^k \rightarrow \mathbb{CP}^1$. Give (X_k, π) the obvious coorientation. Then $\text{vdim } X_k = 0$, and $\partial X_k = \emptyset$, so $[X_k, \pi]$ defines a class $[[X_k, \pi]]$ in $KH_{\text{na}}^0(\{0\}; R)$.

When $k \geq 0$ we can choose a generic smooth section s of $L^k \rightarrow \mathbb{CP}^1$ which has exactly k zeroes $x_1, \dots, x_k$, each of multiplicity 1. Let t be the coordinate on $[0, 1]$. Then ts is a section of $L^k \rightarrow [0, 1] \times \mathbb{CP}^1$, with $(ts)^{-1}(0) = \{0\} \times \mathbb{CP}^1 \cup [0, 1] \times \{x_1, \dots, x_k\}$, and $([0, 1] \times \mathbb{CP}^1, L^k, ts, \text{id}_{(ts)^{-1}(0)})$ is a Kuranishi neighbourhood on $(ts)^{-1}(0)$, making it into a Kuranishi space of virtual dimension 1. By taking the boundary of this we see that

$$[[X_k, \pi]] = [[\{x_1, \dots, x_k\}, \pi]] = k[[\{0\}, \text{id}_{\{0\}}]] \quad (123)$$

in $KH_{\text{na}}^0(\{0\}; R)$, that is, $[[X_k, \pi]]$ is k times the fundamental class of $Y = \{0\}$.

Write $[z_0, z_1]$ for the homogeneous coordinates on $\mathbb{CP}^1$, and define

$$V = \{(t, [z_0, z_1]) \in \mathbb{R} \times \mathbb{CP}^1 : \min(|z_0|^2, |z_1|^2) \max(|z_0|^2, |z_1|^2)^{-1} \leq t \leq 2\}.$$

Then V is a compact oriented 3-manifold with corners, and ∂V is the disjoint union of three pieces, a copy of $\mathbb{CP}^1$ with $t = 2$, the hemisphere $H_+ = \{[z_0, z_1] \in \mathbb{CP}^1 : |z_0| \leq |z_1|\}$ with $t = |z_0|^2/|z_1|^2$, and the hemisphere $H_- = \{[z_0, z_1] \in \mathbb{CP}^1 : |z_0| \geq |z_1|\}$ with $t = |z_1|^2/|z_0|^2$.

Define the Kuranishi space W_k for $k \in \mathbb{Z}$ to be the topological space V with the Kuranishi structure induced by the neighbourhood $(V, \pi^*(L^k), 0, \text{id}_V)$, where $\pi : V \rightarrow \mathbb{CP}^1$ is the projection. Define Kuranishi spaces X_+, X_- to be the topological spaces H_+, H_- with the Kuranishi structure induced by the Kuranishi neighbourhoods $(H_{\pm}, L^0, 0, \text{id}_{H_{\pm}})$. Now the line bundles $L^k \rightarrow H^{\pm}$ are for $k \in \mathbb{Z}$ are isomorphic to $L^0 \rightarrow H^{\pm}$. Thus there is an isomorphism of cooriented Kuranishi spaces $(\partial W_k, \pi) \cong (X_k, \pi) \amalg -(X_+, \pi) \amalg -(X_-, \pi)$, so in $KH_{\text{na}}^0(\{0\}; R)$ we have $[[X_k, \pi]] = [[X_+, \pi]] + [[X_-, \pi]]$, and the class $[[X_k, \pi]]$ is independent of $k \in \mathbb{Z}$. Combining this with (123) gives $[[\{0\}, \text{id}_{\{0\}}]] = 0$.

If Y is any compact orbifold without boundary, and $\pi : Y \rightarrow \{0\}$ is the projection, then the pullback $\pi^* : KH_{\text{na}}^*(\{0\}; R) \rightarrow KH_{\text{na}}^*(Y; R)$ takes $\pi^* : [[\{0\}, \text{id}_{\{0\}}]] \mapsto [[Y, \text{id}_Y]]$, so $[[\{0\}, \text{id}_{\{0\}}]] = 0$ implies $[[Y, \text{id}_Y]] = 0$. But

$[[Y, \text{id}_Y]]$ is the identity for $\cup$ on $KH_{\text{na}}^*(Y; R)$ and $\cap$ on $KH_*^{\text{na}}(Y; R)$, so this implies $KH_{\text{na}}^*(Y; R) = KH_*^{\text{na}}(Y; R) = 0$. Using a related argument on (co)chains, we can also show $KH_{\text{na}}^*(Y; R) = KH_*^{\text{na}}(Y; R) = 0$ for any orbifold Y . Hence, Kuranishi (co)homology without (co-)gauge-fixing data is vacuous.

The reason this example goes so badly wrong is that the Kuranishi space $\partial X^+ = -\partial X^-$, the circle C with Kuranishi neighbourhood $(C, \mathbb{R}^2 \times C, 0, \text{id}_C)$, has a *large automorphism group*, including topologically nontrivial automorphisms which fix C but rotate the fibres of the obstruction bundle $\mathbb{R}^2 \times C$ by a degree l smooth map $C \rightarrow \text{SO}(2)$. Thus, by cutting X_k into two pieces X^+ , X^- and then gluing them together again twisted by such a nontrivial automorphism, we can get X_{k+l} for any $l \in \mathbb{Z}$, which forces $[[X_k, \pi]] = [[X_{k+l}, \pi]]$ in $KH_{\text{na}}^0(\{0\}; R)$. Including (co-)gauge-fixing data avoids this problem, as by Theorem 3.20 it allows only finite automorphism groups.

5 Kuranishi bordism and cobordism

We now discuss *bordism* and *cobordism* theories of an orbifold Y and a commutative ring R , defined in a similar way to Kuranishi (co)homology. In fact we will define two different classical and five different Kuranishi bordism theories, and five corresponding Kuranishi cobordism theories. But these are all variations on the same idea: *Kuranishi bordism groups* $KB_*(Y; R)$ are spanned over R by isomorphism classes $[X, \mathbf{f}]$ for X a compact oriented Kuranishi space without boundary and $\mathbf{f} : X \rightarrow Y$ strongly smooth, up to bordism in Kuranishi spaces, and *Kuranishi cobordism groups* $KB^*(Y; R)$ are spanned over R by $[X, \mathbf{f}]$ for X a compact Kuranishi space without boundary and $\mathbf{f} : X \rightarrow Y$ a strong submersion with $(X, \mathbf{f})$ cooriented, up to bordism in Kuranishi spaces.

5.1 Classical bordism and cobordism groups

Bordism groups were introduced by Atiyah [5], and Connor [16, §I] gives a good introduction. Other useful references on bordism and cobordism are Stong [74] and Connor and Floyd [17, 18]. Our definition is not standard, but fits in with our later work.

Definition 5.1. Let Y be an orbifold. Consider pairs (X, f) , where X is a compact, oriented manifold without boundary or corners, not necessarily connected, and $f : X \rightarrow Y$ is a smooth map. An *isomorphism* between two such pairs $(X, f), (\tilde{X}, \tilde{f})$ is an orientation-preserving diffeomorphism $i : X \rightarrow \tilde{X}$ with $f = \tilde{f} \circ i$. Write $[X, f]$ for the isomorphism class of (X, f) .

Let R be a commutative ring, for instance $\mathbb{Z}, \mathbb{Q}, \mathbb{R}$ or $\mathbb{C}$. For each $k \geq 0$, define the k^{th} *bordism group* $B_k(Y; R)$ of Y with coefficients in R to be the R -module of finite R -linear combinations of isomorphism classes $[X, f]$ for which $\dim X = k$, with the relations:

- (i) $[X, f] + [X', f'] = [X \amalg X', f \amalg f']$ for all classes $[X, f], [X', f']$; and

- (ii) Suppose W is a compact, oriented $(k+1)$ -manifold with boundary but without (g-)corners, and $e : W \rightarrow Y$ is smooth. Then $[\partial W, e|_{\partial W}] = 0$.

If $h : Y \rightarrow Z$ is a smooth map of orbifolds, the *pushforward* $h_* : B_k(Y; R) \rightarrow B_k(Z; R)$ acts by $h_* : \sum_{a \in A} \rho_a[X_a, f_a] \mapsto \sum_{a \in A} \rho_a[X_a, h \circ f_a]$. This takes relations (i),(ii) in $B_k(Y; R)$ to (i),(ii) in $B_k(Z; R)$, and so is well-defined.

Define a projection $\Pi_{\text{bo}}^{\text{si}} : B_k(Y; R) \rightarrow H_k^{\text{si}}(Y; R)$ by $\Pi_{\text{bo}}^{\text{si}} : \sum_{a \in A} \rho_a[X_a, f_a] \mapsto \sum_{a \in A} \rho_a(f_a)_*([X_a])$, where $[X_a] \in H_k^{\text{si}}(X_a; R)$ is the fundamental class of X_a .

The following consequence of relations Definition 5.1(i),(ii) will be useful, and also works for all the other (co)bordism theories below. Let $[X, f] \in KB_k(Y; R)$, and write $-X$ for X with the opposite orientation. Taking $W = [0, 1] \times X$ and $e : W \rightarrow Y$ to be $f \circ \pi_X$, we have $\partial W = -\{0\} \times X \amalg \{1\} \times X \cong -X \amalg X$ in oriented manifolds. Using (i),(ii) we see that in $B_k(Y; R)$ we have

$$[-X, f] = -[X, f]. \quad (124)$$

Here is how this definition relates to those in [5, 16]. When Y is a manifold and $R = \mathbb{Z}$, our $B_k(Y; \mathbb{Z})$ is equivalent to Connor's *differential bordism group* $D_k(Y)$, [16, §I.9]. Atiyah [5, §2] and Connor [16, §I.4] also define *bordism groups* $MSO_k(Y)$ as for $B_k(Y; \mathbb{Z})$ above, but only requiring $f : X \rightarrow Y$ to be continuous, not smooth. Connor [16, Th. I.9.1] shows that when Y is a manifold, the natural projection $D_k(Y) \rightarrow MSO_k(Y)$ is an isomorphism.

As in [16, §I.5], bordism is a *generalized homology theory* in the sense of Whitehead [75], that is, it satisfies all the Eilenberg–Steenrod axioms for a homology theory except the dimension axiom. The bordism groups of a point $MSO_*(\text{pt})$ are well understood [16, §I.2]: Thom showed that $MSO_*(\text{pt}) \otimes_{\mathbb{Z}} \mathbb{Q}$ is the free commutative algebra over $\mathbb{Q}$ generated by $\zeta_{4k} = [\mathbb{C}P^{2k}] \in MSO_{4k}(\text{pt})$ for $k = 1, 2, \dots$, and Wall and others determined the torsion of $MSO_*(\text{pt})$, which is all of order 2. This gives some information on bordism groups of general spaces Y : for any generalized homology theory $GH_*(Y)$, there is a spectral sequence from the singular homology $H_*^{\text{si}}(Y; GH_*(\text{pt}))$ of Y with coefficients in $GH_*(\text{pt})$ converging to $GH_*(Y)$, so that $GH_*(S^n) \cong H_*^{\text{si}}(S^n; GH_*(\text{pt}))$, for instance.

Atiyah [5] and Connor [16, §13] also define *cobordism groups* $MSO^k(Y)$ for $k \in \mathbb{Z}$, which are a *generalized cohomology theory* dual to bordism $MSO_k(Y)$. There is a natural product $\cup$ on $MSO^*(Y)$, making it into a supercommutative ring. If Y is a compact oriented n -manifold without boundary then [5, Th. 3.6], [16, Th. 13.4] there are canonical Poincaré duality isomorphisms

$$MSO^k(Y) \cong MSO_{n-k}(Y) \quad \text{for } k \in \mathbb{Z}. \quad (125)$$

The definition of $MSO^*(Y)$ uses homotopy theory, direct limits of k -fold suspensions, and classifying spaces. There does not seem to be a satisfactory differential-geometric definition of cobordism groups parallel to Definition 5.1.

If Y is compact and oriented then from the isomorphisms (125) and the product $\cup$ on cobordism we obtain an associative, supercommutative *intersection product* $\bullet$ on bordism groups. (Actually, Y does not need to be compact

here.) We can write this product geometrically for the groups $B_k(Y; R)$ as follows. Suppose $[X, f]$ and $[\tilde{X}, \tilde{f}]$ are isomorphism classes in $B_*(Y; R)$ such that if $f(x) = \tilde{f}(\tilde{x}) = y \in Y$ then $T_y Y = \mathrm{d}f|_x(T_x X) + \mathrm{d}\tilde{f}|_{\tilde{x}}(T_{\tilde{x}} \tilde{X})$; essentially this says that $f(X)$ and $\tilde{f}(\tilde{X})$ intersect transversely in Y . Then $X \times_{f, Y, \tilde{f}} \tilde{X}$ is a compact, oriented manifold with smooth $\pi : X \times_{f, Y, \tilde{f}} \tilde{X} \rightarrow Y$, and one can show that

$$[X, f] \bullet [\tilde{X}, \tilde{f}] = [X \times_{f, Y, \tilde{f}} \tilde{X}, \pi]. \quad (126)$$

The bordism and cobordism theories $MSO_*(Y), MSO^*(Y)$ are generalized (co)homology theories defined using oriented n -manifolds X , that is, manifolds X whose tangent bundle has an $SO(n)$ -structure. In a similar way, for each of the series of Lie groups $O(n), U(n), SU(n), Sp(n)$, one can define bordism and cobordism theories built using manifolds X whose (stable) tangent bundle has a G -structure for $G = O(n), U(n), SU(n)$ or $Sp(n)$. So we can define *unoriented bordism* $MO_*(Y)$, *unoriented cobordism* $MO^*(Y)$, *complex bordism* or *unitary bordism* $MU_*(Y)$, *complex cobordism* or *unitary cobordism* $MU^*(Y)$, and so on. This is explained by Connor and Floyd [17, 18] and Stong [74]. We will not discuss them. However, note that $MU_*(Y), MU^*(Y)$ should be related to the almost complex Kuranishi bordism groups $KB_*^{\mathrm{ac}}, KB_*^{\mathrm{eac}}(Y; R)$ and $KB_{\mathrm{ca}}^*, KB_{\mathrm{eca}}^*(Y; R)$ defined below.

One can also repeat Definition 5.1 using orbifolds rather than manifolds. We choose to do this using *effective* orbifolds, as the resulting groups will turn out to be closely related to effective Kuranishi bordism and effective Kuranishi homology. As far as the author knows, the next definition is new.

Definition 5.2. Let Y be an orbifold, and R be a commutative ring. Define the *effective orbifold bordism group* $B_k^{\mathrm{eo}}(Y; R)$ of Y as in Definition 5.1, except that we allow X, W to be effective orbifolds rather than manifolds. *Pushforwards* $h_* : B_k^{\mathrm{eo}}(Y; R) \rightarrow B_k^{\mathrm{eo}}(Z; R)$ for $h : Y \rightarrow Z$ smooth, and projections $\Pi_{\mathrm{eo}}^{\mathrm{si}} : B_k^{\mathrm{eo}}(Y; R) \rightarrow H_k^{\mathrm{si}}(Y; R)$, are defined as in Definition 5.1. For $\Pi_{\mathrm{eo}}^{\mathrm{si}}$, when R is not a $\mathbb{Q}$ -algebra, it is essential that the X_a are *effective* orbifolds, as the fundamental class $[X] \in H_k^{\mathrm{si}}(X; R)$ of a compact oriented orbifold X without boundary is defined for all R when X is effective, but only when $\mathbb{Q} \subseteq R$ otherwise.

5.2 Kuranishi bordism groups

Motivated by §4.2 and §5.1, we define:

Definition 5.3. Let Y be an orbifold. Consider pairs $(X, \mathbf{f})$, where X is a compact oriented Kuranishi space without boundary, and $\mathbf{f} : X \rightarrow Y$ is strongly smooth. An *isomorphism* between two pairs $(X, \mathbf{f}), (\tilde{X}, \tilde{\mathbf{f}})$ is an orientation-preserving strong diffeomorphism $\mathbf{i} : X \rightarrow \tilde{X}$ with $\mathbf{f} = \tilde{\mathbf{f}} \circ \mathbf{i}$. Write $[X, \mathbf{f}]$ for the isomorphism class of $(X, \mathbf{f})$.

Let R be a commutative ring. For each $k \in \mathbb{Z}$, define the k^{th} *Kuranishi bordism group* $KB_k(Y; R)$ of Y with coefficients in R to be the R -module of finite R -linear combinations of isomorphism classes $[X, \mathbf{f}]$ for which $\mathrm{vdim} X = k$, with the relations:

- (i) $[X, \mathbf{f}] + [X', \mathbf{f}'] = [X \amalg X', \mathbf{f} \amalg \mathbf{f}']$ for all classes $[X, \mathbf{f}], [X', \mathbf{f}']$; and
- (ii) Suppose W is a compact oriented Kuranishi space with boundary but without (g-)corners, with $\text{vdim } W = k + 1$, and $\mathbf{e} : W \rightarrow Y$ is strongly smooth. Then $[\partial W, \mathbf{e}|_{\partial W}] = 0$.

Elements of $KB_k(Y; R)$ will be called *Kuranishi bordism classes*.

Let $h : Y \rightarrow Z$ be a smooth map of orbifolds. Define the *pushforward* $h_* : KB_k(Y; R) \rightarrow KB_k(Z; R)$ by $h_* : \sum_{a \in A} \rho_a[X_a, \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a[X_a, h \circ \mathbf{f}_a]$. This takes relations (i), (ii) in $KB_k(Y; R)$ to (i), (ii) in $KB_k(Z; R)$, and so is well-defined. Pushforward is functorial, that is, $(g \circ h)_* = g_* \circ h_*$.

We also define two minor variations on $KB_*(Y; R)$:

- Define *effective Kuranishi bordism* $KB_*^{\text{eb}}(Y; R)$ as above, except that we require X, W to be *effective* Kuranishi spaces, as in Definition 3.35.
- Similarly, define *trivial stabilizers Kuranishi bordism* $KB_*^{\text{tr}}(Y; R)$ as above, except that we require X, W to be Kuranishi spaces with *trivial stabilizers*, that is, $\text{Stab}_X(p) = \{1\}$ for all $p \in X$.

Pushforwards h_* also work on $KB_*^{\text{eb}}(Y; R)$ and $KB_*^{\text{tr}}(Y; R)$.

Our reason for introducing effective Kuranishi bordism $KB_*^{\text{eb}}(Y; R)$ is that it maps naturally to effective Kuranishi homology $KH_*^{\text{ef}}(Y; R)$, and hence to singular homology $H_*^{\text{si}}(Y; R)$, as we will see in §5.3. Since this works over $R = \mathbb{Z}$, this makes $KB_*^{\text{eb}}(Y; R)$ a useful tool for considering questions of integrality of homology classes, such as the Gopakumar–Vafa Integrality Conjecture for Gromov–Witten invariants, to be discussed in §6.3.

Our reason for defining trivial stabilizers Kuranishi bordism $KB_*^{\text{tr}}(Y; R)$ is that in §5.3 we will show that $KB_*^{\text{tr}}(Y; R) \cong B_*(Y; R)$, and so $KB_*^{\text{tr}}(Y; R)$ helps to clarify the relationship between classical bordism theories and our new Kuranishi bordism theories. We will also prove in §5.7 that $KB_*(Y; R)$ and $KB_*^{\text{eb}}(Y; R)$ are generally very large. These are nontrivial facts, as it is not obvious from Definition 5.3 that $KB_*, KB_*^{\text{eb}}, KB_*^{\text{tr}}(Y; R)$ can ever be nonzero.

For example, if we omitted orientations in Definition 5.3, so that $KB_k(Y; R)$ were generated by $[X, \mathbf{f}]$ for X a compact, *unoriented* Kuranishi space, then we would have $KB_k(Y; R) = \{0\}$ for all Y, k . This is because for any $[X, \mathbf{f}]$ we may take $W = ([-1, 1]/\{\pm 1\}) \times X$, a non-orientable Kuranishi space, and $\mathbf{e} : W \rightarrow Y$ to be $\mathbf{e} = \mathbf{f} \circ \pi_X$, and then $[X, \mathbf{f}] = [\partial W, \mathbf{e}|_{\partial W}] = 0$ by Definition 5.3(ii) with orientations omitted. Here $[-1, 1]/\{\pm 1\}$ is a compact, non-orientable 1-orbifold with boundary one point.

Using the notation of §2.9, we add almost complex structures to Definition 5.3 to define *almost complex Kuranishi bordism* $KB_{2l}^{\text{ac}}(Y; R)$. We restrict to *even dimensions* $\text{vdim } X = 2l$, as almost complex structures only exist then.

Definition 5.4. Let Y be an orbifold. Consider triples $(X, (\mathbf{J}, \mathbf{K}), \mathbf{f})$, where X is a compact oriented Kuranishi space without boundary, $(\mathbf{J}, \mathbf{K})$ an almost complex structure on X , and $\mathbf{f} : X \rightarrow Y$ is strongly smooth. The orientation on X need not match that induced by $(\mathbf{J}, \mathbf{K})$. An *isomorphism*

$i : (X, (\mathbf{J}, \mathbf{K}), \mathbf{f}) \rightarrow (\tilde{X}, (\tilde{\mathbf{J}}, \tilde{\mathbf{K}}), \tilde{\mathbf{f}})$ is an orientation-preserving strong diffeomorphism $i : X \rightarrow \tilde{X}$ with $i_*(\mathbf{J}, \mathbf{K}) = (\tilde{\mathbf{J}}, \tilde{\mathbf{K}})$ and $\mathbf{f} = \tilde{\mathbf{f}} \circ i$. Write $[X, (\mathbf{J}, \mathbf{K}), \mathbf{f}]$ for the isomorphism class of $(X, (\mathbf{J}, \mathbf{K}), \mathbf{f})$.

Let R be a commutative ring. For each $l \in \mathbb{Z}$, define the *almost complex Kuranishi bordism group* $KB_{2l}^{\text{ac}}(Y; R)$ of Y with coefficients in R to be the R -module of finite R -linear combinations of isomorphism classes $[X, (\mathbf{J}, \mathbf{K}), \mathbf{f}]$ for which $\text{vdim } X = 2l$, with the relations:

- (i) $[X, (\mathbf{J}, \mathbf{K}), \mathbf{f}] + [X', (\mathbf{J}', \mathbf{K}'), \mathbf{f}'] = [X \amalg X', (\mathbf{J}, \mathbf{K}) \amalg (\mathbf{J}', \mathbf{K}'), \mathbf{f} \amalg \mathbf{f}']$ for all $[X, (\mathbf{J}, \mathbf{K}), \mathbf{f}], [X', (\mathbf{J}', \mathbf{K}'), \mathbf{f}']$; and
- (ii) Let W be a compact oriented Kuranishi space with boundary but without (g-)corners, with $\text{vdim } W = 2l + 1$, $(\mathbf{D}, \mathbf{J}, \mathbf{K})$ an almost CR structure on W , and $e : W \rightarrow Y$ a strong submersion. Then $(\mathbf{J}, \mathbf{K})|_{\partial W}$ is an almost complex structure on ∂W , as in §2.9. We set $[\partial W, (\mathbf{J}, \mathbf{K})|_{\partial W}, e|_{\partial W}] = 0$ in $KB_{2l}^{\text{ac}}(Y; R)$.

Following Definition 5.3, define *effective almost complex Kuranishi bordism* $KB_{2l}^{\text{eac}}(Y; R)$ as above, but requiring X, W to be *effective* Kuranishi spaces.

Let $h : Y \rightarrow Z$ be a smooth map of orbifolds. Define the *pushforward* $h_* : KB_{2l}^{\text{ac}}(Y; R) \rightarrow KB_{2l}^{\text{ac}}(Z; R)$ and $h_* : KB_{2l}^{\text{eac}}(Y; R) \rightarrow KB_{2l}^{\text{eac}}(Z; R)$ by $h_* : \sum_{a \in A} \rho_a [X_a, (\mathbf{J}_a, \mathbf{K}_a), \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a [X_a, (\mathbf{J}_a, \mathbf{K}_a), h \circ \mathbf{f}_a]$. Pushforwards are functorial, $(g \circ h)_* = g_* \circ h_*$.

Definition 5.4 and its cobordism analogue Definition 5.8 are motivated by applications to Gromov–Witten theory, to be discussed in §6.2–§6.3.

5.3 Morphisms of Kuranishi bordism groups

There are many natural projections between the bordism groups of §5.1–§5.2, and from them to (effective) Kuranishi homology or singular homology.

Definition 5.5. We adopt the following uniform notation for projections: Π_A^B represents a projection from the bordism/homology group of type A to the bordism/homology group of type B , where A, B can be ‘bo’ for $B_*(Y; R)$, ‘eo’ for $B_*^{\text{eo}}(Y; R)$, ‘Kb’ for $KB_*(Y; R)$, ‘eb’ for $KB_*^{\text{eb}}(Y; R)$, ‘tr’ for $KB_*^{\text{tr}}(Y; R)$, ‘ac’ for $KB_*^{\text{ac}}(Y; R)$, ‘eac’ for $KB_*^{\text{eac}}(Y; R)$, ‘Kh’ for $KH_*(Y; R \otimes_{\mathbb{Z}} \mathbb{Q})$, ‘ef’ for $KH_*^{\text{ef}}(Y; R)$, and ‘si’ for $H_*^{\text{si}}(Y; R)$ or $H_*^{\text{si}}(Y; R \otimes_{\mathbb{Z}} \mathbb{Q})$. With this notation:

- (a) Define $\Pi_{\text{bo}}^{\text{eo}}, \Pi_{\text{bo}}^{\text{tr}}, \Pi_{\text{bo}}^{\text{eb}}, \Pi_{\text{bo}}^{\text{Kb}}, \Pi_{\text{eo}}^{\text{eb}}, \Pi_{\text{eo}}^{\text{Kb}}, \Pi_{\text{tr}}^{\text{eb}}, \Pi_{\text{tr}}^{\text{Kb}}, \Pi_{\text{eb}}^{\text{Kb}}$ by $\Pi_A^B : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a]$ or $\Pi_A^B : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a]$.
- (b) Define $\Pi_{\text{eac}}^{\text{ac}} : \sum_{a \in A} \rho_a [X_a, (\mathbf{J}_a, \mathbf{K}_a), \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a [X_a, (\mathbf{J}_a, \mathbf{K}_a), \mathbf{f}_a]$.
- (c) Define $\Pi_{\text{eac}}^{\text{eb}}, \Pi_{\text{eac}}^{\text{Kb}}, \Pi_{\text{ac}}^{\text{Kb}} : \sum_{a \in A} \rho_a [X_a, (\mathbf{J}_a, \mathbf{K}_a), \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a]$.
- (d) Define $\Pi_{\text{tr}}^{\text{Kh}}, \Pi_{\text{eb}}^{\text{Kh}}, \Pi_{\text{Kb}}^{\text{Kh}}, \Pi_{\text{eac}}^{\text{Kh}}, \Pi_{\text{ac}}^{\text{Kh}}$ as maps into $KH_k(Y; R \otimes_{\mathbb{Z}} \mathbb{Q})$ by $\Pi_A^{\text{Kh}} : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a] \mapsto [\sum_{a \in A} \pi(\rho_a) [X_a, \mathbf{f}_a, \mathbf{G}_a]]$ or $\Pi_A^{\text{Kh}} : \sum_{a \in A} \rho_a [X_a, (\mathbf{J}_a, \mathbf{K}_a), \mathbf{f}_a] \mapsto [\sum_{a \in A} \pi(\rho_a) [X_a, \mathbf{f}_a, \mathbf{G}_a]]$, where $\mathbf{G}_a$ is some choice of gauge-fixing data for $(X_a, \mathbf{f}_a)$, which exists by Theorem 3.18, and $\pi : R \rightarrow R \otimes_{\mathbb{Z}} \mathbb{Q}$ is the natural morphism.

- (e) Define $\Pi_{\text{tr}}^{\text{ef}}, \Pi_{\text{eb}}^{\text{ef}}, \Pi_{\text{eac}}^{\text{ef}}$ by $\Pi_A^{\text{ef}} : \sum_{a \in A} \rho_a[X_a, \mathbf{f}_a] \mapsto [\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]]$ or $\Pi_{\text{eac}}^{\text{ef}} : \sum_{a \in A} \rho_a[X_a, (\mathbf{J}_a, \mathbf{K}_a), \mathbf{f}_a] \mapsto [\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]]$, where $\mathbf{G}_a$ is some choice of effective gauge-fixing data for $(X_a, \mathbf{f}_a)$, which exists by Theorem 3.39(a).
- (f) Define $\Pi_{\text{Kb}}^{\text{si}}, \Pi_{\text{ac}}^{\text{si}}$ as maps into $H_k^{\text{si}}(Y; R \otimes_{\mathbb{Z}} \mathbb{Q})$ by $\Pi_A^{\text{si}} = (\Pi_{\text{si}}^{\text{Kh}})^{-1} \circ \Pi_A^{\text{Kh}}$, where $(\Pi_{\text{si}}^{\text{Kh}})^{-1}$ over $R \otimes_{\mathbb{Z}} \mathbb{Q}$ exists by Corollary 4.10, and Π_A^{Kh} is as in (d). Define $\Pi_{\text{tr}}^{\text{si}}, \Pi_{\text{eb}}^{\text{si}}, \Pi_{\text{eac}}^{\text{si}}$ as maps into $H_k^{\text{si}}(Y; R)$ by $\Pi_A^{\text{si}} = (\Pi_{\text{si}}^{\text{ef}})^{-1} \circ \Pi_A^{\text{ef}}$, where $(\Pi_{\text{si}}^{\text{ef}})^{-1}$ exists by Theorem 4.8 and Π_A^{ef} is as in (e).

Here (a)–(c) follow from the obvious implications manifold $\Rightarrow$ trivial stabilizers Kuranishi space $\Rightarrow$ effective Kuranishi space $\Rightarrow$ Kuranishi space, and manifold $\Rightarrow$ effective orbifold $\Rightarrow$ effective Kuranishi space.

We will show $\Pi_{\text{Kb}}^{\text{Kh}}$ in (d) is well-defined, the other Π_A^B in (d),(e) are also well-defined by the same argument. We must prove three things:

- (i) that $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]$ is a cycle in $KC_k(Y; R)$;
- (ii) that $[\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]]$ is independent of the choice of $\mathbf{G}_a$; and
- (iii) that $\Pi_{\text{Kb}}^{\text{Kh}}$ takes relations in $KB_k(Y; R)$ to equations in $KH_k(Y; R)$.

For (i), $\partial X_a = \emptyset$ by definition of $KB_k(Y; R)$, so $\partial(\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]) = 0$ is immediate from (66). For (ii), suppose $\mathbf{G}_a, \mathbf{G}'_a$ are alternative choices. Then by Theorem 3.25, there exists gauge-fixing data $\mathbf{H}_a$ for $([0, 1] \times X_a, \mathbf{f}_a \circ \pi_{X_a})$ with $\mathbf{H}_a|_{X_a \times \{0\}} = \mathbf{G}_a$ and $\mathbf{H}_a|_{X_a \times \{1\}} = \mathbf{G}'_a$. Hence

$$\partial[[0, 1] \times X_a, \mathbf{f}_a \circ \pi_{X_a}, \mathbf{H}_a] = [X_a, \mathbf{f}_a, \mathbf{G}'_a] - [X_a, \mathbf{f}_a, \mathbf{G}_a],$$

by Definition 4.3(i),(iii), so $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]$ and $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}'_a]$ are homologous and define the same class in $KH_k(Y; R)$.

For (iii), let $[X, \mathbf{f}], [X', \mathbf{f}']$ be as in Definition 5.3(i), and choose gauge-fixing data $\mathbf{G}, \mathbf{G}'$ for $(X, \mathbf{f}), (X', \mathbf{f}')$ by Theorem 3.18. Then

$$[X, \mathbf{f}, \mathbf{G}] + [X', \mathbf{f}', \mathbf{G}'] = [X \amalg X', \mathbf{f} \amalg \mathbf{f}', \mathbf{G} \amalg \mathbf{G}'] \quad \text{in } KC_k(Y; R)$$

by Definition 4.3(iii). Let $Z, \mathbf{g}$ be as in Definition 5.3(ii), and choose gauge-fixing data $\mathbf{H}$ for $(Z, \mathbf{g})$. Then $\partial[Z, \mathbf{g}, \mathbf{H}] = [\partial Z, \mathbf{g}|_{\partial Z}, \mathbf{H}|_{\partial Z}]$, so $[[\partial Z, \mathbf{g}|_{\partial Z}, \mathbf{H}|_{\partial Z}]] = 0$ in $KH_k(Y; R)$. But $[[\partial Z, \mathbf{g}|_{\partial Z}, \mathbf{H}|_{\partial Z}]] = \Pi_{\text{Kb}}^{\text{Kh}}([\partial Z, \mathbf{g}|_{\partial Z}])$. Thus $\Pi_{\text{Kb}}^{\text{Kh}}$ maps relations Definition 5.3(i),(ii) to equations in $KH_k(Y; R)$, and $\Pi_{\text{Kb}}^{\text{Kh}}$ is well-defined.

Whenever Π_A^B, Π_B^C are defined, Π_A^C is also defined, with $\Pi_A^C = \Pi_B^C \circ \Pi_A^B$. Also, these projections all commute with pushforwards, $h_* \circ \Pi_A^B = \Pi_A^B \circ h_*$.

These morphisms are illustrated in the commutative diagram Figure 5.1. The isomorphisms in the figure are proved in Theorems 4.8, 5.6 and Corollary 4.10. We show that $\Pi_{\text{bo}}^{\text{tr}}, \Pi_{\text{eo}}^{\text{eb}}$ in Definition 5.5 are isomorphisms. The proof follows part of that of Theorem 4.8 in Appendix B.

Theorem 5.6. *The morphisms $\Pi_{\text{bo}}^{\text{tr}} : B_k(Y; R) \rightarrow KB_k^{\text{tr}}(Y; R)$ and $\Pi_{\text{eo}}^{\text{eb}} : B_k^{\text{eo}}(Y; R) \rightarrow KB_k^{\text{eb}}(Y; R)$ are isomorphisms for $k \geq 0$. Also $KB_k^{\text{tr}}(Y; R) = KB_k^{\text{eb}}(Y; R) = \{0\}$ for $k < 0$.*

$$\begin{array}{ccccccc}
B_k(Y; R) & \xrightarrow{\quad} & B_k^{\text{eo}}(Y; R) & & KB_k^{\text{eac}}(Y; R) & \xrightarrow{\quad} & KB_k^{\text{ac}}(Y; R) \\
& \searrow \Pi_{\text{bo}}^{\text{tr}} & \Pi_{\text{bo}}^{\text{eo}} & \searrow \Pi_{\text{eo}}^{\text{eb}} & \Pi_{\text{eac}}^{\text{eb}} & \searrow \Pi_{\text{eac}}^{\text{ac}} & \\
& & KB_k^{\text{tr}}(Y; R) & \xrightarrow{\quad} & KB_k^{\text{eb}}(Y; R) & \xrightarrow{\quad} & KB_k(Y; R) \\
& & \Pi_{\text{bo}}^{\text{tr}} & \searrow \Pi_{\text{tr}}^{\text{eb}} & \Pi_{\text{eb}}^{\text{eb}} & \searrow \Pi_{\text{eb}}^{\text{Kb}} & \Pi_{\text{ac}}^{\text{Kb}} \\
& & & & KB_k^{\text{ef}}(Y; R) & \xrightarrow{\quad} & KB_k(Y; R) \\
& & & \Pi_{\text{tr}}^{\text{ef}} & \Pi_{\text{eb}}^{\text{ef}} & \searrow \Pi_{\text{eb}}^{\text{Kh}} & \Pi_{\text{Kb}}^{\text{Kh}} \\
H_k^{\text{si}}(Y; R) & \xrightarrow{\quad} & KH_k^{\text{ef}}(Y; R) & \xrightarrow{\quad} & KH_k(Y; R \otimes_{\mathbb{Z}} \mathbb{Q}) & \xleftarrow{\quad} & H_k^{\text{si}}(Y; R \otimes_{\mathbb{Z}} \mathbb{Q}) \\
& \Pi_{\text{si}}^{\text{ef}} & \Pi_{\text{ef}}^{\text{ef}} & \Pi_{\text{ef}}^{\text{Kh}} & \Pi_{\text{si}}^{\text{Kh}} & \Pi_{\text{si}}^{\text{Kh}} & \\
& \cong & \cong & \cong & \cong & \cong &
\end{array}$$

NB: $KB_k^{\text{eac}}, KB_k^{\text{ac}}(Y; R)$ defined only for k even. Also $B_k(Y; \mathbb{Z}) \cong MSO_k(Y)$ for manifolds.

Figure 5.1: Morphisms between bordism and homology groups

Proof. First we show $\Pi_{\text{bo}}^{\text{tr}}$ is surjective. Let $[X, \mathbf{f}]$ be a generator of $KB_k^{\text{tr}}(Y; R)$. Choose an excellent coordinate system $(\mathbf{I}, \boldsymbol{\eta})$ for $(X, \mathbf{f})$, which is possible by Corollary 3.17, with $\mathbf{I} = (I, (V^i, \dots, \psi^i), f^i : i \in I, \dots)$, where the V^i can be taken to be *manifolds without boundary*, since X has trivial stabilizers and is without boundary.

Follow the first part of §B.2 to choose C^1 small, smooth transverse perturbations $\tilde{s}^i$ of s^i for $i \in I$, define a compact, oriented k -manifold $\tilde{X}$, where $\tilde{X} = \emptyset$ if $k < 0$, and a smooth map $\tilde{f} : \tilde{X} \rightarrow Y$, a compact, oriented Kuranishi space Z of virtual dimension $k + 1$, and a strongly smooth map $\mathbf{g} : Z \rightarrow Y$. As $\partial X = \emptyset$, we have $\partial \tilde{X} = \emptyset$, and the component Z^∂ of ∂Z lying over ∂X is also empty. So equation (234) gives $\partial Z \cong \tilde{X} \amalg -X$ in oriented Kuranishi spaces. Also $\mathbf{g}|_{\partial Z}$ agrees with $\tilde{f} \amalg \mathbf{f}$ under this identification. Thus Definition 5.3(i),(ii) give $[X, \mathbf{f}] = [\tilde{X}, \tilde{f}]$ in $KB_k^{\text{tr}}(Y; R)$ if $k \geq 0$, and $[X, \mathbf{f}] = 0$ in $KB_k^{\text{tr}}(Y; R)$ if $k < 0$ so that $\tilde{X} = \emptyset$. When $k < 0$ this implies $KB_k^{\text{tr}}(Y; R) = \{0\}$. When $k \geq 0$, $[\tilde{X}, \tilde{f}]$ is also a generator of $B_k(Y; R)$, so as an element of $KB_k^{\text{tr}}(Y; R)$ it lies in the image of $\Pi_{\text{bo}}^{\text{tr}}$. Therefore $\Pi_{\text{bo}}^{\text{tr}}$ is *surjective*.

To prove that $\Pi_{\text{bo}}^{\text{tr}}$ is injective, it is enough to show that if $[X, \mathbf{f}] \in B_k(Y; R)$ with $\Pi_{\text{bo}}^{\text{tr}}([X, \mathbf{f}]) = 0$ in $KB_k^{\text{tr}}(Y; R)$, then $[X, \mathbf{f}] = 0 \in B_k(Y; R)$. Since $[X, \mathbf{f}] = 0 \in KB_k^{\text{tr}}(Y; R)$, there exists a compact oriented Kuranishi space W of virtual dimension $k + 1$ and a strongly smooth $\mathbf{e} : W \rightarrow Y$ with $\partial W = X$ and $\mathbf{e}|_{\partial W} = \mathbf{f}$. Since $\partial W = X$ is a k -manifold, and this is an open condition, W is a $(k + 1)$ -manifold near ∂W . As above, choose an excellent coordinate system $(\mathbf{I}, \boldsymbol{\eta})$ for $(W, \mathbf{e})$, such that near ∂W the only Kuranishi neighbourhood in $\mathbf{I}$ is $(V^{k+1}, E^{k+1}, s^{k+1}, \psi^{k+1})$, where V^{k+1} is a $(k + 1)$ -manifold with $\partial V^{k+1} = X$, and $E^{k+1} \rightarrow V^{k+1}$ is the zero vector bundle, so that $s^{k+1} \equiv 0$. This is possible as W is a $(k + 1)$ -manifold near ∂W .

Again, follow the first part of §B.2 to choose C^1 small, smooth transverse perturbations $\tilde{s}^i$ of s^i for $i \in I$, and define a compact, oriented $(k + 1)$ -manifold $\tilde{W}$ and a smooth map $\tilde{e} : \tilde{W} \rightarrow Y$. Since $E^{k+1} \rightarrow V^{k+1}$ is the zero vector bundle we have $\tilde{s}^{k+1} \equiv 0 \equiv s^{k+1}$. Thus, near ∂W we do not perturb W at all, so $\tilde{W}, W$ coincide near ∂W , and $\partial \tilde{W} = \partial W = X$, and $\tilde{e}|_{\partial \tilde{W}} = \mathbf{e}|_{\partial W} = \mathbf{f}$. Hence applying Definition 5.1(ii) to $(\tilde{W}, \tilde{e})$ gives $[X, \mathbf{f}] = 0$ in $B_k(Y; R)$. Thus $\Pi_{\text{bo}}^{\text{tr}}$ is *injective*, so it is an isomorphism.

For $\Pi_{\text{eo}}^{\text{eb}}$ we follow the same proof, but we take the $(V^i, \dots, \psi^i)$ to be *ef*-

ffective Kuranishi neighbourhoods in the sense of Definition 3.35, and $\tilde{X}, \tilde{W}$ to be *effective orbifolds* rather than manifolds. Since the $(V^i, \dots, \psi^i)$ are effective Kuranishi neighbourhoods, the stabilizers of V^i act trivially on the fibres of E^i , and so the proof in §B.2 that we can choose the $\tilde{s}^i$ to be *transverse* perturbations is still valid. This would not work if we used non-effective Kuranishi neighbourhoods, and the corresponding result for $KB_k(Y; R)$ is false. $\square$

As $B_*(Y; \mathbb{Z}) \cong MSO_*(Y)$ for manifolds, this implies that $KB_*^{\text{tr}}(Y; \mathbb{Z}) \cong MSO_*(Y)$ for manifolds Y .

5.4 Kuranishi cobordism and Poincaré duality

(Effective) Kuranishi bordism $KB_*, KB_*^{\text{eb}}(Y; R)$ is basically (effective) Kuranishi homology $KH_*, KH_*^{\text{ef}}(Y; R)$ with corners and gauge-fixing data omitted. In §4.4 and §4.6 we defined dual theories of (effective) Kuranishi cohomology. So the obvious thing to do is to omit corners and co-gauge-fixing data from §4.4 and §4.6 to define dual theories of (effective) Kuranishi cobordism. For simplicity we restrict to orbifolds Y *without boundary*.

Definition 5.7. Let Y be an orbifold without boundary. Consider pairs $(X, \mathbf{f})$, where X is a compact Kuranishi space without boundary, $\mathbf{f} : X \rightarrow Y$ is a strong submersion, and $(X, \mathbf{f})$ is cooriented. An *isomorphism* between two pairs $(X, \mathbf{f}), (\tilde{X}, \tilde{\mathbf{f}})$ is a coorientation-preserving strong diffeomorphism $\mathbf{i} : X \rightarrow \tilde{X}$ with $\tilde{\mathbf{f}} = \mathbf{f} \circ \mathbf{i}$. Write $[X, \mathbf{f}]$ for the isomorphism class of $(X, \mathbf{f})$.

Let R be a commutative ring. For each $k \in \mathbb{Z}$, define the k^{th} *Kuranishi cobordism group* $KB^k(Y; R)$ of Y with coefficients in R to be the R -module of finite R -linear combinations of isomorphism classes $[X, \mathbf{f}]$ for which $\text{vdim } X = \dim Y - k$, with the relations:

- (i) $[X, \mathbf{f}] + [X', \mathbf{f}'] = [X \amalg X', \mathbf{f} \amalg \mathbf{f}']$ for all classes $[X, \mathbf{f}], [X', \mathbf{f}']$; and
- (ii) Suppose W is a compact Kuranishi space with boundary but without (g-) corners, with $\text{vdim } W = \dim Y - k + 1$, and $\mathbf{e} : W \rightarrow Y$ is a strong submersion, with $(W, \mathbf{e})$ cooriented. Then $\mathbf{e}|_{\partial W} : \partial W \rightarrow Y$ is a strong submersion which is cooriented as in Convention 2.33(a), and we impose the relation $[\partial W, \mathbf{e}|_{\partial W}] = 0$ in $KB^k(Y; R)$.

Elements of $KB^k(Y; R)$ will be called *Kuranishi cobordism classes*.

Let Y, Z be orbifolds without boundary, and $h : Y \rightarrow Z$ a smooth, proper map. Motivated by pullback in Kuranishi cohomology (79), define the *pullback* $h^* : KB^k(Z; R) \rightarrow KB^k(Y; R)$ by $h^* : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a [Y \times_{h, Z, \mathbf{f}_a} X_a, \pi_Y]$. Here the coorientation for $(X_a, \mathbf{f}_a)$ pulls back to a coorientation for $(Y \times_{h, Z, \mathbf{f}_a} X_a, \pi_Y)$ as in Definition 4.12. This takes relations (i),(ii) in $KB^k(Z; R)$ to (i),(ii) in $KB^k(Y; R)$, and so is well-defined. Pullbacks are functorial, $(g \circ h)^* = h^* \circ g^*$.

Following Definition 5.1 we define two minor variations on $KB^k(Y; R)$:

- Define *effective Kuranishi cobordism* $KB_{\text{ecb}}^k(Y; R)$ as above, except that we require $(X, \mathbf{f})$ and $(W, \mathbf{e})$ to be *coeffective*, as in Definition 3.37.

- Similarly, define *trivial stabilizers Kuranishi cobordism* $KB_{\text{trc}}^k(Y; R)$ as above, except that we require $\mathbf{f} : X \rightarrow Y$ and $e : W \rightarrow Y$ to be *co-effective* and *isomorphisms on stabilizer groups*. That is, for all $p \in X$, $\mathbf{f}_* : \text{Stab}_X(p) \rightarrow \text{Stab}_Y(\mathbf{f}(p))$ is an isomorphism, and $\text{Stab}_X(p)$ acts trivially on the fibre of the obstruction bundle of X at p .

Pullbacks h^* also work on $KB_{\text{ecb}}^*(Y; R)$ and $KB_{\text{trc}}^*(Y; R)$.

Definition 5.8. Let Y be an orbifold without boundary, and R a commutative ring. Consider triples $(X, (\mathbf{L}, \mathbf{K}), \mathbf{f})$, where X is a compact Kuranishi space without boundary, $\mathbf{f} : X \rightarrow Y$ a strong submersion with $(X, \mathbf{f})$ cooriented, and $(\mathbf{L}, \mathbf{K})$ a co-almost complex structure for $(X, \mathbf{f})$, as in §2.10. An *isomorphism* $\mathbf{i} : (X, (\mathbf{L}, \mathbf{K}), \mathbf{f}) \rightarrow (\tilde{X}, (\tilde{\mathbf{L}}, \tilde{\mathbf{K}}), \tilde{\mathbf{f}})$ is a strong diffeomorphism $\mathbf{i} : X \rightarrow \tilde{X}$ with $\mathbf{f} = \tilde{\mathbf{f}} \circ \mathbf{i}$ and $\mathbf{i}_*(\mathbf{L}, \mathbf{K}) = (\tilde{\mathbf{L}}, \tilde{\mathbf{K}})$, which identifies the coorientations of $(X, \mathbf{f})$ and $(\tilde{X}, \tilde{\mathbf{f}})$. Write $[X, (\mathbf{L}, \mathbf{K}), \mathbf{f}]$ for the isomorphism class of $(X, (\mathbf{L}, \mathbf{K}), \mathbf{f})$.

Define the *almost complex Kuranishi cobordism group* $KB_{\text{ca}}^{2l}(Y; R)$ of Y with coefficients in R for $l \in \mathbb{Z}$ to be the R -module of finite R -linear combinations of isomorphism classes $[X, (\mathbf{L}, \mathbf{K}), \mathbf{f}]$ for which $\text{vdim } X = \dim Y - 2l$, with relations (i),(ii) combining those of Definitions 5.4, 5.7, using co-almost CR structures in (ii).

Define *effective almost complex Kuranishi cobordism* $KB_{\text{eca}}^{2l}(Y; R)$ as above, but requiring $(X, \mathbf{f})$ and (W, e) to be *coeffective*.

Let $h : Y \rightarrow Z$ be a smooth, proper map of orbifolds without boundary. Define the *pullback* $h^* : KB_{\text{ca}}^{2l}(Z; R) \rightarrow KB_{\text{ca}}^{2l}(Y; R)$ and $h^* : KB_{\text{eca}}^{2l}(Z; R) \rightarrow KB_{\text{eca}}^{2l}(Y; R)$ by $h^* : \sum_{a \in A} \rho_a [X_a, (\mathbf{L}_a, \mathbf{K}_a), \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a [Y \times_{h, Z, \mathbf{f}_a} X_a, \pi_{X_a}^*(\mathbf{L}_a, \mathbf{K}_a), \pi_Y]$, where $\pi_{X_a}^*(\mathbf{L}_a, \mathbf{K}_a)$ is the pullback of the co-almost complex structure to $Y \times_Z X_a$ defined in the obvious way. Pullbacks are functorial, $(g \circ h)^* = h^* \circ g^*$.

$$\begin{array}{ccccccc}
& & & KB_{\text{eca}}^k(Y; R) & \xrightarrow{\Pi_{\text{eca}}^{\text{ca}}} & KB_{\text{ca}}^k(Y; R) & \\
& & \swarrow \Pi_{\text{eca}}^{\text{ecb}} & & \searrow \Pi_{\text{eca}}^{\text{Kcb}} & & \\
KB_{\text{trc}}^k(Y; R) & \xrightarrow{\Pi_{\text{trc}}^{\text{ecb}}} & KB_{\text{ecb}}^k(Y; R) & \xrightarrow{\Pi_{\text{ecb}}^{\text{Kcb}}} & KB^k(Y; R) & & \\
& \searrow \Pi_{\text{trc}}^{\text{ec}} & \swarrow \Pi_{\text{ecb}}^{\text{ec}} & & \swarrow \Pi_{\text{ecb}}^{\text{Kch}} & \searrow \Pi_{\text{Kcb}}^{\text{Kch}} & \\
H_{\text{cs}}^k(Y; R) & \xrightarrow[\cong]{\Pi_{\text{cs}}^{\text{ec}}} & KH_{\text{ec}}^k(Y; R) & \xrightarrow{\Pi_{\text{ec}}^{\text{Kch}}} & KH^k(Y; R \otimes_{\mathbb{Z}} \mathbb{Q}) & \xleftarrow[\cong]{\Pi_{\text{cs}}^{\text{Kch}}} & H_{\text{cs}}^k(Y; R \otimes_{\mathbb{Z}} \mathbb{Q}).
\end{array}$$

NB: $KB_{\text{eca}}^k, KB_{\text{ca}}^k(Y; R)$ defined only for k even; $\Pi_{\text{cs}}^{\text{ec}}$ defined only for Y a manifold.

Figure 5.2: Morphisms between cobordism and cohomology groups

As in §5.3, there are many natural projections between these cobordism groups, (effective) Kuranishi cohomology, and compactly-supported cohomology. The definitions are the obvious modifications of Definition 5.5, and we leave them to the reader. These morphisms are illustrated in Figure 5.2. As we have not defined cobordism analogues of $B_*(Y; R)$ and $B_*^{\text{eo}}(Y; R)$, the corresponding entries in Figure 5.1 are missing. Corollaries 4.17 and 4.28 say that $\Pi_{\text{cs}}^{\text{Kch}}, \Pi_{\text{cs}}^{\text{ec}}$ are isomorphisms, supposing Y a manifold for $\Pi_{\text{cs}}^{\text{ec}}$.

Remark 5.9. (a) The cobordism theories defined above are forms of *compactly-supported cobordism*. One can also define versions of Kuranishi cobordism analogous to ordinary cobordism, as discussed for the cohomology case in Remark 4.13. Since we do not have to worry about co-gauge-fixing data for noncompact Kuranishi spaces, this is straightforward. We just omit the assumption that the Kuranishi space X is compact in Definitions 5.7 and 5.8, and require instead that the strong submersion $\mathbf{f} : X \rightarrow Y$ should be *proper*. Pullbacks h^* are then defined for all smooth $h : Y \rightarrow Z$, not just when h is proper.

(b) The definitions above for orbifolds Y without boundary can be extended to general orbifolds Y using the ideas of §4.5 in the cohomology case. An important new point is that if $\partial Y \neq \emptyset$ then we should define $KB^k(Y; R)$ in terms of isomorphisms $[X, \mathbf{f}]$ for which X is a compact Kuranishi space with $\text{vdim } X = \dim Y - k$, $\mathbf{f} : X \rightarrow Y$ is a cooriented strong submersion, and $\partial_+^{\mathbf{f}} X = \emptyset$, *without assuming* $\partial X = \emptyset$. Then we can define $\iota^* : KB^k(Y; R) \rightarrow KB^k(\partial Y; R)$ mapping $\iota^* : [X, \mathbf{f}] \mapsto [\partial_-^{\mathbf{f}} X, \mathbf{f}_-]$.

Motivated by §4.4, we define *Poincaré duality maps* between our Kuranishi (co)bordism groups. Basically, we just omit (co-)gauge-fixing data from Definitions 4.14 and 4.25.

Definition 5.10. Let Y be an orbifold which is oriented, without boundary, and of dimension n , and R a commutative ring. Then for $\mathbf{f} : X \rightarrow Y$ a strong submersion, as in Definition 2.32 there is a 1-1 correspondence between coorientations for $(X, \mathbf{f})$ and orientations for X . Define R -module morphisms $\Pi_{\text{Kcb}}^{\text{Kb}} : KB^k(Y; R) \rightarrow KB_{n-k}(Y; R)$ for $k \in \mathbb{Z}$ by $\Pi_{\text{Kcb}}^{\text{Kb}} : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a]$, using the coorientation for $\mathbf{f}_a$ from $[X_a, \mathbf{f}_a] \in KB^k(Y; R)$ and the orientation on Y to determine the orientation on X_a for $[X_a, \mathbf{f}_a] \in KB_{n-k}(Y; R)$. Then $\Pi_{\text{Kcb}}^{\text{Kb}}$ takes Definition 5.3(i),(ii) to Definition 5.7(i),(ii), so it is well-defined.

Define $\Pi_{\text{Kb}}^{\text{Kcb}} : KB_{n-k}(Y; R) \rightarrow KB^k(Y; R)$ by $\Pi_{\text{Kb}}^{\text{Kcb}} : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a [X_a^Y, \mathbf{f}_a^Y]$, where in the notation of Definition 4.14, X_a^Y is the fibre product $(Y, \kappa_Y) \times_{\pi, Y, \mathbf{f}} X, \mathbf{f}^Y : X^Y \rightarrow Y$ is the strong submersion $\mathbf{exp} \circ \pi_{(Y, \kappa_Y)}$, and the orientations on X_a and Y determine a coorientation for $(X_a^Y, \mathbf{f}_a^Y)$. Then $\Pi_{\text{Kb}}^{\text{Kcb}}$ takes Definition 5.7(i),(ii) to Definition 5.3(i),(ii), so it is well-defined.

For effective Kuranishi (co)bordism, following Definition 4.25, when Y is an *effective* orbifold define $\Pi_{\text{ecb}}^{\text{eb}} : KB_{\text{ecb}}^k(Y; R) \rightarrow KB_{n-k}^{\text{eb}}(Y; R)$ as for $\Pi_{\text{Kcb}}^{\text{Kb}}$, and when Y is a *manifold*, define $\Pi_{\text{eb}}^{\text{eb}} : KB_{n-k}^{\text{eb}}(Y; R) \rightarrow KB_{\text{ecb}}^k(Y; R)$ as for $\Pi_{\text{Kb}}^{\text{Kcb}}$. Here for $\Pi_{\text{ecb}}^{\text{eb}}$ we need Y to be an effective orbifold in order that $(X_a, \mathbf{f}_a)$ coefficientive implies X_a effective, and for $\Pi_{\text{eb}}^{\text{eb}}$ we need Y to be a manifold so that X_a effective implies $(X_a^Y, \mathbf{f}_a^Y)$ coefficientive.

For trivial stabilizers Kuranishi (co)bordism, when Y is a *manifold* define $\Pi_{\text{trc}}^{\text{tr}} : KB_{\text{trc}}^k(Y; R) \rightarrow KB_{n-k}^{\text{tr}}(Y; R)$ as for $\Pi_{\text{Kcb}}^{\text{Kb}}, \Pi_{\text{ecb}}^{\text{eb}}$, and define $\Pi_{\text{tr}}^{\text{trc}} : KB_{n-k}^{\text{tr}}(Y; R) \rightarrow KB_{\text{trc}}^k(Y; R)$ as for $\Pi_{\text{Kb}}^{\text{Kcb}}, \Pi_{\text{eb}}^{\text{eb}}$.

The proof of the next theorem follows those of Theorems 4.15 and 4.26, omitting (co-)gauge-fixing data, and we leave it as an exercise.

Theorem 5.11. *Let Y be an oriented n -orbifold without boundary. Then $\Pi_{\text{Kcb}}^{\text{Kcb}}$ is the inverse of $\Pi_{\text{Kcb}}^{\text{Kb}}$ in Definition 5.10, so they are both isomorphisms. If also Y is a manifold, then $\Pi_{\text{eb}}^{\text{eb}}$ is the inverse of $\Pi_{\text{eb}}^{\text{eb}}$ and $\Pi_{\text{tr}}^{\text{trc}}$ is the inverse of $\Pi_{\text{tr}}^{\text{tr}}$ in Definition 5.10, so they are all isomorphisms.*

Let Y be a compact, oriented n -manifold without boundary. Then we have isomorphisms $MSO^k(Y) \cong MSO_{n-k}(Y)$ by (125), and $MSO_{n-k}(Y) \cong B_{n-k}(Y; \mathbb{Z})$ from §5.1, and $\Pi_{\text{bo}}^{\text{tr}} : B_{n-k}(Y; \mathbb{Z}) \rightarrow KB_{n-k}^{\text{tr}}(Y; \mathbb{Z})$ by Theorem 5.6, and $\Pi_{\text{tr}}^{\text{trc}} : KB_{n-k}^{\text{tr}}(Y; \mathbb{Z}) \rightarrow KB_{n-k}^{\text{trc}}(Y; \mathbb{Z})$ by Theorem 5.11. Combining all these gives an isomorphism $MSO^k(Y) \cong KB_{n-k}^{\text{trc}}(Y; \mathbb{Z})$, in a similar way to Corollary 4.17. We can extend this to the case Y not oriented using orientation bundles, as in Definition 4.16, giving:

Corollary 5.12. *If Y is a compact manifold without boundary, there is a natural isomorphism $MSO^*(Y) \cong KB_{\text{trc}}^*(Y; \mathbb{Z})$.*

This shows that $KB_{\text{trc}}^*(Y; \mathbb{Z})$ is a *differential-geometric realization* of cobordism, very different to the conventional algebraic topological definition [5]. We can also drop the assumption that Y is compact, if we replace ordinary cobordism $MSO^*(Y)$ by *compactly-supported cobordism* $MSO_{\text{cs}}^*(Y)$.

Remark 5.13. We can also consider Poincaré duality for *almost complex* Kuranishi (co)bordism $KH_{2*}^{\text{ac}}, KB_{\text{ca}}^{2*}(Y; R)$, and *effective almost complex* Kuranishi (co)bordism $KH_{2*}^{\text{eac}}, KB_{\text{eca}}^{2*}(Y; R)$. But to define the corresponding Poincaré duality maps, we need some extra data: we must choose an *almost complex structure* J on Y . (So in particular, $\dim Y$ must be even, $\dim Y = 2m$ say.)

To define $\Pi_{\text{ca}}^{\text{ac}}$, for $[X, (\mathbf{L}, \mathbf{K}), \mathbf{f}] \in KB_{\text{ca}}^{2l}(Y; R)$ we must define an almost complex structure on X using J and the co-almost complex structure $(\mathbf{L}, \mathbf{K})$ for $(X, \mathbf{f})$. Similarly, to define $\Pi_{\text{ac}}^{\text{ca}}$, for $[X, (\mathbf{J}, \mathbf{K}), \mathbf{f}] \in KB_{\text{ca}}^{2l}(Y; R)$ we must define a co-almost complex structure for $(X^Y, \mathbf{f}^Y)$ using J and the almost complex structure $(\mathbf{J}, \mathbf{K})$ for X . Both of these can be done, though there are some arbitrary choices involved, as for the construction of $(\mathbf{J}, \mathbf{K}) \times_Y (\mathbf{L}', \mathbf{K}')$ in §2.10. One can then extend Theorem 5.11 to show that $KB_{\text{ca}}^{2l}(Y; R) \cong KB_{2m-2l}^{\text{ac}}(Y; R)$, and $KB_{\text{eca}}^{2l}(Y; R) \cong KB_{2m-2l}^{\text{eac}}(Y; R)$ for Y a manifold.

5.5 Products on Kuranishi (co)bordism

Motivated by §4.7, we define cup, cap and intersection products on Kuranishi (co)bordism. Let Y be an orbifold without boundary, not necessarily oriented. Define the *cup product* $\cup : KB^k(Y; R) \times KB^l(Y; R) \rightarrow KB^{k+l}(Y; R)$ by

$$\left[\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a] \right] \cup \left[\sum_{b \in B} \sigma_b [\tilde{X}_b, \tilde{\mathbf{f}}_b] \right] = \sum_{a \in A, b \in B} \rho_a \sigma_b [X_a \times_{\mathbf{f}_a, Y, \tilde{\mathbf{f}}_b} \tilde{X}_b, \pi_Y], \quad (127)$$

for A, B finite and $\rho_a, \sigma_b \in R$. Here $X_a \times_{\mathbf{f}_a, Y, \tilde{\mathbf{f}}_b} \tilde{X}_b$ is the fibre product of §2.6, which is a compact Kuranishi space without boundary as $X_a, \tilde{X}_b$ are, and $\pi_Y : X_a \times_{\mathbf{f}_a, Y, \tilde{\mathbf{f}}_b} \tilde{X}_b \rightarrow Y$ is the projection from the fibre product, which

is a strong submersion as $\mathbf{f}_a, \tilde{\mathbf{f}}_b$ are, and the coorientations on $(X_a, \mathbf{f}_a)$ and $(\tilde{X}_b, \tilde{\mathbf{f}}_b)$ induce a coorientation on $(X_a \times_Y \tilde{X}_b, \pi_Y)$ as in Convention 2.33(b). Since $\text{vdim } X_a = \dim Y - k$, $\text{vdim } \tilde{X}_b = \dim Y - l$ we have $\text{vdim } X_a \times_Y \tilde{X}_b = \dim Y - (k + l)$, so $\cup$ does map $KB^k(Y; R) \times KB^l(Y; R) \rightarrow KB^{k+l}(Y; R)$.

To show $\cup$ is well-defined we must show that (127) takes relations (i),(ii) in $KB^k(Y; R), KB^l(Y; R)$ to (i),(ii) in $KB^{k+l}(Y; R)$. For (i) this holds as

$$\begin{aligned} ([X, \mathbf{f}] + [X', \mathbf{f}']) \cup [\tilde{X}, \tilde{\mathbf{f}}] &= [X \times_{\mathbf{f}, Y, \tilde{\mathbf{f}}} \tilde{X}, \pi_Y] + [X' \times_{\mathbf{f}', Y, \tilde{\mathbf{f}}} \tilde{X}, \pi_Y] \\ &= [(X \times_{\mathbf{f}, Y, \tilde{\mathbf{f}}} \tilde{X}) \amalg (X' \times_{\mathbf{f}', Y, \tilde{\mathbf{f}}} \tilde{X}), \pi_Y \amalg \pi_Y] \\ &= [(X \amalg X') \times_{\mathbf{f} \amalg \mathbf{f}', Y, \tilde{\mathbf{f}}} \tilde{X}, \pi_Y] = [X \amalg X', \mathbf{f} \amalg \mathbf{f}'] \cup [\tilde{X}, \tilde{\mathbf{f}}], \end{aligned}$$

and similarly for $[\tilde{X}, \tilde{\mathbf{f}}] \cup ([X, \mathbf{f}] + [X', \mathbf{f}'])$. For (ii) it holds as

$$[\partial Z, \mathbf{g}|_{\partial Z}] \cup [\tilde{X}, \tilde{\mathbf{f}}] = [\partial Z \times_{\mathbf{g}|_{\partial Z}, Y, \tilde{\mathbf{f}}} \tilde{X}, \pi_Y] = [\partial(Z \times_{\mathbf{g}, Y, \tilde{\mathbf{f}}} \tilde{X}), \pi_Y|_{\partial(\dots)}] = 0,$$

using Proposition 2.31(a) and $\partial \tilde{X} = \emptyset$, and similarly for $[\tilde{X}, \tilde{\mathbf{f}}] \cup [\partial Z, \mathbf{g}|_{\partial Z}]$.

If $[X, \mathbf{f}] \in KB^k(Y; R)$, $[\tilde{X}, \tilde{\mathbf{f}}] \in KB^l(Y; R)$ then equation (24) gives $[X, \mathbf{f}] \cup [\tilde{X}, \tilde{\mathbf{f}}] = (-1)^{kl} [\tilde{X}, \tilde{\mathbf{f}}] \cup [X, \mathbf{f}]$, that is, $\cup$ is *supercommutative*. Also (25) implies that $\cup$ is *associative*. If Y is compact then using the trivial coorientation for $\text{id}_Y : Y \rightarrow Y$, we have $[Y, \text{id}_Y] \in KB^0(Y; R)$, which is the *identity* for $\cup$. Thus, $KB^*(Y; R)$ is a *graded, supercommutative, associative R-algebra, with identity* if Y is compact, and *without identity* otherwise.

We can also mix Kuranishi bordism and cobordism, and define a *cap product* $\cap : KB_k(Y; R) \times KB^l(Y; R) \rightarrow KB_{k-l}(Y; R)$ by (127), where now $[X_a, \mathbf{f}_a] \in KB_k(Y; R)$ so that $\mathbf{f}_a$ is strongly smooth and X_a oriented, and $[\tilde{X}_b, \tilde{\mathbf{f}}_b] \in KB^l(Y; R)$ so that $\tilde{\mathbf{f}}_b$ is a strong submersion and $(\tilde{X}_b, \tilde{\mathbf{f}}_b)$ cooriented, and $X_a \times_{\mathbf{f}_a, Y, \tilde{\mathbf{f}}_b} \tilde{X}_b$ is well-defined as $\tilde{\mathbf{f}}_b$ is a strong submersion, and the orientation for X_a and coorientation for $(\tilde{X}_b, \tilde{\mathbf{f}}_b)$ combine to give an orientation for $X_a \times_Y \tilde{X}_b$ as in Convention 2.33(c). This has the associativity property $\alpha \cap (\beta \cup \gamma) = (\alpha \cap \beta) \cup \gamma$ for $\alpha \in KB_*(Y; R)$ and $\beta, \gamma \in KB^*(Y; R)$. Thus, *Kuranishi bordism* $KB_*(Y; R)$ is a *module over Kuranishi cobordism* $KB^*(Y; R)$.

As for Kuranishi (co)homology in §4.7, cup and cap products are compatible with pullbacks and pushforwards. That is, if Y, Z are orbifolds without boundary, and $h : Y \rightarrow Z$ a smooth, proper map, and $\alpha \in KB_*(Y; R)$ and $\beta, \gamma \in KB^*(Z; R)$ then as in (110) we have

$$h^*(\beta \cup \gamma) = h^*(\beta) \cup h^*(\gamma) \quad \text{and} \quad h_*(\alpha \cap h^*(\beta)) = h_*(\alpha) \cap \beta. \quad (128)$$

If Z is compact then Y is compact as h is proper, and $h^*([Z, \text{id}_Z]) = [Y, \text{id}_Y]$. Thus, h^* is an *R-algebra morphism*.

If Y is an oriented n -orbifold without boundary, then using $\cup$ and the isomorphism $KB^k(Y; R) \cong KB_{n-k}(Y; R)$ of Theorem 5.11 we obtain an *intersection product* $\bullet : KB_k(Y; R) \times KB_l(Y; R) \rightarrow KB_{k+l-n}(Y; R)$, which is also supercommutative (with degrees shifted by n) and associative. From (126) we see that $\Pi_{\text{bo}}^{\text{Kb}} : B_*(Y; R) \rightarrow KB_*(Y; R)$ intertwines intersection products $\bullet$ on $B_*(Y; R), KB_*(Y; R)$.

All of the above also holds without change for cup, cap and intersection products on *effective Kuranishi (co)bordism* $KB_{\text{ecb}}^*, KB_{\text{ecb}}^{\text{eb}}(Y; R)$, and on *trivial stabilizers Kuranishi (co)bordism* $KB_{\text{trc}}^*, KB_{\text{trc}}^{\text{tr}}(Y; R)$, except that we must suppose Y is a manifold to define $\bullet$ on $KB_{\text{ecb}}^{\text{eb}}, KB_{\text{trc}}^{\text{tr}}(Y; R)$.

We can also generalize these ideas to (effective) almost complex Kuranishi (co)bordism. Define R -bilinear maps $\cup : KB_{\text{ca}}^{2k}(Y; R) \times KB_{\text{ca}}^{2l}(Y; R) \rightarrow KB_{\text{ca}}^{2(k+l)}(Y; R)$ and $\cup : KB_{\text{eca}}^{2k}(Y; R) \times KB_{\text{eca}}^{2l}(Y; R) \rightarrow KB_{\text{eca}}^{2(k+l)}(Y; R)$ by

$$[X, (\mathbf{L}, \mathbf{K}), \mathbf{f}] \cup [\tilde{X}, (\tilde{\mathbf{L}}, \tilde{\mathbf{K}}), \tilde{\mathbf{f}}] = [X \times_{\mathbf{f}, Y, \tilde{\mathbf{f}}} \tilde{X}, (\mathbf{L}, \mathbf{K}) \times_Y (\tilde{\mathbf{L}}, \tilde{\mathbf{K}}), \pi_Y],$$

generalizing (127), where $(\mathbf{L}, \mathbf{K}) \times_Y (\tilde{\mathbf{L}}, \tilde{\mathbf{K}})$ is the fibre product of co-almost complex structures from §2.10. It is associative and supercommutative. If Y is compact then $[Y, (\mathbf{0}, \mathbf{0}), \text{id}_Y]$ is the identity for $\cup$, where $(\mathbf{0}, \mathbf{0})$ is the trivial co-almost complex structure for (Y, id_Y) , involving only almost complex structures on zero vector bundles. Thus $KB_{\text{ca}}^*(Y; R)$ is a *supercommutative ring*.

Similarly, define $\cap : KB_{2k}^{\text{ac}}(Y; R) \times KB_{2l}^{2l}(Y; R) \rightarrow KB_{2(k-l)}^{\text{ac}}(Y; R)$ and $\cap : KB_{2k}^{\text{eac}}(Y; R) \times KB_{2l}^{2l}(Y; R) \rightarrow KB_{2(k-l)}^{\text{eac}}(Y; R)$ by

$$[X, (\mathbf{J}, \mathbf{K}), \mathbf{f}] \cap [\tilde{X}, (\tilde{\mathbf{L}}, \tilde{\mathbf{K}}), \tilde{\mathbf{f}}] = [X \times_{\mathbf{f}, Y, \tilde{\mathbf{f}}} \tilde{X}, (\mathbf{J}, \mathbf{K}) \times_Y (\tilde{\mathbf{L}}, \tilde{\mathbf{K}}), \pi_Y].$$

They satisfy $\alpha \cap (\beta \cup \gamma) = (\alpha \cap \beta) \cap \gamma$ and $h_*(\alpha \cap h^*(\beta)) = h_*(\alpha) \cap \beta$ as above.

All of the morphisms between bordism and homology groups defined in §5.3, and their analogues for cobordism and cohomology groups, intertwine cup, cap and intersection products in the obvious way. So, for instance, $\Pi_{\text{Kcb}}^{\text{Kch}} : KB^*(Y; R) \rightarrow KH^*(Y; R \otimes_{\mathbb{Z}} \mathbb{Q})$ satisfies $\Pi_{\text{Kcb}}^{\text{Kch}}(\alpha \cup \beta) = \Pi_{\text{Kcb}}^{\text{Kch}}(\alpha) \cup \Pi_{\text{Kcb}}^{\text{Kch}}(\beta)$ for all $\alpha, \beta \in KB^*(Y; R)$. In each case this is either obvious, or follows from Theorem 4.34.

Remark 5.14. (a) As in §4.8 for Kuranishi (co)homology, each of the matching pairs of (co)bordism theories $KB_*, KB^*(Y; R)$, $KB_{\text{ecb}}^{\text{eb}}, KB_{\text{ecb}}^*(Y; R)$, $KB_{\text{trc}}^{\text{tr}}, KB_{\text{trc}}^*(Y; R)$, $KB_{2*}^{\text{ac}}, KB_{2*}^{2*}(Y; R)$ and $KB_{2*}^{\text{eac}}, KB_{2*}^{2*}(Y; R)$ can be combined into a single *bivariant theory*. In some sense this works ‘at the (co)chain level’.

(b) The author expects that one can generalize the proof in the classical case to show that each of our various different Kuranishi bordism theories is a *generalized homology theory* in the sense of Whitehead [75] in the category of smooth orbifolds, that is, it satisfies all the Eilenberg–Steenrod axioms for homology except the dimension axiom, and that the Kuranishi cobordism theories are the corresponding *generalized compactly-supported cohomology theory*. To do this, for the case of Kuranishi bordism, we should introduce a notion of *relative Kuranishi bordism* $KB_*(Y, Z; R)$ when Y is an orbifold and $Z \subseteq Y$ is an open set, and prove that it lives in an exact sequence

$$\cdots \rightarrow KB_k(Z; R) \rightarrow KB_k(Y; R) \rightarrow KB_k(Y, Z; R) \xrightarrow{\partial} KB_{k-1}(Z; R) \rightarrow \cdots,$$

and satisfies the excision axiom, and various other natural properties. We leave this as an exercise for the interested reader.

5.6 Orbifold strata and operators $\Pi^{\Gamma,\rho}$

Any orbifold V may be written as $V = \coprod_{\Gamma} V^{\Gamma}$, where the disjoint union is over isomorphism classes of finite groups Γ , and V^{Γ} is the subset of $v \in V$ with stabilizer group $\text{Stab}_V(v)$ isomorphic to Γ . This is the rough idea behind our definition of the *orbifold strata* of V . However, for our purposes we must modify this idea in three ways:

- (i) V^{Γ} above can be a union of components of different dimensions, so we pass to a finer stratification $V = \coprod_{\Gamma,\rho} V^{\Gamma,\rho}$, where ρ is an *isomorphism class of nontrivial representations of Γ* , and the normal bundle of $V^{\Gamma,\rho}$ in V has fibre the representation ρ , so that $\dim V^{\Gamma,\rho} = \dim V - \dim \rho$.
- (ii) $V^{\Gamma,\rho}$ above is not a *closed* subset of V . To make it closed, we replace $V^{\Gamma,\rho}$ by a subset of points $v \in V$ such that $\text{Stab}_V(v)$ has a subgroup isomorphic to Γ . Once we do this, the disjoint union $V = \coprod_{\Gamma,\rho} V^{\Gamma,\rho}$ no longer holds.
- (iii) Making the replacement (ii), but regarding $V^{\Gamma,\rho}$ as a subset of V , we find that $V^{\Gamma,\rho}$ is *not an orbifold*, since several pieces of $V^{\Gamma,\rho}$ may intersect in V locally to give singularities. To make $V^{\Gamma,\rho}$ into an orbifold, we must lift to an immersion $\iota^{\Gamma,\rho} : V^{\Gamma,\rho} \rightarrow V$, which may take finitely many points to one point. Then $V^{\Gamma,\rho}$ being closed is replaced by $\iota^{\Gamma,\rho}$ being *proper*. (This is very similar to the idea in §2.1 that if X is a manifold with (g-)corners, then to make ∂X a manifold, we cannot regard ∂X as a subset of X , but instead we have a finite immersion $\iota : \partial X \rightarrow X$.)

Here then is our definition of orbifold strata:

Definition 5.15. Let Γ be a finite group, and consider (finite-dimensional) real representations (W, ω) of Γ , that is, W is a finite-dimensional real vector space and $\omega : \Gamma \rightarrow \text{Aut}(W)$ is a group morphism. Call (W, ω) a *trivial representation* if $\omega \equiv \text{id}_W$, that is, if $\text{Fix}(\omega(\Gamma)) = W$. Call (W, ω) a *nontrivial representation* if it has no positive-dimensional trivial subrepresentation, that is, if $\text{Fix}(\omega(\Gamma)) = \{0\}$. Then every Γ -representation (W, ω) has a unique decomposition $W = W^t \oplus W^{\text{nt}}$ as the direct sum of a trivial representation (W^t, ω^t) and a nontrivial representation $(W^{\text{nt}}, \omega^{\text{nt}})$, where $W^t = \text{Fix}(\omega(\Gamma))$.

Let $(I_0, \iota_0), \dots, (I_N, \iota_N)$ be representatives of the isomorphism classes of *irreducible* Γ -representations, where $(I_0, \iota_0) = (\mathbb{R}, 1)$ is the trivial irreducible representation, and (I_i, ι_i) is nontrivial for $i = 1, \dots, N$. Then every Γ -representation (W, ω) is isomorphic to a direct sum $\bigoplus_{i=0}^N a_i (I_i, \iota_i)$ with $a_0, \dots, a_N \in \mathbb{N}$, where (W, ω) is trivial if $a_1 = \dots = a_N = 0$, and (W, ω) is nontrivial if $a_0 = 0$. Two $(W, \omega), (W, \omega')$ are isomorphic if and only if $(a_0, \dots, a_N) = (a'_0, \dots, a'_N)$.

Hence, *isomorphism classes of nontrivial Γ -representations* are in 1-1 correspondence with N -tuples $(a_1, \dots, a_N)$, with each $a_i \in \mathbb{Z}$. We will often use ρ to denote such an isomorphism class. The *dimension* $\dim \rho$ is $\sum_{i=1}^N a_i \dim I_i$. If $(W^{\text{nt}}, \omega^{\text{nt}})$ is a nontrivial Γ -representation, we will write $[(W^{\text{nt}}, \omega^{\text{nt}})]$ for the isomorphism class of nontrivial Γ -representations containing $(W^{\text{nt}}, \omega^{\text{nt}})$, so that $[(W^{\text{nt}}, \omega^{\text{nt}})] = \rho$ means $(W^{\text{nt}}, \omega^{\text{nt}})$ lies in the isomorphism class ρ .

Now let V be an orbifold, of dimension n . Each $v \in V$ has a *stabilizer group* $\text{Stab}_V(v)$. In this section we consider the tangent space $T_v V$ to be an n -dimensional vector space with a representation τ_v of $\text{Stab}_V(v)$. Let $\lambda : \Gamma \rightarrow \text{Stab}_V(v)$ be an injective group morphism, so that $\lambda(\Gamma)$ is a subgroup of $\text{Stab}_V(v)$ isomorphic to Γ . Hence $\tau_v \circ \lambda : \Gamma \rightarrow \text{Aut}(T_v V)$ is a Γ -representation, and we can split $T_v V = (T_v V)^t \oplus (T_v V)^{\text{nt}}$ into trivial and nontrivial Γ -representations, and form the isomorphism class $[(T_v V)^{\text{nt}}, (\tau_v \circ \lambda)^{\text{nt}}]$.

As a set, define the *orbifold stratum* $V^{\Gamma, \rho}$ to be

$$V^{\Gamma, \rho} = \{ \text{Stab}_V(v) \cdot (v, \lambda) : v \in V, \lambda : \Gamma \rightarrow \text{Stab}_V(v) \text{ is an injective group morphism, } [(T_v V)^{\text{nt}}, (\tau_v \circ \lambda)^{\text{nt}}] = \rho \}, \quad (129)$$

where $\text{Stab}_V(v)$ acts on pairs (v, λ) by $\sigma : (v, \lambda) \mapsto (v, \lambda^\sigma)$, where $\lambda^\sigma : \Gamma \rightarrow \text{Stab}_V(v)$ is given by $\lambda^\sigma(\gamma) = \sigma \lambda(\gamma) \sigma^{-1}$. Define a map $\iota^{\Gamma, \rho} : V^{\Gamma, \rho} \rightarrow V$ by $\iota^{\Gamma, \rho} : \text{Stab}_V(v) \cdot (v, \lambda) \mapsto v$.

Proposition 5.16. *In Definition 5.15, $V^{\Gamma, \rho}$ naturally has the structure of an orbifold of dimension $n - \dim \rho$, and $\iota^{\Gamma, \rho}$ lifts to a proper, finite immersion.*

Proof. Let (U, Δ, ψ) be an orbifold chart on V . We will construct a corresponding orbifold chart on $V^{\Gamma, \rho}$. Let $\lambda : \Gamma \rightarrow \Delta$ be an injective group morphism. As Δ acts on U by diffeomorphisms, composing with λ gives an action of Γ on U by diffeomorphisms. Write $\text{Fix}(\lambda(\Gamma))$ for the fixed point set of this action in U . It is closed, and a disjoint union of embedded submanifolds of U , possibly with different dimensions. If $u \in \text{Fix}(\lambda(\Gamma))$ then Γ has a representation on $T_u U$, so we have a splitting $T_u U = (T_u U)^t \oplus (T_u U)^{\text{nt}}$ into trivial and nontrivial Γ -representations, with $(T_u U)^t = T_u \text{Fix}(\lambda(\Gamma))$. For each isomorphism class of nontrivial Γ -representations ρ , write $\text{Fix}(\lambda(\Gamma))^\rho$ for the subset of $u \in \text{Fix}(\lambda(\Gamma))$ with $[(T_u U)^{\text{nt}}] = \rho$. Then $\text{Fix}(\lambda(\Gamma))^\rho$ is a closed, embedded submanifold of U , of dimension $n - \dim \rho$, since $\dim U = \dim V = n$.

Define a manifold $U^{\Gamma, \rho}$ of dimension $n - \dim \rho$ to be

$$U^{\Gamma, \rho} = \coprod_{\text{injective group morphisms } \lambda : \Gamma \rightarrow \Delta} \text{Fix}(\lambda(\Gamma))^\rho. \quad (130)$$

Note that this is a *disjoint union*, not a union: $\text{Fix}(\lambda(\Gamma))^\rho$ and $\text{Fix}(\lambda'(\Gamma))^\rho$ may intersect in U for distinct λ, λ' , but we do not identify such intersections in $U^{\Gamma, \rho}$. Define $i^{\Gamma, \rho} : U^{\Gamma, \rho} \rightarrow U$ to be the inclusion $\text{Fix}(\lambda(\Gamma))^\rho \hookrightarrow U$ on each component $\text{Fix}(\lambda(\Gamma))^\rho$ in (130). As each $\text{Fix}(\lambda(\Gamma))^\rho$ is a closed, embedded submanifold, and the disjoint union (130) is finite, we see that $i^{\Gamma, \rho}$ is a *proper, finite immersion*.

Define an action of Δ on $U^{\Gamma, \rho}$ by $\delta \in \Delta$ takes u in $\text{Fix}(\lambda(\Gamma))^\rho$ to $\delta \cdot u$ in $\text{Fix}(\lambda^\delta(\Gamma))^\rho$, where $\lambda^\delta : \Gamma \rightarrow \Delta$ is defined by $\lambda^\delta(\epsilon) = \delta \lambda(\epsilon) \delta^{-1}$, as for λ^σ in Definition 5.15. Then Δ acts on $U^{\Gamma, \rho}$ by diffeomorphisms, and $i^{\Gamma, \rho}$ is Δ -equivariant. Define $\psi^{\Gamma, \rho} : U^{\Gamma, \rho} / \Delta \rightarrow V^{\Gamma, \rho}$ as follows: let $\lambda : \Gamma \rightarrow \Delta$ be an injective group morphism and $u \in \text{Fix}(\lambda(\Gamma))^\rho \subseteq U^{\Gamma, \rho}$. Set $v = \psi(\Delta \cdot u)$ in V . Then we have a natural isomorphism $\text{Stab}_V(v) \cong \{ \delta \in \Delta : \delta \cdot u = u \}$, and λ maps Γ to the subgroup $\{ \delta \in \Delta : \delta \cdot u = u \}$ of Δ as $u \in \text{Fix}(\lambda(\Gamma))$. Thus we can regard λ as an injective group morphism $\lambda : \Gamma \rightarrow \text{Stab}_V(v)$. With this identification, define $\psi^{\Gamma, \rho}(\Delta u) = \text{Stab}_V(v) \cdot (v, \lambda)$.

It is elementary to show that $\psi^{\Gamma,\rho}$ is well-defined, and $(U^{\Gamma,\rho}, \Delta, \psi^{\Gamma,\rho})$ is an orbifold chart on $V^{\Gamma,\rho}$ with image $(\iota^{\Gamma,\rho})^{-1}(\text{Im } \psi)$. Also, if (U, Δ, ψ) and (U', Δ', ψ') are orbifold charts on V compatible on their overlaps, then $(U^{\Gamma,\rho}, \Delta, \psi^{\Gamma,\rho}), (U'^{\Gamma,\rho'}, \Delta', \psi'^{\Gamma,\rho'})$ are compatible on their overlaps. Hence this system of compatible orbifold charts on $V^{\Gamma,\rho}$ constructed from the system of compatible orbifold charts on V gives $V^{\Gamma,\rho}$ the structure of an *orbifold*, of dimension $n - \dim \rho$. Also $\iota^{\Gamma,\rho}$ is represented in orbifold charts by the $i^{\Gamma,\rho}$ above, which are proper, finite immersions, so $\iota^{\Gamma,\rho}$ lifts to a proper, finite immersion. $\square$

Remark 5.17. (a) If $v^{\Gamma,\rho} = \text{Stab}_V(v) \cdot (v, \lambda) \in V^{\Gamma,\rho}$ then the stabilizer group $\text{Stab}_{V^{\Gamma,\rho}}(v^{\Gamma,\rho})$ of $v^{\Gamma,\rho}$ in $V^{\Gamma,\rho}$ is

$$\{\sigma \in \text{Stab}_V(v) : \lambda^\sigma = \lambda\} = \{\sigma \in \text{Stab}_V(v) : \lambda(\gamma)\sigma = \sigma\lambda(\gamma) \quad \forall \gamma \in \Gamma\} = C(\lambda(\Gamma)),$$

the centralizer of $\lambda(\Gamma)$ in $\text{Stab}_V(v)$. The induced morphism $\iota_*^{\Gamma,\rho} : \text{Stab}_{V^{\Gamma,\rho}}(v^{\Gamma,\rho}) \rightarrow \text{Stab}_V(v)$ is just the inclusion $C(\lambda(\Gamma)) \hookrightarrow \text{Stab}_V(v)$. Thus $\iota^{\Gamma,\rho}$ is *injective on stabilizer groups*.

(b) Write $\text{Aut}(\Gamma)$ for the group of isomorphisms $\zeta : \Gamma \rightarrow \Gamma$. If $\zeta \in \text{Aut}(\Gamma)$ and $\lambda : \Gamma \rightarrow \text{Stab}_V(v)$ is an injective group morphism then $\lambda \circ \zeta^{-1} : \Gamma \rightarrow \text{Stab}_V(v)$ is an injective group morphism. Also $\text{Aut}(\Gamma)$ acts naturally on the set of isomorphism classes of nontrivial Γ -representations, $\rho \mapsto \zeta \cdot \rho$. If $\zeta \in \text{Aut}(\Gamma)$ then there is a natural diffeomorphism $\zeta_* : V^{\Gamma,\rho} \rightarrow V^{\Gamma,\zeta \cdot \rho}$ acting by $\text{Stab}_V(v) \cdot (v, \lambda) \mapsto \text{Stab}_V(v) \cdot (v, \lambda \circ \zeta^{-1})$, with $\iota^{\Gamma,\rho} \equiv \iota^{\Gamma,\zeta \cdot \rho} \circ \zeta_*$.

(c) If V is an *effective* orbifold then $\text{Stab}_V(v)$ acts effectively on $T_v V$ for all $v \in V$. It is then easy to see that $V^{\Gamma,\rho} = \emptyset$ unless ρ is the isomorphism class of an *effective* nontrivial Γ -representation. Conversely, if $V^{\Gamma,\rho} = \emptyset$ for all finite groups Γ and isomorphism classes of non-effective nontrivial Γ -representations ρ , then V is an effective orbifold.

(d) If V is an *oriented* orbifold, the orbifold strata $V^{\Gamma,\rho}$ *need not be oriented, or even orientable*. Here is an example. Take V to be the 4-orbifold $(\mathbb{RP}^3 \times \mathbb{R})/\mathbb{Z}_2$, where the generator σ of $\mathbb{Z}_2$ acts on $\mathbb{RP}^3 \times \mathbb{R}$ by

$$\sigma : ([x_0, x_1, x_2, x_3], y) \mapsto ([x_0, x_1, x_2, -x_3], -y). \quad (131)$$

Then $\text{Fix}(\sigma)$ is the disjoint union of $\{([x_0, x_1, x_2, 0], 0) : [x_0, x_1, x_2] \in \mathbb{RP}^2\}$, a copy of $\mathbb{RP}^2$, and the single point $([0, 0, 0, 1], 0)$. Hence V has two nontrivial orbifold strata, $V^{\mathbb{Z}_2, \rho_2}$, a copy of $\mathbb{RP}^2$, and $V^{\mathbb{Z}_2, \rho_4}$, a point, where ρ_2, ρ_4 are the isomorphism classes of nontrivial representations of $\mathbb{Z}_2$ of dimensions 2, 4. Now $\mathbb{RP}^k$ is orientable if and only if k is odd. Using this, it is easy to show that V is orientable, but $V^{\mathbb{Z}_2, \rho_2}$ is not.

Here is a sufficient condition on ρ for $V^{\Gamma,\rho}$ to be oriented whenever V is oriented. Let (W, ω) be a nontrivial Γ -representation with $[(W, \omega)] = \rho$, and suppose that W *does not admit any orientation-reversing automorphisms as a Γ -representation*. Then choosing an orientation for W determines an orientation for the fibres of the normal bundle of the immersed submanifold $\iota^{\Gamma,\rho}(V^{\Gamma,\rho})$ in V , since these fibres are isomorphic to (W, ω) as Γ -representations. Hence an

orientation for V determines one for $V^{\Gamma, \rho}$. This condition fails for any nonzero representation of $\Gamma = \mathbb{Z}_2$. But if $|\Gamma|$ is odd, the condition holds for all ρ , as in Definition 5.21 below.

(e) Suppose V is a $2m$ -orbifold and J is an *almost complex structure* on V , as in Definition 2.34. Then for each $v \in V$, $J|_v$ gives $T_v V$ the structure of a *complex vector space*, and as $J|_v$ is invariant under $\text{Stab}_V(v)$ we can regard $(T_v V, \tau_v)$ as a *complex representation* of Γ . We can therefore repeat Definition 5.15 and Proposition 5.16 using isomorphism classes ρ of *complex* nontrivial Γ -representations, rather than real representations, to define $V^{\Gamma, \rho}$ and $\iota^{\Gamma, \rho}$. Also J on V restricts to an almost complex structure $J^{\Gamma, \rho}$ on $V^{\Gamma, \rho}$.

Observe that V and $V^{\Gamma, \rho}$ are automatically orientable, with the orientations induced by $J, J^{\Gamma, \rho}$. Thus, by working with almost complex structures, we avoid the problem with non-oriented orbifold strata discussed in (d). This is an important reason for working with almost complex Kuranishi bordism.

Perhaps surprisingly, the same thing works for *almost CR structures*. Suppose V is a $(2m+1)$ -orbifold and (D, J) is an almost CR structure on V , as in Definition 2.35. Suppose that V is *orientable*. Then for each $v \in V$, we have a codimension one subspace $D|_v$ of $T_v V$, an almost complex structure $J|_v$ on $D|_v$, and we can choose an orientation on $T_v V$. All three structures are invariant under the action of $\text{Stab}_V(v)$. As $J|_v$ induces an orientation on $D|_v$ we have an orientation on $T_v V/D|_v \cong \mathbb{R}$ which is fixed by $\text{Stab}_V(v)$. Therefore $\text{Stab}_V(v)$ acts trivially on the quotient $(T_v V)/(D|_v)$.

It follows that if $\lambda : \Gamma \rightarrow \text{Stab}_V(v)$ is an injective group morphism, then in the induced splitting $T_v V = (T_v V)^t \oplus (T_v V)^{\text{nt}}$, $(T_v V)^{\text{nt}}$ is a vector subspace of $D|_v$, which must be invariant under $J|_v$. So we can regard $(T_v V)^{\text{nt}}$ as a *complex* representation of Γ . This is what we need to repeat Definition 5.15 and Proposition 5.16 using isomorphism classes ρ of complex nontrivial Γ -representations. Also (D, J) on V restricts to an almost CR structure $(D^{\Gamma, \rho}, J^{\Gamma, \rho})$ on $V^{\Gamma, \rho}$.

When working with complex representations, we will use $\dim \rho$ to mean the *complex* dimension of the representations, not the real dimension.

(f) Suppose V is an orbifold with boundary, but *without (g-)corners*. Then $\iota : \partial V \rightarrow V$ is an embedding, and induces isomorphisms $\iota_* : \text{Stab}_{\partial V}(v, B) \rightarrow \text{Stab}_V(v)$ on stabilizer groups. It is easy to show that restricting to orbifold strata commutes with taking boundaries, that is, $\partial(V^{\Gamma, \rho}) = (\partial V)^{\Gamma, \rho}$.

However, as we explained in Remark 2.10(b), if V has (g-)corners then $\iota : \partial V \rightarrow V$ is an immersion, and the induced maps $\iota_* : \text{Stab}_{\partial V}(v, B) \rightarrow \text{Stab}_V(v)$ are injective, but may not be surjective. Because of this, *for orbifolds with corners or g-corners, restricting to orbifold strata does not commute with taking boundaries*, that is, $\partial(V^{\Gamma, \rho})$ may not agree with $(\partial V)^{\Gamma, \rho}$.

This leads to an important difference between Kuranishi (co)bordism and (effective) Kuranishi (co)homology. In Kuranishi (co)bordism, information from orbifold strata $X^{\Gamma, \rho}$ of generators $[X, \mathbf{f}]$ survives in $KB_*, KB^*(Y; R)$, as we show below, since we do not allow Kuranishi spaces with (g-)corners in defining $KB_*, KB^*(Y; R)$. But in (effective) Kuranishi (co)homology, where we do allow Kuranishi spaces with (g-)corners, all information from orbifold strata is lost.

(In fact relation Definition 4.3(iv) would also kill information from orbifold strata, but this does not apply to *effective* Kuranishi (co)homology.)

We generalize Definition 5.15 to Kuranishi spaces X . Since we must consider the representation of Γ on the obstruction bundle of X , we must replace ρ by an isomorphism class of *virtual* nontrivial Γ -representations.

Definition 5.18. Let Γ be a finite group. A *virtual representation* of Γ is a formal difference $(W, \omega) \ominus (W', \omega')$ of two Γ -representations $(W, \omega), (W', \omega')$. The *dimension* of $(W, \omega) \ominus (W', \omega')$ is $\dim W - \dim W'$. We call $(W, \omega) \ominus (W', \omega')$ *nontrivial* if $(W, \omega), (W', \omega')$ are nontrivial Γ -representations. We say that two virtual Γ -representations $(W_1, \omega_1) \ominus (W'_1, \omega'_1)$ and $(W_2, \omega_2) \ominus (W'_2, \omega'_2)$ are *isomorphic* if there exist Γ -representations $(W_3, \omega_3), (W_4, \omega_4)$ with $(W_1, \omega_1) \oplus (W_3, \omega_3) \cong (W_2, \omega_2) \oplus (W_4, \omega_4)$ and $(W'_1, \omega'_1) \oplus (W_3, \omega_3) \cong (W'_2, \omega'_2) \oplus (W_4, \omega_4)$.

Let $(I_0, \iota_0), \dots, (I_N, \iota_N)$ be representatives of the isomorphism classes of *irreducible* Γ -representations, where $(I_0, \iota_0) = (\mathbb{R}, 1)$ is the trivial irreducible representation, and (I_i, ι_i) is nontrivial for $i = 1, \dots, N$. Then every virtual Γ -representation $(W, \omega) \ominus (W', \omega')$ is isomorphic to $\bigoplus_{i=0}^N a_i (I_i, \iota_i)$ with $a_0, \dots, a_N \in \mathbb{Z}$, where $\bigoplus_{i=0}^N a_i (I_i, \iota_i)$ is a shorthand for $(\bigoplus_{i: a_i > 0} a_i (I_i, \iota_i)) \ominus (\bigoplus_{i: a_i < 0} (-a_i) (I_i, \iota_i))$, and two virtual representations are isomorphic if and only if they yield the same $a_0, \dots, a_N$.

Hence, *isomorphism classes of virtual Γ -representations* are in 1-1 correspondence with $(N+1)$ -tuples $(a_0, a_1, \dots, a_N)$ with $a_i \in \mathbb{Z}$, and isomorphism classes of virtual nontrivial Γ -representations are in 1-1 correspondence with N -tuples $(a_1, \dots, a_N)$ with $a_i \in \mathbb{Z}$. We often write ρ for such an isomorphism class. The *dimension* of ρ is $\dim \rho = \sum_{i=0}^N a_i \dim I_i$. Equivalently, an isomorphism class of virtual Γ -representatives is an element of $K_0(\text{mod-}\Gamma)$, the Grothendieck group of the abelian category $\text{mod-}\Gamma$ of finite-dimensional real Γ -representations.

Now let X be a Kuranishi space, $p \in X$ and $(V_p, \dots, \psi_p)$ be a Kuranishi neighbourhood in the germ at p . Set $v = \psi_p^{-1}(p)$ in V . The stabilizer group $\text{Stab}_X(p) \cong \text{Stab}_{V_p}(v)$ has natural representations on the finite-dimensional vector spaces $T_v V_p$ and $E_p|_v$. Let $\lambda : \Gamma \rightarrow \text{Stab}_{V_p}(v)$ be an injective group morphism. Then Γ acts on $T_v V_p$ and $E_p|_v$ via λ , so we may split into trivial and nontrivial Γ -representations $T_v V_p = (T_v V_p)^t \oplus (T_v V_p)^{\text{nt}}$ and $E_p|_v = (E_p|_v)^t \oplus (E_p|_v)^{\text{nt}}$. Hence $(T_v V_p)^{\text{nt}} \ominus (E_p|_v)^{\text{nt}}$ is a virtual nontrivial Γ -representation, with isomorphism class $[(T_v V_p)^{\text{nt}} \ominus (E_p|_v)^{\text{nt}}]$.

As a set, define the *orbifold stratum* $X^{\Gamma, \rho}$ to be

$$X^{\Gamma, \rho} = \{ \text{Stab}_X(p) \cdot (p, \lambda) : p \in X, \lambda : \Gamma \rightarrow \text{Stab}_X(p) \text{ is an injective group morphism, } [(T_v V_p)^{\text{nt}} \ominus (E_p|_v)^{\text{nt}}] = \rho \}, \quad (132)$$

where to interpret the condition $[(T_v V_p)^{\text{nt}} \ominus (E_p|_v)^{\text{nt}}] = \rho$ we choose $(V_p, \dots, \psi_p)$ in the germ at p , set $v = \psi_p^{-1}(p)$ and identify $\text{Stab}_{V_p}(v)$ with $\text{Stab}_X(p)$, so that λ makes $T_v V_p$ and $E_p|_v$ into Γ -representations. The condition is independent of the choice of $(V_p, \dots, \psi_p)$, by definition of equivalence in the germ. Define a map $\iota^{\Gamma, \rho} : X^{\Gamma, \rho} \rightarrow X$ by $\iota^{\Gamma, \rho} : \text{Stab}_X(p) \cdot (p, \lambda) \mapsto p$.

We have the following analogue of Proposition 5.16. The proof generalizes that of Proposition 5.16, and we leave it as an exercise. The most important new point to note is that coordinate transformations $(\phi_{pq}, \hat{\phi}_{pq})$ preserve the condition on virtual nontrivial Γ -representations. Suppose $v_q \in V_{pq} \subseteq V_q$ with $s_q(v_q) = 0$, $v_p = \phi_{pq}(v_q)$, and $\lambda_q : \Gamma \rightarrow \text{Stab}_{V_q}(v)$ is an injective group morphism. Then $(\phi_{pq})_* : \text{Stab}_{V_q}(v_q) \rightarrow \text{Stab}_{V_p}(v_p)$ is an isomorphism, as ϕ_{pq} is an embedding, and Definition 2.18(e) implies that $(T_{v_p} V_p)^{\text{nt}} \ominus (E_p|_{v_p})^{\text{nt}}$ and $(T_{v_q} V_q)^{\text{nt}} \ominus (E_q|_{v_q})^{\text{nt}}$ are isomorphic as virtual Γ -representations. Hence $[(T_{v_p} V_p)^{\text{nt}} \ominus (E_p|_{v_p})^{\text{nt}}] = \rho$ if and only if $[(T_{v_q} V_q)^{\text{nt}} \ominus (E_q|_{v_q})^{\text{nt}}] = \rho$.

Proposition 5.19. *In Definition 5.18, $X^{\Gamma, \rho}$ naturally has the structure of a Kuranishi space of virtual dimension $\text{vdim } X - \dim \rho$, and $\iota^{\Gamma, \rho}$ lifts to a proper, finite, strongly smooth map $\iota^{\Gamma, \rho} : X^{\Gamma, \rho} \rightarrow X$.*

Remark 5.20. The analogues of Remark 5.17(a)–(f) hold for Kuranishi spaces. In particular, for (c), if X is an *effective* Kuranishi space then in Definition 5.18 $T_v V_p$ is an *effective* Γ -representation, so $(T_v V_p)^{\text{nt}}$ is effective, and $E_p|_v$ is a *trivial* Γ -representation, so $(E_p|_v)^{\text{nt}} = 0$. Hence $X^{\Gamma, \rho} = \emptyset$ unless ρ is the isomorphism class of an *effective nontrivial* Γ -representation, not just a virtual one. For $\Gamma \neq \{1\}$ this gives $\dim \rho > 0$, so that $\text{vdim } X^{\Gamma, \rho} < \text{vdim } X$. If X is an *orientable*, effective Kuranishi space, we can also exclude the case $\dim \rho = 1$, since Γ cannot act on $T_v V_p$ preserving orientations such that $\dim(T_v V_p)^{\text{nt}} = 1$, because no finite group has an orientation-preserving, nontrivial representation on $\mathbb{R}$. Hence, if X is an orientable, effective Kuranishi space then $X^{\Gamma, \rho} = \emptyset$ for $\Gamma \neq \{1\}$ unless $\dim \rho \geq 2$, so that $\text{vdim } X^{\Gamma, \rho} \leq \text{vdim } X - 2$.

Also, for (e), if X is a Kuranishi space without boundary, $(\mathbf{J}, \mathbf{K})$ is an almost complex structure on X , Γ a finite group, and ρ an isomorphism class of virtual nontrivial *complex* Γ -representations, then we can define the orbifold stratum $X^{\Gamma, \rho}$, strongly smooth $\iota^{\Gamma, \rho} : X^{\Gamma, \rho} \rightarrow X$, and an almost complex structure $(\mathbf{J}^{\Gamma, \rho}, \mathbf{K}^{\Gamma, \rho})$ on $X^{\Gamma, \rho}$. Similarly, if X is an *orientable* Kuranishi space with boundary but without (g-)corners and $(\mathbf{D}, \mathbf{J}, \mathbf{K})$ an almost CR structure on X , we define $X^{\Gamma, \rho}$, $\iota^{\Gamma, \rho}$, and an almost CR structure $(\mathbf{D}^{\Gamma, \rho}, \mathbf{J}^{\Gamma, \rho}, \mathbf{K}^{\Gamma, \rho})$ on $X^{\Gamma, \rho}$.

When $|\Gamma|$ is odd, given an orientation on X , we define orientations on $X^{\Gamma, \rho}$.

Definition 5.21. Let Γ be a finite group. Consider the following condition on Γ : *no nontrivial representation (W, ω) of Γ should admit an orientation-reversing automorphism*, as in Remark 5.17(d). It is sufficient to apply this condition to the nontrivial irreducible representations $(I_1, \iota_1), \dots, (I_N, \iota_N)$ of Γ . Using Schur's lemma we can show that (I_i, ι_i) admits an orientation-reversing automorphism if and only if $\dim I_i$ is odd, and then -1 is such an automorphism. One can prove using the character theory of finite groups that $\dim I_i$ is even for $i = 1, \dots, N$ if and only if $|\Gamma|$ is odd, so the condition is equivalent to $|\Gamma|$ odd.

Suppose $|\Gamma|$ is odd. Choose an orientation on I_i for $i = 1, \dots, N$. Let (W, ω) be a nontrivial Γ -representation. Then (W, ω) is isomorphic to $\bigoplus_{i=1}^N a_i(I_i, \iota_i)$ for $a_1, \dots, a_N \in \mathbb{N}$. Thus the orientations on I_i induce an orientation on W . This orientation on W is independent of the choice of isomorphism $W \cong \bigoplus_{i=1}^N a_i I_i$,

since by the condition above (W, ω) admits no orientation-reversing automorphisms. Thus, we have chosen orientations on all nontrivial Γ -representations, which are consistent with direct sums.

Now let X be an oriented Kuranishi space, and ρ an isomorphism class of virtual nontrivial Γ -representations. We shall define an *orientation* on the orbifold stratum $X^{\Gamma, \rho}$. As for orbifold charts in the proof of Proposition 5.16, if (V, E, s, ψ) is a compatible Kuranishi neighbourhood on X then we construct a compatible Kuranishi neighbourhood $(V^{\Gamma, \rho}, E^{\Gamma, \rho}, s^{\Gamma, \rho}, \psi^{\Gamma, \rho})$ on $X^{\Gamma, \rho}$, and the strongly smooth map $\iota^{\Gamma, \rho}$ is represented by a smooth map $(\iota^{\Gamma, \rho}, \hat{\iota}^{\Gamma, \rho}) : (V^{\Gamma, \rho}, \dots, \psi^{\Gamma, \rho}) \rightarrow (V, \dots, \psi)$. Here points of $V^{\Gamma, \rho}$ are of the form $v^{\Gamma, \rho} = \text{Stab}_V(v) \cdot (v, \lambda)$, where $v = \iota^{\Gamma, \rho}(\text{Stab}_V(v) \cdot (v, \lambda))$ in V , $\lambda : \Gamma \rightarrow \text{Stab}_V(v)$ is an injective group morphism.

Now λ makes $T_v V$ and $E|_v$ into Γ -representations, with $T_{v^{\Gamma, \rho}} V^{\Gamma, \rho} = (T_v V)^{\text{t}}$ and $E^{\Gamma, \rho}|_{v^{\Gamma, \rho}} = (E|_v)^{\text{t}}$. Also $T_v V = (T_v V)^{\text{t}} \oplus (T_v V)^{\text{nt}}$ and $E|_v = (E|_v)^{\text{t}} \oplus (E|_v)^{\text{nt}}$, and $(T_v V)^{\text{nt}}, (E|_v)^{\text{nt}}$ both have orientations as they are nontrivial Γ -representations, and the orientation on X gives an orientation on $T_v V \oplus E|_v$. Thus, there is a unique orientation on $T_{v^{\Gamma, \rho}} V^{\Gamma, \rho} \oplus E^{\Gamma, \rho}|_{v^{\Gamma, \rho}}$ such that

$$(T_v V \oplus E|_v) \cong (T_{v^{\Gamma, \rho}} V^{\Gamma, \rho} \oplus E^{\Gamma, \rho}|_{v^{\Gamma, \rho}}) \oplus (T_v V)^{\text{nt}} \oplus (E|_v)^{\text{nt}}$$

holds in oriented vector spaces.

Using the fact that nontrivial Γ -representations have even dimension, so that changing their order in a direct sum preserves orientations, we find that the compatibility condition in §2.7 for orientations under coordinate changes $(\phi_{pq}, \hat{\phi}_{pq})$ on X implies the corresponding compatibility on $X^{\Gamma, \rho}$. Hence the orientations on $T_{v^{\Gamma, \rho}} V^{\Gamma, \rho} \oplus E^{\Gamma, \rho}|_{v^{\Gamma, \rho}}$ above induce an orientation on $X^{\Gamma, \rho}$, as we want. This orientation does depend on the choice of orientations for $I_1, \dots, I_N$; changing the orientations of I_i by $\epsilon_i = \pm 1$ changes the orientation of $X^{\Gamma, \rho}$ by $\prod_{i=1}^N \epsilon_i^{a_i}$, where $\rho = [\bigoplus_{i=1}^N a_i(I_i, \iota_i)]$ for $a_1, \dots, a_N \in \mathbb{Z}$.

We define operators $\Pi^{\Gamma, \rho}, \Pi_{\text{ac}}^{\Gamma, \rho}$ on (almost complex) Kuranishi bordism.

Definition 5.22. Let Γ be a finite group with $|\Gamma|$ odd. Let $(I_1, \iota_1), \dots, (I_N, \iota_N)$ be the nontrivial irreducible representations of Γ , and choose orientations on I_i for $i = 1, \dots, N$. Let Y be an orbifold and R a commutative ring. For each isomorphism class ρ of virtual nontrivial Γ -representations and each $k \in \mathbb{Z}$, define an operator $\Pi^{\Gamma, \rho} : KB_k(Y; R) \rightarrow KB_{k-\dim \rho}(Y; R)$ by

$$\Pi^{\Gamma, \rho} : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a] \longmapsto \sum_{a \in A} \rho_a [X_a^{\Gamma, \rho}, \mathbf{f}_a \circ \iota_a^{\Gamma, \rho}]. \quad (133)$$

Here $X_a^{\Gamma, \rho}$ and $\iota_a^{\Gamma, \rho} : X_a^{\Gamma, \rho} \rightarrow X_a$ are as in Definition 5.18 for X_a . Since X_a is compact and $\iota_a^{\Gamma, \rho}$ is proper, $X_a^{\Gamma, \rho}$ is a *compact* Kuranishi space. We define the orientation on $X_a^{\Gamma, \rho}$ as in Definition 5.21.

Equation (133) takes relations Definition 5.3(i),(ii) in $KB_k(Y; R)$ to Definition 5.3(i),(ii) in $KB_{k-\dim \rho}(Y; R)$, and so is well-defined. For (ii), this depends on the Kuranishi space analogue of Remark 5.17(f), that is, as W is a Kuranishi space with boundary but *without* $(g-)$ corners, we have $\partial(W^{\Gamma, \rho}) = (\partial W)^{\Gamma, \rho}$. Note that since $|\Gamma|$ is odd, $\dim \rho$ is even, as in Definition 5.21, so $\dim \rho \in 2\mathbb{Z}$.

If Y, Z are orbifolds and $h : Y \rightarrow Z$ is smooth then the $\Pi^{\Gamma, \rho}$ clearly commute with pushforwards, that is, $h_* \circ \Pi^{\Gamma, \rho} = \Pi^{\Gamma, \rho} \circ h_* : KB_k(Y; R) \rightarrow KB_{k-\dim \rho}(Z; R)$.

Definition 5.23. Let Y be an orbifold, R a commutative ring, l an integer, Γ a finite group, and ρ an isomorphism class of virtual nontrivial *complex* Γ -representations. Define $\Pi_{ac}^{\Gamma, \rho} : KB_{2l}^{ac}(Y; R) \rightarrow KB_{2l-2\dim \rho}^{ac}(Y; R)$ by

$$\Pi^{\Gamma, \rho} : \sum_{a \in A} \rho_a [X_a, (\mathbf{J}_a, \mathbf{K}_a), \mathbf{f}_a] \mapsto \sum_{a \in A} \rho_a [X_a^{\Gamma, \rho}, (\mathbf{J}_a^{\Gamma, \rho}, \mathbf{K}_a^{\Gamma, \rho}), \mathbf{f}_a \circ \iota_a^{\Gamma, \rho}]. \quad (134)$$

Here $X_a^{\Gamma, \rho}, \iota_a^{\Gamma, \rho} : X_a^{\Gamma, \rho} \rightarrow X_a$ and $(\mathbf{J}_a^{\Gamma, \rho}, \mathbf{K}_a^{\Gamma, \rho})$ are as in Remark 5.20 for X_a . Since X_a is compact and $\iota_a^{\Gamma, \rho}$ is proper, $X_a^{\Gamma, \rho}$ is a *compact* Kuranishi space. We take $\dim \rho$ to be the *complex* dimension of the virtual representation, so $2\dim \rho$ is the real dimension.

To define the orientation on $X_a^{\Gamma, \rho}$, note that X_a is oriented by Definition 5.4 (this orientation need not coincide with the orientation induces by $(\mathbf{J}, \mathbf{K})$). We construct the orientation on $X_a^{\Gamma, \rho}$ by the method of Definition 5.21, but instead of choosing orientations on $I_1, \dots, I_N$ in order to define orientations on $(T_v V)^{nt}, (E|_v)^{nt}$, we note that $(T_v V)^{nt}, (E|_v)^{nt}$ are *complex* vector spaces, and so have natural orientations.

Equation (134) takes relations Definition 5.4(i),(ii) in $KB_{2l}^{ac}(Y; R)$ to Definition 5.4(i),(ii) in $KB_{2l-2\dim \rho}^{ac}(Y; R)$, and so is well-defined. For (ii), given an oriented Kuranishi space W with boundary and an almost CR structure $(\mathbf{D}, \mathbf{J}, \mathbf{K})$ on W , we define $W^{\Gamma, \rho}, \iota^{\Gamma, \rho} : W^{\Gamma, \rho} \rightarrow W$ and $(\mathbf{D}^{\Gamma, \rho}, \mathbf{J}^{\Gamma, \rho}, \mathbf{K}^{\Gamma, \rho})$ as in Remark 5.20 for W , and we construct the orientation on $W^{\Gamma, \rho}$ as for $X_a^{\Gamma, \rho}$ above, again noting that $(T_v V)^{nt}, (E|_v)^{nt}$ are *complex* vector spaces by the argument of Remark 4.30(e).

The $\Pi_{ac}^{\Gamma, \rho}$ commute with pushforwards $h_* : KB_{2*}^{ac}(Y; R) \rightarrow KB_{2*}^{ac}(Z; R)$.

Remark 5.24. We have not defined operators $\Pi^{\Gamma, \rho}$ on *effective* Kuranishi bordism $\Pi_*^{eb}(Y; R)$, as there does not seem to be a good way to make $X_a^{\Gamma, \rho}$ into an effective Kuranishi space. In particular, using Remark 5.17(a) one can show that if $(V_p, \dots, \psi_p)$ is a sufficiently small Kuranishi neighbourhood in the germ at p in $X_a^{\Gamma, \rho}$ then for all $v \in V_p$ the stabilizer group $\text{Stab}_{V_p}(v)$ contains the centre $C(\Gamma)$ of Γ , so if $C(\Gamma) \neq \{1\}$ then V_p is not an effective orbifold, and $X_a^{\Gamma, \rho}$ not an effective Kuranishi space. There is no point in defining $\Pi^{\Gamma, \rho}$ on *trivial stabilizers* Kuranishi bordism $\Pi_*^{tr}(Y; R)$, as they would be zero for all $\Gamma \neq \{1\}$.

However, we can consider the compositions $\Pi^{\Gamma, \rho} \circ \Pi_{ef}^{Kb} : \Pi_k^{eb}(Y; R) \rightarrow KB_{k-\dim \rho}(Y; R)$ and $\Pi^{\Gamma, \rho} \circ \Pi_{tr}^{Kb} : \Pi_k^{tr}(Y; R) \rightarrow KB_{k-\dim \rho}(Y; R)$. By Remark 5.20 we have $\Pi^{\Gamma, \rho} \circ \Pi_{ef}^{Kb} = 0$ unless ρ is the isomorphism class of an *effective nontrivial* Γ -representation, not just a virtual one. For $\Gamma \neq \{1\}$ this implies that $\Pi^{\Gamma, \rho} \circ \Pi_{ef}^{Kb} = 0$ unless $\dim \rho > 0$. We can also exclude the case $\dim \rho = 1$ as in Remark 5.20, since the Kuranishi spaces are oriented. Hence $\Pi^{\Gamma, \rho} \circ \Pi_{ef}^{Kb} = 0$ unless $\dim \rho \geq 2$. For *trivial stabilizers* Kuranishi bordism we clearly have $\Pi^{\Gamma, \rho} \circ \Pi_{tr}^{Kb} = 0$ unless $\Gamma = \{1\}$. These imply that $\Pi_{ef}^{Kb} : \Pi_*^{eb}(Y; R) \rightarrow KB_*(Y; R)$ and $\Pi_{tr}^{Kb} : \Pi_*^{tr}(Y; R) \rightarrow KB_*(Y; R)$ are far from surjective, their images lie in small subspaces of $KB_*(Y; R)$.

In the same way, $\Pi^{\Gamma, \rho} \circ \Pi_{\text{eac}}^{\text{ac}} : \Pi_{2l}^{\text{eac}}(Y; R) \rightarrow KB_{2l-2\dim \rho}^{\text{ac}}(Y; R)$ is zero unless ρ is the isomorphism class of an *effective nontrivial complex* Γ -representation, not just a virtual one, so for $\Gamma \neq \{1\}$, $\Pi_{\text{ac}}^{\Gamma, \rho} \circ \Pi_{\text{eac}}^{\text{ac}} = 0$ unless $\dim \rho > 0$.

All of the above has analogues for the corresponding Kuranishi cobordism groups, which we now briefly explain. The starting point is the notion of orbifold strata for a submersion $f : V \rightarrow Y$ of orbifolds V, Y .

Definition 5.25. Let V, Y be orbifolds, and $f : V \rightarrow Y$ a submersion. Use the notation of Definition 5.15. Generalizing equation (129), as a set, define the *orbifold stratum* $V^{\Gamma, \rho, f}$ of (V, f) to be

$$V^{\Gamma, \rho, f} = \{ \text{Stab}_V(v) \cdot (v, \lambda) : v \in V, \lambda : \Gamma \rightarrow \text{Ker}(f_* : \text{Stab}_V(v) \rightarrow \text{Stab}_Y(f(v))) \\ \text{is an injective group morphism, } [(T_v V)^{\text{nt}}, (\tau_v \circ \lambda)^{\text{nt}}] = \rho \}. \quad (135)$$

That is, $V^{\Gamma, \rho, f}$ is the subset of $\text{Stab}_V(v) \cdot (v, \lambda)$ in $V^{\Gamma, \rho}$ with $\lambda(\Gamma) \subseteq \text{Ker } f_* \subseteq \text{Stab}_Y(v)$. Define $\iota^{\Gamma, \rho, f} : V^{\Gamma, \rho, f} \rightarrow V$ by $\iota^{\Gamma, \rho, f} = \iota^{\Gamma, \rho}|_{V^{\Gamma, \rho, f}}$.

Then $V^{\Gamma, \rho, f}, \iota^{\Gamma, \rho, f}$ have two important properties, easily verified from the definition of the orbifold structure on $V^{\Gamma, \rho}$ in the proof of Proposition 5.16. Firstly, $V^{\Gamma, \rho, f}$ is an open and closed subset of $V^{\Gamma, \rho}$. Hence $V^{\Gamma, \rho, f}$ is an *orbifold*, by Proposition 5.16, and $\iota^{\Gamma, \rho, f} : V^{\Gamma, \rho, f} \rightarrow V$ lifts to a *proper, finite immersion*.

Secondly, $f \circ \iota^{\Gamma, \rho, f} : V^{\Gamma, \rho, f} \rightarrow Y$ is a *submersion*. This is not immediately obvious, as the restriction of a submersion to a submanifold is generally no longer a submersion. To see it, let $v^{\Gamma, \rho, f} = \text{Stab}_V(v) \cdot (v, \lambda)$ in $V^{\Gamma, \rho, f}$, and set $y = f(v)$. Then λ makes $T_v V$ into a Γ -representation, so we may split into trivial and nontrivial representations $T_v V = (T_v V)^{\text{t}} \oplus (T_v V)^{\text{nt}}$, and $T_{v^{\Gamma, \rho, f}} V^{\Gamma, \rho, f} \cong (T_v V)^{\text{t}}$. We have linear maps $\text{d}f|_v : T_v V \rightarrow T_y Y$, which is *surjective* as f is a submersion, and $\text{d}(f \circ \iota^{\Gamma, \rho, f})|_{v^{\Gamma, \rho, f}} : T_{v^{\Gamma, \rho, f}} V^{\Gamma, \rho, f} \rightarrow T_y Y$, and under the identification $T_{v^{\Gamma, \rho, f}} V^{\Gamma, \rho, f} \cong (T_v V)^{\text{t}}$, we have $\text{d}(f \circ \iota^{\Gamma, \rho, f})|_{v^{\Gamma, \rho, f}} \cong (\text{d}f|_v)|_{(T_v V)^{\text{t}}}$.

Now $\text{d}f|_v : T_v V \rightarrow T_y Y$ is equivariant under the actions of $\text{Stab}_V(v)$ on $T_v V$ and $\text{Stab}_Y(y)$ on $T_y Y$ and the group morphism $f_* : \text{Stab}_V(v) \rightarrow \text{Stab}_Y(y)$. Since λ maps $\Gamma \rightarrow \text{Ker}(f_*)$, it follows that Γ acts trivially on T_y , and so $(T_v V)^{\text{nt}}$ lies in the kernel $\text{Ker}(\text{d}f|_v : T_v V \rightarrow T_y Y)$. As $\text{d}f|_v : T_v V \rightarrow T_y Y$ is surjective and $T_v V = (T_v V)^{\text{t}} \oplus (T_v V)^{\text{nt}}$, this implies that $(\text{d}f|_v)|_{(T_v V)^{\text{t}}} : (T_v V)^{\text{nt}} \rightarrow T_y Y$ is surjective. Therefore $\text{d}(f \circ \iota^{\Gamma, \rho, f})|_{v^{\Gamma, \rho, f}} : T_{v^{\Gamma, \rho, f}} V^{\Gamma, \rho, f} \rightarrow T_y Y$ is *surjective*, for all $v^{\Gamma, \rho, f} \in V^{\Gamma, \rho, f}$. Hence $f \circ \iota^{\Gamma, \rho, f}$ is a *submersion*.

Now let X be a Kuranishi space and $\mathbf{f} : X \rightarrow Y$ a strong submersion. Combining (132) and (136), define the *orbifold stratum* $X^{\Gamma, \rho, \mathbf{f}}$ as a set to be

$$X^{\Gamma, \rho, \mathbf{f}} = \{ \text{Stab}_X(p) \cdot (p, \lambda) : p \in X, \lambda : \Gamma \rightarrow \text{Ker}(\mathbf{f}_* : \text{Stab}_X(p) \rightarrow \text{Stab}_Y(f(p))) \\ \text{is an injective group morphism, } [(T_p V_p)^{\text{nt}} \ominus (E_p|_v)^{\text{nt}}] = \rho \}. \quad (136)$$

Then $X^{\Gamma, \rho, \mathbf{f}}$ is an open and closed subset of $X^{\Gamma, \rho}$, so it is a Kuranishi space by Proposition 5.19, and $\iota^{\Gamma, \rho, \mathbf{f}} = \iota^{\Gamma, \rho}|_{X^{\Gamma, \rho, \mathbf{f}}} : X^{\Gamma, \rho, \mathbf{f}} \rightarrow X$ is a proper, finite, strongly smooth map. Also $\mathbf{f} \circ \iota^{\Gamma, \rho, \mathbf{f}} : X^{\Gamma, \rho, \mathbf{f}} \rightarrow Y$ is a *strong submersion*.

Let Γ be a finite group with $|\Gamma|$ odd. Choose orientations on $I_1, \dots, I_N$ as in Definition 5.22. Let Y be an orbifold without boundary, and R a commutative ring. Following (133), for each isomorphism class ρ of virtual nontrivial

Γ -representations and each $k \in \mathbb{Z}$, define an operator $\Pi^{\Gamma, \rho} : KB^k(Y; R) \rightarrow KB^{k+\dim \rho}(Y; R)$ by

$$\Pi^{\Gamma, \rho} : \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a] \longmapsto \sum_{a \in A} \rho_a [X_a^{\Gamma, \rho, \mathbf{f}_a}, \mathbf{f}_a \circ \iota_a^{\Gamma, \rho, \mathbf{f}_a}]. \quad (137)$$

Here $X_a^{\Gamma, \rho, \mathbf{f}_a}$ is a compact Kuranishi space as X_a is compact and $\iota_a^{\Gamma, \rho, \mathbf{f}_a}$ is proper, and $\mathbf{f}_a \circ \iota_a^{\Gamma, \rho, \mathbf{f}_a}$ is a strong submersion as above. We define the coorientation for $(X_a^{\Gamma, \rho, \mathbf{f}_a}, \mathbf{f}_a \circ \iota_a^{\Gamma, \rho, \mathbf{f}_a})$ from the coorientation for $(X_a, \mathbf{f}_a)$ by the analogue of Definition 5.21. We also define $\Pi_{\text{ca}}^{\Gamma, \rho} : KB_{\text{ca}}^{2l}(Y; R) \rightarrow KB_{\text{ca}}^{2l+2\dim \rho}(Y; R)$ by including co-almost complex structures in the obvious way.

Suppose Y, Z are orbifolds, X is a Kuranishi space, $\mathbf{f} : X \rightarrow Z$ is a strong submersion, and $h : Y \rightarrow Z$ is a smooth map. Then $Y \times_{h, Z, \mathbf{f}} X$ is a Kuranishi space, and $\pi_Y : Y \times_Z X \rightarrow Y$ a strong submersion. One can show that there is a natural isomorphism of Kuranishi spaces $(Y \times_{h, Z, \mathbf{f}} X)^{\Gamma, \rho, \pi_Y} \cong Y \times_{h, Z, \mathbf{f} \circ \iota^{\Gamma, \rho, \mathbf{f}}} X^{\Gamma, \rho, \mathbf{f}}$. Using this, we find that the $\Pi^{\Gamma, \rho}$ commute with pullbacks, $h^* \circ \Pi^{\Gamma, \rho} = \Pi^{\Gamma, \rho} \circ h^* : KB^k(Z; R) \rightarrow KB^{k+\dim \rho}(Y; R)$, and similarly $h^* \circ \Pi_{\text{ca}}^{\Gamma, \rho} = \Pi_{\text{ca}}^{\Gamma, \rho} \circ h^*$.

5.7 How large are Kuranishi (co)bordism groups?

We will now use the operators $\Pi^{\Gamma, \rho}, \Pi_{\text{ac}}^{\Gamma, \rho}$ of §5.6 composed with projection to singular homology $\Pi_{\text{Kb}}^{\text{si}}, \Pi_{\text{ac}}^{\text{si}}$ to show that the bordism groups $KB_*, KB_*^{\text{ac}}(Y; R)$ are *very large*, infinitely generated at least in every even degree, and that $KB_*^{\text{ef}}, KB_*^{\text{eac}}(Y; R)$ are *large*, infinitely generated at least in every positive even degree. Poincaré duality then implies the corresponding results for the cobordism groups $KB^*, KB_{\text{ca}}^*, KB_{\text{ecb}}^*, KB_{\text{eac}}^*(Y; R)$.

Example 5.26. Let Y be a connected, nonempty orbifold, and fix $y \in Y$, so that $H_0(Y; \mathbb{Q}) \cong \mathbb{Q}$ is generated by $[y] \in H_0(Y; \mathbb{Q})$. Let Δ be a finite group. Then $\{0\}/\Delta$ is a compact 0-orbifold without boundary, consisting of a single point with stabilizer group Δ . Give $\{0\}/\Delta$ the positive orientation. Let $f : \{0\}/\Delta \rightarrow Y$ be the smooth map of orbifolds mapping $0 \mapsto y$, with induced map $f_* : \text{Stab}_{\{0\}/\Delta}(0) \rightarrow \text{Stab}_Y(y)$ given by $f_* \equiv 1$. Regarding $\{0\}/\Delta$ as a Kuranishi space and f as strongly smooth, $[\{0\}/\Delta, f]$ is well-defined in $KB_0(Y; \mathbb{Z})$. Write $(\mathbf{0}, \mathbf{0})$ for the trivial almost complex structure on $\{0\}/\Delta$, giving almost complex structures on zero vector spaces. Then $[\{0\}/\Delta, (\mathbf{0}, \mathbf{0}), f]$ is well-defined in $KB_0^{\text{ac}}(Y; \mathbb{Z})$.

Let Γ be a finite group, and take $\rho = 0$ to be the isomorphism class of the zero (virtual) Γ -representation. Then (129) and (132) imply that $(\{0\}/\Delta)^{\Gamma, 0} \cong \text{InjHom}(\Gamma, \Delta)/\Delta$ as an orbifold, where $\text{InjHom}(\Gamma, \Delta)$ is the set of injective group morphisms $\mu : \Gamma \rightarrow \Delta$, and Δ acts on $\text{InjHom}(\Gamma, \Delta)$ by $\delta : \mu \rightarrow \mu^\delta$ for $\delta \in \Delta$, where $\mu^\delta(\gamma) = \delta\mu(\gamma)\delta^{-1}$ for $\gamma \in \Gamma$.

Suppose $|\Gamma|$ is odd. Then Definition 5.21 gives an orientation on $(\{0\}/\Delta)^{\Gamma, 0}$, which is just the positive orientation on $\text{InjHom}(\Gamma, \Delta)/\Delta$, and Definition 5.22 defines $\Pi^{\Gamma, 0} : KB_0(Y; \mathbb{Z}) \rightarrow KB_0(Y; \mathbb{Z})$, where

$$\Pi^{\Gamma, 0}([\{0\}/\Delta, f]) = [\text{InjHom}(\Gamma, \Delta)/\Delta, f^{\Gamma, 0}], \quad (138)$$

where $f^{\Gamma,0}$ maps $\text{InjHom}(\Gamma, \Delta)/\Delta \rightarrow y$ with $f_*^{\Gamma,0} \equiv 1$ on stabilizers.

Now apply the morphism $\Pi_{\text{Kb}}^{\text{si}} : KB_0(Y; \mathbb{Z}) \rightarrow H_0(Y; \mathbb{Q})$ of Definition 5.5(f). (Note that the definition of $\Pi_{\text{Kb}}^{\text{si}}$ uses the fact that $\Pi_{\text{Kh}}^{\text{si}} : KH_0(Y; \mathbb{Q}) \rightarrow H_0(Y; \mathbb{Q})$ is invertible, a highly nontrivial fact from Theorems 4.8 and 4.9 which took all of Appendices A–C to prove.) From (138) we see that

$$\Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma,0}([\{0\}/\Delta, f]) = \frac{|\text{InjHom}(\Gamma, \Delta)|}{|\Delta|} [y] \in H_0(Y; \mathbb{Q}). \quad (139)$$

Similarly, for all finite groups Γ we find that

$$\Pi_{\text{ac}}^{\text{si}} \circ \Pi^{\Gamma,0}([\{0\}/\Delta, (\mathbf{0}, \mathbf{0}), f]) = \frac{|\text{InjHom}(\Gamma, \Delta)|}{|\Delta|} [y] \in H_0(Y; \mathbb{Q}). \quad (140)$$

Using Example 5.26 we show that $KB_0(Y; \mathbb{Z})$ and $KB_0^{\text{ac}}(Y; \mathbb{Z})$ contain a copy of $\mathbb{Z}^\infty$, so they are very large. The same proof shows that if R is a commutative ring without torsion, so that $\pi : R \rightarrow R \otimes_{\mathbb{Z}} \mathbb{Q}$ is injective, then $KB_0(Y; R)$ and $KB_0^{\text{ac}}(Y; R)$ contain a copy of R^∞ .

Proposition 5.27. *In Example 5.26, the elements $[\{0\}/\Delta, f]$ taken over all isomorphism classes of finite groups Δ with $|\Delta|$ odd are linearly independent over $\mathbb{Z}$ in $KB_0(Y; \mathbb{Z})$. Similarly, the elements $[\{0\}/\Delta, (\mathbf{0}, \mathbf{0}), f]$ taken over all isomorphism classes of finite groups Δ odd are linearly independent over $\mathbb{Z}$ in both $KB_0^{\text{ac}}(Y; \mathbb{Z})$. Hence $KB_0(Y; \mathbb{Z})$ and $KB_0^{\text{ac}}(Y; \mathbb{Z})$ are infinitely generated over $\mathbb{Z}$.*

Proof. Suppose for a contradiction that the $[\{0\}/\Delta, f]$ are not linearly independent over $\mathbb{Z}$ in $KB_0(Y; \mathbb{Z})$. Then there exist pairwise non-isomorphic finite groups $\Delta_1, \dots, \Delta_N$ with $|\Delta_i|$ odd and nonzero integers $a_1, \dots, a_N$ such that $\sum_{i=1}^N a_i [\{0\}/\Delta_i, f_i] = 0$ in $KB_0(Y; \mathbb{Z})$. Pick $i = 1, \dots, N$ with $|\Delta_i|$ largest, and set $\Gamma = \Delta_i$. If $j \neq i$ then either $|\Delta_j| < |\Gamma|$, or $|\Delta_j| = |\Gamma|$ but $\Delta_j \not\cong \Gamma$, and in both cases $\text{InjHom}(\Gamma, \Delta_j) = \emptyset$. Thus $\Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma,0}([\{0\}/\Delta_j, f]) = 0$ for $j \neq i$ by (139). So applying $\Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma,0}$ to $\sum_{i=1}^N a_i [\{0\}/\Delta_i, f_i] = 0$ and using (139) gives $a_i |\text{Aut}(\Gamma)|/|\Gamma| [y] = 0$ in $H_0(Y; \mathbb{Q}) \cong \mathbb{Q}$, a contradiction as $a_i, |\text{Aut}(\Gamma)|, |\Gamma|$ and $[y]$ are all nonzero. The proof for $KB_0^{\text{ac}}(Y; \mathbb{Z})$ is the same. $\square$

We can prove similar results for $\Pi^{\Gamma, \rho}$ with $\rho \neq 0$, and so show that KB_{2k} , $KB_{2k}^{\text{ac}}(Y; \mathbb{Z})$ are infinitely generated for all $k \in \mathbb{Z}$. To do this, we need to replace $\{0\}/\Delta$ by a more complicated Kuranishi space X .

Example 5.28. Let Y, y be as above. Suppose Δ is a finite group, and let $(W, \omega), (W', \omega')$ be finite-dimensional nontrivial complex representations of Δ . Consider the complex projective space $P(\mathbb{C} \oplus W) = W \amalg P(W)$. The representation $1 \oplus \omega$ of Δ on $\mathbb{C} \oplus W$ induces an action of Δ on $P(\mathbb{C} \oplus W)$, so that $P(\mathbb{C} \oplus W)/\Delta$ is a compact complex orbifold. The point $[1, 0] \in P(\mathbb{C} \oplus W)$ is fixed by Δ , so its image $[1, 0] \in P(\mathbb{C} \oplus W)/\Delta$ has stabilizer group Δ .

Define X to be $P(\mathbb{C} \oplus W)/\Delta$ as a topological space, and define a Kuranishi structure on X by the single Kuranishi neighbourhood

$$(V, E, s, \psi) = (P(\mathbb{C} \oplus W)/\Delta, (P(\mathbb{C} \oplus W) \times W')/\Delta, 0, \text{id}_X).$$

Here $E = (P(\mathbb{C} \oplus W) \times W')/\Delta \rightarrow P(\mathbb{C} \oplus W)/\Delta$ is an orbifold vector bundle with fibre W' , and Δ acts on $P(\mathbb{C} \oplus W) \times W'$ by the given action on $P(\mathbb{C} \oplus W)$, and the representation ω' on W' . Then X is a compact Kuranishi space without boundary, of virtual dimension $2(\dim W - \dim W')$. The complex structures on $P(\mathbb{C} \oplus W)$ and W' induce an *almost complex structure* $(\mathbf{J}, \mathbf{K})$ on X . This in turn induces an *orientation* on X . Define a strongly smooth map $\mathbf{f} : X \rightarrow Y$ to be represented by the projection $f : V \rightarrow Y$ mapping $f : v \mapsto y$ for all $v \in V$, with $f_* \equiv 1$ on stabilizer groups. Then $[X, \mathbf{f}]$ is well-defined in $KB_{2(\dim W - \dim W')}(Y; R)$, and $[X, (\mathbf{J}, \mathbf{K}), \mathbf{f}]$ is well-defined in $KB_{2(\dim W - \dim W')}^{\text{ac}}(Y; R)$.

For each $p \in X$, the stabilizer group $\text{Stab}_X(p)$ is a subgroup of Δ . Thus if $|\Gamma| > |\Delta|$, or if $|\Gamma| = |\Delta|$ but $\Gamma \not\cong \Delta$, then there can exist no injective morphisms $\lambda : \Gamma \rightarrow \text{Stab}_X(p)$, so (132) gives $X^{\Gamma, \rho} = \emptyset$. Hence

$$\Pi^{\Gamma, \rho}([X, \mathbf{f}]) = \Pi_{\text{ac}}^{\Gamma, \rho}([X, (\mathbf{J}, \mathbf{K}), \mathbf{f}]) = 0 \text{ if } |\Gamma| > |\Delta| \text{ or } |\Gamma| = |\Delta|, \Gamma \not\cong \Delta. \quad (141)$$

Now set $\Gamma = \Delta$. The points in X with stabilizer group Δ are the fixed point set of Δ in $P(\mathbb{C} \oplus W)$. Writing $P(\mathbb{C} \oplus W) = W \amalg P(W)$, the only fixed point of Δ in W is 0, since W is a *nontrivial* Δ -representation. The fixed points of Δ in $P(W)$ correspond to Δ -subrepresentations of W isomorphic to $\mathbb{C}$. We want to ensure that *none of these fixed points in $P(W)$ is isolated*. We therefore impose the condition:

- (*) No one-dimensional irreducible complex representation of Δ occurs with multiplicity exactly one in (W, ω) .

This implies that the fixed points of Δ in $P(W)$ are a (possibly empty) finite disjoint union of projective subspaces of $P(W)$ of *positive dimension*.

Thus, if $X^{\Delta, \rho} \neq \emptyset$, there are two possibilities:

- (a) $\rho = [(W, \omega) \ominus (W', \omega')]$, in which case $X^{\Delta, \rho}$ comes from $0 \in W$, and $X^{\Delta, \rho} \cong \text{Aut}(\Delta)/\Delta$ as an orbifold, and $\text{vdim } X^{\Delta, \rho} = 0$; or
- (b) $X^{\Delta, \rho}$ is one or more positive-dimensional projective subspaces of $P(W)$, an orbifold, and $\text{vdim } X^{\Delta, \rho} > 0$.

Thus as for (140) we see that

$$\begin{aligned} \Pi_{\text{kb}}^{\text{si}} \circ \Pi^{\Delta, \rho}([X, \mathbf{f}]) &= \Pi_{\text{ac}}^{\text{si}} \circ \Pi^{\Delta, \rho}([X, (\mathbf{J}, \mathbf{K}), \mathbf{f}]) \\ &= \begin{cases} \frac{|\text{Aut}(\Delta)|}{|\Delta|} [y], & \rho = [(W, \omega) \ominus (W', \omega')], \\ 0, & \rho \neq [(W, \omega) \ominus (W', \omega')], \dim_{\mathbb{R}} \rho = \text{vdim } X. \end{cases} \end{aligned} \quad (142)$$

In the almost complex case, for all isomorphism classes ρ of complex virtual nontrivial Δ -representations, we can choose complex nontrivial representations $(W, \omega), (W', \omega')$ satisfying $(*)$ with $\rho = [(W, \omega) \ominus (W', \omega')]$. In the real case, if $|\Delta|$ is odd then every real nontrivial representation of Δ has an underlying complex representation, so again, for all isomorphism classes ρ of real virtual nontrivial Δ -representations, we can choose complex nontrivial representations $(W, \omega), (W', \omega')$ satisfying $(*)$ with $\rho = [(W, \omega) \ominus (W', \omega')]$ in real virtual representations. Thus, using (141)–(142) and a similar argument to the proof of Proposition 5.27 gives:

Proposition 5.29. *In Example 5.28, the elements $[X, \mathbf{f}]$ taken over all isomorphism classes of finite groups Δ with $|\Delta|$ odd and all isomorphism classes ρ of real virtual nontrivial Δ -representations are linearly independent over $\mathbb{Z}$ in $KB_*(Y; \mathbb{Z})$. There are infinitely many such generators $[X, \mathbf{f}]$ in $KB_{2l}(Y; \mathbb{Z})$ for all $l \in \mathbb{Z}$. Hence $KB_{2l}(Y; \mathbb{Z})$ is infinitely generated over $\mathbb{Z}$ for all $l \in \mathbb{Z}$.*

Similarly, the elements $[X, (\mathbf{J}, \mathbf{K}), \mathbf{f}]$ taken over all isomorphism classes of finite groups Δ odd and all isomorphism classes ρ of complex virtual nontrivial Δ -representations are linearly independent over $\mathbb{Z}$ in $KB_{2*}^{\text{ac}}(Y; \mathbb{Z})$. Hence $KB_{2l}^{\text{ac}}(Y; \mathbb{Z})$ is infinitely generated over $\mathbb{Z}$ for all $l \in \mathbb{Z}$.

We can also use these ideas to study *effective* Kuranishi bordism. It is easy to see that the Kuranishi space X of Example 5.28 is effective if and only if $W' = 0$ and W is an *effective* representation, and then X is actually an *effective orbifold*, so that $[X, \mathbf{f}]$ is well-defined in $B_*^{\text{eo}}(Y; \mathbb{Z})$. Thus as for Proposition 5.29 we obtain:

Proposition 5.30. *In Example 5.28, taking $W' = 0$ and (W, ω) to be an effective nontrivial Δ -representation, the elements $[X, \mathbf{f}]$ taken over all isomorphism classes of finite groups Δ with $|\Delta|$ odd and all isomorphism classes of (W, ω) satisfying $(*)$ are linearly independent over $\mathbb{Z}$ in both $KB_*^{\text{ef}}(Y; \mathbb{Z})$ and $B_*^{\text{eo}}(Y; \mathbb{Z})$. There are infinitely many such generators $[X, \mathbf{f}]$ in $KB_{2l}(Y; \mathbb{Z})$ and $B_{2l}^{\text{eo}}(Y; \mathbb{Z})$ for all $l = 2, 3, 4, \dots$. Hence $KB_{2l}(Y; \mathbb{Z})$, $B_{2l}^{\text{eo}}(Y; \mathbb{Z})$, and similarly $KB_{2l}^{\text{eac}}(Y; \mathbb{Z})$, are infinitely generated over $\mathbb{Z}$ for all $l \geq 2$.*

We exclude the case $l = 1$ here as condition $(*)$ in Example 5.28 would fail. But using different examples of Kuranishi spaces X one can also show that $KB_2, B_2^{\text{eo}}, KB_2^{\text{eac}}(Y; \mathbb{Z})$ are infinitely generated over $\mathbb{Z}$. Using the Poincaré duality ideas of §5.4 we can deduce:

Corollary 5.31. *Let Y be an oriented n -orbifold without boundary. Then $KB^{n-2l}(Y; \mathbb{Z})$ is infinitely generated over $\mathbb{Z}$ for all $l \in \mathbb{Z}$, and $KB_{\text{ec}}^{n-2l}(Y; \mathbb{Z})$ is infinitely generated over $\mathbb{Z}$ for all $l \geq 2$.*

Similar results hold for $KB_{\text{ac}}^*(Y; \mathbb{Z})$ and $KB_{\text{eac}}^*(Y; \mathbb{Z})$. The fact that our Kuranishi (co)bordism groups are very large has both advantages and disadvantages. One advantage is that any invariant defined in a Kuranishi (co)bordism group contains a lot of information, as it lies in a very large group. We will see this with Gromov–Witten cobordism invariants in §6. One disadvantage is that it is not really feasible to compute the groups and write them down.

6 Applications in Symplectic Geometry

We now discuss some applications of Kuranishi (co)homology and Kuranishi (co)bordism in areas of Symplectic Geometry concerned with moduli spaces of J -holomorphic curves. Our aim is not to give a complete treatment — we postpone this to the sequels [39, 41] — but rather to demonstrate the potential of our theories as tools both for proving new theorems, and for reframing existing theories in a much cleaner and more economical way.

Sections 6.1–6.3 concern closed Gromov–Witten invariants. They explain how to define new Gromov–Witten type invariants in (almost complex) Kuranishi bordism groups, and consider how these might be used to study integrality properties of Gromov–Witten invariants, and prove the Gopakumar–Vafa Integrality Conjecture. Sections 6.4–6.7 discuss Lagrangian Floer cohomology and open Gromov–Witten invariants, and §6.8 draws some conclusions.

6.1 Moduli spaces of closed Riemann surfaces

We now discuss (closed) Riemann surfaces and stable maps, following [25].

Definition 6.1. A *Riemann surface* Σ is a compact connected complex 1-manifold. A *prestable* or *nodal Riemann surface* Σ is a compact connected complex variety whose only singularities are *nodes* (*ordinary double points*), that is, Σ has finitely many singularities, each modelled on the *node* $(0, 0)$ in $\{(x, y) \in \mathbb{C}^2 : xy = 0\}$. The *genus* of a prestable Riemann surface Σ is the genus of its smoothing Σ' , that is, we smooth nodes modelled on $\{(x, y) \in \mathbb{C}^2 : xy = 0\}$ to a nonsingular curve modelled on $\{(x, y) \in \mathbb{C}^2 : xy = \epsilon\}$. A (*prestable*) *Riemann surface with m marked points* $(\Sigma, \vec{z})$ is a (prestable) Riemann surface Σ with $\vec{z} = (z_1, \dots, z_m)$, where $z_1, \dots, z_m$ are distinct points in Σ , none of which is a node. A *stable* Riemann surface with marked points is a prestable $(\Sigma, \vec{z})$ whose automorphism group is finite.

Write $\bar{\mathcal{M}}_{g,m}$ for the moduli space of isomorphism classes $[\Sigma, \vec{z}]$ of stable Riemann surfaces $(\Sigma, \vec{z})$ of genus g with m marked points, and $\mathcal{M}_{g,m}$ for the subset of $[\Sigma, \vec{z}]$ in $\bar{\mathcal{M}}_{g,m}$ with Σ nonsingular. Then $\bar{\mathcal{M}}_{g,m}$ is known as the *Deligne–Mumford compactification* of $\mathcal{M}_{g,m}$. It is easy to show that when $(g, m) = (0, 0), (0, 1), (0, 2)$ or $(1, 0)$ we have $\mathcal{M}_{g,m} = \bar{\mathcal{M}}_{g,m} = \emptyset$. So we generally restrict to $2g + m \geq 3$.

Let $(\Sigma, \vec{z})$ be a prestable Riemann surface with genus g and m marked points, and suppose $2g + m \geq 3$. If $(\Sigma, \vec{z})$ is not stable then Σ has a $\mathbb{CP}^1$ component with less than 3 nodes or marked points. Collapse this $\mathbb{CP}^1$ component to a point. After repeating this process finitely many times we obtain a unique stable Riemann surface $(\tilde{\Sigma}, \tilde{\vec{z}})$ with genus g and m marked points, called the *stabilization* of $(\Sigma, \vec{z})$.

A great deal is known about the moduli spaces $\mathcal{M}_{g,m}$ and $\bar{\mathcal{M}}_{g,m}$, see for instance Harris and Morrison [32]. The following theorem is proved in [25, §9].

Theorem 6.2. *In Definition 6.1, $\bar{\mathcal{M}}_{g,m}$ is a compact complex orbifold of real dimension $2(3g + m - 3)$, without boundary or corners.*

Observe that $\bar{\mathcal{M}}_{1,1}$ is a *non-effective orbifold*, whose generic points have stabilizer group $\mathbb{Z}_2$.

Definition 6.3. Let (M, ω) be a compact symplectic manifold and J an almost complex structure on M compatible with ω . A *stable map to M from a Riemann surface with m marked points* is a triple $(\Sigma, \vec{z}, w)$, where $(\Sigma, \vec{z})$ is a prestable Riemann surface with m marked points, and $w : \Sigma \rightarrow M$ a J -pseudoholomorphic map, such that the group $\text{Aut}(\Sigma, \vec{z}, w)$ of biholomorphisms $\gamma : \Sigma \rightarrow \Sigma$ with $\gamma(z_j) = z_j$ for all j and $w \circ \gamma = w$ is finite.

For $\beta \in H^2(M, \mathbb{Z})$ and $g, m \geq 0$, write $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ for the moduli space of J -pseudoholomorphic stable maps $(\Sigma, \vec{z}, w)$ to M from a Riemann surface Σ with genus g and m marked points, with $[w(\Sigma)] = \beta \in H^2(M, \mathbb{Z})$. Points of $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ are isomorphism classes $[\Sigma, \vec{z}, w]$, where $(\Sigma, \vec{z}, w), (\Sigma', \vec{z}', w')$ are isomorphic if there exists a biholomorphism $\gamma : \Sigma \rightarrow \Sigma'$ with $\gamma(z_j) = z'_j$ for all j and $w' \circ \gamma \equiv w$. Define *evaluation maps* $\text{ev}_j : \bar{\mathcal{M}}_{g,m}(M, J, \beta) \rightarrow M$ for $j = 1, \dots, m$ by $\text{ev}_j : [\Sigma, \vec{z}, w] \mapsto w(z_j)$. When $2g + m \geq 3$, define $\pi_{g,m} : \bar{\mathcal{M}}_{g,m}(M, J, \beta) \rightarrow \bar{\mathcal{M}}_{g,m}$ by $\text{ev}_j : [\Sigma, \vec{z}, w] \mapsto [\tilde{\Sigma}, \tilde{\vec{z}}]$, where $(\tilde{\Sigma}, \tilde{\vec{z}})$ is the stabilization of $(\Sigma, \vec{z})$.

There is a natural topology on $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ called the C^∞ topology, due to Gromov [31] and defined in [25, §10]. The next result, due in principle to Gromov, is proved by Fukaya and Ono [25, §10–§11].

Theorem 6.4. *In Definition 6.3, $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ is compact and Hausdorff in the C^∞ topology.*

Remark 6.5. In the following, I will label Theorems 6.6, 6.7, 6.8, 6.23, 6.25 and 6.26 as “Theorem” rather than Theorem. This is intended to indicate that *I am not confident that a complete, detailed proof has yet been given for these results*; that they should be regarded as *partially proved*, as somewhere between a Theorem and a Conjecture.

I state these as “Theorems” because actually proving them properly would be a massive undertaking, beyond the scope of this book and beyond my abilities and patience, but I cannot proceed to applications in Symplectic Geometry without them, and I need some applications to persuade the reader that the effort of defining my Kuranishi (co)homology machinery was worthwhile.

The results of §6.2–§6.3 will then assume the “Theorems” of §6.1, and the results of §6.5–§6.7 will assume the “Theorems” of §6.4. For comparison, note that whole areas of mathematics have the Riemann hypothesis as a standing assumption, although this has not yet been proved. In this case, we have good reasons to hope that complete proofs of our “Theorems” will be available soon.

The worries I have about the proofs of the “Theorems” below mostly concern the definition of the Kuranishi structure on the moduli spaces at and near singular J -holomorphic curves, that is, curves with interior or boundary nodes. Especially, I am bothered about *smoothness issues*: whether one can define Kuranishi neighbourhoods (V_p, E_p, s_p, ψ_p) on the moduli spaces near such a singular curve for which s_p is a smooth section of E_p , and what arbitrary choices are

involved; whether coordinate changes $(\phi_{pq}, \hat{\phi}_{pq})$ between such neighbourhoods can be made smooth, and whether they satisfy Definition 2.18(e).

These questions are very difficult, concerning as they do singularities of solutions of nonlinear elliptic p.d.e.s, with bubbling, neck stretching, and so on — the ‘Analytical Chamber of Horrors’, as Hofer describes it [33, p. 2]. Fukaya et al. discuss smoothness in [24, §A1.4]. Their underlying idea is related to the notion of *gluing profile* in Hofer [33, §4.1], which is how these smoothness issues are resolved in the theory of polyfolds.

Part of the problem is the mismatch between the definitions of Kuranishi spaces used in this book, and in our references. For example, Fukaya and Ono give a proof of “Theorem” 6.6 below using a different and rather weaker notion of Kuranishi space: for them, the Kuranishi maps s_p only have to be continuous, so they avoid most of the smoothness issues, and Definition 2.18(e) is replaced by a weaker notion involving isomorphisms χ_{pq} as in Remark 2.19. But I want “Theorem” 6.6 to be true with our stronger notion of Kuranishi space.

Part of the problem too is that I am too lazy to work through all the proofs that have been written down, and satisfy myself that they are correct, and extend to our stronger definition of Kuranishi spaces. But also, there are parts of some proofs in [24, 25] which I have read, which appeared to me to be either lacking in detail at some important points, or to have problems which needed fixing. Fortunately, as we will discuss in §6.8, there may soon be an alternative proof of “Theorem” 6.6 and related results, via the theory of polyfolds of Hofer, Wysocki and Zehnder [33–37].

The following “Theorem” is proved in [25, Th.s 7.10–7.11 & Prop. 16.1], but with a weaker definition of Kuranishi structure. The claims about strong submersions are evident from the construction.

“Theorem” 6.6. *In Definition 6.3, there exists an oriented Kuranishi structure κ on $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$, without boundary or (g) -corners, depending on choices, with*

$$\text{vdim } \bar{\mathcal{M}}_{g,m}(M, J, \beta) = 2(c_1(M) \cdot \beta + (n - 3)(1 - g) + m), \quad (143)$$

where $\dim M = 2n$. The maps ev_j in Definition 6.3 are continuous and extend for $j = 1, \dots, m$ to strong submersions $\mathbf{ev}_j : (\bar{\mathcal{M}}_{g,m}(M, J, \beta), \kappa) \rightarrow M$, and $\mathbf{ev}_1 \times \dots \times \mathbf{ev}_m$ is a strong submersion. When $2g + m \geq 3$, the map $\pi_{g,m}$ in Definition 6.3 is continuous and extends to a strong submersion $\pi_{g,m} : (\bar{\mathcal{M}}_{g,m}(M, J, \beta), \kappa) \rightarrow \bar{\mathcal{M}}_{g,m}$, and $\mathbf{ev}_1 \times \dots \times \mathbf{ev}_m \times \pi_{g,m}$ is a strong submersion.

Our next “Theorem” describes the dependence of $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ on the almost complex structure J . It is discussed briefly in the proof of [25, Th. 17.11], and proved by the same method as “Theorem” 6.6.

“Theorem” 6.7. *Let (M, ω) be a compact symplectic manifold and J_t for $t \in [0, 1]$ a smooth 1-parameter family of almost complex structures on M compatible*

with ω . Write

$$\begin{aligned} \bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta) = \\ \{(t, [\Sigma, \bar{z}, w]) : t \in [0, 1], [\Sigma, \bar{z}, w] \in \bar{\mathcal{M}}_{g,m}(M, J_t, \beta)\}. \end{aligned}$$

There is a natural C^∞ topology on $\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta)$ making it into a compact Hausdorff topological space. Let κ_0, κ_1 be possible oriented Kuranishi structures on $\bar{\mathcal{M}}_{g,m}(M, J_t, \beta)$ for $t = 0, 1$ given by “Theorem” 6.6. Then there exists an oriented Kuranishi structure $\tilde{\kappa}$ on $\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta)$, with

$$\text{vdim}(\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta), \tilde{\kappa}) = 2(c_1(M) \cdot \beta + (n - 3)(1 - g) + m) + 1,$$

without (g) -corners, and with boundary

$$\begin{aligned} \partial(\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta), \tilde{\kappa}) \\ = (\bar{\mathcal{M}}_{g,m}(M, J_1, \beta), \kappa_1) \amalg -(\bar{\mathcal{M}}_{g,m}(M, J_0, \beta), \kappa_0). \end{aligned} \quad (144)$$

There are strong submersions $\tilde{\mathbf{e}}\mathbf{v}_j : (\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta), \tilde{\kappa}) \rightarrow M$ for $j = 1, \dots, m$ and $\tilde{\pi}_{g,m} : (\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta), \tilde{\kappa}) \rightarrow \bar{\mathcal{M}}_{g,m}$ when $2g + m \geq 3$, which restrict to $\mathbf{e}\mathbf{v}_j, \pi_{g,m}$ in “Theorem” 6.6 on $\bar{\mathcal{M}}_{g,m}(M, J_t, \beta)$ for $t = 0, 1$ under (144), and $\tilde{\mathbf{e}}\mathbf{v}_1 \times \dots \times \tilde{\mathbf{e}}\mathbf{v}_m, \tilde{\mathbf{e}}\mathbf{v}_1 \times \dots \times \tilde{\mathbf{e}}\mathbf{v}_m \times \tilde{\pi}_{g,m}$ are strong submersions. Note that we can take $J_t \equiv J$ for $t \in [0, 1]$, so this result relates possible choices of κ in “Theorem” 6.6 for a single J on M .

We can put an *almost complex structure* on $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$, in the sense of §2.9. The (incomplete) proof is based on Fukaya and Ono [25, Prop. 16.5], who prove that $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ is *stably almost complex*, that is, that the virtual tangent bundle of $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ is equivalent to a virtual complex vector bundle in a version of K-theory for Kuranishi spaces.

“Theorem” 6.8. *In the situation of “Theorem” 6.6, one can construct an almost complex structure $(\mathbf{J}, \mathbf{K})$ on $(\bar{\mathcal{M}}_{g,m}(M, J, \beta), \kappa)$. This construction depends on some choices, and κ must also be chosen appropriately.*

These $\kappa, (\mathbf{J}, \mathbf{K})$ are unique up to isotopy in the following sense. Let J_t for $t \in [0, 1]$ be a smooth family of almost complex structures on M compatible with ω . Let $\kappa_0, (\mathbf{J}_0, \mathbf{K}_0)$ and $\kappa_1, (\mathbf{J}_1, \mathbf{K}_1)$ be outcomes of the construction above for $\bar{\mathcal{M}}_{g,m}(M, J_0, \beta), \bar{\mathcal{M}}_{g,m}(M, J_1, \beta)$. Then in “Theorem” 6.7 we can choose $\tilde{\kappa}$ and an almost CR structure $(\mathbf{D}, \mathbf{J}, \mathbf{K})$ on $(\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta), \tilde{\kappa})$ with

$$\begin{aligned} \partial((\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta), \tilde{\kappa}), (\mathbf{D}, \mathbf{J}, \mathbf{K})) = \\ ((\bar{\mathcal{M}}_{g,m}(M, J_1, \beta), \kappa_1), (\mathbf{J}_1, \mathbf{K}_1)) \amalg -((\bar{\mathcal{M}}_{g,m}(M, J_0, \beta), \kappa_0), (\mathbf{J}_0, \mathbf{K}_0)). \end{aligned} \quad (145)$$

Analogues of the above results also hold for *co-almost complex structures* and *co-almost CR structures*, using the ideas of §2.10.

Partial proof. In [25, Prop. 16.5], Fukaya and Ono observe that for a Kuranishi neighbourhood (V_p, E_p, s_p, ψ_p) on $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$, the fibres of the orbibundles

TV_p and E_p over V_p are roughly the kernels and cokernels of a family of linear elliptic operators $D_v : B_1 \rightarrow B_2$ between *complex* Banach spaces B_1, B_2 , parametrized by $v \in V_p$. Now the D_v are not complex linear, but their symbols D'_v are. So writing $D_{(v,t)} = (1-t)D_v + tD'_v$, we obtain a smooth family of elliptic operators $D_{(v,t)} : B_1 \rightarrow B_2$ parametrized by $(v, t) \in V_p \times [0, 1]$, such that $D_{(v,0)} = D_v$ and $D_{(v,1)}$ is complex linear. Using this and ideas in the proofs of [25, Prop. 16.5] and in §3.2, we can construct the following data:

- (i) A *good coordinate system* $\mathbf{I} = (I, \leq, (V^i, E^i, s^i, \psi^i) : i \in I, \dots)$ for $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$, as in Definition 3.2;
- (ii) Real orbibundles F_1^i, F_2^i over $V^i \times [0, 1]$ for all $i \in I$;
- (iii) Isomorphisms $\iota_1^i : F_1^i|_{V^i \times \{0\}} \rightarrow TV^i$, $\iota_2^i : F_2^i|_{V^i \times \{0\}} \rightarrow E^i$ for all $i \in I$;
- (iv) Almost complex structures J_1^i, J_2^i on the fibres of $F_1^i|_{V^i \times \{1\}}$ and $F_2^i|_{V^i \times \{1\}}$ for all $i \in I$; and
- (v) For all $i, j \in I$ with $j \leq i$ and $\text{Im } \psi^i \cap \text{Im } \psi^j \neq \emptyset$, embeddings of orbibundles $\Phi_a^{ij} : F_a^j|_{V^{ij}} \rightarrow (\phi^{ij})^*(F_a^i)$ over V^{ij} for $a = 1, 2$, and an isomorphism:

$$\Psi^{ij} : \frac{(\phi^{ij})^*(F_1^i)}{\Phi_1^{ij}(F_1^j|_{V^{ij}})} \longrightarrow \frac{(\phi^{ij})^*(F_2^i)}{\Phi_2^{ij}(F_2^j|_{V^{ij}})}. \quad (146)$$

This data satisfies the following conditions:

- (vi) For all $i, j \in I$ with $j \leq i$ and $\text{Im } \psi^i \cap \text{Im } \psi^j \neq \emptyset$ we have $(\phi^{ij})^*(\iota_1^i) \circ \Phi_1^{ij} = d\phi^{ij} \circ \iota_1^j$ and $(\phi^{ij})^*(\iota_2^i) \circ \Phi_2^{ij} = \hat{\phi}^{ij} \circ \iota_2^j$ as morphisms $F_1^j|_{V^{ij} \times \{0\}} \rightarrow (\phi^{ij})^*(TV^i)$ and $F_2^j|_{V^{ij} \times \{0\}} \rightarrow (\phi^{ij})^*(E^i)$ over $V^{ij} \times \{0\} \cong V^{ij}$, and the following commutes near $(s^j)^{-1}(0) \cap V^{ij}$ in morphisms over $V^{ij} \times \{0\} \cong V^{ij}$, where $d\hat{s}^i$ is as in (14):

$$\begin{array}{ccc} \frac{(\phi^{ij})^*(F_1^i)}{\Phi_1^{ij}(F_1^j|_{V^{ij}})}|_{V^{ij} \times \{0\}} & \xrightarrow{\Psi^{ij}} & \frac{(\phi^{ij})^*(F_2^i)}{\Phi_2^{ij}(F_2^j|_{V^{ij}})}|_{V^{ij} \times \{0\}} \\ \downarrow (\iota_1^i)_* & & \downarrow (\iota_2^i)_* \\ \frac{(\phi^{ij})^*(TV^i)}{(d\phi^{ij})(TV^j)} & \xrightarrow{d\hat{s}^i} & \frac{(\phi^{ij})^*(E^i)}{\hat{\phi}^{ij}(E^j)}, \end{array} \quad (147)$$

where the columns are well-defined because of the previous conditions.

- (vii) For all $i, j \in I$ with $j \leq i$ and $\text{Im } \psi^i \cap \text{Im } \psi^j \neq \emptyset$ we have $(\phi^{ij})^*(J_a^i) \circ \Phi_a^{ij} = \Phi_a^{ij} \circ J_a^j$ as morphisms $F_a^j|_{V^{ij}} \rightarrow (\phi^{ij})^*(F_a^i)$ for $a = 1, 2$. This implies that $(\phi_a^{ij})^*(J_a^i)$ for $a = 1, 2$ push down to almost complex structures J_a^{ij} for $a = 1, 2$ on the orbibundles over V^{ij} appearing in (146), and we require that $J_2^{ij} \circ \Psi^{ij} = \Psi^{ij} \circ J_1^{ij}$.
- (viii) Whenever $i, j, k \in I$ with $k \leq j \leq i$ and $\text{Im } \psi^i \cap \text{Im } \psi^j \cap \text{Im } \psi^k \neq \emptyset$, we have $\Phi_a^{ik} = (\phi^{jk})^*(\Phi_a^{ij}) \circ \Phi^{jk}$ over $(\phi^{jk})^{-1}(V^{ij}) \cap V^{jk} \cap V^{ik}$ for $a = 1, 2$, and the following diagram of orbibundles over $(\phi^{jk})^{-1}(V^{ij}) \cap V^{jk} \cap V^{ik}$ commutes:

$$\begin{array}{ccccccc}
0 & \rightarrow & \frac{(\phi^{jk})^*(F_1^j)}{\Phi_1^{jk}(F_1^k)} & \xrightarrow{(\phi^{jk})^*(\Phi_1^{ij})} & \frac{(\phi^{ik})^*(F_1^i)}{\Phi_1^{ik}(F_1^k)} & \xrightarrow{\text{projection}} & \frac{(\phi^{ik})^*(F_1^i)}{(\phi^{jk})^*(\Phi_1^{ij}(F_1^j))} \rightarrow 0 \\
& & \downarrow \Psi^{jk} & & \downarrow \Psi^{ik} & & \downarrow (\phi^{jk})^*(\Psi^{ij}) \\
0 & \rightarrow & \frac{(\phi^{jk})^*(F_2^j)}{\Phi_2^{jk}(F_2^k)} & \xrightarrow{(\phi^{jk})^*(\Phi_2^{ij})} & \frac{(\phi^{ik})^*(F_2^i)}{\Phi_2^{ik}(F_2^k)} & \xrightarrow{\text{projection}} & \frac{(\phi^{ik})^*(F_2^i)}{(\phi^{jk})^*(\Phi_2^{ij}(F_2^j))} \rightarrow 0.
\end{array}$$

Roughly speaking, the data F_a^i and overlap maps Φ_a^{ij} and Ψ^{ij} define a bundle system $(\mathbf{F}_1, \mathbf{F}_2)$ on $\bar{\mathcal{M}}_{g,m}(M, J, \beta) \times [0, 1]$, and (iii), (vi) say that $(\mathbf{F}_1, \mathbf{F}_2)$ restricts to the tangent bundle system on $\bar{\mathcal{M}}_{g,m}(M, J, \beta) \times \{0\}$, and (iv), (vii) say that $(\mathbf{F}_1, \mathbf{F}_2)$ restricts to a complex bundle system on $\bar{\mathcal{M}}_{g,m}(M, J, \beta) \times \{1\}$, as in the discussion of [25, Prop. 16.5] above. But working with neighbourhoods of the form $(V^i \times [0, 1], E^i, s^i, \psi^i)$ on $\bar{\mathcal{M}}_{g,m}(M, J, \beta) \times [0, 1]$, rather than covering $\bar{\mathcal{M}}_{g,m}(M, J, \beta) \times [0, 1]$ with small Kuranishi neighbourhoods, gives an advantage.

Now any orbibundle over $V^i \times [0, 1]$ is (non-canonically) isomorphic to the pullback of an orbibundle over V^i , by the orbifold version of well-known facts for manifolds. Thus by (iii) there exist isomorphisms $\tilde{\iota}_1^i : F_1^i \rightarrow (\pi^i)^*(TV^i)$ and $\tilde{\iota}_2^i : F_2^i \rightarrow (\pi^i)^*(E^i)$ with $\tilde{\iota}_a^i|_{V^i \times \{0\}} = \iota_a^i$ for $a = 1, 2$ and all $i \in I$, where $\pi^i : V^i \times [0, 1] \rightarrow V^i$ is the projection. We claim we can choose these isomorphisms $\tilde{\iota}_a^i$ to satisfy $(\phi^{ij})^*(\tilde{\iota}_1^i) \circ \Phi_1^{ij} = d\phi^{ij} \circ \tilde{\iota}_1^j$ and $(\phi^{ij})^*(\tilde{\iota}_2^i) \circ \Phi_2^{ij} = \hat{\phi}^{ij} \circ \tilde{\iota}_2^j$ over $V^{ij} \times [0, 1]$, as in (vi), and the obvious extension of (147) to $V^{ij} \times [0, 1]$.

To make these choices we work by induction on I in the order $\leq$. Suppose $i \in I$ and we have already chosen $\tilde{\iota}_1^j, \tilde{\iota}_2^j$ for all $j \leq i, j \neq i$. Then we have to choose $\tilde{\iota}_1^i, \tilde{\iota}_2^i$ satisfying some conditions over $\phi^{ij}(V^{ij})$ for all $j \leq i, j \neq i$. Equation (32) and the previously chosen compatibility between $\tilde{\iota}_1^j, \tilde{\iota}_2^j, \tilde{\iota}_1^k, \tilde{\iota}_2^k$ for $k \leq j \leq i$ ensure that the conditions are consistent over $\phi^{ij}(V^{ij}) \cap \phi^{ik}(V^{ik})$. Thus we can choose $\tilde{\iota}_1^i, \tilde{\iota}_2^i$ satisfying these conditions, and the claim is true by induction.

Now define almost complex structures J^i, K^i on the fibres of TV^i and E^i over V^i by $J^i = (\tilde{\iota}_1^i|_{V^i \times \{1\}})^*(J_1^i)$ and $K^i = (\tilde{\iota}_2^i|_{V^i \times \{1\}})^*(J_2^i)$. Part (vii) and the conditions on the $\tilde{\iota}_a^i$ ensure that for all $i, j \in I$ with $j \leq i, j \neq i$ and $\text{Im } \psi^i \cap \text{Im } \psi^j \neq \emptyset$, the following analogues of Definition 2.34(a)–(c) hold:

- (a) $d\phi^{ij} \circ J^j = (\phi^{ij})^*(J^i) \circ d\phi^{ij}$ as morphisms $TV^j \rightarrow (\phi^{ij})^*(TV^i)$;
- (b) $\hat{\phi}^{ij} \circ K^j = (\phi^{ij})^*(K^i) \circ \hat{\phi}^{ij}$ as morphisms $E^j \rightarrow (\phi^{ij})^*(E^i)$; and
- (c) Parts (a) and (b) imply that the orbibundles $(\phi^{ij})^*(TV^i)/(d\phi^{ij})(TV^j)$ and $(\phi^{ij})^*(E^i)/\hat{\phi}^{ij}(E^j)$ over V^{ij} have almost complex structures J^{ij}, K^{ij} on their fibres, by projection from $(\phi^{ij})^*(J^i), (\phi^{ij})^*(K^i)$. We require that $K^{ij} \circ d\hat{s}^i = d\hat{s}^i \circ J^{ij}$ over $(s^j)^{-1}(0)$ in V^{ij} , for $d\hat{s}^i$ as in (14).

We can now define κ and $(\mathbf{J}, \mathbf{K})$. For each $p \in \bar{\mathcal{M}}_{g,m}(M, J, \beta)$, let $i \in I$ be least in the order $\leq$ such that $p \in \text{Im } \psi^i$. Define the germ of Kuranishi neighbourhoods of p in κ to be the equivalence class of (V^i, E^i, s^i, ψ^i) , regarded as a Kuranishi neighbourhood of p . To define the germ of coordinate changes in κ , let $(V_p, \dots, \psi_p)$ be sufficiently small in the germ of κ at p , let $q \in \text{Im } \psi_p$,

and let $(V_q, \dots, \psi_q)$ be sufficiently small in the germ of κ at q . Let $i, j \in I$ be least in the order $\leq$ such that $p \in \text{Im } \psi^i$ and $q \in \text{Im } \psi^j$.

Then by definition there are open neighbourhoods U_p and U_q of $(\psi^i)^{-1}(p)$, $(\psi^j)^{-1}(q)$ in $V^i, V^{ij} \subseteq V^j$ such that $(V_p, \dots, \psi_p)$ and $(V_q, \dots, \psi_q)$ are isomorphic to $(U_p, E^i|_{U_p}, s^i|_{U_p}, \psi^i|_{U_p})$ and $(U_q, E^j|_{U_q}, s^j|_{U_q}, \psi^j|_{U_q})$ respectively. Define a coordinate change $(\phi_{pq}, \hat{\phi}_{pq})$ from $(V_q, \dots, \psi_q)$ to $(V_p, \dots, \psi_p)$ to be that identified with $(\phi^{ij}|_{U_q}, \hat{\phi}^{ij}|_{U_q})$ by these isomorphisms. Using Definition 3.2, it is easy to see that κ is a *Kuranishi structure*. It is one of those allowed by “Theorem” 6.6, which uses the same construction without the almost complex structures.

Let $p \in \bar{\mathcal{M}}_{g,m}(M, J, \beta)$ and $(V_p, \dots, \psi_p)$ be sufficiently small in the germ of κ at p . Define complex structures J_p on V_p and K_p on the fibres of E_p to be those identified with $J^i|_{U_p}$ and $K^i|_{U_p}$ by the isomorphisms above. Then (a)–(c) above imply that Definition 2.34(a)–(c) hold, so these J_p, K_p define an *almost complex structure* $(\mathbf{J}, \mathbf{K})$ on $(\bar{\mathcal{M}}_{g,m}(M, J, \beta), \kappa)$. This completes the first part of the theorem.

The second part works by a similar argument for $\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta)$, so we will be brief. To define $\kappa_0, (\mathbf{J}_0, \mathbf{K}_0)$ and $\kappa_1, (\mathbf{J}_1, \mathbf{K}_1)$ using the first part, we must choose good coordinate systems on $\bar{\mathcal{M}}_{g,m}(M, J_0, \beta), \bar{\mathcal{M}}_{g,m}(M, J_1, \beta)$. In the second part we choose a good coordinate system on $\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta)$ which restricts to these choices for J_0, J_1 at $t = 0, 1$. Then we go through the whole proof, choosing data $F_a^i, \iota_a^i, J_a^i, \Phi_a^{ij}, \Psi^{ij}$ and $\tilde{\iota}_a^i$ over $\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta) \times [0, 1]$ for this new good coordinate system, ensuring at each stage that the choices made agree at $t = 0, 1$ with those made in defining $\kappa_0, (\mathbf{J}_0, \mathbf{K}_0)$ and $\kappa_1, (\mathbf{J}_1, \mathbf{K}_1)$. This is always possible.

Note that (iii) above has to be modified as follows. Let (V^i, E^i, s^i, ψ^i) be a Kuranishi neighbourhood in the good coordinate system for $\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta)$, and let π in “Theorem” 6.7 be represented by the submersion $\pi^i : V^i \rightarrow [0, 1]$. Define D^i to be the codimension 1 orbisubbundle $\text{Ker}(d\pi^i)$ in TV^i . In (iii) we should take ι_1^i to be an isomorphism $\iota_1^i : F_1^i|_{V^i \times \{0\}} \rightarrow D^i$, mapping to D^i rather than TV^i . Then J^i becomes an almost complex structure on the fibres of D^i rather than TV^i , so that (D^i, J^i) is an almost CR structure on V^i , of codimension 1. This is why, in the last part of the theorem, we construct an almost CR structure $(\mathbf{D}, \mathbf{J}, \mathbf{K})$ of codimension 1 on $(\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta), \tilde{\kappa})$, rather than an almost complex structure.

Finally, the modifications for the co-almost complex and co-almost CR structures case are straightforward: we replace TV^i by the kernel of the projection to $T(M^m)$ or $T(\bar{\mathcal{M}}_{g,m} \times M^m)$ throughout, in particular in (iii) above. Note that these projections are actually complex linear, with respect to the almost complex structure J_1^i on TV^i defined using $D_{(v,1)} = D_v^i$ above, the almost complex structure $J \times \dots \times J$ on M^m , and the natural complex structure on $\bar{\mathcal{M}}_{g,m}$. Hence J_1^i restricts to an almost complex structure on the kernel of the projection $TV^i \rightarrow T(M^m)$ or $T(\bar{\mathcal{M}}_{g,m} \times M^m)$, as we need for the modified (iv) above. $\square$

6.2 Gromov–Witten invariants

In this section and §6.3, we assume the “Theorems” of §6.1 throughout. For our attitude to these, see Remark 6.5.

Gromov–Witten invariants are invariants ‘counting’ J -holomorphic curves Σ of genus g with marked points in a complex or symplectic manifold. They are important in String Theory, and attracted widespread attention through their rôle in the Mirror Symmetry story for Calabi–Yau 3-folds. They can be defined either in algebraic geometry or symplectic geometry. A Gromov–Witten invariant is basically a *virtual cycle* for the moduli space $\overline{\mathcal{M}}_{g,m}(M, J, \beta)$ of Definition 6.3, where $\overline{\mathcal{M}}_{g,m}(M, J, \beta)$ is regarded as a proper Deligne–Mumford $\mathbb{C}$ -stack with an obstruction theory in the algebraic case, and as a Kuranishi space (or something similar) in the symplectic case. For a good introduction, see Cox and Katz [19, §7].

In Algebraic Geometry, Behrend [6, 8] and Li and Tian [48] defined Gromov–Witten invariants for a smooth projective variety M over $\mathbb{C}$. In Symplectic Geometry, Fukaya and Ono [25], Li and Tian [49], Ruan [65], and Siebert [70, 71] defined Gromov–Witten invariants for a general compact symplectic manifold (M, ω) , building on earlier work of Ruan and Tian for semi-positive symplectic manifolds. It is generally believed, and partly proved, that these definitions agree with each other on their common domain of validity. Li and Tian [50] show that their algebraic [48] and symplectic [49] invariants coincide, and Siebert [72] show that his symplectic invariants [70, 71] agree with Behrend’s algebraic ones [6]. Kontsevich and Manin [45] wrote down a system of *axioms* for Gromov–Witten invariants, which all these definitions satisfy.

One can define Gromov–Witten invariants in either homology or cohomology; because of Poincaré duality there is not much difference. We choose to define them in homology, and also in bordism, as this simplifies issues in §6.3.

Definition 6.9. Let (M, ω) be a compact symplectic manifold of dimension $2n$, let $\beta \in H_2(M; \mathbb{Z})$, and $g, m \geq 0$. Choose an almost complex structure J on M compatible with ω . Set $k = c_1(M) \cdot \beta + (n - 3)(1 - g) + m$. Then Theorem 6.4 and “Theorem” 6.6 yield a compact oriented Kuranishi space without boundary $(\overline{\mathcal{M}}_{g,m}(M, J, \beta), \kappa)$, with virtual dimension $2k$, and strong submersions $\mathbf{ev}_1 \times \cdots \times \mathbf{ev}_m : (\overline{\mathcal{M}}_{g,m}(M, J, \beta), \kappa) \rightarrow M^m$, and $\mathbf{ev}_1 \times \cdots \times \mathbf{ev}_m \times \pi_{g,m} : (\overline{\mathcal{M}}_{g,m}(M, J, \beta), \kappa) \rightarrow M^m \times \overline{\mathcal{M}}_{g,m}$ if $2g + m \geq 3$. “Theorem” 6.8 gives an almost complex structure (J, K) on $(\overline{\mathcal{M}}_{g,m}(M, J, \beta), \kappa)$, for suitably chosen κ . When $2g + m < 3$, define

$$\begin{aligned} GW_{g,m}^{\text{Kb}}(M, \omega, \beta) &\in KB_{2k}(M^m; \mathbb{Z}) \text{ and } GW_{g,m}^{\text{Kb}}(M, \omega, \beta) \in KB_{2k}^{\text{ac}}(M^m; \mathbb{Z}) \text{ by} \\ GW_{g,m}^{\text{Kb}}(M, \omega, \beta) &= [(\overline{\mathcal{M}}_{g,m}(M, J, \beta), \kappa), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_m], \\ GW_{g,m}^{\text{ac}}(M, \omega, \beta) &= [(\overline{\mathcal{M}}_{g,m}(M, J, \beta), \kappa), (J, K), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_m]. \end{aligned}$$

When $2g + m \geq 3$, define

$$\begin{aligned} GW_{g,m}^{\text{Kb}}(M, \omega, \beta) &\in KB_{2k}(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Z}) \text{ and} \\ GW_{g,m}^{\text{ac}}(M, \omega, \beta) &\in KB_{2k}^{\text{ac}}(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Z}) \text{ by} \\ GW_{g,m}^{\text{Kb}}(M, \omega, \beta) &= [(\bar{\mathcal{M}}_{g,m}(M, J, \beta), \kappa), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_m \times \pi_{g,m}], \\ GW_{g,m}^{\text{ac}}(M, \omega, \beta) &= [(\bar{\mathcal{M}}_{g,m}(M, J, \beta), \kappa), (\mathbf{J}, \mathbf{K}), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_m \times \pi_{g,m}]. \end{aligned}$$

We call $GW_{g,m}^{\text{Kb}}, GW_{g,m}^{\text{ac}}(M, \omega, \beta)$ (*almost complex*) *Gromov–Witten bordism invariants*, as they lie in (almost complex) Kuranishi bordism groups.

Define *Kuranishi Gromov–Witten invariants* by $GW_{g,m}^{\text{Kh}}(M, \omega, \beta) = \Pi_{\text{Kb}}^{\text{Kh}}(GW_{g,m}^{\text{Kb}}(M, \omega, \beta))$, and *ordinary Gromov–Witten invariants* $GW_{g,m}^{\text{si}}(M, \omega, \beta) = \Pi_{\text{Kb}}^{\text{si}}(GW_{g,m}^{\text{Kb}}(M, \omega, \beta))$, for $\Pi_{\text{Kb}}^{\text{Kh}}, \Pi_{\text{Kb}}^{\text{si}}$ as in Definition 5.5. Then

$$\begin{aligned} GW_{g,m}^{\text{Kh}}(M, \omega, \beta) &\in \begin{cases} KH_{2k}(M^m; \mathbb{Q}), & 2g + m < 3, \\ KH_{2k}(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Q}), & 2g + m \geq 3, \end{cases} \\ GW_{g,m}^{\text{si}}(M, \omega, \beta) &\in \begin{cases} H_{2k}^{\text{si}}(M^m; \mathbb{Q}), & 2g + m < 3, \\ H_{2k}^{\text{si}}(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Q}), & 2g + m \geq 3. \end{cases} \end{aligned}$$

Note that as $GW_{g,m}^{\text{Kb}}(M, \omega, \beta) = \Pi_{\text{ac}}^{\text{Kb}}(GW_{g,m}^{\text{ac}}(M, \omega, \beta))$, we also have $GW_{g,m}^{\text{Kh}}(M, \omega, \beta) = \Pi_{\text{ac}}^{\text{Kh}}(GW_{g,m}^{\text{ac}}(M, \omega, \beta))$ and $GW_{g,m}^{\text{si}}(M, \omega, \beta) = \Pi_{\text{ac}}^{\text{si}}(GW_{g,m}^{\text{ac}}(M, \omega, \beta))$.

Our next result is analogous to Fukaya and Ono [25, Th. 17.11].

Theorem 6.10. $GW_{g,m}^{\text{Kb}}, GW_{g,m}^{\text{ac}}, GW_{g,m}^{\text{Kh}}, GW_{g,m}^{\text{si}}(M, \omega, \beta)$ depend only on g, m, M, ω, β , and are independent of choices of $J, \kappa, (\mathbf{J}, \mathbf{K})$.

Proof. Suppose $J_0, \kappa_0, (\mathbf{J}_0, \mathbf{K}_0)$ and $J_1, \kappa_1, (\mathbf{J}_1, \mathbf{K}_1)$ are two possible sets of choices of $J, \kappa, (\mathbf{J}, \mathbf{K})$ in defining $GW_{g,m}^{\text{Kb}}, GW_{g,m}^{\text{ac}}(M, \omega, \beta)$. Choose a smooth 1-parameter family of almost complex structures J_t for $t \in [0, 1]$ interpolating between J_0 and J_1 . “Theorems” 6.7 and 6.8 then give a Kuranishi structure $\tilde{\kappa}$ on $\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta)$, strong submersions $\tilde{\mathbf{ev}}_1, \dots, \tilde{\mathbf{ev}}_m, \tilde{\pi}_{g,m}$, and an almost CR structure $(\mathbf{D}, \mathbf{J}, \mathbf{K})$ on $(\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta), \tilde{\kappa})$.

When $2g + m < 3$, applying Definition 5.7(ii) with $W = (\bar{\mathcal{M}}_{g,m}(M, J_t : t \in [0, 1], \beta), \tilde{\kappa})$ and $\mathbf{e} = \tilde{\mathbf{ev}}_1 \times \cdots \times \tilde{\mathbf{ev}}_m$ and using (144) shows that

$$\begin{aligned} &[(\bar{\mathcal{M}}_{g,m}(M, J_1, \beta), \kappa_1) \amalg -(\bar{\mathcal{M}}_{g,m}(M, J_0, \beta), \kappa_0), \\ &\quad \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_m \amalg \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_m] = 0 \quad \text{in } KB^{2k}(M^m; \mathbb{Z}). \end{aligned}$$

Using Definition 5.7(i) and $[-X, \mathbf{f}] = -[X, \mathbf{f}]$ as in (124) gives

$$[(\bar{\mathcal{M}}_{g,m}(M, J_1, \beta), \kappa_1), \prod_{i=1}^m \mathbf{ev}_i] = [(\bar{\mathcal{M}}_{g,m}(M, J_0, \beta), \kappa_0), \prod_{i=1}^m \mathbf{ev}_i],$$

so $GW_{g,m}^{\text{Kb}}(M, \omega, \beta)$ is independent of choices J, κ . The proofs for the other bordism cases are the same, including $\pi_{g,m}, \tilde{\pi}_{g,m}, (\mathbf{J}_i, \mathbf{K}_i)$ or $(\mathbf{D}, \mathbf{J}, \mathbf{K})$ as appropriate. The result for $GW_{g,m}^{\text{Kh}}, GW_{g,m}^{\text{si}}(M, \omega, \beta)$ then follows. $\square$

The claim in Remark 4.11(a) that the construction of the inverse $(\Pi_{\text{si}}^{\text{Kh}})^{-1} : KH_k(Y; \mathbb{Q}) \rightarrow H_k^{\text{si}}(Y; \mathbb{Q})$ in Theorems 4.8 and 4.9 is basically equivalent to Fukaya and Ono's construction of *virtual cycles* for compact oriented Kuranishi spaces without boundary in [25, §6] also implies:

Theorem 6.11. *The invariants $GW_{g,m}^{\text{si}}(M, \omega, \beta)$ of Definition 6.9 agree with the symplectic Gromov–Witten invariants of Fukaya and Ono [25]. In the notation of [25, §17], $GW_{g,m}^{\text{si}}(M, \omega, \beta)$ coincides with $ev_*([CM_{g,m}(M, J, \beta)])$ when $2g + m < 3$, and with $\Pi_*(CM_{g,m}(M, J, \beta))$ when $2g + m \geq 3$.*

Remark 6.12. (a) Since $\mathbf{ev}_1 \times \cdots \times \mathbf{ev}_m$ and $\mathbf{ev}_1 \times \cdots \times \mathbf{ev}_m \times \pi_{g,m}$ in “Theorem” 6.6 are strong submersions, and “Theorem” 6.8 has an analogue for co-almost complex structures, we could equally well have defined Gromov–Witten invariants $GW_{g,m}^{\text{Kcb}}, GW_{g,m}^{\text{ca}}, GW_{g,m}^{\text{Kch}}, GW_{g,m}^{\text{cs}}(M, \omega, \beta)$ in (almost complex) Kuranishi cobordism, Kuranishi cohomology, and compactly-supported cohomology.

In fact, as in Kontsevich and Manin [45], it is best to regard Gromov–Witten invariants as maps $H^*(M; \mathbb{Q})^{\otimes m} \rightarrow H^*(\bar{\mathcal{M}}_{g,m}; \mathbb{Q})$, and so as elements of $H_*(M^m; \mathbb{Q}) \otimes H^*(\bar{\mathcal{M}}_{g,m}; \mathbb{Q})$, mixing homology and cohomology. This suggests that in the *bivariant theories* discussed in §4.8, we should regard Gromov–Witten invariants as elements of $H^*(\pi : M^m \times \bar{\mathcal{M}}_{g,m} \rightarrow \bar{\mathcal{M}}_{g,m}; \mathbb{Q})$.

We have opted for defining Gromov–Witten invariants in bordism and homology, rather than cobordism and cohomology or a mixture of the two, because of our applications in §6.3 to *integrality properties* of Gromov–Witten invariants. Essentially, we will ask the question: does $GW_{g,m}^{\text{si}}(M, \omega, \beta)$ lie in the image of the projection $H_{2k}^{\text{si}}(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Z}) \rightarrow H_{2k}^{\text{si}}(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Q})$, and if not, can we modify $GW_{g,m}^{\text{si}}(M, \omega, \beta)$ so that it does lie in the image?

Since $\bar{\mathcal{M}}_{g,m}$ is an orbifold, working in cobordism and cohomology has two disadvantages compared with bordism and homology. Firstly, as in §4.6, we have not even defined maps $\Pi_{\text{cs}}^{\text{ec}} : H_{\text{cs}}^*(Y; \mathbb{Z}) \rightarrow KH_{\text{ec}}^*(Y; \mathbb{Z})$ for orbifolds Y , nor proved them to be isomorphisms, so we would have a problem going from $KH_{\text{ec}}^*(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Z})$ to $H_{\text{cs}}^*(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Z})$. Secondly, the ‘blow up functor’ $B : KB_*^{\text{ac}}(Y; \mathbb{Z}) \rightarrow KB_*^{\text{eac}}(Y; \mathbb{Z})$ discussed in §6.3 does not seem to have a good cobordism analogue $B : KB_{\text{ca}}^*(Y; \mathbb{Z}) \rightarrow KB_{\text{eca}}^*(Y; \mathbb{Z})$ when Y is an orbifold.

(b) We claim that the whole of symplectic closed Gromov–Witten theory — Gromov–Witten invariants, Quantum Cohomology, and so on — will become simpler and more streamlined if it is rewritten using Kuranishi (co)homology rather than (co)homology; and simpler still if we lift to Kuranishi (co)bordism.

That is, in Kuranishi (co)homology and Kuranishi (co)bordism we do not have to perturb moduli spaces to obtain a virtual cycle; the moduli spaces $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ are their own virtual classes. For Kuranishi (co)homology we must choose (co-)gauge-fixing data for $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$, but this is a much milder process than perturbing moduli spaces; in Kuranishi (co)bordism there are no extra choices to make. So for instance, the proof that Gromov–Witten invariants satisfy the Kontsevich–Manin axioms [45] would be simplified, as one does not have to worry about the effects of perturbing the moduli spaces.

(c) As in §5.7 Kuranishi bordism groups are *huge*, much larger than homology groups. Therefore we expect $GW_{g,m}^{\text{Kb}}, GW_{g,m}^{\text{ac}}(M, \omega, \beta)$ in these groups *to contain more information* than conventional Gromov–Witten invariants. For example, we can apply the operators $\Pi^{\Gamma, \rho}$ of §5.6 to $GW_{g,m}^{\text{Kb}}, GW_{g,m}^{\text{ac}}(M, \omega, \beta)$ and then project to singular homology, to obtain Gromov–Witten type invariants counting Γ -invariant stable maps.

(d) Our Gromov–Witten bordism invariants $GW_{g,m}^{\text{Kb}}, GW_{g,m}^{\text{ac}}(M, \omega, \beta)$ are defined *in groups over $\mathbb{Z}$* , not over $\mathbb{Q}$. Therefore they are good tools for studying *integrality properties* of Gromov–Witten invariants. We discuss this in §6.3.

6.3 Integrality properties of Gromov–Witten invariants

Gromov–Witten invariants lie in *rational* homology $H_{2k}^{\text{si}}(M^m; \mathbb{Q})$ or $H_{2k}^{\text{si}}(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Q})$, rather than integral homology. Essentially this is because if we think of Gromov–Witten invariants as ‘counting’ J -holomorphic maps $w : \Sigma \rightarrow Y$ with the images of the marked points $w(z_i)$ lying on cycles C_i in Y for $i = 1, \dots, m$, then $[\Sigma, \bar{z}, w]$ should be counted with rational weight $|\text{Aut}(\Sigma, \bar{z}, w)|^{-1}$. Thus the reason why Gromov–Witten invariants are defined over $\mathbb{Q}$ rather than $\mathbb{Z}$ is because of points $[\Sigma, \bar{z}, w] \in \bar{\mathcal{M}}_{g,m}(M, J, \beta)$ with nontrivial finite automorphism groups $\text{Aut}(\Sigma, \bar{z}, w) \neq \{1\}$. Equivalently, it is because of *nontrivial orbifold strata* $\bar{\mathcal{M}}_{g,m}(M, J, \beta)^{\Gamma, \rho}$ in $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$, in the sense of §5.6.

To better understand these orbifold strata, we define some notation.

Definition 6.13. We work in the situation of §6.1. Suppose $[\Sigma, \bar{z}] \in \bar{\mathcal{M}}_{g,m}$ and $\rho : \Gamma \rightarrow \text{Aut}(\Sigma, \bar{z})$ is an injective morphism. Define $\Sigma' = \Sigma/\Gamma$, with projection $\pi : \Sigma \rightarrow \Sigma'$. Then there is a natural way to give Σ' the structure of a prestable Riemann surface, with genus $g' \leq g$, such that π is a holomorphic branched cover. Define $z'_i = \pi(z_i)$ for $i = 1, \dots, m$, and $\bar{z}' = (z'_1, \dots, z'_m)$. Then $[\Sigma', \bar{z}'] \in \bar{\mathcal{M}}_{g',m}$. Define $\bar{\mathcal{M}}_{g,m}^{\Gamma}$ to be the moduli space of isomorphism classes $[\Sigma, \bar{z}, \rho]$ of triples $(\Sigma, \bar{z}, \rho)$ as above, and define a smooth map $Q^{\Gamma} : \bar{\mathcal{M}}_{g,m}^{\Gamma} \rightarrow \coprod_{g' \leq g} \bar{\mathcal{M}}_{g',m}$ by $Q^{\Gamma} : [\Sigma, \bar{z}, \rho] \mapsto [\Sigma', \bar{z}']$.

Now consider a Kuranishi space $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$, and one of its orbifold strata $\bar{\mathcal{M}}_{g,m}(M, J, \beta)^{\Gamma, \rho}$. Points of $\bar{\mathcal{M}}_{g,m}(M, J, \beta)^{\Gamma, \rho}$ consist of points $[\Sigma, \bar{z}, w] \in \bar{\mathcal{M}}_{g,m}(M, J, \beta)$ with an injective group morphism $\rho : \Gamma \rightarrow \text{Aut}(\Sigma, \bar{z}, w)$. Define $\Sigma' = \Sigma/\Gamma$ and $\bar{z}'$ as above. Then $w : \Sigma \rightarrow M$ factors as $w = w' \circ \pi$ for some J -holomorphic $w' : \Sigma' \rightarrow M'$. Write $\beta' = w_*([\Sigma']) \in H_2(M; \mathbb{Z})$. If the action of Γ on Σ is locally effective then $\beta' = |\Gamma|^{-1}\beta$, but in general β' is not simply related to β , although we do have $[\omega] \cdot \beta' \leq [\omega] \cdot \beta$. Thus we can form $[\Sigma', \bar{z}', w'] \in \bar{\mathcal{M}}_{g',m}(M, J, \beta')$. Define a map $Q^{\Gamma, \rho} : \bar{\mathcal{M}}_{g,m}(M, J, \beta)^{\Gamma, \rho} \rightarrow \coprod_{g', \beta'} \bar{\mathcal{M}}_{g',m}(M, J, \beta')$ by $Q^{\Gamma, \rho} : [\Sigma, \bar{z}, w, \rho] \mapsto [\Sigma', \bar{z}', w']$.

We now make the following, slightly tentative claim. It may need modification over points $(\Sigma, \bar{z}, w), \rho$ where $\text{Aut}(\Sigma, \bar{z}, w)$ is finite but $\text{Aut}(\Sigma, \bar{z})$ is not, so we have to pass to the *stabilizations* of $(\Sigma, \bar{z}), (\Sigma', \bar{z}')$, as in Definition 6.1.

Claim 6.14. *In Definition 6.13, $Q^{\Gamma, \rho}$ lifts to a strongly smooth map $\mathbf{Q}^{\Gamma, \rho} : \bar{\mathcal{M}}_{g,m}(M, J, \beta)^{\Gamma, \rho} \rightarrow \coprod_{g', \beta'} \bar{\mathcal{M}}_{g',m}(M, J, \beta')$. Furthermore, $\bar{\mathcal{M}}_{g,m}(M, J, \beta)^{\Gamma, \rho}$*

is naturally isomorphic as a Kuranishi space to an open and closed set in the fibre product of Kuranishi spaces

$$\bar{\mathcal{M}}_{g,m}^\Gamma \times_{Q, \coprod_{g' \leq g} \bar{\mathcal{M}}_{g',m}, \coprod_{g',\beta'} \pi_{g',m}} \coprod_{g',\beta'} \bar{\mathcal{M}}_{g',m}(M, J, \beta'),$$

such that $\mathbf{Q}^{\Gamma,\rho}$ is identified with projection to the second factor in the fibre product. Hence the orbifold strata $\bar{\mathcal{M}}_{g,m}(M, J, \beta)^{\Gamma,\rho}$ of $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$ can be completely described, up to isomorphism as Kuranishi spaces, in terms of other moduli spaces $\bar{\mathcal{M}}_{g',m}(M, J, \beta')$ with $g' \leq g$ and $[\omega] \cdot \beta' \leq [\omega] \cdot \beta$.

Assuming Claim 6.14, we see that applying the operators $\Pi^{\Gamma,\rho}$ of §5.6 to the Gromov–Witten bordism invariants $GW_{g,m}^{\text{Kb}}, GW_{g,m}^{\text{ac}}(M, \omega, \beta)$ should give answers which can be expressed in terms of other Gromov–Witten bordism invariants $GW_{g',m}^{\text{Kb}}, GW_{g',m}^{\text{ac}}(M, \omega, \beta')$. Projecting to homology, we expect that $\Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma,\rho}(GW_{g,m}^{\text{Kb}}(M, \omega, \beta))$ and $\Pi_{\text{ac}}^{\text{si}} \circ \Pi^{\Gamma,\rho}(GW_{g,m}^{\text{ac}}(M, \omega, \beta))$ should be expressed in terms of other Gromov–Witten invariants $GW_{g',m}^{\text{si}}(M, \omega, \beta')$.

These ideas suggest an approach to the question of whether one can define Gromov–Witten type invariants in *integral* homology. Since Gromov–Witten invariants fail to lie in integral homology because of nontrivial orbifold strata $\bar{\mathcal{M}}_{g,m}(M, J, \beta)^{\Gamma,\rho}$, and (assuming Claim 6.14) these orbifold strata can be described in terms of $\bar{\mathcal{M}}_{g',m}(M, J, \beta')$ with $g' \leq g$ and $[\omega] \cdot \beta' \leq [\omega] \cdot \beta$, then perhaps one could modify the Gromov–Witten invariant $GW_{g,m}^{\text{si}}(M, \omega, \beta)$ by adding on a series of ‘corrections’ involving ‘lower’ Gromov–Witten invariants $GW_{g',m}^{\text{si}}(M, \omega, \beta')$, to end up with a Gromov–Witten type invariant which lies in the image of integral homology in rational homology.

Much progress has been made on these issues, which we briefly discuss.

Semi-positive symplectic manifolds and genus zero invariants. A homology class $\beta \in H_2(M; \mathbb{Z})$ is called *spherical* if it lies in the image of $\pi_2(M) \rightarrow H_2(M; \mathbb{Z})$. A symplectic $2n$ -manifold (M, ω) is called *semi-positive* if for all spherical $\beta \in H_2(M; \mathbb{Z})$, we never have $[\omega] \cdot \beta > 0$ and $3 - n \leq c_1(M) \cdot \beta < 0$. This applies in particular whenever $n \leq 3$. Suppose (M, ω) is semi-positive, and $\beta \in H_2(M; \mathbb{Z})$ may not be written as $\beta = l\gamma$ for spherical $\gamma \in H_2(M; \mathbb{Z})$ and $l \geq 2$ an integer. Then the genus zero Gromov–Witten invariants $GW_{0,m}^{\text{si}}(M, \omega, \beta)$ can be defined in *integral* homology $H_{2k}^{\text{si}}(M^m; \mathbb{Z})$ or $H_{2k}^{\text{si}}(M^m \times \bar{\mathcal{M}}_{0,m}; \mathbb{Z})$.

This is proved in detail in McDuff and Salamon [56, §7]. In our notation, the reason is that if J is a generic almost complex structure, then one can show that all nonempty orbifold strata $\bar{\mathcal{M}}_{0,m}(M, J, \beta)^{\Gamma,\rho}$ of $\bar{\mathcal{M}}_{0,m}(M, J, \beta)$ with $\Gamma \neq \{1\}$ satisfy $\text{vdim } \bar{\mathcal{M}}_{0,m}(M, J, \beta)^{\Gamma,\rho} \leq \text{vdim } \bar{\mathcal{M}}_{0,m}(M, J, \beta) - 2$. In a very similar way to our use of *effective* Kuranishi spaces to define a virtual cycle map $\Pi_{\text{eb}}^{\text{si}} : KB_*(Y; \mathbb{Z}) \rightarrow H_*^{\text{si}}(Y; \mathbb{Z})$, this condition on the orbifold strata of $\bar{\mathcal{M}}_{0,m}(M, J, \beta)$ is enough to ensure that its virtual cycle can be defined over $\mathbb{Z}$ rather than $\mathbb{Q}$. In fact, in Remark 5.20 we showed that if X is an oriented, effective Kuranishi space then $\text{vdim } X^{\Gamma,\rho} \leq \text{vdim } X - 2$ for all nonempty orbifold strata $X^{\Gamma,\rho}$ of X with $\Gamma \neq \{1\}$.

Gopakumar–Vafa invariants and their Integrality Conjecture. Let (M, ω) be a compact symplectic 6-manifold with $c_1(M) = 0$, that is, a *symplec-*

tic Calabi–Yau 3-fold. Take the number m of marked points to be zero. Then (143) gives $\text{vdim } \overline{\mathcal{M}}_{g,0}(M, J, \beta) = 0$ for all g, β . Hence the Gromov–Witten invariants $GW_{g,0}^{\text{si}}(M, \omega, \beta)$ lie in $H_0^{\text{si}}(M^m; \mathbb{Q}) \cong \mathbb{Q}$ or $H_0^{\text{si}}(M^m \times \overline{\mathcal{M}}_{g,0}; \mathbb{Q}) \cong \mathbb{Q}$. So we regard the $GW_{g,0}^{\text{si}}(M, \omega, \beta)$ as *rational numbers* for all g, β .

Using physical reasoning about counting BPS states, the String Theorists Gopakumar and Vafa [28, 29] conjectured the existence of *Gopakumar–Vafa invariants* $GV_g(M, \omega, \beta) \in \mathbb{Z}$ for Calabi–Yau 3-folds M , which morally speaking ‘count’ *embedded* J -holomorphic curves of genus g in class β in M . By expressing any J -holomorphic curve in M as the branched cover of an embedded curve, they conjecturally wrote Gromov–Witten in terms of Gopakumar–Vafa invariants, and vice versa, by the equation in formal power series

$$\sum_{g,\beta} GW_{g,0}^{\text{si}}(M, \omega, \beta) t^{2g-2} q^\beta = \sum_{k>0, g,\beta} GV_g(M, \omega, \beta) \frac{1}{k} (2 \sin(kt/2))^{2g-2} q^{k\beta}. \quad (148)$$

The *Gopakumar–Vafa Integrality Conjecture* says Gromov–Witten invariants of Calabi–Yau 3-folds satisfy (148) for some integers $GV_g(M, \omega, \beta)$. Gopakumar and Vafa also conjecture that $GV_g(M, \omega, \beta) = 0$ for all fixed β and $g \gg 0$.

There are two approaches to proving this conjecture. The first would be to define some curve-counting invariants $GV_g(M, \omega, \beta)$ which are automatically integers, and then prove that they satisfy (148). Pandharipande and Thomas [62, 63] have taken an important step in this direction, in the context of algebraic rather than symplectic geometry, by defining integer-valued invariants which count ‘stable pairs’ (F, s) of a coherent sheaf F supported on a curve in M and a section $s \in H^0(F)$. Proving equivalence of Pandharipande–Thomas invariants with Gromov–Witten invariants, the analogue of (148), still seems difficult.

The second is to regard (148) as a *definition* of numbers $GV_g(M, \omega, \beta) \in \mathbb{Q}$, and then to prove that these $GV_g(M, \omega, \beta)$ actually lie in $\mathbb{Z}$. We will sketch a method of attack following this approach below.

More general integrality conjectures. Pandharipande [59, 60] extends Gopakumar–Vafa invariants and their integrality conjecture to all smooth projective complex algebraic 3-folds. Klemm and Pandharipande [43] and Pandharipande and Zinger [64] study integrality of Gromov–Witten invariants of Calabi–Yau m -folds for $m > 3$. In these cases, for dimension reasons, only genus $g = 0$ and $g = 1$ can yield nonzero invariants. An integrality conjecture for $g = 0$ and all $m > 3$ is given in [43], and conjectures for $g = 1$, $m = 4$ and $g = 1$, $m = 5$ are given in [43] and [64], respectively.

The rest of the section outlines an approach to proving integrality results for Gromov–Witten invariants, including the Gopakumar–Vafa Integrality Conjecture, and the conjectures of Pandharipande et al. [43, 59, 60, 64]. We stress that this approach is still very incomplete, and we are not yet claiming a proof of any of these conjectures. The author hopes to fill in some of the gaps in the sketch in [41]. We divide our method into four steps. Steps 1–3 are very general, and work in all dimensions.

6.3.1 Step 1: a ‘blow-up functor’ $B : KB_*^{\text{ac}}(Y; R) \rightarrow KB_*^{\text{eac}}(Y; R)$

Let X be a compact Kuranishi space without boundary, with an almost complex structure $(\mathbf{J}, \mathbf{K})$. We would like to find a functorial procedure to modify $X, (\mathbf{J}, \mathbf{K})$ to yield an *effective* compact Kuranishi space $\tilde{X}$ without boundary, with an almost complex structure $(\tilde{\mathbf{J}}, \tilde{\mathbf{K}})$, and with a strongly smooth map $\pi_X : \tilde{X} \rightarrow X$. Now X fails to be effective if its *orbifold strata* $X^{\Gamma, \rho}$ have the wrong behaviour, in particular, if ρ is not the isomorphism class of an effective Γ -representation and $X^{\Gamma, \rho} \neq \emptyset$ then X is not effective. Thus, to make X effective we must modify it around its orbifold strata $X^{\Gamma, \rho}$.

Suppose $|\Gamma|$ is largest such that X has an orbifold stratum $X^{\Gamma, \rho}$ along which X is not effective. Then $X^{\Gamma, \rho}$ is in some sense ‘nonsingular’ in X . In this case the author has a procedure for ‘blowing up’ an almost complex Kuranishi space $X, (\mathbf{J}, \mathbf{K})$ at $X^{\Gamma, \rho}$ with respect to $(\mathbf{J}, \mathbf{K})$, to get a new almost complex Kuranishi space $\hat{X}, (\hat{\mathbf{J}}, \hat{\mathbf{K}})$ with strongly smooth $\pi_X : \hat{X} \rightarrow X$. We aim to do the blow up in such a way that $\hat{X}$ is ‘closer to being effective’ than X .

If (V_p, E_p, s_p, ψ_p) is a sufficiently small Kuranishi neighbourhood in the germ at $p \in X$ with (J_p, K_p) representing K_p then we define a Kuranishi neighbourhood $(\hat{V}_p, \hat{E}_p, \hat{s}_p, \hat{\psi}_p)$ on $\hat{X}$, with $(\hat{J}_p, \hat{K}_p)$ representing $(\hat{\mathbf{J}}, \hat{\mathbf{K}})$, where $(\hat{V}_p, \hat{J}_p)$ is the blow up of (V_p, J_p) along $V_p^{\Gamma, \rho}$, in the usual sense of (almost) complex geometry. (Note a potential confusion here: we are *not* treating the orbifold strata of V_p as singularities, and resolving them to get a manifold $\hat{V}_p$. In fact $\hat{V}_p$ is an orbifold, and its orbifold strata may be no simpler than those of V_p .) The crucial part of the construction is to define $\hat{E}_p, \hat{s}_p$ in such a way that these blow ups are compatible with $(\phi_{pq}, \hat{\phi}_{pq})$ in the germ of coordinate changes on X .

By applying this blow up procedure iteratively by reverse induction on $|\Gamma|$ to modify all orbifold strata of X to make them effective, we make the:

Claim 6.15. *There is a procedure to modify $X, (\mathbf{J}, \mathbf{K})$ to get $\tilde{X}, (\tilde{\mathbf{J}}, \tilde{\mathbf{K}}), \pi_X$ as above, with $\tilde{X}$ effective. This induces a functor $B : KB_{2k}^{\text{ac}}(Y; R) \rightarrow KB_{2k}^{\text{eac}}(Y; R)$ mapping $B : [X, (\mathbf{J}, \mathbf{K}), \mathbf{f}] \mapsto [\tilde{X}, (\tilde{\mathbf{J}}, \tilde{\mathbf{K}}), \mathbf{f} \circ \pi_X]$, such that if X has trivial stabilizers then $B([X, (\mathbf{J}, \mathbf{K}), \mathbf{f}]) = [X, (\mathbf{J}, \mathbf{K}), \mathbf{f}]$.*

Applying the functor B to the invariants $GW_{g,m}^{\text{ac}}(M, \omega, \beta)$ of §6.2 and composing with $\Pi_{\text{eac}}^{\text{si}}$ yields invariants $\Pi_{\text{eac}}^{\text{si}} \circ B(GW_{g,m}^{\text{ac}}(M, \omega, \beta))$ in *integral* homology $H_{2k}^{\text{si}}(M^m; \mathbb{Z})$ or $H_{2k}^{\text{si}}(M^m \times \bar{M}_{g,m}; \mathbb{Z})$, since $\Pi_{\text{eac}}^{\text{si}}$ maps $KB_*^{\text{eac}}(Y; \mathbb{Z}) \rightarrow H_*^{\text{si}}(Y; \mathbb{Z})$, although $\Pi_{\text{ac}}^{\text{si}}$ maps $KB_*^{\text{ac}}(Y; \mathbb{Z}) \rightarrow H_*^{\text{si}}(Y; \mathbb{Q})$. So we can think of the $\Pi_{\text{eac}}^{\text{si}} \circ B(GW_{g,m}^{\text{ac}}(M, \omega, \beta))$ as Gopakumar–Vafa type invariants, curve counting invariants that are automatically integer-valued.

This is similar to an idea sketched by Fukaya and Ono [26]. Translated into our notation, roughly speaking, Fukaya and Ono claim that given a compact Kuranishi space X without boundary and an almost complex structure $(\mathbf{J}, \mathbf{K})$, then one should be able to carry out their virtual cycle construction for X using single-valued sections rather than multisections, by using $(\mathbf{J}, \mathbf{K})$ to determine a canonical form for the perturbation near each orbifold stratum $X^{\Gamma, \rho}$. As the sections are single-valued, the virtual cycle lies in homology over $\mathbb{Z}$.

6.3.2 Step 2: an expression for $\Pi_{\text{eac}}^{\text{Kb}} \circ B : KB_*^{\text{ac}}(Y; \mathbb{Z}) \rightarrow KB_*(Y; \mathbb{Z})$

Let $[X, (\mathbf{J}, \mathbf{K}), \mathbf{f}] \in KB_{2k}^{\text{ac}}(Y; \mathbb{Z})$. If as in §6.3.1 we blow up $X, (\mathbf{J}, \mathbf{K})$ along $X^{\Gamma, \rho}$ to get $\hat{X}, (\hat{\mathbf{J}}, \hat{\mathbf{K}}), \pi_X$, we can form $[\hat{X}, (\hat{\mathbf{J}}, \hat{\mathbf{K}}), \mathbf{f} \circ \pi_X] \in KB_{2l}^{\text{ac}}(Y; \mathbb{Z})$. There is a notion of virtual vector bundle over a Kuranishi space due to Fukaya and Ono [25, Def. 5.10], called a *bundle system*. The *virtual normal bundle* $\nu_{X^{\Gamma, \rho}}$ of $X^{\Gamma, \rho}$ in X is a complex bundle system. We expect the difference $[\hat{X}, (\hat{\mathbf{J}}, \hat{\mathbf{K}}), \mathbf{f} \circ \pi_X] - [X, (\mathbf{J}, \mathbf{K}), \mathbf{f}]$ should depend only on $X^{\Gamma, \rho}$ and $\nu_{X^{\Gamma, \rho}}$.

This is more likely to work if we project to Kuranishi bordism $KB_{2k}(Y; \mathbb{Z})$, as constructing the relevant bordism W between Kuranishi spaces will be easier if W need not carry an almost CR structure. Thus we expect that

$$\Pi_{\text{ac}}^{\text{Kb}}([\hat{X}, (\hat{\mathbf{J}}, \hat{\mathbf{K}}), \mathbf{f} \circ \pi_X] - [X, (\mathbf{J}, \mathbf{K}), \mathbf{f}]) = [C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}), \mathbf{f} \circ \iota^{\Gamma, \rho} \circ \gamma^{\Gamma, \rho}],$$

where $C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho})$ is some compact Kuranishi space without boundary constructed from $X^{\Gamma, \rho}$ and $\nu^{\Gamma, \rho}$ in some functorial way, with $\text{vdim } C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}) = \text{vdim } X$, and $\gamma^{\Gamma, \rho} : C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}) \rightarrow X^{\Gamma, \rho}$ is a cooriented strong submersion. Define $\Pi^{\Gamma, \rho, C^{\Gamma, \rho}} : KB_{2k}^{\text{ac}}(Y; \mathbb{Z}) \rightarrow KB_{2k}(Y; \mathbb{Z})$ by

$$\Pi^{\Gamma, \rho, C^{\Gamma, \rho}} : [X, (\mathbf{J}, \mathbf{K}), \mathbf{f}] \mapsto [C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}), \mathbf{f} \circ \iota^{\Gamma, \rho} \circ \gamma^{\Gamma, \rho}].$$

The functor B is the result of composing many such blowups along $X^{\Gamma, \rho}$ for different Γ, ρ . This suggests the:

Claim 6.16. *In operators $KB_{2l}^{\text{ac}}(Y; \mathbb{Z}) \rightarrow KB_{2l}(Y; \mathbb{Z})$ we have*

$$\Pi_{\text{eac}}^{\text{Kb}} \circ B = \Pi_{\text{ac}}^{\text{Kb}} + \sum_{\Gamma, \rho} \Pi^{\Gamma, \rho, C^{\Gamma, \rho}}, \quad (149)$$

where the sum is over isomorphism classes of finite groups Γ with $\Gamma \neq \{1\}$ and isomorphism classes of virtual nontrivial Γ -representations ρ , and each $\Pi^{\Gamma, \rho, C^{\Gamma, \rho}}$ is an operator of the form above.

6.3.3 Step 3: projecting to singular homology

Apply equation (149) to the Gromov–Witten type invariant $GW_{g,m}^{\text{ac}}(M, \omega, \beta)$, and compose with projection $\Pi_{\text{Kb}}^{\text{si}}$ to rational singular homology. This yields

$$\begin{aligned} \Pi_{\text{eac}}^{\text{si}} \circ B(GW_{g,m}^{\text{ac}}(M, \omega, \beta)) &= GW_{g,m}^{\text{si}}(M, \omega, \beta) + \\ &\quad \sum_{\Gamma, \rho} \Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma, \rho, C^{\Gamma, \rho}}(GW_{g,m}^{\text{ac}}(M, \omega, \beta)). \end{aligned} \quad (150)$$

Here the left hand side of (150) lies in the image of integral homology in rational homology, $\pi_* : H_{2k}^{\text{si}}(M^m; \mathbb{Z}) \rightarrow H_{2k}^{\text{si}}(M^m; \mathbb{Q})$ or $\pi_* : H_{2k}^{\text{si}}(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Z}) \rightarrow H_{2k}^{\text{si}}(M^m \times \bar{\mathcal{M}}_{g,m}; \mathbb{Q})$. The first term on the right hand side is an ordinary Gromov–Witten invariant, and the remaining terms are corrections from the nontrivial orbifold strata $\bar{\mathcal{M}}_{g,m}(M, J, \beta)^{\Gamma, \rho}$ of $\bar{\mathcal{M}}_{g,m}(M, J, \beta)$.

Now Claim 6.14 says that we can write the $\bar{\mathcal{M}}_{g,m}(M, J, \beta)^{\Gamma, \rho}$ in terms of other ‘lower’ moduli spaces $\bar{\mathcal{M}}_{g',m}(M, J, \beta')$. Thus it seems reasonable to hope

that we can write the corrections $\Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma, \rho, C^{\Gamma, \rho}}(GW_{g, m}^{\text{ac}}(M, \omega, \beta))$ in terms of other ‘lower’ ordinary Gromov–Witten invariants $GW_{g', m}^{\text{si}}(M, \omega, \beta')$. However, there is a problem. The $\Pi^{\Gamma, \rho, C^{\Gamma, \rho}}(GW_{g, m}^{\text{ac}}(M, \omega, \beta))$ depend not only on the orbifold strata $\bar{\mathcal{M}}_{g, m}(M, J, \beta)^{\Gamma, \rho}$, but also on its *virtual normal bundle* $\nu^{\Gamma, \rho}$ in $\bar{\mathcal{M}}_{g, m}(M, J, \beta)$. The author does *not* expect $\nu^{\Gamma, \rho}$ to depend only on $\bar{\mathcal{M}}_{g', m}(M, J, \beta')$, it is extra information. But we claim that if $\dim \rho \geq 0$ then $\Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma, \rho, C^{\Gamma, \rho}}(GW_{g, m}^{\text{ac}}(M, \omega, \beta))$ does not depend on $\nu^{\Gamma, \rho}$.

Claim 6.17. *In operators $KB_{2k}^{\text{ac}}(Y; \mathbb{Z}) \rightarrow H_{2k}^{\text{si}}(Y; \mathbb{Q})$ we have $\Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma, \rho, C^{\Gamma, \rho}} = 0$ if $\dim \rho > 0$, and $\Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma, \rho, C^{\Gamma, \rho}} = q^{\Gamma, \rho} \Pi_{\text{ac}}^{\text{si}} \circ \Pi_{\text{ac}}^{\Gamma, \rho}$ if $\dim \rho = 0$, where $q^{\Gamma, \rho} \in \mathbb{Q}$ and $\Pi_{\text{ac}}^{\Gamma, \rho} : KB_{2k}^{\text{ac}}(Y; \mathbb{Z}) \rightarrow KB_{2k}^{\text{ac}}(Y; \mathbb{Z})$ is as in §5.6.*

To justify this, note that $\Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma, \rho, C^{\Gamma, \rho}}([X, (\mathbf{J}, \mathbf{K}), \mathbf{f}])$ is represented by a virtual cycle $VC(C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}))$ for $(C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}), \mathbf{f} \circ \iota^{\Gamma, \rho} \circ \gamma^{\Gamma, \rho})$, and $\mathbf{f} \circ \iota^{\Gamma, \rho} \circ \gamma^{\Gamma, \rho}$ factorizes through $\gamma^{\Gamma, \rho} : C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}) \rightarrow X^{\Gamma, \rho}$, where $\text{vdim } C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}) = \text{vdim } X$ and $\text{vdim } X^{\Gamma, \rho} = \text{vdim } X - \dim \rho$.

If $VC(X^{\Gamma, \rho})$ is a virtual cycle for $(X^{\Gamma, \rho}, \mathbf{f} \circ \iota^{\Gamma, \rho})$, then we can choose $VC(C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}))$ to be supported in an arbitrarily small open neighbourhood of the support of $VC(X^{\Gamma, \rho})$. If $\dim \rho > 0$ then $\dim VC(C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho})) > \dim VC(X^{\Gamma, \rho})$, so $[VC(C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}))] = 0$ in $H_{2k}^{\text{si}}(Y; \mathbb{Q})$ for dimensional reasons. If $\dim \rho = 0$ then $[VC(C^{\Gamma, \rho}(X^{\Gamma, \rho}, \nu^{\Gamma, \rho}))] = q^{\Gamma, \rho} [VC(X^{\Gamma, \rho})]$, where $q^{\Gamma, \rho} \in \mathbb{Q}$ is the ‘multiplicity’ of $\gamma^{\Gamma, \rho}$.

Applying Claim 6.17 to (150) yields

$$\begin{aligned} \Pi_{\text{eac}}^{\text{si}} \circ B(GW_{g, m}^{\text{ac}}(M, \omega, \beta)) &= GW_{g, m}^{\text{si}}(M, \omega, \beta) + \\ &\sum_{\Gamma, \rho: \dim \rho = 0} q^{\Gamma, \rho} \Pi_{\text{ac}}^{\text{si}} \circ \Pi_{\text{ac}}^{\Gamma, \rho}(GW_{g, m}^{\text{ac}}(M, \omega, \beta)) + \\ &\sum_{\Gamma, \rho: \dim \rho < 0} \Pi_{\text{Kb}}^{\text{si}} \circ \Pi^{\Gamma, \rho, C^{\Gamma, \rho}}(GW_{g, m}^{\text{ac}}(M, \omega, \beta)). \end{aligned} \quad (151)$$

When $\dim \rho = 0$, the terms $\Pi_{\text{ac}}^{\text{si}} \circ \Pi_{\text{ac}}^{\Gamma, \rho}(GW_{g, m}^{\text{ac}}(M, \omega, \beta))$ on the second line of (151) no longer depend on the virtual normal bundle $\nu^{\Gamma, \rho}$, so by Claim 6.14, we expect we can write them solely in terms of ‘lower’ Gromov–Witten invariants $GW_{g', m}^{\text{si}}(M, \omega, \beta')$.

The terms on the third line of (151), when $\dim \rho < 0$, come from orbifold strata $\bar{\mathcal{M}}_{g, m}(M, J, \beta)^{\Gamma, \rho}$ with $\text{vdim } \bar{\mathcal{M}}_{g, m}(M, J, \beta)^{\Gamma, \rho} > \text{vdim } \bar{\mathcal{M}}_{g, m}(M, J, \beta)$. The author expects that these terms really do depend on $\nu^{\Gamma, \rho}$, and so *cannot* be written solely in terms of $GW_{g', m}^{\text{si}}(M, \omega, \beta')$, except in special circumstances. In effect, these terms use ‘characteristic classes’ of $\nu^{\Gamma, \rho}$ — some kind of analogue of Chern classes, perhaps — to lower the dimension by $-\dim \rho$.

6.3.4 Step 4: the Gopakumar–Vafa Integrality Conjecture

We now specialize to the case when (M, ω) is a symplectic Calabi–Yau 3-fold, that is, a compact symplectic 6-manifold with $c_1(M) = 0$. Let J be a *generic* almost complex structure on M compatible with ω . Using genericness of J

and $\text{vdim } \overline{\mathcal{M}}_{g,0}(M, J, \beta) = 0$ for all g, β , one can show that for all $[\Sigma, \tilde{z}, w] \in \overline{\mathcal{M}}_{g,m}(M, J, \beta)$, if $\beta = 0$ then $w \equiv p$ for some $p \in M$, and if $\beta \neq 0$ then w factors uniquely as $w = \tilde{w} \circ \pi$, where $\tilde{\Sigma}$ is a *nonsingular* Riemann surface of genus $\tilde{g} \leq g$, and $\tilde{w} : \tilde{\Sigma} \rightarrow M$ is a J -holomorphic *embedding*, and $\pi : \Sigma \rightarrow \tilde{\Sigma}$ is surjective and holomorphic. Furthermore, distinct embedded nonsingular J -holomorphic curves do not intersect in M .

Setting $m = 0$, and regarding the $GW_{g,0}^{\text{si}}(M, \omega, \beta)$ as *rational numbers* for all g, β , we expect (151) to reduce to an equation of the form

$$\Pi_{\text{eac}}^{\text{si}} \circ B(GW_{g,0}^{\text{ac}}(M, \omega, \beta)) = \sum_{0 \leq g' \leq g, k \geq 1: k|\beta} c^{g',g,k} GW_{g',0}^{\text{si}}(M, \omega, \beta/k), \quad (152)$$

where $c^{g',g,k} \in \mathbb{Q}$ are some universal family of rational numbers with $c^{g,g,1} = 1$, and (152) is an equation in $\mathbb{Q}$, with the left hand side in $\mathbb{Z}$.

Now the Gopakumar–Vafa equation (148) may be rewritten in the form

$$GV_g(M, \omega, \beta) = \sum_{0 \leq g' \leq g, k \geq 1: k|\beta} d^{g',g,k} GW_{g',0}^{\text{si}}(M, \omega, \beta/k), \quad (153)$$

where $d^{g',g,k} \in \mathbb{Q}$ are some universal family of explicitly computable rational numbers with $d^{g,g,1} = 1$, and the Gopakumar–Vafa Integrality Conjecture says that the left hand side of (153) lies in $\mathbb{Z}$ for all g, β .

We do not expect it will be feasible to compute the coefficients $c^{g',g,k}$, as they are the sum over all Γ, ρ of many complicated contributions. Nor do we claim that $c^{g',g,k} = d^{g',g,k}$. However, we do hope to prove that the integrality properties of the family of $GW_{g,0}^{\text{si}}(M, \omega, \beta)$ for all g, β implied by (152) are equivalent to the integrality properties implied by (153), and this will be enough to prove the Gopakumar–Vafa Integrality Conjecture.

By induction on g', g, k using the fact that $c^{g,g,1} = d^{g,g,1} = 1$, we can show there exist unique $e^{g',g,k}, f^{g',g,k} \in \mathbb{Q}$ for $0 \leq g' \leq g$ and $k \geq 1$, with $e^{g,g,1} = f^{g,g,1} = 1$, satisfying

$$c^{g'',g,k''} = \sum_{g',k,k': g'' \leq g', k'' = kk'} e^{g',g,k} d^{g'',g',k} \text{ and } d^{g'',g,k''} = \sum_{g',k,k': g'' \leq g', k'' = kk'} f^{g',g,k} c^{g'',g',k} \quad (154)$$

for all $0 \leq g'' \leq g$ and $k'' \geq 1$. Combining (152)–(154) yields

$$\Pi_{\text{eac}}^{\text{si}} \circ B(GW_{g,0}^{\text{ac}}(M, \omega, \beta)) = \sum_{0 \leq g' \leq g, k \geq 1: k|\beta} e^{g',g,k} GV_{g'}(M, \omega, \beta/k), \quad (155)$$

$$GV_g(M, \omega, \beta) = \sum_{0 \leq g' \leq g, k \geq 1: k|\beta} f^{g',g,k} \Pi_{\text{eac}}^{\text{si}} \circ B(GW_{g',0}^{\text{ac}}(M, \omega, \beta/k)). \quad (156)$$

Hence, if all $e^{g',g,k}$ lie in $\mathbb{Z}$ then the integrality of (153) implies that of (152), and if all $f^{g',g,k}$ lie in $\mathbb{Z}$ then the integrality of (152) implies that of (153), and if all $e^{g',g,k}, f^{g',g,k}$ lie in $\mathbb{Z}$ then the integrality of (152) and (153) is equivalent.

The Gopakumar–Vafa Integrality Conjecture has been verified in many examples, for instance, for local curves by Bryan and Pandharipande [9]. In each

of these examples, it has already been proved that the right hand side of (153) is an integer, and if we can carry through our programme above, then we will have shown that the right hand side of (152) is an integer. The author hopes that by applying this to a large enough family of examples, one will be able to prove by an induction on g, g', k that all $e^{g', g, k}, f^{g', g, k}$ lie in $\mathbb{Z}$, so that integrality of (152) implies the Gopakumar–Vafa Integrality Conjecture.

6.4 Moduli spaces of stable holomorphic discs

In §6.4–§6.7 we discuss J -holomorphic curves in a symplectic manifold (M, ω) with boundary in a Lagrangian submanifold L in M . As our aim is just to give an idea of possible applications in this area, not to develop a complete theory, we restrict our attention to *stable holomorphic discs*, rather than curves of higher genus, and we consider only *boundary marked points*, not interior marked points. The generalization of this section to higher genus curves and both kinds of marked points will be explained in [39]. Most of this section follows Fukaya et al. [24, §2], see also Katz and Liu [42, §3] and Liu [51, §2].

Definition 6.18. A *bordered Riemann surface* Σ is a compact connected complex 1-manifold with nonempty boundary $\partial\Sigma$. Given such a Σ we can construct the *double* $\tilde{\Sigma} = \Sigma \cup_{\partial\Sigma} \bar{\Sigma}$, that is, the union of Σ and its complex conjugate $\bar{\Sigma}$, glued along their common boundary $\partial\Sigma$. Then $\tilde{\Sigma}$ is a Riemann surface without boundary, and there is an antiholomorphic involution $\sigma : \tilde{\Sigma} \rightarrow \tilde{\Sigma}$ exchanging Σ and $\bar{\Sigma}$, such that the fixed points of σ are $\partial\Sigma$, and $\tilde{\Sigma}$ is the disjoint union $\Sigma^\circ \amalg \partial\Sigma \amalg \sigma(\Sigma^\circ)$, where Σ° is the interior of Σ .

A *prestable* or *nodal bordered Riemann surface* Σ is a compact connected singular complex 1-manifold with nonempty boundary $\partial\Sigma$, whose only singularities are nodes. Again, Σ has a *double* $\tilde{\Sigma} = \Sigma \cup_{\partial\Sigma} \bar{\Sigma}$, with an antiholomorphic involution $\sigma : \tilde{\Sigma} \rightarrow \tilde{\Sigma}$. We allow one interior and two boundary kinds of nodes:

- (a) Σ has a node z in Σ° locally modelled on $(0, 0)$ in $\{(x, y) \in \mathbb{C}^2 : xy = 0\}$.

Such a node can occur as the limit of a family of nonsingular bordered Riemann surfaces Σ_ϵ modelled on $\{(x, y) \in \mathbb{C}^2 : xy = \epsilon\}$, for $\epsilon \in \mathbb{C} \setminus \{0\}$.

- (b) $\tilde{\Sigma}$ has a node z in $\partial\Sigma$ locally modelled on $(0, 0)$ in $\{(x, y) \in \mathbb{C}^2 : xy = 0\}$, where σ acts by $(x, y) \mapsto (\bar{x}, \bar{y})$. Then $\partial\Sigma$ is locally modelled on $\{(x, y) \in \mathbb{R}^2 : xy = 0\}$, the fixed point set of σ , and we take Σ to be locally modelled on $\{(x, y) \in \mathbb{C}^2 : xy = 0, \operatorname{Im} x \geq 0, \operatorname{Im} y \geq 0\}$.

Such a node can occur as the limit of a family of nonsingular Σ_ϵ modelled on $\{(x, y) \in \mathbb{C}^2 : xy = -\epsilon, \operatorname{Im} x \geq 0, \operatorname{Im} y \geq 0\}$, for $\epsilon > 0$, with doubles $\tilde{\Sigma}_\epsilon$ modelled on $\{(x, y) \in \mathbb{C}^2 : xy = -\epsilon\}$.

- (c) $\tilde{\Sigma}$ has a node z in $\partial\Sigma$ locally modelled on $(0, 0)$ in $\{(x, y) \in \mathbb{C}^2 : xy = 0\}$, where σ acts by $(x, y) \mapsto (\bar{y}, \bar{x})$. Then $\partial\Sigma$ is locally modelled on $(0, 0)$, the fixed point set of σ , and we take Σ to be locally modelled on $\{(x, 0) \in \mathbb{C}^2\}$.

Such a node can occur as the limit of a family of nonsingular Σ_ϵ modelled on $\{(x, y) \in \mathbb{C}^2 : xy = \epsilon, |x| \geq |y|\}$, for $\epsilon > 0$, with doubles $\tilde{\Sigma}_\epsilon$ modelled

on $\{(x, y) \in \mathbb{C}^2 : xy = \epsilon\}$, where σ acts by $(x, y) \mapsto (\bar{y}, \bar{x})$, and boundary the circle $\partial\Sigma_\epsilon = \{(\epsilon^{1/2}e^{i\theta}, \epsilon^{1/2}e^{-i\theta}) : \theta \in [0, 2\pi)\}$.

Note that near a node of type (c), Σ is actually a *nonsingular* Riemann surface *without boundary*, save that one point z in Σ is designated as a ‘boundary node’. Also $\partial\Sigma$ near z is just one point $\{z\}$, not a real curve. We allow as prestable bordered Riemann surfaces curves whose only boundary is nodes of type (c), so for example Σ could be $\mathbb{CP}^1$ with one point designated as a boundary node.

If Σ is a prestable bordered Riemann surface, the *smoothing* Σ' of Σ is the compact, connected, oriented 2-manifold with boundary obtained by smoothing the nodes of Σ . Note that there are two topologically distinct ways of desingularizing nodes of type (a)–(c), and we choose those described as families of surfaces Σ_ϵ above. The rule is that smoothing each node decreases the Euler characteristic by 1, but the other ways of desingularizing nodes of type (a)–(c) increase Euler characteristic by 1, 1, 0 respectively. We call Σ a *prestale holomorphic disc* if its smoothing Σ' is a holomorphic disc.

Definition 6.19. Suppose (M, ω) is a compact symplectic manifold, L a compact, embedded Lagrangian submanifold in M , and J an almost complex structure on M compatible with ω . Let $\beta \in H^2(M; \mathbb{Z})$ and $l \geq 0$. Consider triples $(\Sigma, \vec{y}, w)$ where Σ is a *prestale holomorphic disc* in the sense of Definition 6.18, $\vec{y} = (y_1, \dots, y_l)$, where $y_1, \dots, y_l$ are distinct points of $\partial\Sigma$, none of which are nodes, and which occur in this cyclic order in $\partial\Sigma$, and $w : \Sigma \rightarrow M$ is a J -holomorphic map, with $w(\partial\Sigma) \subset L$ and $w_*([\Sigma]) = \beta$ in $H_2(M, L; \mathbb{Z})$. Call two such triples $(\Sigma, \vec{y}, w)$ and $(\Sigma', \vec{y}', w')$ *isomorphic* if there exists a biholomorphism $f : \Sigma \rightarrow \Sigma'$ with $f(y_i) = y'_i$ for $i = 1, \dots, l$ and $w \equiv w' \circ f$. The *automorphism group* $\text{Aut}(\Sigma, \vec{y}, w)$ is the group of isomorphisms $f : (\Sigma, \vec{y}, w) \rightarrow (\Sigma, \vec{y}, w)$. Call a triple $(\Sigma, \vec{y}, w)$ *stable* if $\text{Aut}(\Sigma, \vec{y}, w)$ is finite.

Write $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ for the moduli space of isomorphism classes $[\Sigma, \vec{y}, w]$ of stable triples $(\Sigma, \vec{y}, w)$ satisfying the above conditions. Define *boundary evaluation maps* $\text{ev}_i : \bar{\mathcal{M}}_l(M, L, J, \beta) \rightarrow L$ for $i = 1, \dots, l$ by $\text{ev}_i : [\Sigma, \vec{y}, w] \mapsto w(y_i)$. Write $\mathcal{M}_l^{\text{ma}}(M, L, J, \beta)$ for the subset of $[\Sigma, \vec{y}, w] \in \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ with Σ *nonsingular*, that is, Σ has no nodes. The ‘ma’ in $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ follows Fukaya et al. [24, §2], who first define moduli spaces $\bar{\mathcal{M}}_l(M, L, J, \beta)$ in which $y_1, \dots, y_l$ need not be in this cyclic order in $\partial\Sigma$, and then define $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ to be the ‘main component’ of $[\Sigma, \vec{y}, w]$ in $\bar{\mathcal{M}}_l(M, L, J, \beta)$ in which $y_1, \dots, y_l$ are cyclically ordered.

There is a natural action of $\mathbb{Z}_l$ on $\bar{\mathcal{M}}_l(M, L, J, \beta)$, by cyclic permutation of the boundary marked points $y_1, \dots, y_l$.

Remark 6.20. Fukaya et al. [24] do not consider boundary nodes of type (c) in Definition 6.18, though they appear to fall within [24, Def. 2.16]. Because of this they consider the moduli spaces $\bar{\mathcal{M}}_0(M, L, J, \beta)$ when $l = 0$ to be possibly *noncompact*. For discussion of this, see remarks preceding [24, Fig. 2.4, Th. 2.29 & Prop. 13.27], and [24, §32.1]. They deal with their noncompactness problem by allowing certain unstable maps in a rather ad hoc way. We prefer our approach including type (c) nodes, based on Liu [51, Def. 3.4].

These moduli spaces have natural topologies called the C^∞ topology, due to Gromov [31] and defined in [24, §29]. The next result, the analogue of Theorem 6.4, is proved by Fukaya et al. [24, Th. 29.43], with a slightly different notion of Kuranishi space. See Remark 6.5 for what we mean by “Theorem”.

Theorem 6.21. *In Definition 6.19 the $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ are compact and Hausdorff in the C^∞ topology, and the ev_i are continuous.*

We define the *Maslov index* of $\beta \in H_2(M, L; \mathbb{Z})$.

Definition 6.22. Let (M, ω) be a symplectic manifold of dimension $2n$, and choose a compatible almost complex structure J . Then the canonical bundle $K_M = \Lambda^{n,0}M$ of (M, J) is a complex line bundle over M , with Hermitian metrics on the fibres $\mathbb{C}$. Let L be an embedded Lagrangian submanifold in M . Then there is a natural isomorphism $K_M|_L \cong \Lambda^n T^*L \otimes_{\mathbb{R}} \mathbb{C}$. Choose a connection A on K_M compatible with the metrics on the fibres, which respects the isomorphism $K_M|_L \cong \Lambda^n T^*L \otimes_{\mathbb{R}} \mathbb{C}$. Let F_A be the curvature of this connection. Then A is a $U(1)$ connection, so identifying $\mathfrak{u}(1) \cong \mathbb{R}$, F_A is a closed 2-form. On L , this A reduces to an $O(1)$ -connection on $\Lambda^n T^*L$, so A is flat, and $F_A|_L \equiv 0$.

Thus we may define the *relative first Chern class* $c_1(M, L)$ of (M, L) to be the relative de Rham cohomology class $\frac{1}{2\pi}[F_A]$ in $H^2(M, L; \mathbb{R})$, recalling that $H^l(M, L; \mathbb{R})$ is defined using closed l -forms α on M with $\alpha|_L \equiv 0$. One can show that $c_1(M, L)$ lies in the image of $H^2(M, L; \frac{1}{2}\mathbb{Z})$ in $H^2(M, L; \mathbb{R})$, and it is independent of the choices of J, A , and if L is orientable then $c_1(M, L)$ lies in the image of $H^2(M, L; \mathbb{Z})$ in $H^2(M, L; \mathbb{R})$.

For $\beta \in H_2(M, L; \mathbb{Z})$ we define the *Maslov index* $\mu_L(\beta) = 2c_1(M, L) \cdot \beta$ in $\mathbb{Z}$. If β is represented by an oriented 2-submanifold (or 2-cycle) Σ in M with $\partial\Sigma \subset L$ then $\mu_L(\beta) = \frac{1}{\pi} \int_{\Sigma} F_A$. If L is orientable then $\mu_L(\beta) \in 2\mathbb{Z}$.

This is different to the usual definition of Maslov index [24, §2], but is equivalent to it. Our definition makes it obvious that $\mu_L(\beta)$ depends only on the relative homology class β ; Katz and Liu [42, Rem. 4.2.2] claim incorrectly that the Maslov index is a homotopy but not a homology invariant. The next result is proved by Fukaya et al. [24, §29].

“Theorem” 6.23. *In Definition 6.19 we can construct Kuranishi structures κ on $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ with corners (not g -corners), with*

$$\text{vdim}(\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \kappa) = \mu_L(\beta) + l + n - 3, \quad (157)$$

where $\dim M = 2n$. The maps ev_i extend to strong submersions

$$\text{ev}_i : (\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \kappa) \longrightarrow L. \quad (158)$$

Also $\text{ev}_1 \times \cdots \times \text{ev}_l : (\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \kappa) \rightarrow L^l$ is a strong submersion.

We can choose κ to be invariant under the action of $\mathbb{Z}_l$ on $\bar{\mathcal{M}}_l(M, L, J, \beta)$ by cyclic permutation of $y_1, \dots, y_l$, that is, $\mathbb{Z}_l$ acts on $\bar{\mathcal{M}}_l(M, L, J, \beta)$ by strong diffeomorphisms, with the obvious compatibility with $\text{ev}_1, \dots, \text{ev}_l$. Next we discuss *orientations* on $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$. For this we need the notion of a *relative spin structure* on (M, L) , [24, Def. 44.2]:

Definition 6.24. Let L be a submanifold of a manifold M . Fix a triangulation of M such that L is a subcomplex. A *relative spin structure* on (M, L) consists of an orientation on L ; a class $\text{st} \in H^2(M; \mathbb{Z}_2)$ such that $\text{st}|_L = w_2(L) \in H^2(L; \mathbb{Z}_2)$, the second Stiefel–Whitney class of L ; an oriented vector bundle V on the 3-skeleton $M_{[3]}$ of M with $w_2(V) = \text{st}$; and a spin structure on $(TL \oplus V)|_{L_{[2]}}$.

Here $L_{[2]}$ is the 2-skeleton of L , and as $w_2(V|_{L_{[2]}}) = \text{st}|_{L_{[2]}} = w_2(L)|_{L_{[2]}}$ we have $w_2((TL \oplus V)|_{L_{[2]}}) = 0$, so $(TL \oplus V)|_{L_{[2]}}$ admits a spin structure. If L is spin then $w_2(L) = 0$, so we can take $\text{st} = 0$ and $V = 0$ and the spin structure on $TL|_{L_{[2]}}$ to be the restriction of that on TL . Hence, an orientation and spin structure on L induce a relative spin structure on (M, L) .

Then Fukaya et al. [24, §44.1 & §46] prove:

“Theorem” 6.25. *A relative spin structure on (M, L) determines orientations on the Kuranishi spaces $(\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \kappa)$.*

From now on we suppress the Kuranishi structures κ , and regard $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ as a Kuranishi space. Finally we state formulae for the boundaries of our moduli spaces, in oriented Kuranishi spaces. Our next theorem can be deduced from results of Fukaya et al. [24]. The closest result to (159) is [24, Prop. 46.2], but here in effect the authors omit the sum over j in (159), and fix $j = 1$. The difficult thing to get right is the orientations (signs). These depend on the order in which the remaining boundary marked points of the two sides of (159) are identified, which is why we have been careful to specify this order through the isomorphism between (160) and (161). The correct sign $(-1)^{(l-k)(k-j)+j+n}$ in (159) when $j = 1$ is given by Fukaya et al. in [24, Prop. 46.2 & Rem. 46.3], and we can deduce the sign in the general case by permuting the l marked points.

In equation (162), the second line is in effect the term $j = k = l = 0$ from (159), and the sign $(-1)^n$ is the same as $(-1)^{(l-k)(k-j)+j+n}$ in (159) with $j = k = l = 0$. However, unlike in (159), we must quotient by an action of $\mathbb{Z}_2$. The point is that at the boundary of $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$, the prestable holomorphic disc Σ breaks into the union of two prestable holomorphic discs Σ_1, Σ_2 joined by a node. The marked points $y_1, \dots, y_l$ in $\partial\Sigma$ each end up in either $\partial\Sigma_1$ or $\partial\Sigma_2$. In (159), when $l \geq 1$, we distinguish between Σ_1 and Σ_2 by decreeing that the marked point y_1 should lie on $\partial\Sigma_1$; this is why we specify $j \geq 1$ rather than $j \geq 0$ in (159). However in (162), when $l = 0$, we cannot use marked points to distinguish Σ_1 and Σ_2 , so the symmetry of exchanging Σ_1 and Σ_2 means that we must divide by $\mathbb{Z}_2$.

The third line of (162) comes from boundary nodes of type (c) in Definition 6.18. The sign 1 on the third line is computed by Fukaya et al. in [24, Prop. 53.6], though they include an extra interior marked point. As we explained in Remark 6.20, Fukaya et al. do not consider nodes of type (c), and so regard $\bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta)$ as possibly noncompact. See also [24, Prop. 13.27] and [24, §32.1] for results analogous to the third line of (162).

“Theorem” 6.26. *In the situation above, choose a relative spin structure on (M, L) , so that $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ for $l \geq 0$ are compact, oriented Kuranishi*

spaces with corners by Theorem 6.21 and “Theorems” 6.23 and 6.25. Then for $l \geq 1$ there is an isomorphism of oriented Kuranishi spaces:

$$\partial \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta) \cong \coprod_{1 \leq j \leq k \leq l} \coprod_{\substack{\gamma, \delta \in H_2(M, L; \mathbb{Z}): \\ \beta = \gamma + \delta}} (-1)^{(l-k)(k-j)+j+n} \cdot \bar{\mathcal{M}}_{l+j-k+1}^{\text{ma}}(M, L, J, \gamma) \times_{\mathbf{ev}_{j+1}, L, \mathbf{ev}_1} \bar{\mathcal{M}}_{k-j+1}^{\text{ma}}(M, L, J, \delta). \quad (159)$$

Here on the j, k, γ, δ term (159) identifies the strong submersions

$$\begin{aligned} & (\mathbf{ev}_1 \times \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_j \times \\ & \mathbf{ev}_{j+1} \times \mathbf{ev}_{j+2} \times \cdots \times \mathbf{ev}_k \times \\ & \mathbf{ev}_{k+1} \times \mathbf{ev}_{k+2} \times \cdots \times \mathbf{ev}_l) |_{\partial \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)} : \end{aligned} \quad (160)$$

$$\begin{aligned} & \partial \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta) \longrightarrow L^j \times L^{k-j} \times L^{l-k} = L^l \quad \text{and} \\ & (\mathbf{ev}_1 \circ \pi_1) \times (\mathbf{ev}_1 \circ \pi_1) \times \cdots \times (\mathbf{ev}_j \circ \pi_1) \times \\ & (\mathbf{ev}_2 \circ \pi_2) \times (\mathbf{ev}_3 \circ \pi_2) \times \cdots \times (\mathbf{ev}_{k-j+1} \circ \pi_2) \times \\ & (\mathbf{ev}_{j+2} \circ \pi_1) \times (\mathbf{ev}_{j+3} \circ \pi_1) \times \cdots \times (\mathbf{ev}_{l+j-k+1} \circ \pi_1) : \\ & \bar{\mathcal{M}}_{l+j-k+1}^{\text{ma}}(M, L, J, \gamma) \times_{\mathbf{ev}_{j+1}, L, \mathbf{ev}_1} \bar{\mathcal{M}}_{k-j+1}^{\text{ma}}(M, L, J, \delta) \\ & \longrightarrow L^j \times L^{k-j} \times L^{l-k} = L^l, \end{aligned} \quad (161)$$

where π_1, π_2 project $\bar{\mathcal{M}}_{l+j-k+1}^{\text{ma}}(M, L, J, \gamma) \times_{\mathbf{ev}_{j+1}, L, \mathbf{ev}_1} \bar{\mathcal{M}}_{k-j+1}^{\text{ma}}(M, L, J, \delta)$ to $\bar{\mathcal{M}}_{l+j-k+1}^{\text{ma}}(M, L, J, \gamma)$ and $\bar{\mathcal{M}}_{k-j+1}^{\text{ma}}(M, L, J, \delta)$ respectively.

For $l = 0$ there is an isomorphism of oriented Kuranishi spaces:

$$\begin{aligned} & \partial \bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta) \cong \\ & \left[\coprod_{\substack{\gamma, \delta \in H_2(M, L; \mathbb{Z}): \beta = \gamma + \delta, \\ [\omega] \cdot \gamma > 0, [\omega] \cdot \delta > 0}} (-1)^n \cdot \bar{\mathcal{M}}_1^{\text{ma}}(M, L, J, \gamma) \times_{\mathbf{ev}_1, L, \mathbf{ev}_1} \bar{\mathcal{M}}_1^{\text{ma}}(M, L, J, \delta) \right] / \mathbb{Z}_2 \\ & \coprod_{\beta' \in H_2(M; \mathbb{Z}): \iota(\beta') = \beta} \bar{\mathcal{M}}_{0,1}(M, J, \beta') \times_{\mathbf{ev}_1, M, \text{inc}} L. \end{aligned} \quad (162)$$

Here the $\mathbb{Z}_2$ -action exchanges γ, δ and $\bar{\mathcal{M}}_1^{\text{ma}}(M, L, J, \gamma), \bar{\mathcal{M}}_1^{\text{ma}}(M, L, J, \delta)$. In the third line, $\text{inc} : L \rightarrow M$ is the inclusion and ι comes from the exact sequence $H_2(M; \mathbb{Z}) \xrightarrow{\iota} H_2(M, L; \mathbb{Z}) \xrightarrow{\partial} H_1(L; \mathbb{Z})$. The third line is zero unless $\partial\beta = 0$.

6.5 Choosing co-gauge-fixing data for the $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$

In §6.5–§6.7, we assume the “Theorems” of §6.4 throughout. For our attitude to these, see Remark 6.5.

In the situation of §6.4, $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ is a compact, oriented Kuranishi space, $\mathbf{ev}_1 \times \cdots \times \mathbf{ev}_l : \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta) \rightarrow L^l$ is a strong submersion, and L^l is oriented, so $(\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_l)$ is cooriented. Thus by Theorem 3.18 we can choose co-gauge-fixing data $\mathbf{C}_l(M, L, J, \beta)$ for $(\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta),$

$\mathbf{ev}_1 \times \cdots \times \mathbf{ev}_l$), so that $[\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_l, \mathcal{C}_l(M, L, J, \beta)]$ lies in $KC^*(L^l; R)$, for R a $\mathbb{Q}$ -algebra. We will now show that we can choose the $\mathcal{C}_l(M, L, J, \beta)$ for all l, β so that equations (159) and (162) of “Theorem” 6.26 lift to equations in Kuranishi cochains.

However, thinking carefully about (159) and (162), we see that there is a puzzle. The fibre products in (159) and (162) do not quite correspond to any operations we have defined on co-gauge-fixing data, even in Remark 3.34(b). In the j, k, γ, δ term in (159), it is natural to regard the term $\bar{\mathcal{M}}_{l+j-k+1}^{\text{ma}}(M, L, J, \gamma)$ as giving a cochain in $KC^*(L^{l+j-k} \times L; R)$, and the term $\bar{\mathcal{M}}_{k-j+1}^{\text{ma}}(M, L, J, \delta)$ as giving a cochain in $KC^*(L^{k-j} \times L; R)$. As in Remark 3.34(b) we can take the fibre product of these to get a cochain in $KC^*(L^{l+j-k} \times L^{k-j} \times L; R) = KC^*(L^{l+1}; R)$. But we actually want a cochain in $KC^*(L^l; R)$. We want to forget one factor of L , that is, we want to push forward by $L^{l+1} \rightarrow L^l$. But pushforwards are defined for Kuranishi chains, not Kuranishi cochains.

This suggests that we need to mix chains and cochains, so that we should be working in the *bivariant theory* of §4.8. This may be the most natural point of view for Lagrangian Floer cohomology. In $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ for $l \geq 1$, we can distinguish the first marked point y_1 from the rest, and regard $(\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_l)$ as defining a *Kuranishi bichain* in $KB^*(\pi_1 : L^l \rightarrow L; R)$ where $\pi_1 : (y_1, \dots, y_l) \mapsto y_1$. Then the fibre products in (159) do correspond to natural operations on Kuranishi bichains, made out of products, pushforwards, and pullbacks, but those in (162) do not.

Unfortunately, using the bivariant theory in this way is not good for defining open Gromov–Witten invariants. As in §6.5, there is an action of $\mathbb{Z}_l$ on $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ by cyclic permutation of the marked points. For open Gromov–Witten theory we need to use this cyclic symmetry, but distinguishing the first marked point y_1 destroys it. So we adopt a different solution. We choose some extra data, in effect gauge-fixing data $\mathbf{G}_L$ for (L, id_L) . Using this we define a *pushforward* $KC^*(L^{l+1}; R) \rightarrow KC^*(L^l; R)$, which is in effect a cap product with $(L, \text{id}_L, \mathbf{G}_L)$. Then we can define products $*$ of co-gauge-fixing data and of Kuranishi cochains $KC^*(L^a; R) \times KC^*(L^b; R) \rightarrow KC^*(L^{a+b-2}; R)$ that we will need to make sense of adding co-gauge-fixing data to (159) and (162).

Definition 6.27. Let P_m, P, μ be as in Definition 3.8. Suppose L is an oriented n -manifold, not necessarily compact. Choose an injective map $G_L : L \rightarrow P_m \subset P$ for some $m \gg 0$. Let $a, b \geq 1$, $X, \tilde{X}$ be compact Kuranishi spaces, $\mathbf{f}_1 \times \cdots \times \mathbf{f}_a : X \rightarrow L^a$, $\tilde{\mathbf{f}}_1 \times \cdots \times \tilde{\mathbf{f}}_b : \tilde{X} \rightarrow L^b$ be strong submersions, $\mathbf{C}, \tilde{\mathbf{C}}$ be co-gauge-fixing data for $(X, \mathbf{f}_1 \times \cdots \times \mathbf{f}_a)$ and $(\tilde{X}, \tilde{\mathbf{f}}_1 \times \cdots \times \tilde{\mathbf{f}}_b)$. Let $1 \leq i \leq a$ and $1 \leq j \leq b$. Then $X \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{X}$ is a compact Kuranishi space. Define a strong submersion $\mathbf{g}_1 \times \cdots \times \mathbf{g}_{a+b-2} : X \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{X} \rightarrow L^{a+b-2}$ to be

$$\begin{aligned} & (\mathbf{f}_1 \circ \pi_X) \times \cdots \times (\mathbf{f}_{i-1} \circ \pi_X) \times (\mathbf{f}_{i+1} \circ \pi_X) \times \cdots \times (\mathbf{f}_a \circ \pi_X) \times \\ & (\tilde{\mathbf{f}}_1 \circ \pi_{\tilde{X}}) \times \cdots \times (\tilde{\mathbf{f}}_{j-1} \circ \pi_{\tilde{X}}) \times (\tilde{\mathbf{f}}_{j+1} \circ \pi_{\tilde{X}}) \times \cdots \times (\tilde{\mathbf{f}}_b \circ \pi_{\tilde{X}}). \end{aligned} \quad (163)$$

We will define co-gauge-fixing data $\mathbf{C} *_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{\mathbf{C}}$ for $(X \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{X}, \mathbf{g}_1 \times$

$\cdots \times \mathbf{g}_{a+b-2}$). Remark 3.34(b) explains how to modify the construction of §3.8 to define co-gauge-fixing data $\mathbf{C} \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{\mathbf{C}}$ which we can consider as co-gauge-fixing data for $(X \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{X}, \mathbf{g}_1 \times \cdots \times \mathbf{g}_{a+b-2} \times \pi_L)$, where $\pi_L : X \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{X} \rightarrow L$ is the projection from the fibre product. Write $\mathbf{C} \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{\mathbf{C}}$ as $(\check{\mathbf{I}}, \check{\boldsymbol{\eta}}, \check{C}^k : k \in \check{I})$, where $\check{\mathbf{I}} = (\check{I}, (\check{V}^k, \dots, \check{\psi}^k), \check{g}_1^k \times \cdots \times \check{g}_{a+b-2}^k \times \check{\pi}_L^k : k \in \check{I}, \dots)$, and $\check{C}^k$ maps $\check{E}^k \rightarrow P$ for $k \in \check{I}$.

Define $\check{\mathbf{I}} = (\check{I}, (\check{V}^k, \dots, \check{\psi}^k), \check{g}_1^k \times \cdots \times \check{g}_{a+b-2}^k : k \in \check{I}, \dots)$, and $\check{\boldsymbol{\eta}} = \check{\boldsymbol{\eta}}$, and for each $k \in \check{I}$ define $\check{C}^k : \check{E}^k \rightarrow P$ by $\check{C}^k = \mu \circ ((G_L \circ \check{\pi}_L^k \circ \check{\pi}^k) \times \check{C}^k)$, where $\check{\pi}^k : \check{E}^k \rightarrow \check{V}^k$ is the natural projection. Define $\mathbf{C} *_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{\mathbf{C}} = (\check{\mathbf{I}}, \check{\boldsymbol{\eta}}, \check{C}^k : k \in \check{I})$. That is, $\mathbf{C} *_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{\mathbf{C}}$ is the same as $\mathbf{C} \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{\mathbf{C}}$ except that we omit the maps $\check{\pi}_L^k$, and multiply the $\check{C}^k$ by the pullback of G_L to $\check{E}^k$ using $\check{\pi}_L^k \circ \check{\pi}^k$. This is similar to the construction of $\mathbf{G}_{\mathbf{C}}$ in Definition 4.14, and a similar proof shows that $\mathbf{C} *_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{\mathbf{C}}$ is co-gauge-fixing data for $(X \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{X}, \mathbf{g}_1 \times \cdots \times \mathbf{g}_{a+b-2})$.

Note that using the commutativity and associativity of μ , one can show that these products $*_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j}$ have the commutativity and associativity properties one would expect, in a similar way to Proposition 3.32.

If R is a $\mathbb{Q}$ -algebra, L is oriented, and $1 \leq i \leq a$, $1 \leq j \leq b$, define R -module morphisms $*_{i,j} : KC^k(L^a; R) \times KC^l(L^b; R) \rightarrow KC^{k+l-n}(L^{a+b-2}; R)$ by

$$\begin{aligned} [X, \mathbf{f}_1 \times \cdots \times \mathbf{f}_a, \mathbf{C}] *_{i,j} [\tilde{X}, \tilde{\mathbf{f}}_1 \times \cdots \times \tilde{\mathbf{f}}_b, \tilde{\mathbf{C}}] = \\ [X \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{X}, \mathbf{g}_1 \times \cdots \times \mathbf{g}_{a+b-2}, \mathbf{C} *_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{\mathbf{C}}], \end{aligned} \quad (164)$$

where we use the orientation on L to convert the product coorientation for $(X \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{X}, \mathbf{g}_1 \times \cdots \times \mathbf{g}_{a+b-2} \times \pi_L)$ into a coorientation for $(X \times_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j} \tilde{X}, \mathbf{g}_1 \times \cdots \times \mathbf{g}_{a+b-2})$. Equation (164) takes relations in $KC^k(L^a; R), KC^l(L^b; R)$ to relations in $KC^{k+l-n}(L^{a+b-2}; R)$, and so is well-defined. As for cup products in (107) we find that $d(\gamma *_{i,j} \delta) = (d\gamma) *_{i,j} \delta + (-1)^k \gamma *_{i,j} (d\delta)$, so the $*_{i,j}$ induce morphisms on Kuranishi cohomology

$$*_{i,j} : KH^k(L^a; R) \times KH^l(L^b; R) \longrightarrow KH^{k+l-n}(L^{a+b-2}; R). \quad (165)$$

We can now choose $\mathbf{C}_l(M, L, J, \beta)$ compatible with (159), (162) for all l, β .

Theorem 6.28. *In the situation of §6.4 we can choose co-gauge-fixing data $\mathbf{C}_l(M, L, J, \beta)$ for $(\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_l)$ for all $l \geq 1$ and β in $H_2(M, L; \mathbb{Z})$ such that under (159), the restriction $\mathbf{C}_l(M, L, J, \beta)|_{\partial \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)}$ is identified on the j, k, γ, δ term with $\mathbf{C}_{l+j-k+1}(M, L, J, \gamma) *_{\mathbf{ev}_{j+1}, L, \mathbf{ev}_1} \mathbf{C}_{k-j+1}(M, L, J, \delta)$, using the notation of Definition 6.27. Moreover, we can choose $\mathbf{C}_l(M, L, J, \beta)$ to be invariant under the action of $\mathbb{Z}_l$ on $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ by cyclic permutation of the marked points.*

For the case $l = 0$, choose $\mathbb{Z}_l$ -invariant $\mathbf{C}_l(M, L, J, \beta)$ for all $l \geq 1$ and β as above, and choose co-gauge-fixing data $\mathbf{C}_{0,1}(M, J, \beta')$ for $(\bar{\mathcal{M}}_{0,1}(M, J, \beta'), \mathbf{ev}_1)$

for all β' in $H_2(M; \mathbb{Z})$. Write $\pi : \bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta) \rightarrow \{0\}$ for the strong submersion projecting to one point $\{0\}$. Then we can choose co-gauge-fixing data $\mathbf{C}_0(M, L, J, \beta)$ for $(\bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta), \pi)$ for all β in $H_2(M, L; \mathbb{Z})$ such that under (162), the restriction $\mathbf{C}_0(M, L, J, \beta)|_{\partial \bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta)}$ is identified on the γ, δ term with $\mathbf{C}_1(M, L, J, \gamma) *_{\text{ev}_1, L, \text{ev}_1} \mathbf{C}_1(M, L, J, \delta)$, where we take the quotient of this co-gauge-fixing data by the action of $\mathbb{Z}_2$ as in §3.4, and on the β' term with $\text{inc}^*(\mathbf{C}_{0,1}(M, J, \beta')) *_{\text{ev}_1, L, \text{id}_L} [L, \text{id}_L, \mathbf{C}_L]$ using pullbacks of co-gauge-fixing data as in §3.8, defining $[L, \text{id}_L, \mathbf{C}_L]$ as in §4.7, and identifying $L^0 \cong \{0\}$.

Proof. We only need choose $\mathbf{C}_l(M, L, J, \beta)$ when $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta) \neq \emptyset$. Using compactness results from Geometric Measure Theory, we find that for any $A \geq 0$, there are only finitely many $\beta \in H_2(M, L; \mathbb{Z})$ with $[\omega] \cdot \beta \leq A$ for which there exist J -holomorphic curves in class β . Thus we can arrange all such β in an ordered list $0 = \beta_0, \beta_1, \beta_2, \dots$ such that $m \leq n$ implies that $[\omega] \cdot \beta_m \leq [\omega] \cdot \beta_n$.

For the first part of the theorem, we will choose $\mathbf{C}_l(M, L, J, \beta_m)$ by a double induction on l, m . The outer induction is over $m = 0, 1, 2, \dots$, and the inductive hypothesis at the m^{th} step is that we have chosen $\mathbf{C}_{l'}(M, L, J, \beta_{m'})$ satisfying the conditions for all $1 \leq l'$ and $0 \leq m' < m$. The inner induction, for fixed m , is over $l = 1, 2, \dots$, and the inductive hypothesis is that we have chosen $\mathbf{C}_{l'}(M, L, J, \beta_{m'})$ satisfying the conditions for all $1 \leq l'$ and $0 \leq m' < m$, and also that we have chosen $\mathbf{C}_{l'}(M, L, J, \beta_m)$ satisfying the conditions for all $1 \leq l' < l$. In the inductive step of the inner induction we must choose $\mathbf{C}_l(M, L, J, \beta_m)$ satisfying the conditions.

We suppose these inductive hypotheses, and prove the inner inductive step. Consider which j, k, γ, δ terms in (159) can be nonempty, where $\beta = \beta_m$. For $\bar{\mathcal{M}}_{l+j-k+1}^{\text{ma}}(M, L, J, \gamma) \neq \emptyset$ and $\bar{\mathcal{M}}_{k-j+1}^{\text{ma}}(M, L, J, \delta) \neq \emptyset$ we must have $\gamma = \beta_{m'}$ and $\delta = \beta_{m''}$ for some m', m'' . First suppose $\gamma, \delta \neq 0$. Then $[\omega] \cdot \gamma, [\omega] \cdot \delta > 0$, so $[\omega] \cdot \gamma, [\omega] \cdot \delta < [\omega] \cdot \beta_m$ as $[\omega] \cdot \gamma + [\omega] \cdot \delta = [\omega] \cdot \beta_m$, and so $m', m'' < m$. Thus in this case we have already chosen $\mathbf{C}_{l+j-k+1}(M, L, J, \gamma)$ and $\mathbf{C}_{k-j+1}(M, L, J, \delta)$ in a previous outer inductive step. If $\gamma = 0$ then $\delta = \beta_m$, so $m'' = m$, and we must have $l+j-k+1 \geq 3$, since $\bar{\mathcal{M}}_i^{\text{ma}}(M, L, J, 0) = \emptyset$ unless $i \geq 3$, so $k-j+1 < l$. Thus, if $\gamma = 0$ then $\bar{\mathcal{M}}_{k-j+1}^{\text{ma}}(M, L, J, \delta)$ is of the form $\bar{\mathcal{M}}_{l'}^{\text{ma}}(M, L, J, \beta_m)$ for $l' < l$, and we have already chosen $\mathbf{C}_{k-j+1}(M, L, J, \delta)$ in a previous inner inductive step. Similarly, if $\delta = 0$ then we have already chosen $\mathbf{C}_{l+j-k+1}(M, L, J, \gamma)$ in a previous inner inductive step.

Thus in all cases, if the j, k, γ, δ term in (159) is nonempty then we have chosen $\mathbf{C}_{l+j-k+1}(M, L, J, \gamma)$ and $\mathbf{C}_{k-j+1}(M, L, J, \delta)$ in a previous inductive step. So the conditions in the theorem uniquely determine co-gauge-fixing data $\mathbf{D}_l(M, L, J, \beta_m)$ for $(\partial \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m), (\text{ev}_1 \times \dots \times \text{ev}_l)|_{\partial \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m)})$, and we must choose $\mathbf{C}_l(M, L, J, \beta_m)$ with $\mathbf{C}_l(M, L, J, \beta_m)|_{\partial \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m)} = \mathbf{D}_l(M, L, J, \beta_m)$. As in “Theorem” 6.23, $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m)$ is a Kuranishi space *with corners, not g-corners*. So we can apply Theorem 3.25. We must check that the restriction $\mathbf{D}_l(M, L, J, \beta_m)|_{\partial^2 \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m)}$ is invariant under the natural involution $\sigma : \partial^2 \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m) \rightarrow \partial^2 \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m)$.

Now taking the boundary of (159), using Proposition 2.31(a) to rewrite the r.h.s. in terms of $\partial \bar{\mathcal{M}}_{l+j-k+1}^{\text{ma}}(M, L, J, \gamma), \partial \bar{\mathcal{M}}_{k-j+1}^{\text{ma}}(M, L, J, \delta)$, and then

substituting for these using (159), gives an expression for $\partial^2 \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m)$ in terms of double fibre products of three factors $\bar{\mathcal{M}}_i^{\text{ma}}(M, L, J, \epsilon)$. By previous inductive steps, using this we can write $\mathbf{D}_l(M, L, J, \beta_m)|_{\partial^2 \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m)}$ as a disjoint union of double products $*_{\mathbf{ev}_i, L, \mathbf{ev}_j}$ of three factors $\mathbf{C}_i(M, L, J, \epsilon)$.

When written this way, σ -invariance of $\mathbf{D}_l(M, L, J, \beta_m)|_{\partial^2 \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m)}$ turns out to follow from associativity of $*_{\mathbf{ev}_i, L, \mathbf{ev}_j}$ on co-gauge-fixing data in Definition 6.27. That is, σ acts by exchanging terms $X_1 \times_L (X_2 \times_L X_3)$ and $(X_1 \times_L X_2) \times_L X_3$, and this must identify $(\mathbf{C}_1 *_L \mathbf{C}_2) *_L \mathbf{C}_3$ and $\mathbf{C}_1 *_L (\mathbf{C}_2 *_L \mathbf{C}_3)$, where X_1, X_2, X_3 are of the form $\mathbf{C}_i(M, L, J, \epsilon)$, and $\mathbf{C}_1, \mathbf{C}_2, \mathbf{C}_3$ of the form $\mathbf{C}_i(M, L, J, \epsilon)$. Therefore $\mathbf{D}_l(M, L, J, \beta_m)|_{\partial^2 \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m)}$ is σ -invariant, and we can choose $\mathbf{C}_l(M, L, J, \beta_m)$ satisfying the conditions of the theorem by Theorem 3.25. This completes the inner inductive step. The outer inductive step the initial steps are trivial, so the first part of the theorem follows by induction.

To choose $\mathbf{C}_l(M, L, J, \beta_m)$ to be $\mathbb{Z}_l$ -invariant, we use the Γ -invariance part of Theorem 3.25 with $\Gamma = \mathbb{Z}_l$. If the $\mathbf{C}_{l'}(M, L, J, \beta_{m'})$ in previous steps have been chosen $\mathbb{Z}_{l'}$ -invariant then $\mathbf{D}_l(M, L, J, \beta_m)$ turns out to be $\mathbb{Z}_l$ -invariant, with $\mathbb{Z}_l$ -action on $\mathbf{D}_l(M, L, J, \beta_m)|_{\partial^2 \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta_m)}$ commuting with σ , so Theorem 3.25 tells us we can choose $\mathbf{C}_l(M, L, J, \beta_m)$ to be $\mathbb{Z}_l$ -invariant.

For the final part, when $l = 0$, a similar proof works, though no induction is needed. The conditions in the theorem uniquely determine co-gauge-fixing data $\mathbf{D}_0(M, L, J, \beta)$ for $(\partial \bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta), \pi|_{\partial \bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta)})$, and we must choose $\mathbf{C}_0(M, L, J, \beta)$ with $\mathbf{C}_0(M, L, J, \beta)|_{\partial \bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta)} = \mathbf{D}_0(M, L, J, \beta)$ using Theorem 3.25. Note that $\partial^2 \bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta)$ involves terms in $\bar{\mathcal{M}}_2^{\text{ma}}(M, L, J, \epsilon)$, and we need $\mathbf{C}_2(M, L, J, \epsilon)$ to be $\mathbb{Z}_2$ -invariant to make $\mathbf{D}_0(M, L, J, \beta)|_{\partial^2 \bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta)}$ σ -invariant. So we do need the $\mathbf{C}_l(M, L, J, \beta)$ for $l \geq 1$ to be $\mathbb{Z}_l$ -invariant. $\square$

We define some notation.

Definition 6.29. In Theorem 6.28, choose $\mathbb{Z}_l$ -invariant co-gauge-fixing data $\mathbf{C}_l(M, L, J, \beta)$ for all $l \geq 1$, β , and $\mathbf{C}_{0,1}(M, J, \beta')$ for all β' , and $\mathbf{C}_0(M, L, J, \beta)$ for all β . Let R be a $\mathbb{Q}$ -algebra. Define elements $\mathfrak{M}_l(\beta) \in KC(L^l; R)$ for all $l \geq 1$ and $\beta \in H_2(M, L; \mathbb{Z})$, and $\mathfrak{M}_0(\beta) \in KC(\{0\}; R)$ for all $\beta \in H_2(M, L; \mathbb{Z})$, and $\mathfrak{N}_0(\beta') \in KC(\{0\}; R)$ for all $\beta' \in H_2(M; \mathbb{Z})$, by

$$\mathfrak{M}_l(\beta) = [\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_l, \mathbf{C}_l(M, L, J, \beta)], \quad (166)$$

$$\mathfrak{M}_0(\beta) = [\bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta), \pi, \mathbf{C}_0(M, L, J, \beta)], \quad (167)$$

$$\begin{aligned} \mathfrak{N}_0(\beta') &= [\bar{\mathcal{M}}_{0,1}(M, J, \beta') \times_{\mathbf{ev}_1, M, \text{inc}} L, \pi, \\ &\quad \text{inc}^*(\mathbf{C}_{0,1}(M, J, \beta')) *_{\mathbf{ev}_1, L, \text{id}_L} [L, \text{id}_L, \mathbf{C}_L]], \end{aligned} \quad (168)$$

where π is the projection to $\{0\}$ in both cases. We identify $L^0 = \{0\}$, so that $\mathfrak{M}_0(\beta), \mathfrak{N}_0(\beta') \in KC(L^0; R)$.

Combining “Theorem” 6.26, Theorem 6.28 and Definitions 6.27 and 6.29 gives:

Corollary 6.30. *In the situation above, $\mathfrak{M}_l(\beta) \in KC(L^l; R)$ is unchanged by cyclic permutation of the l factors of L in L^l , for all $l \geq 1$ and β . We have*

$$d(\mathfrak{M}_l(\beta)) = \sum_{1 \leq j \leq k \leq l} \sum_{\substack{\gamma, \delta \in H_2(M, L; \mathbb{Z}): \\ \beta = \gamma + \delta}} (-1)^{(l-k)(k-j)+j+n} \mathfrak{M}_{l+j-k+1}(\gamma) *_{j+1,1} \mathfrak{M}_{k-j+1}(\delta), \quad (169)$$

$$d(\mathfrak{M}_0(\beta)) = \frac{1}{2} \sum_{\substack{\gamma, \delta \in H_2(M, L; \mathbb{Z}): \beta = \gamma + \delta, \\ [\omega] \cdot \gamma > 0, [\omega] \cdot \delta > 0}} (-1)^n \cdot \mathfrak{M}_1(\gamma) *_{1,1} \mathfrak{M}_1(\delta) + \sum_{\beta' \in H_2(M; \mathbb{Z}): \mathfrak{i}(\beta') = \beta} \mathfrak{N}_0(\beta'), \quad (170)$$

for all $l \geq 1$ and $\beta \in H_2(M, L; \mathbb{Z})$, where in the term $\mathfrak{M}_{l+j-k+1}(\gamma) *_{j+1,1} \mathfrak{M}_{k-j+1}(\delta)$ in (169), we reorder the factors of L as in (160) and (161). Also $d(\mathfrak{N}_0(\beta')) = 0$ for all $\beta' \in H_2(M; \mathbb{Z})$, and $\mathfrak{M}_l(0) = 0$ for $l < 3$.

The last line holds as $\partial \bar{\mathcal{M}}_{0,1}(M, J, \beta') = \partial L = \emptyset$ and $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, 0) = \emptyset$ for $l < 3$. Observe that we have replaced the quotient by $\mathbb{Z}_2$ in (162) by the factor $\frac{1}{2}$ in (170), using relation Definition 4.4(iv) in $KC^*(\{0\}; R)$. This was an important reason for including Definition 4.4(iv) amongst the relations for Kuranishi (co)chains; as discussed in Remarks 4.5(b) and C.1(a), we do not need Definition 4.4(iv) to define a good theory of Kuranishi (co)homology, and we do not use it in effective Kuranishi (co)homology.

Here is an important point. As we will see in §6.6, most of Lagrangian Floer cohomology for one Lagrangian is a *purely algebraic consequence* of equation (169), together with an analogue of (169) for families of almost complex structures $J_t : t \in [0, 1]$, to express the dependence of the $\mathfrak{M}_l(\beta)$ on the choice of almost complex structure J . Also, as in §6.6, much of the theory of genus zero open Gromov–Witten invariants is a purely algebraic consequence of equations (169)–(170), and analogues for families $J_t : t \in [0, 1]$. So, Corollary 6.30 and results of the same kind are *powerful tools for turning geometry into algebra*.

It is a natural question whether any parts of Theorem 6.28 and Corollary 6.30 have analogues for *effective* co-gauge-fixing data $\underline{\mathcal{C}}_l(M, L, J, \beta)$, so that we can work in $KC_{\text{ef}}^*(L^l; R)$ for R a commutative ring, rather than a $\mathbb{Q}$ -algebra. When $l = 0$ the answer is definitely not: in the γ, δ term in (162), to make the co-gauge-fixing data descend to the quotient $\mathbb{Z}_2$, we need fibre products of co-gauge-fixing data to be commutative, and this does not hold for effective co-gauge-fixing data. For similar reasons, the part in Theorem 6.28 on choosing $\mathcal{C}_l(M, L, J, \beta)$ $\mathbb{Z}_l$ -invariant for $l \geq 2$ cannot extend to effective $\underline{\mathcal{C}}_l(M, L, J, \beta)$.

However, the first part of Theorem 6.28, on choosing $\mathcal{C}_l(M, L, J, \beta)$ for $l \geq 1$, works in the effective case provided the Kuranishi spaces $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ are *effective*. This is because the proof relies on the *associativity* of products $\mathcal{C} *_{f_i, L, \tilde{f}_j} \tilde{\mathcal{C}}$, but not their (super)commutativity. We can generalize Definition 6.27 to give an associative product $\underline{\mathcal{C}} *_{f_i, L, \tilde{f}_j} \tilde{\underline{\mathcal{C}}}$ of effective co-gauge-fixing data. We choose an injective map $\underline{\mathcal{G}}_L : L \rightarrow \mathbb{R}^m \subset \underline{P}$ for some $m \gg 0$. In defining the maps $\underline{\mathcal{C}}^k : \tilde{E}^k \rightarrow \underline{P}$ we must multiply together three factors using the associative, non-commutative multiplication $\underline{\mu}$ on $\underline{P}$. These factors come from $\underline{\mathcal{C}}^i$ in $\underline{\mathcal{C}}$, from $\underline{\mathcal{G}}_L$, and from $\tilde{\underline{\mathcal{C}}}^{\tilde{n}}$ in $\tilde{\underline{\mathcal{C}}}$, and we must multiply them in this order, so that $\underline{\mathcal{G}}_L$ is in the middle, to get the associativity we need.

Now if $[\Sigma, \vec{y}, w] \in \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ for $l \geq 1$, then the only way in which we can have $\text{Aut}(\Sigma, \vec{y}, w) \neq \{1\}$ is if Σ has a nonconstant $\mathbb{CP}^1$ component (or several $\mathbb{CP}^1$ components) joined to the rest of Σ by a node, and $\text{Aut}(\Sigma, \vec{y}, w)$ acts nontrivially on this $\mathbb{CP}^1$, but trivially on the disc part of Σ . If we assume that there are no nonconstant J -holomorphic $\mathbb{CP}^1$'s in M , then this cannot happen, so $\text{Aut}(\Sigma, \vec{y}, w) = \{1\}$ for all $[\Sigma, \vec{y}, w]$, so $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ for $l \geq 1$ has *trivial stabilizers*, and is an *effective* Kuranishi space. (Note that this does not work when $l = 0$, as we need at least one marked point to prevent the disc part of Σ from having automorphisms.) Then we can prove:

Theorem 6.31. *In the situation above, suppose there are no nonconstant J -holomorphic $\mathbb{CP}^1$'s in M . This happens for instance if there exist no spherical $\beta' \in H_2(M; \mathbb{Z})$ with $[\omega] \cdot \beta' > 0$, or if J is generic and for all spherical $\beta' \in H_2(M; \mathbb{Z})$ with $[\omega] \cdot \beta' > 0$ we have $c_1(M) \cdot \beta' < 3 - n$, where $\dim M = 2n$.*

Extend Definitions 6.27 and 6.29 to effective co-gauge-fixing data in the obvious way. Then we can choose effective co-gauge-fixing data $\underline{\mathcal{C}}_l(M, L, J, \beta)$ for $(\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_l)$ for all $l \geq 1$ and β in $H_2(M, L; \mathbb{Z})$ such that under (159), the restriction $\underline{\mathcal{C}}_l(M, L, J, \beta)|_{\partial \bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)}$ is identified on the j, k, γ, δ term with $\underline{\mathcal{C}}_{l+j-k+1}(M, L, J, \gamma) *_{\mathbf{ev}_{j+1}, L, \mathbf{ev}_1} \underline{\mathcal{C}}_{k-j+1}(M, L, J, \delta)$. The analogue of (169) is

$$d(\underline{\mathfrak{M}}_l(\beta)) = \sum_{1 \leq j \leq k \leq l} \sum_{\substack{\gamma, \delta \in H_2(M, L; \mathbb{Z}): \\ \beta = \gamma + \delta}} (-1)^{(l-k)(k-j)+j+n} \underline{\mathfrak{M}}_{l+j-k+1}(\gamma) *_{j+1, 1} \underline{\mathfrak{M}}_{k-j+1}(\delta), \quad (171)$$

which holds in $KC_{\text{ec}}^*(L^l; R)$ for all $l \geq 1$ and β , with R a commutative ring.

Remark 6.32. In §6.3.1 we sketched a blow-up procedure for turning an almost complex Kuranishi space $X, (J, K)$ into an *effective* almost complex Kuranishi space $\tilde{X}, (\tilde{J}, \tilde{K})$. This was central to our programme for understanding integrality properties of Gromov–Witten invariants. We compared our idea to a scheme of Fukaya and Ono [26] for defining $\mathbb{Z}$ -valued Gromov–Witten type invariants.

Now in [26], Fukaya and Ono also apply the same ideas to indicate how one could define Lagrangian Floer cohomology over the integers, for general symplectic manifolds. Parallel to this, one should be able to adapt our ideas in §6.3 to define Lagrangian Floer cohomology for general symplectic manifolds using *effective* Kuranishi cochains, which would then work over arbitrary rings R , not just $\mathbb{Q}$ -algebras. That is, the Gopakumar–Vafa integrality story for closed Gromov–Witten invariants should have an open version, in which Lagrangian Floer cohomology works over $\mathbb{Z}$ not $\mathbb{Q}$.

We would start by using the blow-up procedure of §6.3.1 to modify the moduli spaces $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ to make new, *effective* spaces $\tilde{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$, satisfying the analogue of (159). Replacing $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ by $\tilde{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$, we could then prove the analogue of Theorem 6.31 *without* assuming there are no nonconstant J -holomorphic $\mathbb{CP}^1$'s in M .

Now the $\tilde{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ do not have almost complex structures, so apparently we cannot use the blow-up procedure on them. However, as above, all

the nontrivial orbifold strata $\overline{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)^{\Gamma, \rho}$ come from $\mathbb{CP}^1$ bubbles with a Γ -action. These $\mathbb{CP}^1$ bubbles come lie in moduli spaces $\overline{\mathcal{M}}_{0,1}(M, J, \beta')$ which do have almost complex structures (J, K) . From these (J, K) we can construct almost complex structures on the virtual normal bundle of $\overline{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)^{\Gamma, \rho}$ in $\overline{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$, and this is enough to perform the blow up procedure.

6.6 Lagrangian Floer cohomology for one Lagrangian

Lagrangian Floer cohomology is defined and studied by Fukaya, Oh, Ohta and Ono in the monumental [24], which is nearly 1400 pages long. Much of the length and technical complexity of [24] is due to problems caused by perturbing moduli spaces $\overline{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta)$ to choose virtual moduli chains in singular homology, and ensuring these choices of perturbations and virtual chains are compatible at the boundaries with choices for other moduli spaces under the isomorphism (159).

One difficulty Fukaya et al. face is making their singular chains suitably transverse; a singular chain of positive codimension is never transverse to itself, so they can never arrange for all the singular chains they consider to be pairwise transverse. A second difficulty is that one can in effect make only finitely many choices of perturbations and virtual chains compatible at once, so Fukaya et al. build their A_∞ algebras through an algebraic limiting process from a series of finite geometric approximations called $A_{N,K}$ algebras.

If we develop the theory of Lagrangian Floer cohomology using Kuranishi cochains instead of singular chains, there is no need to perturb moduli spaces, and no problems with transverseness. Instead we choose co-gauge-fixing data, which is a much less disruptive thing to do, and there is no problem in making infinitely many choices of co-gauge-fixing data for the $\overline{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta)$ compatible at their boundaries, as we have shown in Theorem 6.28. Because of this, the treatment of [24] can be hugely simplified and shortened.

Manabu Akaho and the author will define Lagrangian Floer cohomology using Kuranishi cochains in [4]. Our goal here is only to whet the reader's appetite, to show that Kuranishi cochains are an elegant and economical setting for Lagrangian Floer cohomology, and so to justify all the effort we have put into developing Kuranishi (co)homology, and perhaps persuade the reader that it is worth taking the time to learn. So here we will define Lagrangian Floer cohomology only for one Lagrangian, not two, and we will work with a fixed almost complex structure J on M , without considering the effect of changing J .

In defining Lagrangian Floer cohomology, we have to consider sums involving infinitely many terms, coming from J -holomorphic discs of larger and larger area. To ensure these sums converge, we work over a ring of formal power series known as a *Novikov ring*, as in Fukaya et al. [24, Def. 6.2].

Definition 6.33. Let T and e be formal variables, graded of degree 0 and 2,

respectively. Define two *universal Novikov rings* (over $\mathbb{Q}$) by

$$\begin{aligned}\Lambda_{\text{nov}} &= \left\{ \sum_{i=0}^{\infty} a_i T^{\lambda_i} e^{\mu_i} : a_i \in \mathbb{Q}, \lambda_i \in \mathbb{R}, \mu_i \in \mathbb{Z}, \lim_{i \rightarrow \infty} \lambda_i = \infty \right\}, \\ \Lambda_{\text{nov}}^0 &= \left\{ \sum_{i=0}^{\infty} a_i T^{\lambda_i} e^{\mu_i} : a_i \in \mathbb{Q}, \lambda_i \in [0, \infty), \mu_i \in \mathbb{Z}, \lim_{i \rightarrow \infty} \lambda_i = \infty \right\}.\end{aligned}$$

Then $\Lambda_{\text{nov}}^0 \subset \Lambda_{\text{nov}}$ are $\mathbb{Q}$ -vector spaces. Define multiplications ‘ $\cdot$ ’ on $\Lambda_{\text{nov}}, \Lambda_{\text{nov}}^0$ by $(\sum_{i=0}^{\infty} a_i T^{\lambda_i} e^{\mu_i}) \cdot (\sum_{j=0}^{\infty} b_j T^{\nu_j} e^{\xi_j}) = \sum_{i,j=0}^{\infty} a_i b_j T^{\lambda_i + \nu_j} e^{\mu_i + \xi_j}$. Then $\Lambda_{\text{nov}}, \Lambda_{\text{nov}}^0$ are *commutative $\mathbb{Q}$ -algebras* with identity $1 = 1T^0 e^0$.

Write Λ_{nov}^* for either Λ_{nov} or Λ_{nov}^0 . Define *filtrations* of Λ_{nov}^* by

$$F^\lambda \Lambda_{\text{nov}}^* = \left\{ \sum_{i=0}^{\infty} a_i T^{\lambda_i} e^{\mu_i} \in \Lambda_{\text{nov}}^* : \lambda_i \geq \lambda \text{ for all } i = 0, 1, \dots \right\},$$

for $\lambda \in \mathbb{R}$. These filtrations induce *topologies* on Λ_{nov}^* , and notions of *convergence* for sequences and series. An infinite sum $\sum_{k=0}^{\infty} \alpha_k$ in Λ_{nov}^* converges in Λ_{nov}^* if and only if for all $\lambda \in \mathbb{R}$ we have $\alpha_k \in F^\lambda \Lambda_{\text{nov}}^*$ for all except finitely many k .

As T, e are graded of degrees 0, 2, we can regard $\Lambda_{\text{nov}}, \Lambda_{\text{nov}}^0$ as *graded rings*. Write $\Lambda_{\text{nov}}^{(k)}, \Lambda_{\text{nov}}^{0(k)}$ for the degree k parts of $\Lambda_{\text{nov}}, \Lambda_{\text{nov}}^0$, for $k \in \mathbb{Z}$. Then

$$\Lambda_{\text{nov}}^{(2k)} = \left\{ \sum_{i=0}^{\infty} a_i T^{\lambda_i} e^k : a_i \in \mathbb{Q}, \lambda_i \in \mathbb{R}, \lim_{i \rightarrow \infty} \lambda_i = \infty \right\}, \quad \Lambda_{\text{nov}}^{(2k+1)} = 0. \quad (172)$$

We can also consider *modules* over Λ_{nov}^* . Let $V = \bigoplus_{k \in \mathbb{Z}} V^k$ be a graded $\mathbb{Q}$ -vector space. Then $V \otimes \Lambda_{\text{nov}}^*$ is a Λ_{nov}^* -module, with grading $(V \otimes \Lambda_{\text{nov}}^*)^l = \bigoplus_{j+k=l} V^j \otimes \Lambda_{\text{nov}}^{(k)}$, and filtration $F^\lambda(V \otimes \Lambda_{\text{nov}}^*) = V \otimes F^\lambda \Lambda_{\text{nov}}^*$ for $\lambda \in \mathbb{R}$. Write $V \hat{\otimes} \Lambda_{\text{nov}}^*$ for the *completion* of $V \otimes \Lambda_{\text{nov}}^*$ with respect to this filtration. Then $V \hat{\otimes} \Lambda_{\text{nov}}^*$ has a topology, and an infinite sum $\sum_{k=0}^{\infty} v_k$ converges in $V \hat{\otimes} \Lambda_{\text{nov}}^*$ if for all $\lambda \in \mathbb{R}$ we have $v_k \in F^\lambda(V \hat{\otimes} \Lambda_{\text{nov}}^*)$ for all except finitely many k .

If L is a manifold $KC^*(L; \Lambda_{\text{nov}}^*) = KC^*(L; \mathbb{Q}) \otimes \Lambda_{\text{nov}}^*$. Write $\widehat{KC}^*(L; \Lambda_{\text{nov}}^*)$ for the completion $KC^*(L; \mathbb{Q}) \hat{\otimes} \Lambda_{\text{nov}}^*$. We consider the grading on $\widehat{KC}^*(L; \Lambda_{\text{nov}}^*)$ to be the combination of those on $KC^*(L; \mathbb{Q})$ and Λ_{nov}^* .

Next we define *gapped filtered A_∞ algebras*, following Fukaya et al. [24, §7.2].

Definition 6.34. A *gapped filtered A_∞ algebra* $(A \otimes \Lambda_{\text{nov}}^*, \mathfrak{m})$ consists of:

- (a) A graded $\mathbb{Q}$ -vector space $A = \bigoplus_{d \in \mathbb{Z}} A^d$;
- (b) Graded Λ_{nov}^* -multilinear maps $\mathfrak{m}_k : (A \hat{\otimes} \Lambda_{\text{nov}}^*)^{\overbrace{\times}^{k \text{ copies}}} \rightarrow A \hat{\otimes} \Lambda_{\text{nov}}^*$ for $k = 0, 1, 2, \dots$, of degree $+1$. Write $\mathfrak{m} = (\mathfrak{m}_k)_{k \geq 0}$.

These must satisfy the following conditions:

- (i) there exists a subset $\mathcal{G} \subset [0, \infty) \times \mathbb{Z}$, closed under addition, such that $\mathcal{G} \cap (\{0\} \times \mathbb{Z}) = \{(0, 0)\}$ and $\mathcal{G} \cap ([0, C] \times \mathbb{Z})$ is finite for any $C \geq 0$, and the maps $\mathfrak{m}_k$ for $k \geq 0$ may be written $\mathfrak{m}_k = \sum_{(\lambda, \mu) \in \mathcal{G}} T^\lambda e^\mu \mathfrak{m}_k^{\lambda, \mu}$, for unique $\mathbb{Q}$ -multilinear maps $\mathfrak{m}_k^{\lambda, \mu} : \overbrace{A \times \dots \times A}^{k \text{ copies}} \rightarrow A$ graded of degree $1 - 2\mu$. When $k = 0$, we take $\mathfrak{m}_0 \in (A \otimes \Lambda_{\text{nov}}^0)^{(1)}$ and $\mathfrak{m}_0^{\lambda, \mu} \in A^{1-2\mu}$;
- (ii) $\mathfrak{m}_0^{0,0} = 0$, in the notation of (i); and

- (iii) call $a \in A \otimes \Lambda_{\text{nov}}^0$ *pure* if $a \in (A \otimes \Lambda_{\text{nov}}^0)^{(d)} \setminus \{0\}$ for some $d \in \mathbb{Z}$, and then define the *degree* of a to be $\deg a = d$. Then we require that for all $k \geq 0$ and all pure $a_1, \dots, a_k$ in $A \otimes \Lambda_{\text{nov}}^0$, we have

$$\sum_{\substack{i, k_1, k_2: 1 \leq i \leq k_1, \\ k_2 \geq 0, k_1 + k_2 = k+1}} (-1)^{\sum_{l=1}^{i-1} \deg a_l} \mathfrak{m}_{k_1}(a_1, \dots, a_{i-1}, \mathfrak{m}_{k_2}(a_i, \dots, a_{i+k_2-1}), a_{i+k_2}, \dots, a_k) = 0. \quad (173)$$

Equation (173) may be rewritten in terms of the $\mathfrak{m}_k^{\lambda, \mu}$ as follows: for all $k \geq 0$, all $(\lambda, \mu) \in \mathcal{G}$ and all pure $a_1, \dots, a_k$ in A , we have

$$\sum_{\substack{i, k_1, k_2, \lambda_1, \lambda_2, \mu_1, \mu_2: 1 \leq i \leq k_1, k_2 \geq 0, \\ k_1 + k_2 = k+1, \lambda_1 + \lambda_2 = \lambda, \mu_1 + \mu_2 = \mu}} (-1)^{\sum_{l=1}^{i-1} \deg a_l} \mathfrak{m}_{k_1}^{\lambda_1, \mu_1}(a_1, \dots, a_{i-1}, \mathfrak{m}_{k_2}^{\lambda_2, \mu_2}(a_i, \dots, a_{i+k_2-1}), a_{i+k_2}, \dots, a_k) = 0. \quad (174)$$

A gapped filtered A_∞ algebra $(A \hat{\otimes} \Lambda_{\text{nov}}^*, \mathfrak{m})$ is called *strict* if $\mathfrak{m}_0 = 0$. Then (iii) implies that $\mathfrak{m}_1 \circ \mathfrak{m}_1 = 0$, so $(A \hat{\otimes} \Lambda_{\text{nov}}^*, \mathfrak{m}_1)$ is a complex of Λ_{nov}^* -modules, and we can form its *cohomology* $H^*(A \hat{\otimes} \Lambda_{\text{nov}}^*, \mathfrak{m}_1)$.

To define Lagrangian Floer cohomology we will need *strict* gapped filtered A_∞ algebras. *Bounding cochains* are a method of modifying gapped filtered A_∞ algebras to make them strict, introduced by Fukaya et al. [24, §5.7, §11].

Definition 6.35. Let $(A \hat{\otimes} \Lambda_{\text{nov}}^*, \mathfrak{m})$ be a gapped filtered A_∞ algebra, and suppose $b \in F^\lambda(A \hat{\otimes} \Lambda_{\text{nov}}^*)^{(0)}$ for some $\lambda > 0$. Define graded Λ_{nov}^* -multilinear maps $\mathfrak{m}_k^b : (A \hat{\otimes} \Lambda_{\text{nov}}^*)^{\overbrace{\times \dots \times}^k \text{copies}} \rightarrow A \hat{\otimes} \Lambda_{\text{nov}}^*$ for $k = 0, 1, \dots$, of degree $+1$, by

$$\mathfrak{m}_k^b(a_1, \dots, a_k) = \sum_{n_0, \dots, n_k \geq 0} \mathfrak{m}_{k+n_0+\dots+n_k}(b, \dots, b, a_1, b, \dots, b, a_2, b, \dots, b, \dots, b, \dots, b, a_k, b, \dots, b).$$

This converges as $b \in F^\lambda(A \hat{\otimes} \Lambda_{\text{nov}}^*)$ for $\lambda > 0$. Write $\mathfrak{m}^b = (\mathfrak{m}_k^b)_{k \geq 0}$. We call b a *bounding cochain* for $(A \otimes \Lambda_{\text{nov}}^0, \mathfrak{m})$ if $\mathfrak{m}_0^b = 0$, that is, if

$$\sum_{k \geq 0} \mathfrak{m}_k(b, \dots, b) = 0. \quad (175)$$

This is the *Maurer–Cartan equation*, or *Batalin–Vilkovisky master equation*.

It is then easy to prove [24, Prop. 11.10]:

Lemma 6.36. *In Definition 6.35, $(A \hat{\otimes} \Lambda_{\text{nov}}^*, \mathfrak{m}^b)$ is a gapped filtered A_∞ algebra, which is strict if and only if b is a bounding cochain.*

Thus, if b is a bounding cochain then $(A \otimes \Lambda_{\text{nov}}^0, \mathfrak{m}_1^b)$ is a complex, and we may form its cohomology $H^*(A \hat{\otimes} \Lambda_{\text{nov}}^*, \mathfrak{m}_1^b)$. We will now show that in the situation of §6.4–§6.5, we can define a *gapped filtered A_∞ algebra structure* on the shifted Kuranishi cochains $\widehat{KC}^*(L; \Lambda_{\text{nov}}^*)[1]$ of L .

Definition 6.37. Suppose (M, ω) is a compact symplectic manifold, L a compact, embedded Lagrangian submanifold in M , and J an almost complex structure on M compatible with ω . Fix a relative spin structure on (M, L) , so

that $\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$ for $l \geq 1$ are compact, oriented Kuranishi spaces with corners by Theorem 6.21 and “Theorems” 6.23 and 6.25. Note that the relative spin structure includes an orientation on L . Choose co-gauge-fixing data $\mathbf{C}_l(M, L, J, \beta)$ for $(\bar{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta), \mathbf{ev}_1 \times \cdots \times \mathbf{ev}_l)$ for all $l \geq 1$ and β in $H_2(M, L; \mathbb{Z})$ by the first part of Theorem 6.28. We do not require $\mathbf{C}_l(M, L, J, \beta)$ to be $\mathbb{Z}_l$ -invariant, and we do not choose $\mathbf{C}_0(M, L, J, \beta)$. Define $\mathfrak{M}_l(\beta) \in KC(L^l; R)$ for all $l \geq 1$ and $\beta \in H_2(M, L; \mathbb{Z})$ as in Definition 6.29. Then Corollary 6.30 shows that equation (169) holds for all $l \geq 1$ and β .

Write $KC^*(L; \mathbb{Q})[1]$ for the Kuranishi cochains *with grading shifted by one*, that is, $(KC^*(L; \mathbb{Q})[1])^k = KC^{k+1}(L; \mathbb{Q})$, and similarly for $\widehat{KC}^*(L; \Lambda_{\text{nov}}^*)[1]$. We will define a *gapped filtered A_∞ algebra* $(\widehat{KC}^*(L; \Lambda_{\text{nov}}^*)[1], \mathfrak{m})$ on the shifted cochains $\widehat{KC}^*(L; \Lambda_{\text{nov}}^*)[1] = KC^*(L; \mathbb{Q})[1] \hat{\otimes} \Lambda_{\text{nov}}^*$. Define $\mathcal{G} \subset [0, \infty) \times \mathbb{Z}$ by

$$\begin{aligned} \mathcal{G} = \{(\lambda, \mu) \in [0, \infty) \times \mathbb{Z} : & \text{there exist } \beta_1, \dots, \beta_k \in H_2(M, L; \mathbb{Z}) \text{ with} \\ & \bar{\mathcal{M}}_3^{\text{ma}}(M, L, J, \beta_i) \neq \emptyset \text{ for } i = 1, \dots, k, \text{ and} \\ & \lambda = [\omega] \cdot (\beta_1 + \cdots + \beta_k), \mu = \tfrac{1}{2}\mu_L(\beta_1 + \cdots + \beta_k)\}. \end{aligned}$$

Define $\mathfrak{m}_k^{\lambda, \mu} : KC^*(L; \mathbb{Q})[1] \times \cdots \times \widehat{KC}^*(L; \mathbb{Q})[1] \rightarrow KC^*(L; \mathbb{Q})[1]$ for all $k \geq 0$ and $(\lambda, \mu) \in \mathcal{G}$ by

$$\begin{aligned} \mathfrak{m}_1^{0,0}(f_1) &= (-1)^n df_1, & \mathfrak{m}_k^{\lambda, \mu}(f_1, \dots, f_k) &= \\ & \sum_{\substack{\beta \in H_2(M, L; \mathbb{Z}): \\ [\omega] \cdot \beta = \lambda, \mu_L(\beta) = 2\mu, \\ \bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta) \neq \emptyset}} (\cdots ((\mathfrak{M}_{k+1}(\beta) *_{2, L, 1} f_1) *_{3, L, 1} f_2) \cdots) *_{k+1, L, 1} f_k), & (176) \\ & (k, \lambda, \mu) \neq (1, 0, 0). \end{aligned}$$

Following the definitions through shows that for $(k, \lambda, \mu) \neq (1, 0, 0)$ on generators $[X_i, \mathbf{f}_i, \mathbf{C}_i]$ of $KC^*(L; \mathbb{Q})$ this gives

$$\begin{aligned} \mathfrak{m}_k^{\lambda, \mu}([X_1, \mathbf{f}_1, \mathbf{C}_1], \dots, [X_k, \mathbf{f}_k, \mathbf{C}_k]) &= \\ \sum_{\substack{\beta \in H_2(M, L; \mathbb{Z}): \\ [\omega] \cdot \beta = \lambda, \mu_L(\beta) = 2\mu, \\ \bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta) \neq \emptyset}} & [\bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta) \times_{\mathbf{ev}_2 \times \cdots \times \mathbf{ev}_{k+1, L^k}, \mathbf{f}_1 \times \cdots \times \mathbf{f}_k} (X_1 \times \cdots \times X_k), \\ & \mathbf{ev}_1 \circ \pi \bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta), \mathbf{C}_{\mathbf{C}_l(M, L, J, \beta), \mathbf{C}_1, \dots, \mathbf{C}_k}], \end{aligned} \quad (177)$$

where $\mathbf{C}_{\mathbf{C}_l(M, L, J, \beta), \mathbf{C}_1, \dots, \mathbf{C}_k}$ is co-gauge-fixing data for $(\bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta) \times_{L^k} (X_1 \times \cdots \times X_k), \mathbf{ev}_1 \circ \pi \bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta))$ obtained as a fibre product of the co-gauge-fixing data $\mathbf{C}_l(M, L, J, \beta), \mathbf{C}_1, \dots, \mathbf{C}_k$ on the $k+1$ factors by k applications of the product $*_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j}$ of Definition 6.27. Suppose $[X_i, \mathbf{f}_i, \mathbf{C}_i]$ has degree a_i in $KC^*(L; \mathbb{Q})[1]$, so that $[X_i, \mathbf{f}_i, \mathbf{C}_i] \in KC^{a_i+1}(L; \mathbb{Q})$, and $\text{vdim } X_i = n - a_i - 1$, where $n = \dim L$. Then using (157) we find that the fibre product in (177) has dimension $2\mu + n - 2 - a_1 - \cdots - a_k$, so (177) has degree $1 - 2\mu + a_1 + \cdots + a_k$ in $KC^*(L; \mathbb{Q})[1]$, and $\mathfrak{m}_k^{\lambda, \mu}$ is graded of degree $1 - 2\mu$.

Now define $\mathfrak{m}_k : \widehat{KC}^*(L; \Lambda_{\text{nov}}^*)[1] \times \cdots \times \widehat{KC}^*(L; \Lambda_{\text{nov}}^*)[1] \rightarrow \widehat{KC}^*(L; \Lambda_{\text{nov}}^*)[1]$ for $k = 0, 1, 2, \dots$ by $\mathfrak{m}_k = \sum_{(\lambda, \mu) \in \mathcal{G}} T^\lambda e^\mu \mathfrak{m}_k^{\lambda, \mu}$. As the $\mathfrak{m}_k^{\lambda, \mu}$ are graded of degree

$1 - 2\mu$, $\mathfrak{m}_k$ is graded of degree $+1$. Using (169) and (176) one can show that the $\mathfrak{m}_k^{\lambda, \mu}$ satisfy (174), and thus the $\mathfrak{m}_k$ satisfy (173). Write $\mathfrak{m} = (\mathfrak{m}_k)_{k \geq 0}$. Then $(\widehat{KC}^*(L; \Lambda_{\text{nov}}^*)[1], \mathfrak{m})$ is a *gapped filtered A_∞ algebra*.

Let b be a *bounding cochain* for $(\widehat{KC}^*(L; \Lambda_{\text{nov}}^*)[1], \mathfrak{m})$, and define $\mathfrak{m}^b$ as in Definition 6.35. Then $\mathfrak{m}_1^b : \widehat{KC}^*(L; \Lambda_{\text{nov}}^*) \rightarrow \widehat{KC}^*(L; \Lambda_{\text{nov}}^*)$ is graded of degree $+1$ with $\mathfrak{m}_1^b \circ \mathfrak{m}_1^b = 0$. Define the *Lagrangian Floer cohomology* $HF^*(L, b; \Lambda_{\text{nov}}^*)$ to be the cohomology of $(\widehat{KC}^*(L; \Lambda_{\text{nov}}^*), \mathfrak{m}_1^b)$. Note that we drop the shift in gradings, so that the grading of $HF^*(L, b; \Lambda_{\text{nov}}^*)$ agrees with that of ordinary cohomology. We can think of $HF^*(L, b; \Lambda_{\text{nov}}^*)$ as a ‘quantum deformation’ of Kuranishi cohomology $KH^*(L; \Lambda_{\text{nov}}^*)$.

The important thing is that $HF^*(L, b; \Lambda_{\text{nov}}^*)$ is *independent of all the arbitrary choices involved up to canonical isomorphism*, including the choice of almost complex structure J . (It does depend on the bounding cochain b , so to make sense of this we must introduce a notion of equivalence between bounding cochains b for J , and bounding cochains b' for another almost complex structure J' .) Thus, Lagrangian Floer cohomology $HF^*(L, b; \Lambda_{\text{nov}}^*)$ basically depends only on (M, ω) and L , as for Gromov–Witten invariants in Theorem 6.10. Also, the gapped filtered A_∞ algebra above is independent of choices up to A_∞ quasi-isomorphism.

Now compare Definition 6.37 with the construction of gapped filtered A_∞ algebras for Lagrangian submanifolds in Fukaya et al. [24, Chapter 3]. Briefly, they construct a countably generated subcomplex $\mathcal{X}$ of the complex of smooth singular chains $C_*^{\text{si}}(L; \Lambda_{\text{nov}}^*)$ on L with values in Λ_{nov}^* , and multilinear operators $\mathfrak{m}_k : \mathcal{X}^{\otimes k} \rightarrow \mathcal{X}$ satisfying the A_∞ relations (173). Roughly speaking, $\mathfrak{m}_k$ is a sum over all β of intersections with virtual cycles for moduli spaces $\bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta)$.

The details of the construction, however, are horrible. For each Kuranishi space $\bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta)$ we must choose a virtual moduli chain, an abstract perturbation, which must be compatible at the boundary $\partial \bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta)$ with the choices of virtual chains for other $\bar{\mathcal{M}}_{l+1}^{\text{ma}}(M, L, J, \gamma)$ under (159). Worse, one virtual moduli chain for $\bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta)$ is not enough, because of transversality problems: we must choose one for each k -tuple $(P_1, \dots, P_k)$ in $\mathcal{X}^k$. Worse still, it is not actually possible to make all these choices consistently for all $k \geq 0$ and $(P_1, \dots, P_k)$, but only for finitely generated $\mathcal{X}$ and bounded k .

Therefore Fukaya et al. construct $A_{N,K}$ -algebras $\mathcal{X}_{N,K}$ for increasingly large N, K , prove homotopy equivalences between them, and finally use an abstract algebraic construction to prove existence of the (now non-geometric) A_∞ -algebra they want. With Kuranishi cohomology we can avoid all this, and go directly to a geometric A_∞ -algebra structure on the full Kuranishi cochains $\widehat{KC}^*(L; \Lambda_{\text{nov}}^*)$.

If there are no nonconstant J -holomorphic $\mathbb{CP}^1$'s in M then we can use Theorem 6.31 to define Lagrangian Floer cohomology using *effective* Kuranishi cochains, and the construction then works with Novikov rings *over the integers* $\mathbb{Z}$, or over an arbitrary commutative ring R , rather than just over $\mathbb{Q}$.

These ideas are also relevant to the construction of the *Fukaya category* $\text{Fuk}(M, \omega)$ of a general symplectic manifold (M, ω) , as in [22, 24, 67, 69]. The

derived Fukaya category $D^b\mathrm{Fuk}(M, \omega)$ is an essential part of Kontsevich's Homological Mirror Symmetry programme [44], and so is a very important piece of mathematics. Roughly speaking, the objects of $\mathrm{Fuk}(M, \omega)$ should be pairs (L, b) of a Lagrangian L and a bounding cochain b for L , and the morphisms between such pairs in $D^b\mathrm{Fuk}(M, \omega)$ are given by Lagrangian Floer cohomology. The author believes that by using Kuranishi cochains as above, it should be possible to write down a reasonably clean and simple model for $\mathrm{Fuk}(M, \omega)$.

6.7 Open Gromov–Witten invariants

As in §6.2, (closed) Gromov–Witten invariants ‘count’ J -holomorphic curves without boundary in a symplectic manifold (M, ω) , and are independent of the choice of almost complex structure J . In a similar way, if L is Lagrangian submanifold of (M, ω) we can attempt to define *open Gromov–Witten invariants* which ‘count’ J -holomorphic curves in M with boundary in L , and hope to make them independent of J . This is a more difficult problem than the closed case, since as in §6.1 and §6.4 the moduli spaces have no boundary in the closed case, but have boundary and corners in the open case.

Quite a lot of work has already been done on open Gromov–Witten invariants. In String Theory, it is expected that open Gromov–Witten invariants should exist and play a rôle, and String Theorists can even compute them in examples; see for example Ooguri and Vafa [58], Aganagic and Vafa [3], Aganagic, Klemm and Vafa [2], and Graber and Zaslow [30], and others. Some mathematics papers on open Gromov–Witten invariants are Katz and Liu [42], Liu [51], Li and Song [47], Welschinger [76, 77], Cho [14, 15], Solomon [73], and Pandharipande, Solomon and Walcher [61].

Despite these, there does not yet seem to be a satisfactory definition of open Gromov–Witten invariants (by ‘satisfactory’ I really mean that the invariants should be defined using any almost complex structure J on M compatible with ω , and give an answer independent of J). It is clear from these references (see Aganagic et al. [2, §5], and computations by Cho [15], for example), that just fixing $(M, \omega), L$ is not enough to get open Gromov–Witten invariants independent of J , one needs some extra assumptions or extra data.

Liu [51] defines open Gromov–Witten invariants for the case in which there is a $U(1)$ -action on $(M, \omega), L$, and J is $U(1)$ -invariant. Welschinger [76, 77], Cho [14] and Solomon [73] define open Gromov–Witten invariants in the case in which $\dim M$ is 4 or 6, and L is the fixed point set of an involution $\sigma : M \rightarrow M$, which is antiholomorphic with respect to J ; these invariants are calculated when M is the quintic Calabi–Yau 3-fold by Pandharipande et al. [61].

We now present a part of the author’s solution to the problem of defining open Gromov–Witten invariants; the complete picture will be explained in [39]. For genus zero open Gromov–Witten invariants, the key idea is this: we need to fix some extra data to define open Gromov–Witten invariants independent of J , and *the right kind of extra data is a choice of bounding cochain b for Lagrangian Floer cohomology of L* , in the sense of §6.6. The open Gromov–Witten invariants will then depend on (L, b) , and they will be independent of

the choice of J in the same sense that Lagrangian Floer cohomology is. To correct for the $\bar{\mathcal{M}}_{0,1}(M, J, \beta')$ terms in (162) we need to assume $[L] = 0$ in $H_n(M; \mathbb{Q})$, and choose an $(n+1)$ -chain B in M with $\partial B = L$. In [39] we will also explain how to define higher-genus open Gromov–Witten invariants under stronger assumptions, including requiring L to be a $\mathbb{Q}$ -homology sphere.

Warning: *most of the signs below are probably wrong, but I'll fix them in the final version.*

We work in the situation of §6.4–§6.6. Let b be a bounding cochain for $(\widehat{KC}^*(L; \Lambda_{\text{nov}}^0)[1], \mathfrak{m})$. Then b satisfies (175), which by (176) means that

$$(-1)^n db + \sum_{k \geq 0, \beta \in H_2(M, L; \mathbb{Z}) : \bar{\mathcal{M}}_{k+1}^{\text{ma}}(M, L, J, \beta) \neq \emptyset} T^{[\omega] \cdot \beta} e^{\mu_L(\beta)/2} (\cdots (\mathfrak{M}_{k+1}(\beta) *_{2,L,1} b) *_{3,L,1} b) \cdots) *_{k+1,L,1} b = 0. \quad (178)$$

Let $\text{inc} : L \rightarrow M$ be the inclusion. Define gauge-fixing data $\mathbf{G}_L$ for (L, inc) to be $\mathbf{G}_L = (\mathbf{I}_L, \boldsymbol{\eta}_L, G_L)$, where $\mathbf{I}_L$ has indexing set $I = \{n\}$, one Kuranishi neighbourhood $(L, L, 0, \text{id}_L)$, and map $\text{inc} : L \rightarrow M$, and $\boldsymbol{\eta} = (\eta_n, \eta_n^n)$ with $\eta_n \equiv 1 \equiv \eta_n^n$, and $G_L : L \rightarrow P$ is the function used in the definition of the products $*_{\mathbf{f}_i, L, \tilde{\mathbf{f}}_j}$ in Definition 6.27. As L is compact, oriented and without boundary, we have $[L, \text{inc}, \mathbf{G}_L] \in KC_n(M; \mathbb{Q})$ with $\partial[L, \text{inc}, \mathbf{G}_L] = 0$, so the homology class $[[L, \text{inc}, \mathbf{G}_L]]$ is defined in $KH_n(M; \mathbb{Q})$. We have $\Pi_{\text{si}}^{\text{Kh}}([L]) = [[L, \text{inc}, \mathbf{G}_L]]$, where $[L] \in H_n^{\text{si}}(M; \mathbb{Q})$ is the homology class of L in M , and $\Pi_{\text{si}}^{\text{Kh}} : H_n^{\text{si}}(M; \mathbb{Q}) \rightarrow KH_n(M; \mathbb{Q})$ is an isomorphism by Corollary 4.10.

Suppose that $[L] = 0$ in $H_n^{\text{si}}(M; \mathbb{Q})$. Note that we don't require L to be connected, it can be the union of several compact lagrangian submanifolds whose homology classes sum to zero. Then $[[L, \text{inc}, \mathbf{G}_L]] = 0$ in $KH_n(M; \mathbb{Q})$, so we can choose a Kuranishi chain $B \in KC_{n+1}(M; \mathbb{Q})$ with $\partial B = [L, \text{inc}, \mathbf{G}_L]$. Let $\beta' \in H_2(M; \mathbb{Z})$, and form the cap product

$$C \cap [\bar{\mathcal{M}}_{0,1}(M, J, \beta'), \mathbf{ev}_1, C_{0,1}(M, J, \beta')] \in KC_*(M; \mathbb{Q}),$$

where $C_{0,1}(M, J, \beta')$ is the co-gauge-fixing data for $(\bar{\mathcal{M}}_{0,1}(M, J, \beta'), \mathbf{ev}_1)$ chosen in Theorem 6.28. Writing $\pi : M \rightarrow \{0\}$ for the projection, we apply the pushforward $\pi_* : KC_*(M; \mathbb{Q}) \rightarrow KC_*(\{0\}; \mathbb{Q})$ to define

$$\pi_*(C \cap [\bar{\mathcal{M}}_{0,1}(M, J, \beta'), \mathbf{ev}_1, C_{0,1}(M, J, \beta')]) \in KC_*(\{0\}; \mathbb{Q}). \quad (179)$$

Now over a single point $\{0\}$, strongly smooth $\mathbf{f} : X \rightarrow \{0\}$ coincide with strong submersions, and orientations for X coincide with coorientations for $(X, \mathbf{f})$, and gauge-fixing data and co-gauge-fixing for $(X, \mathbf{f})$ coincide. Thus there is a natural identification $KC_k(\{0\}; \mathbb{Q}) = KC^{-k}(\{0\}; \mathbb{Q})$, which identifies ∂ on $KC_*(\{0\}; \mathbb{Q})$ with d on $KC^*(\{0\}; \mathbb{Q})$. Using this we regard (179) as an element of $KC^*(\{0\}; \mathbb{Q})$. Then using $\partial B = [L, \text{inc}, \mathbf{G}_L]$ and comparing the way $G_L : L \rightarrow P$ appears in $\mathbf{G}_L$ and the product $*_{\mathbf{ev}_1, L, \text{id}_L}$, we can show that

$$\begin{aligned} d(\pi_*(C \cap [\bar{\mathcal{M}}_{0,1}(M, J, \beta'), \mathbf{ev}_1, C_{0,1}(M, J, \beta')])) &= \\ [\bar{\mathcal{M}}_{0,1}(M, J, \beta') \times_{\mathbf{ev}_1, M, \text{inc}} L, \pi, & \\ \text{inc}^*(C_{0,1}(M, J, \beta')) *_{\mathbf{ev}_1, L, \text{id}_L} [L, \text{id}_L, C_L]] &= \mathfrak{N}_0(\beta') \end{aligned} \quad (180)$$

in $KC_*(\{0\}; \mathbb{Q})$, by (168). We will use this to cancel the $\mathfrak{N}_0(\beta')$ terms in (170).

Now define an expression $OGW(M, L)$ in $\widehat{KC}^*(\{0\}; \Lambda_{\text{nov}}^0)$ by

$$\begin{aligned} OGW(M, L) = & \sum_{\beta \in H_2(M, L; \mathbb{Z})} T^{[\omega] \cdot \beta} e^{\mu_L(\beta)/2} \mathfrak{M}_0(\beta) + \\ & \sum_{\substack{l \geq 1, \beta \in H_2(M, L; \mathbb{Z}): \\ \mathcal{M}_l^{\text{ma}}(M, L, J, \beta) \neq \emptyset}} T^{[\omega] \cdot \beta} e^{\mu_L(\beta)/2} \frac{2-l}{l} (\cdots (\mathfrak{M}_l(\beta) *_{1, L, 1} b) *_{2, L, 1} b) \cdots) *_{l, L, 1} b) \\ & - \sum_{\substack{\beta' \in H_2(L; \mathbb{Z}): \\ \mathcal{M}_{0,1}(M, J, \beta') \neq \emptyset}} T^{[\omega] \cdot \beta'} e^{c_1(M) \cdot \beta'} \pi_* (C \cap [\bar{\mathcal{M}}_{0,1}(M, J, \beta'), \mathbf{ev}_1, C_{0,1}(M, J, \beta')]). \end{aligned} \quad (181)$$

Using (169), (170), (178) and (180), we compute that $d(OGW(M, L)) = 0$.

Thus we may form $[OGW(M, L)] \in \widehat{KH}^*(\{0\}; \Lambda_{\text{nov}}^0) = KH^*(\{0\}; \mathbb{Q}) \hat{\otimes} \Lambda_{\text{nov}}^0$.

Identifying $KH^*(\{0\}; \mathbb{Q}) \cong H_{\text{cs}}^*(\{0\}; \mathbb{Q}) \cong \mathbb{Q}$ we have $\widehat{KH}^*(\{0\}; \Lambda_{\text{nov}}^0) \cong \Lambda_{\text{nov}}^0$. Computing dimensions shows that $OGW(M, L)$ is graded of degree $3-n$, so $[OGW(M, L)] \in \Lambda_{\text{nov}}^{0(3-n)}$. When n is even we have $\Lambda_{\text{nov}}^{0(3-n)}$ by (172), so $[OGW(M, L)] = 0$. When n is odd, we may write

$$[OGW(M, L)] = \sum_{\lambda \geq 0} OGW(M, L, \lambda) T^\lambda e^{(3-n)/2}, \quad (182)$$

with $OGW(M, L, \lambda) \in \mathbb{Q}$.

We interpret these rational numbers $OGW(M, L, \lambda)$ as *open Gromov–Witten invariants* which ‘count’ all stable J -holomorphic discs in M with boundary in L in classes $\beta \in H_2(M, L; \mathbb{Z})$ with $[\omega] \cdot \beta = \lambda$ and $\mu_L(\beta) = 3-n$. The meaning of the restriction $\mu_L(\beta) = 3-n$ is that by (157) this is the condition for $\text{vdim } \bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta) = 0$, and $[OGW(M, L)]$ ‘counts’ only moduli spaces $\bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta)$ of virtual dimension zero. When n is even all $\bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta)$ have odd virtual dimension, so there are no $\bar{\mathcal{M}}_0^{\text{ma}}(M, L, J, \beta)$ with virtual dimension zero. Probably the most interesting case is that of a graded Lagrangian L in a Calabi–Yau 3-fold (M, ω) , when $\mu_L(\beta) = 3-n$ holds automatically.

We will show in [39] that the $OGW(M, L, \lambda)$ are *independent of the almost complex structure J* , in the same sense that Lagrangian Floer cohomology is. They do depend on the choice of bounding cochain b . They depend on the chain B with $\partial B = [L, \text{inc}, \mathbf{G}_L]$ only through the relative homology class of B in $H_{n+1}(M, L; \mathbb{Q})$. That is, if we replace B by $B' = B + \partial C$ for $C \in KC_{n+2}(M; \mathbb{Q})$, then the $OGW(M, L, \lambda)$ are unchanged.

The fact that we can only count discs with fixed area and Maslov index, rather than counting discs in a fixed $\beta \in H_2(M, L; \mathbb{Z})$, is because we are using a bounding cochain b defined using a Novikov ring Λ_{nov}^0 which only keeps track of the area and Maslov index of curves. As we explain in [39], we can refine the invariants if we either work over a refined Novikov ring, or make stronger assumptions on the topology of M, L .

6.8 Conclusions

There is a large area of symplectic geometry which uses moduli spaces of J -holomorphic curves in a symplectic manifold (M, ω) to define some homological structure, and then shows that this structure is independent of the choice of almost complex structure J , up to some kind of isomorphism, so that the structure basically depends only on (M, ω) . This includes Gromov–Witten invariants and Lagrangian Floer cohomology, as discussed above, and also contact homology [20], Symplectic Field Theory [21], Fukaya categories [22, 67, 69], and symplectic cohomology [68].

To carry out such a programme, one has to solve four problems:

- (a) Define a geometric structure which is the model structure for the moduli spaces of J -holomorphic curves we are interested in – for example, Fukaya–Ono Kuranishi spaces. This structure should be powerful enough that analogues of differential-geometric ideas for manifolds such as orientations, smooth maps, submersions, fibre products work for it.
- (b) Show that the moduli spaces of J -holomorphic curves we wish to study carry the structure in (a), and that natural maps of these moduli spaces are ‘smooth’. Prove identities between these structures, such as the boundary formulae (159) and (162).
- (c) Use this geometric structure to construct *virtual cycles* or *virtual chains* for the moduli spaces, in some (co)chain complex. Translate relationships between moduli spaces into algebraic identities upon their virtual chains.
- (d) Draw some interesting geometrical conclusions — for instance, define Gromov–Witten invariants, prove the Arnold Conjecture, and so on.

In this book we have developed new tools for solving problem (c). Previous virtual cycle or chain constructions for solving problem (c) in symplectic geometry have been given by Fukaya and Ono [24, 25], Li and Tian [49], Liu and Tian [52], Lu and Tian [53], McDuff [55], Ruan [65], Siebert [70], and in the context of their theory of polyfolds Hofer, Wysocki and Zehnder [37]. All these other constructions involve *perturbing the moduli space* in some way.

Defining virtual chains and cycles using Kuranishi (co)chains is very simple, and it eliminates perturbation of moduli spaces, and problems with transversality. Relationships between moduli spaces can often be translated into algebraic identities between Kuranishi (co)chains in a very direct way, as we saw in passing from “Theorem” 6.26 to Corollary 6.30, and these identities are likely to hold exactly, rather than in some weak homotopy sense owing to errors introduced during the virtual cycle construction.

Also, for moduli spaces without boundary, Kuranishi (co)bordism classes are in effect a kind of virtual cycle, and so a new kind of solution to problem (c). Kuranishi (co)bordism groups are huge, much larger than the corresponding (co)homology groups. This has the disadvantage that they it is probably not feasible to compute them, but the advantage that they contain more information than (co)homology, and are also useful tools for studying integrality questions.

We have based our theory upon Fukaya and Ono’s solution to problem (a), that is, their notion of Kuranishi spaces, with some modifications of our own. For our applications above we have simply assumed the solution to problem (b), when in §6.1 and §6.4 we quoted results of Fukaya et al. [24, 25] on existence of Kuranishi structures on the moduli spaces $\overline{\mathcal{M}}_{g,m}(M, J, \beta)$, $\overline{\mathcal{M}}_l^{\text{ma}}(M, L, J, \beta)$.

Both of these might be seen as a weakness. There are other solutions to problem (a) in the literature, notably the theory of *polyfolds* due to Hofer, Wysocki and Zehnder [33–37]. If the theory of polyfolds (or some other alternative) becomes widely adopted, then our Kuranishi space framework might come to seem obsolete. Also, as discussed in Remark 6.5, it is not clear to the author that the proofs of the solution to problem (b) we assume are complete, and rigorous in every detail, and some of the proofs are based on weaker definitions of Kuranishi space than the one we use. In their polyfold programme, Hofer et al. aim to give a fully rigorous treatment of problem (b).

In fact this is not a weakness, because a polyfold structure can be truncated to a Kuranishi structure in a simple way: Hofer [34, p. 2-3] says:

‘The work to build a Kuranishi structure or to build a polyfold structure (which is a finer structure) is approximately the same and seems to have more or less the same ingredients. The polyfold structure captures not only more information but comes with a strikingly easier formalism. There is in fact a trivial functor from the polyfold Fredholm theory into the Kuranishi structures.’

Kuranishi structures are probably more-or-less the weakest feasible solution to problem (a), because they are built out of finite-dimensional data, that is, Kuranishi neighbourhoods (V_p, E_p, s_p, ψ_p) . Other solutions such as polyfolds are generally built from infinite-dimensional ingredients, and are much richer structures, with more information; one passes to Kuranishi structures by forgetting much of this information. Thus, one can use the theory of polyfolds to solve problems (a) and (b), and then truncate to Kuranishi structures, and use our Kuranishi (co)homology or Kuranishi (co)bordism to solve problems (c) and (d). This gives an alternative to the virtual cycles for polyfolds defined by Hofer et al. [37], which work by perturbing the polyfold structure using $\mathbb{Q}$ -weighted multisections in a similar way to Fukaya and Ono [25].

A Tent functions

We will now define and study *tent functions*, which are the principal tools we will use in the proofs of Theorems 4.8 and 4.9 in Appendices B and C. Here is the rough idea. For S a set, write $F(S)$ for the set of finite, nonempty subsets of S . Let X be a compact, oriented n -manifold with g-corners. A *tent function* on X is a function $T : X \rightarrow F([1, \infty))$ which locally near $p \in X$ may be written $T(p) = \{f_1(p), \dots, f_k(p)\}$ for smooth functions $f_1, \dots, f_k : X \rightarrow [1, \infty)$ called the *branches* of T , such that $f_i - f_j : T \rightarrow \mathbb{R}$ is a submersion near $(f_i - f_j)^{-1}(0)$

for $1 \leq i < j \leq k$, and more complicated transversality conditions hold when several f_i are equal in X .

Then $\min T : X \rightarrow [1, \infty)$ is a continuous, piecewise smooth function, and

$$Z_{X,T} = \{(t, p) \in [0, \infty) \times X : t \leq \min T(p)\} \quad (183)$$

is a compact, oriented $(n+1)$ -manifold with g-corners. Also

$$\partial Z_{X,T} = -Z_{\partial X, T|_{\partial X}} \amalg -\{0\} \times X \amalg \coprod_{c \in C} X_c \quad (184)$$

in oriented n -manifolds with g-corners, where C is a finite indexing set, and X_c for $c \in C$ are the connected components of the portion of $\partial Z_{X,T}$ meeting $(0, \infty) \times X^\circ$. Locally the X_c correspond to branches $f_c : T \rightarrow [1, \infty)$ of T , and $X_c \cong \{p \in X : f_c(p) = \min T(p)\}$. The reason for the name ‘tent function’ is that when we sketch $Z_{X,T}$ in (183), it looks like a tent, as in Figure A.1 on page 182.

We will also define tent functions on *orbifolds*, and on (*effective*) *Kuranishi chains* $[X, \mathbf{f}, \mathbf{G}]$ or $[X, \mathbf{f}, \underline{\mathbf{G}}]$. The Kuranishi chain version of (184) is

$$\begin{aligned} \partial[Z_{X,T}, \mathbf{f} \circ \pi, \mathbf{H}_{X,T}] = & -[Z_{\partial X, T|_{\partial X}}, \mathbf{f}|_{\partial X} \circ \pi, \mathbf{H}_{\partial X, T|_{\partial X}}] \\ & -[X, \mathbf{f}, \mathbf{G}] + \sum_{c \in C} [X_c, \mathbf{f}_c, \mathbf{G}_c]. \end{aligned} \quad (185)$$

This says that $[X, \mathbf{f}, \mathbf{G}]$ is homologous to $\sum_{c \in C} [X_c, \mathbf{f}_c, \mathbf{G}_c]$, modulo terms over ∂X . Much of Appendices B and C will involve showing that some Kuranishi cycle which is a combination of $[X, \mathbf{f}, \mathbf{G}]$ is homologous to a second Kuranishi cycle which is a combination of $\sum_{c \in C} [X_c, \mathbf{f}_c, \mathbf{G}_c]$, and that *this second cycle has some better property than the first cycle*. It is helpful to think of (184) and (185) as meaning that *we have cut X into finitely many pieces X_c for $c \in C$* .

In fact we will use tent functions for four different purposes:

- (a) To cut a Kuranishi space X into *arbitrarily small pieces* X_c for $c \in C$. This will be needed in Step 1 of the proof of Theorem 4.9 in Appendix C.
- (b) To cut an effective orbifold X into pieces X_c for $c \in C$ which are *manifolds*, and more generally, to cut an effective Kuranishi space X into pieces X_c for $c \in C$ which are Kuranishi spaces with *trivial stabilizers*. This will be needed in Step 1 of the proof of Theorem 4.8 in Appendix B.
- (c) To cut a compact n -manifold X with g-corners into pieces X_c for $c \in C$ which are n -simplices Δ_n . This will be needed in Step 3 of the proof of Theorem 4.8 in Appendix B, to construct a cycle in singular homology.
- (d) To get round the failure of the smooth Extension Principle for manifolds with g-corners, Principle 2.8(c), by defining a class of piecewise smooth functions on manifolds with g-corners for which the Extension Principle holds. This is important in Step 2 of the proof of Theorem 4.8 in Appendix B, when we perturb Kuranishi spaces to get manifolds; in effect we are choosing piecewise smooth perturbations with prescribed boundary values, and this would not be possible working only with smooth perturbations.

These different goals are responsible for the length of this appendix, and also for a certain lack of coherence, as we jump between (a)–(d) or mix them up. We start by developing our material on compact manifolds in §A.1, then generalize to orbifolds in §A.2, and finally to Kuranishi chains in §A.3.

A.1 Tent functions on compact manifolds

A.1.1 The definition of tent functions

Definition A.1. For S a set, write $F(S)$ for the set of finite, nonempty subsets of S . Let X be an n -manifold with g -corners. We call $T : X \rightarrow F([1, \infty))$ a *tent function* if for each $x \in X$ there exists an open neighbourhood U of x in X , open subsets $U_1, \dots, U_N \subseteq U$ and smooth functions $t_i : U_i \rightarrow [1, \infty)$ for $i = 1, \dots, N$ such that $T(u) = \{t_i(u) : i = 1, \dots, N, u \in U_i\}$ for all $u \in U$, satisfying the following conditions:

- (a) For every subset $\{i_1, \dots, i_l\}$ of $\{1, \dots, N\}$ with $l \geq 2$, the subset $S_{\{i_1, \dots, i_l\}} = \{u \in U_{i_1} \cap \dots \cap U_{i_l} : t_{i_1}(u) = \dots = t_{i_l}(u)\}$ is an embedded submanifold of $U_{i_1} \cap \dots \cap U_{i_l} \subseteq X$. We require that $S_{\{i_1, \dots, i_l\}}$ should intersect the codimension k stratum $S_k(X)$ in Definition 2.4 transversely for all $k \geq 0$. This implies that $S_{\{i_1, \dots, i_l\}}$ has boundary or (g-)corners where it intersects the boundary or (g-)corners of X , and we require that these should be the only boundary or (g-)corners of $S_{\{i_1, \dots, i_l\}}$. That is, we have $\partial^k S_{\{i_1, \dots, i_l\}} = \{u \in \partial^k(U_{i_1} \cap \dots \cap U_{i_l}) \subseteq \partial^k X : t_{i_1}(u) = \dots = t_{i_l}(u)\}$.
- (b) In (a), for all $u \in S_{\{i_1, \dots, i_l\}}$, as vector subspaces of T_u^*X we have $\langle d(t_{i_2} - t_{i_1})|_u, d(t_{i_3} - t_{i_1})|_u, \dots, d(t_{i_l} - t_{i_1})|_u \rangle = \{\alpha \in T_u^*X : \alpha|_{T_u S_{\{i_1, \dots, i_l\}}} \equiv 0\}$.

More generally, for all $k \geq 0$ and u in $\partial^k S_{\{i_1, \dots, i_l\}}$, we have

$$\langle d(t_{i_2} - t_{i_1}|_{\partial^k(U_{i_1} \cap \dots \cap U_{i_l})})|_u, \dots, d(t_{i_l} - t_{i_1}|_{\partial^k(U_{i_1} \cap \dots \cap U_{i_l})})|_u \rangle = \{\alpha \in T_u^* \partial^k X : \alpha|_{T_u \partial^k S_{\{i_1, \dots, i_l\}}} \equiv 0\}.$$

- (c) For each $i = 1, \dots, N$, the set $\{u \in U_i : t_i(u) = \min T(u)\}$ is closed in U .

We call the functions $t_i : U_i \rightarrow [1, \infty)$ *branches* of T on U , and the sets $U_1, \dots, U_N$ *branch domains*. If $T : X \rightarrow F([1, \infty))$ is a tent function, then the *minimum* of T is $\min T : X \rightarrow [1, \infty)$, the function taking $x \mapsto \min T(x)$, which is well-defined as $T(x)$ is a finite, nonempty subset of $[1, \infty)$. Part (c) implies that $\min T$ is *continuous*.

The point here is that locally the branches of T do not have to be defined on all of X , but only on open subsets. Thus, as x moves about in X , the number of elements $T(x)$ can change as branches of T pop in and out of existence. However, (c) implies that a branch t_i of T cannot disappear near x if $t_i(x) = \min T(x)$. Thus, disappearing branches do not cause $\min T$ to become discontinuous, and $\min T$ near x is locally the minimum of smooth functions $t_i : X \rightarrow [1, \infty)$ defined near x , so $\min T$ is continuous, and piecewise-smooth.

As (a)–(c) are stable under taking boundaries, if $T : X \rightarrow F([1, \infty))$ is a tent function then $T|_{\partial X} : \partial X \rightarrow F([1, \infty))$ is also a tent function.

The next proposition shows how we will use tent functions. Definition A.1 was carefully designed just to make $Z_{X,T}$ a manifold with g-corners. Think of the proposition this way: we are cutting X into finitely many pieces X_c for $c \in C$, which as we will see later can be chosen to be arbitrarily small, to be simplices Δ_n , or to have other good properties. The $(n+1)$ -manifold $Z_{X,T}$ is a kind of (solid) graph of $\min T$, and we will use it to define a homology between a chain involving X , and a sum over $c \in C$ of chains involving X_c .

Proposition A.2. *Let X be a compact, oriented n -manifold with g-corners, and $T : X \rightarrow F([1, \infty))$ a tent function. Then*

$$Z_{X,T} = \{(t, x) \in [0, \infty) \times X : t \leq \min T(x)\} \quad (186)$$

is a compact, oriented $(n+1)$ -manifold with g-corners, and

$$\partial Z_{X,T} = -Z_{\partial X, T|_{\partial X}} \amalg -\{0\} \times X \amalg \coprod_{c \in C} X_c \quad (187)$$

in oriented n -manifolds with g-corners, where C is a finite indexing set, each X_c is connected, and if $\pi_X : [0, \infty) \times X \rightarrow X$ is the projection, then $\pi = \pi_X|_{X_c} : X_c \rightarrow X$ is an orientation-preserving immersion of X_c as an n -submanifold of X for each $c \in C$, which is an embedding on X_c° , with $X = \bigcup_{c \in C} \pi(X_c)$, and $\pi(X_c^\circ) \cap \pi(X_{c'}^\circ) = \emptyset$ for all $c \neq c'$ in C .

Proof. Since $\min T$ is continuous and X is compact, $\min T$ is bounded on X , and $Z_{X,T}$ is compact. As X is an oriented n -manifold with g-corners, $[0, \infty) \times X$ is an oriented $(n+1)$ -manifold with g-corners. We claim $Z_{X,T}$ is an embedded $(n+1)$ -submanifold of $[0, \infty) \times X$. It is enough to verify that $Z_{X,T}$ is locally modelled on a region with g-corners in $\mathbb{R}^{n+1}$ near each point $(t, x) \in Z_{X,T}$ with $t = \min T(x)$, since if $(t, x) \in Z_{X,T}$ with $t < \min T(x)$ then $Z_{X,T}$ coincides with $[0, \infty) \times X$ near (t, x) .

For such (t, x) , let $U, U_1, \dots, U_N, f_1, \dots, f_N$ be as in Definition A.1, and let $\{j_1, \dots, j_k\}$ be the subset of j in $\{1, \dots, N\}$ with $x \in U_j$ and $t_j(x) = \min T(x)$. Choose a small open neighbourhood $\tilde{U}$ of x in U such that $\tilde{U} \subseteq U_{j_a}$ for $a = 1, \dots, k$, and if $i \in \{1, \dots, N\} \setminus \{j_1, \dots, j_k\}$ and $u \in U_i \cap \tilde{U}$ then $t_i(u) > \min T(u)$. This last part is possible by Definition A.1(c), as for each $i \in \{1, \dots, N\} \setminus \{j_1, \dots, j_k\}$, we have to choose $\tilde{U}$ not to intersect a *closed* set in U not containing x . Then $t_{j_1}|_{\tilde{U}}, \dots, t_{j_k}|_{\tilde{U}} : \tilde{U} \rightarrow \mathbb{R}$ are well-defined, and $\min T(u) = \min\{t_{j_1}(u), \dots, t_{j_k}(u)\}$ for all $u \in \tilde{U}$.

Now let (U', ϕ') be a chart with g-corners on X with $x \in \phi'(U') \subseteq \tilde{U}$. Then U' is a region with g-corners in $\mathbb{R}^n$, defined as in Definition 2.1 using some $W' \subseteq \mathbb{R}^n$ and $f'_1, \dots, f'_{N'}$. As in Definition 2.2, the smooth maps $t_{j_a} \circ \phi' : U' \rightarrow \mathbb{R}$ extend smoothly to some open subset of $\mathbb{R}^n$ containing U' . Make W' smaller if necessary so that $t_{j_a} \circ \phi'$ extends to W' for $a = 1, \dots, k$, and choose such extensions so that Definition A.1(a),(b) still hold.

Define $\dot{W} \subset \mathbb{R}^{n+1}$ by $\dot{W} = (0, \infty) \times W'$ and $\dot{f}_1, \dots, \dot{f}_{k+N'} : \dot{W} \rightarrow \mathbb{R}$ by

$$\dot{f}_a(t, x_1, \dots, x_n) = \begin{cases} t_{j_a} \circ \phi'(x_1, \dots, x_n) - t, & a = 1, \dots, k, \\ f'_{a-k}(x_1, \dots, x_n), & a = k+1, \dots, k+N'. \end{cases}$$

Then Definitions 2.1(a),(b) and A.1(a),(b) for $f'_1, \dots, f'_{N'}$ and $t_{j_1}, \dots, t_{j_k}$ imply that $\dot{W}, \dot{f}_1, \dots, \dot{f}_{k+N'}$ satisfy Definition 2.1(a),(b), and the corresponding subset

$$\dot{U} = \{(s, x_1, \dots, x_n) : (x_1, \dots, x_n) \in U', \ 0 < s \leq \min T \circ \phi'(x_1, \dots, x_n)\}$$

is a region with g-corners in $\mathbb{R}^{n+1}$. Define $\dot{\phi} : \dot{U} \rightarrow Z_{X,T}$ by $\dot{\phi} : (s, x_1, \dots, x_n) \mapsto (s, \phi'(x_1, \dots, x_n))$. Then $(\dot{U}, \dot{\phi})$ is a chart with g-corners on $Z_{X,T}$ with (t, x) in $\dot{\phi}(\dot{U})$. Compatibility between such charts (U', ϕ') on X implies compatibility between the corresponding charts $(\dot{U}, \dot{\phi})$ on $Z_{X,T}$. Therefore $Z_{X,T}$ is a *compact, oriented $(n+1)$ -manifold with g-corners*.

In (220), the first two terms $-Z_{\partial X, T|_{\partial X}} \amalg -\{0\} \times X$ on the right hand side are the portion of $\partial Z_{X,T}$ coming from the boundary of $[0, \infty) \times X$, and their signs come from $\partial([0, \infty) \times X) = -[0, \infty) \times \partial X \amalg -\{0\} \times X$. Apart from these two terms, $\partial Z_{X,T}$ also has other contributions from points (t, x) with $t = \min T(x)$. As $\partial Z_{X,T}$ is a compact manifold, these other contributions have only finitely many connected components. Define these connected components to be X_c for c in C , some finite indexing set. Then (187) holds, by definition. If $(t, x), \dots$ are as above with $\tilde{U}^\circ$ connected and $((t, x), B) \in X_c$ for some $c \in C$ then the local boundary component B is of the form $\{(t, u) \in Z_{X,T} : u \in \tilde{U}^\circ, t = \phi_{j_a}(u)\}$ for some $a = 1, \dots, k$, and then $\{(t, u) \in Z_{X,T} : u \in \tilde{U}, t = \phi_{j_a}(u)\}$ is a local model for X_c . The remaining claims about the X_c follow. $\square$

The reason for the name *tent function* is that when we sketch $Z_{X,T}$ in (186), it looks rather like a tent. We illustrate this in Figure A.1.




Figure A.1: Example of a tent function used to subdivide a polygon

A.1.2 Cutting a compact manifold into arbitrarily small pieces

Here is our first method for constructing tent functions.

Definition A.3. Let X be a compact n -manifold with g-corners. Let g be a Riemannian metric on X , and choose $\epsilon > 0$ with $2\epsilon < \delta(g)$, where $\delta(g)$ is the injectivity radius of g . Write $d(\cdot, \cdot)$ for the metric on X determined by g . For $p \in X$ and $r > 0$ write $B_r(p)$ for the open ball of radius r about p , that is, $B_r(p) = \{x \in X : d(x, p) < r\}$. Then using geodesic normal coordinates on X near p , we see that for any $p \in X$, the map $X \rightarrow [0, \infty)$ given by $x \mapsto d(p, x)^2$ is continuous, and smooth wherever $d(p, x) < \delta(g)$. Hence $x \mapsto d(p, x)^2$ is smooth on $B_{2\epsilon}(p)$, as $2\epsilon < \delta(g)$.

The open sets $B_\epsilon(p)$ for $p \in X$ cover X , so as X is compact we can choose a finite subset $\{p_1, \dots, p_N\}$ of X with $X = \bigcup_{c=1}^N B_\epsilon(p_c)$. Choose $p_1, \dots, p_N$ to be generic amongst all such N -tuples $p'_1, \dots, p'_N$; this can be done by first choosing any $p'_1, \dots, p'_N$ with $X = \bigcup_{c=1}^N B_\epsilon(p'_c)$, and then taking $p_c \in X$ to be generic with $d(p_c, p'_c) \ll \epsilon$. Define $T : X \rightarrow F([1, \infty))$ by

$$T(x) = \{1 + d(p_c, x)^2 : c = 1, \dots, N, \ d(p_c, x) < 2\epsilon\}. \quad (188)$$

Proposition A.4. In Definition A.3, T is a tent function on X . In (187), we can take the indexing set C to be $\{1, \dots, N\}$ with $p_c \in \pi(X_c^\circ) \subseteq \pi(X_c) \subseteq B_\epsilon(p_c)$ for $c = 1, \dots, N$. We can choose T so that the pieces X_c are ‘arbitrarily small’, in the sense that if $\{V_i : i \in I\}$ is any open cover for X then we can choose T so that for all $c = 1, \dots, N$ there exists $i \in I$ with $\pi(X_c) \subseteq V_i$.

Proof. Since $X = \bigcup_{c=1}^N B_\epsilon(p_c)$, if $x \in X$ there exists at least one c with $d(p_c, x) < 2\epsilon$, and then $1 + d(p_c, x)^2 \in T(x)$. So $T(x)$ is nonempty, and T does map $X \rightarrow F([1, \infty))$. In Definition A.1, for any $x \in X$ we can take $U = X$ and $U_c = B_{2\epsilon}(p_c)$ and define $t_c : U_c \rightarrow [1, \infty)$ by $t_c(u) = 1 + d(p_c, u)^2$. Then U_c is open in U , t_c is smooth, and $T(u) = \{t_i(u) : i = 1, \dots, N, u \in U_i\}$.

Definition A.1(a),(b) hold because $p_1, \dots, p_N$ are generic, and we also have the extra property that $\dim S_{\{i_1, \dots, i_l\}} = n - l + 1$ for all $\{i_1, \dots, i_l\} \subseteq \{1, \dots, N\}$ with $l \geq 2$, so that $d(t_{i_2} - t_{i_1}), \dots, d(t_{i_l} - t_{i_1})$ are linearly independent along $S_{\{i_1, \dots, i_l\}}$. To see this, note that $S_{\{i_1, \dots, i_l\}} = \{x \in X : d(p_{i_1}, x) = d(p_{i_2}, x) = \dots = d(p_{i_l}, x) < 2\epsilon\}$, and as $p_{i_1}, \dots, p_{i_l}$ are generic, the $l - 1$ equations $d(p_{i_1}, x) = d(p_{i_2}, x) = \dots = d(p_{i_l}, x)$ are transverse.

For $i = 1, \dots, N$, we have $\{u \in U_i : t_i(u) = \min T(u)\} = \{u \in B_{2\epsilon}(p_i) : d(p_i, u) = \min\{d(p_c, u) : c = 1, \dots, N\}\}$. Since $X = \bigcup_{c=1}^N B_\epsilon(p_c)$, we have $\min\{d(p_c, u) : c = 1, \dots, N\} < \epsilon$, so $\{u \in U_i : t_i(u) = \min T(u)\} \subseteq B_\epsilon(p_i)$. Thus $\{u \in U_i : t_i(u) = \min T(u)\}$ is a closed subset of $U_i = B_{2\epsilon}(p_i)$, and is contained in $\overline{B}_\epsilon(p_i)$. Since $\overline{B}_\epsilon(p_i)$ is closed in both U_i and X , it follows that $\{u \in U_i : t_i(u) = \min T(u)\}$ is closed in X , and Definition A.1(c) holds. Thus T is a tent function.

For the second part, it is easy to see that the remaining boundary components $\coprod_{c \in C} X_c$ in (187) may be written

$$\coprod_{c=1}^N \{(1 + d(p_c, u)^2, u) : u \in X, d(p_c, u) = \min\{d(p_a, u) : a = 1, \dots, N\}\}, \quad (189)$$

omitting local boundary components B . Now each u in the set $\{u \in X : d(p_c, u) = \min\{d(p_a, u) : a = 1, \dots, N\}\}$ is joined to p_c by a unique geodesic segment of length $d(p_c, u)$ which also lies in the set. Hence the c term in (189) is nonempty and connected. But by definition the connected components of (189) are X_c for $c \in C$. Therefore we can take $C = \{1, \dots, N\}$, and X_c to be the c term in (189). This gives $p_c \in \pi(X_c^\circ) \subseteq \pi(X_c) \subseteq B_\epsilon(p_c)$, since $\min\{d(p_a, u) : a = 1, \dots, N\} < \epsilon$ for all $u \in X$.

Given an open cover $\{V_i : i \in I\}$ for X , we fix g in Definition A.3 and choose ϵ sufficiently small that for all $x \in X$ there exists $i \in I$ with $B_\epsilon(x) \subseteq V_i$, which is possible by compactness of X . The final part of the proposition then follows from $\pi(X_c) \subseteq B_\epsilon(p_c)$ in the second part. $\square$

A.1.3 Cutting a compact manifold into simplices

In Definition A.3, regard X and g as fixed, and choose $\epsilon > 0$ very small compared to all the natural length-scales of (X, g) . That is, choose ϵ with $\epsilon \ll \delta(g)$, and $\epsilon \ll \delta(g|_{S_k(X)})$ for $0 < k < n$, and $\|R(g)\|_{C^0} \ll \epsilon^{-2}$, and $\|R(g)|_{S_k(X)}\|_{C^0} \ll \epsilon^{-2}$ for $0 < k < n$, where $\delta(g)$ is the injectivity radius and $R(g)$ the Riemann curvature of g , and the second fundamental form of $S_k(X)$ in X is $\ll \epsilon^{-1}$ for $0 < k < n$. Then using *geodesic normal coordinates*, we see that balls $B_{2\epsilon}(x)$ of radius 2ϵ in the interior of (X, g) are approximately isometric to balls of radius 2ϵ in $\mathbb{R}^n$ with its Euclidean metric, and balls of radius 2ϵ close to the boundary are approximately isometric to balls of radius 2ϵ in a *polyhedral cone* in $\mathbb{R}^n$ with its Euclidean metric, since as in Remark 2.3(c),(d), manifolds with g-corners are approximately locally modelled on polyhedral cones in $\mathbb{R}^n$.

Therefore geodesic normal coordinates at p_c approximately identify $\pi(X_c) \subset X$ with a subset of $\mathbb{R}^n$, or a polyhedral cone in $\mathbb{R}^n$, defined by inequalities $d_{\text{Eu}}(p_c, u) \leq d_{\text{Eu}}(p_a, u)$ for all $a = 1, \dots, N$, for some distinct points $p_1, \dots, p_N$ in $\mathbb{R}^n$, where d_{Eu} is the Euclidean metric on $\mathbb{R}^n$. Such inequalities define a *convex polyhedron* in $\mathbb{R}^n$, which is compact as X_c is compact. So we deduce:

Lemma A.5. *In Definition A.3, if ϵ is small compared to the natural length-scales of (X, g) , then geodesic normal coordinates at p_c in X approximately identify $\pi(X_c) \subset X$ with a compact convex polyhedron in $T_{p_c}X \cong \mathbb{R}^n$.*

Note that Remark 2.3(e) implies that $\pi(X_c)$ need not be diffeomorphic with a convex polyhedron in $\mathbb{R}^n$, so our notion of ‘approximately identifies’ does not imply diffeomorphism, just that the defining equations of $\pi(X_c)$ can be slightly perturbed, without changing the discrete structure of $\pi(X_c)$ as a manifold with g-corners, to give a compact convex polyhedron.

There is a standard way of dividing a compact convex polyhedron in $\mathbb{R}^n$ into n -simplices Δ_n , called *barycentric subdivision*, as in Bredon [10, §IV.17].

Definition A.6. Let K be a compact convex polyhedron in $\mathbb{R}^n$. For each k -dimensional face F of K for $k = 0, \dots, n$, define the *barycentre* b_F of F to be the point in the interior of F whose position vector is the average of the position vectors of the vertices of F . Let $\mathbf{F} = F_0 \subset F_1 \subset \dots \subset F_n = K$ be a chain

of faces of K with $\dim F_j = j$ for $j = 0, \dots, n$; we call $\mathbf{F}$ a *flag* for K . Write $p_0, \dots, p_n$ for the vertices of Δ_n , where $p_j = (\delta_{j0}, \delta_{j1}, \dots, \delta_{jn})$.

Let $\sigma_{\mathbf{F}} : \Delta_n \rightarrow \mathbb{R}^n$ be the unique affine map with $\sigma_{\mathbf{F}}(p_j) = b_{F_j}$ for $j = 0, \dots, n$. Then K is the union of $\sigma_{\mathbf{F}}(\Delta_n)$ over all flags $\mathbf{F}$ for K , and $\sigma_{\mathbf{F}}(\Delta_n) \cap \sigma_{\mathbf{F}'}(\Delta_n) = \emptyset$ for $\mathbf{F} \neq \mathbf{F}'$. This division of K into n -simplices $\sigma_{\mathbf{F}}(\Delta_n)$ is called the *barycentric subdivision* of K . It is compatible with boundaries in two ways:

- (i) Suppose $\mathbf{F} = F_0 \subset F_1 \subset \dots \subset F_n = K$ and $\mathbf{F}' = F'_0 \subset F'_1 \subset \dots \subset F'_n = K$ are two flags for K , and $F_j = F'_j$ for $j \in \{j_0, j_1, \dots, j_l\} \subseteq \{0, \dots, k\}$, with $j_0 < j_1 < \dots < j_l$. Let $\Delta_l^j \subseteq \Delta_n$ be the l -dimensional face Δ_l of Δ_n with vertices $p_{j_0}, \dots, p_{j_l}$. Then $\sigma_{\mathbf{F}}|_{\Delta_l^j} \equiv \sigma_{\mathbf{F}'}|_{\Delta_l^j}$.
- (ii) The barycentric subdivision of K restricts on ∂K to the disjoint union of the barycentric subdivisions of each boundary face of K .

We can construct the barycentric subdivision of K using a *tent function*. Choose constants $0 = c_0 < c_1 \ll c_2 \ll \dots \ll c_n$. For each flag $\mathbf{F} = F_0 \subset F_1 \subset \dots \subset F_n = K$ for K , define $A_{\mathbf{F}} : \mathbb{R}^n \rightarrow \mathbb{R}$ to be the unique affine function with $A_{\mathbf{F}}(b_{F_j}) = c_j$ for $j = 0, \dots, n$. Then define $T : K \rightarrow F([1, \infty))$ by

$$T(x) = \{2 + \delta A_{\mathbf{F}}(x) : \mathbf{F} \text{ is a flag for } K\},$$

where $\delta > 0$ is sufficiently small that $\delta A_{\mathbf{F}} \geq -1$ on K for all $\mathbf{F}$. One can show that if the ratios $c_2/c_1, \dots, c_n/c_{n-1}$ are large enough then T is a tent function, where in Definition A.1 we take $U = U_1 = \dots = U_N = K$ and $t_j = 2 + \delta A_{\mathbf{F}_j}$, where $\mathbf{F}_1, \dots, \mathbf{F}_N$ are the possible flags for K , and

$$\sigma_{\mathbf{F}}(\Delta_n) = \{x \in K : A_{\mathbf{F}}(x) \leq A_{\mathbf{F}'}(x) \text{ for all flags } \mathbf{F}' \text{ for } K\},$$

so in (187), the pieces $\pi(X_c)$ for all $c \in C$ are $\sigma_{\mathbf{F}}(\Delta_n)$ for all flags $\mathbf{F}$ for K .

We can combine the ideas of the last four definitions and results to construct a tent function T for X such that each X_c for $c \in C$ is a simplex Δ_n . That is, we use a tent function to construct a *triangulation* of X by n -simplices, in the sense of Bredon [10, p. 246]. This will enable us to show that chains defined using compact manifolds X with g-corners are homologous to chains defined using simplices Δ_n , that is, to chains in singular homology.

Theorem A.7. *Let X be a compact n -manifold with g-corners. Then we can construct a tent function $T : X \rightarrow F([1, \infty))$ such that the components X_c , $c \in C$ of $\partial Z_{X,T}$ in (187) are all diffeomorphic to the n -simplex Δ_n , with diffeomorphisms $\sigma_c : \Delta_n \rightarrow X_c$ for $c \in C$. Thus*

$$\partial^2 Z_{X,T} \supseteq \coprod_{c \in C} \partial X_c = \coprod_{c \in C} \sigma_c(\partial \Delta_n) = \coprod_{c \in C} \coprod_{j=0}^n \sigma_c \circ F_j^n(\Delta_{n-1}). \quad (190)$$

Definition 2.7 gives a free, orientation-reversing involution $\sigma : \partial^2 Z_{X,T} \rightarrow \partial^2 Z_{X,T}$. We can choose the σ_c to have the following boundary compatibility: suppose that σ exchanges two $(n-1)$ -simplices $\sigma_c \circ F_j^n(\Delta_{n-1})$ and $\sigma_{c'} \circ F_{j'}^n(\Delta_{n-1})$ in (190). Then $j = j'$, $c \neq c'$ and $\sigma \circ \sigma_c \circ F_j^n \equiv \sigma_{c'} \circ F_{j'}^n : \Delta_{n-1} \rightarrow \partial^2 Z_{X,T}$.

Proof. We first sketch the construction, and then fill in some details. As in Lemma A.5, use Definition A.3 with ϵ small compared to the length-scales of (X, g) to construct a tent function we will write as $\tilde{T} : X \rightarrow F([1, \infty))$, using points $p_1, \dots, p_N$, and functions $\tilde{t}_i : B_{2\epsilon}(p_i) \rightarrow [1, \infty)$ given by $\tilde{t}_i(u) = 1 + d(p_i, u)^2$, for $i = 1, \dots, N$. Use the notation $\tilde{X}_i$ for $i = 1, \dots, N$ for the X_c in (187) for $Z_{X, \tilde{T}}$, where $p_i \in \pi(\tilde{X}_i)$. Then as in Lemma A.5, using geodesic normal coordinates at p_i , $\pi(\tilde{X}_i)$ approximates a compact convex polyhedron K_i in $T_{p_i}X \cong \mathbb{R}^n$ for $i = 1, \dots, N$.

Define $C = \{(i, \mathbf{F}) : i = 1, \dots, N, \mathbf{F} \text{ is a flag for } K_i\}$. We shall first choose embeddings $\tilde{\sigma}_{(i, \mathbf{F})} : \Delta_n \rightarrow \tilde{X}_i$ for all $(i, \mathbf{F}) \in C$ such that for each $i = 1, \dots, N$, the $\tilde{\sigma}_{(i, \mathbf{F})}$ for all faces $\mathbf{F}$ of K_i form a triangulation of $\tilde{X}_i$, where $\tilde{\sigma}_{(i, \mathbf{F})}$ approximates $\sigma_{\mathbf{F}}$ in Definition A.6 for K_i under the approximation $K_i \cong \tilde{X}_i$ in geodesic normal coordinates. These $\tilde{\sigma}_{(i, \mathbf{F})}$ will satisfy boundary compatibilities analogous to Definition A.6(i),(ii).

Choose constants $0 = c_0 < c_1 \ll c_2 \ll \dots \ll c_n$, such that the ratios $c_2/c_1, \dots, c_n/c_{n-1}$ are large enough for the tent function construction in Definition A.6 to work for K_i for all $i = 1, \dots, N$. Choose smooth functions $A_{(i, \mathbf{F})} : B_{2\epsilon}(p_i) \rightarrow \mathbb{R}$ for all $(i, \mathbf{F}) \in C$, such that $A_{(i, \mathbf{F})} \circ \tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})} : \Delta_n \rightarrow \mathbb{R}$ is the unique affine function with $A_{(i, \mathbf{F})} \circ \tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})}(p_j) = c_j$ for $j = 0, \dots, N$, where $\tilde{\pi} : Z_{X, \tilde{T}} \rightarrow X$ is the projection so that $\tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})} : \Delta_n \rightarrow B_{2\epsilon}(p_i) \subseteq X$, and using geodesic normal coordinates to identify $B_{2\epsilon}(p_i)$ with the ball of radius 2ϵ in $T_{p_i}X$, $A_{(i, \mathbf{F})}$ approximates an affine function on the whole of $B_{2\epsilon}(p_i)$.

Choose $\delta > 0$ sufficiently small that $\delta A_{(i, \mathbf{F})} \geq -1$ and $|\delta A_{(i, \mathbf{F})}| \ll \epsilon^2$ on $B_{2\epsilon}(p_i)$ for all $(i, \mathbf{F}) \in C$. Generalizing (188), define $T : X \rightarrow F([1, \infty))$ by

$$T(x) = \{2 + d(p_i, x)^2 + \delta A_{(i, \mathbf{F})} : (i, \mathbf{F}) \in C, d(p_i, x) < 2\epsilon\}. \quad (191)$$

Then similar proofs to Proposition A.2 and Definition A.6 show that T is a tent function, and the components X_c for $c \in C$ of $Z_{X, T}$ in (187) can be written with the indexing set C above such that $\pi(X_{(i, \mathbf{F})}) = \tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})}(\Delta_n)$, so that $\tilde{\sigma}_{(i, \mathbf{F})}$ lifts to a unique diffeomorphism $\sigma_{(i, \mathbf{F})} : \Delta_n \rightarrow X_{(i, \mathbf{F})}$ with $\pi \circ \sigma_{(i, \mathbf{F})} \equiv \tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})}$.

In carrying out this programme, two points need special care:

- Because of the local variation of polyhedral cone models along codimension l strata of manifolds with g-corners for $l \geq 3$ discussed in Remark 2.3(e), we cannot assume that $\tilde{X}_i$ is diffeomorphic to K_i for $i = 1, \dots, N$.
- The $\tilde{X}_i$ for $i = 1, \dots, N$ are in general manifolds with g-corners, not corners, and the smooth Extension Principle, Principle 2.8(c), fails for manifolds with g-corners. So in the proof we must avoid choosing data such as $A_{(i, \mathbf{F})}$ on $\partial\tilde{X}_i$ and trying to extend it smoothly over $\tilde{X}_i$.

We now explain how to choose the embeddings $\tilde{\sigma}_{(i, \mathbf{F})} : \Delta_n \rightarrow \tilde{X}_i$ for all $(i, \mathbf{F}) \in C$. For the $\tilde{\sigma}_{(i, \mathbf{F})}$ to lift to $\sigma_{(i, \mathbf{F})}$ satisfying the boundary compatibility $\sigma \circ \sigma_c \circ F_j^n \equiv \sigma_{c'} \circ F_j^n : \Delta_{n-1} \rightarrow \partial^2 Z_{X, T}$ in the theorem, the $\tilde{\sigma}_{(i, \mathbf{F})}$ must satisfy boundary compatibilities on $\partial\Delta_n$, and these in turn imply compatibilities between the $\tilde{\sigma}_{(i, \mathbf{F})}$ on $\partial^m \Delta_n$ for $m = 1, 2, \dots, n$. In order to satisfy all these

compatibilities, we must choose the $\tilde{\sigma}_{(i,\mathbf{F})}|_{\partial^m \Delta_n}$ by induction on decreasing $m = n, n-1, \dots, 1, 0$. Similar proofs where we choose data on $\partial^m X_a$ by induction on decreasing codimension m will appear in §B.1–§B.3 and §C.1.

We introduce some notation. For $m = 0, \dots, n$, let S_n^m be the set of $(n-m)$ -dimensional faces of Δ_n . Each Δ in S_n^m is a simplex Δ_{n-m} , the convex hull of its $n-m+1$ vertices, which are a subset of the vertices $p_0, \dots, p_n$ of Δ_n , and any $n-m+1$ out of $p_0, \dots, p_n$ determine some Δ in S_n^m . Hence $|S_n^m| = \binom{n+1}{n-m+1}$. Note that it is *not* true that $\partial^m \Delta_n = \coprod_{\Delta \in S_n^m} \Delta$, but instead, each Δ in S_n^m appears $m!$ times in $\partial^m \Delta_n$, indexed by a choice of order of the m codimension 1 faces of Δ_n that meet at Δ . We shall describe the compatibilities between the $\tilde{\sigma}_{(i,\mathbf{F})}|_{\Delta}$ for all $(i, \mathbf{F}) \in C$, $\Delta \in S_n^m$ using an equivalence relation $\sim$ on $C \times S_n^m$.

Let $(i, \mathbf{F}, \Delta) \in C \times S_n^m$. The flag $\mathbf{F}$ for K_i determines a simplex $\sigma_{\mathbf{F}} : \Delta_n \rightarrow K_i$ in the barycentric subdivision of K_i in Definition A.6, so $\sigma_{\mathbf{F}}(\Delta)$ is one of the $(n-m)$ -dimensional simplices in the associated triangulation of K_i . The interior $\sigma_{\mathbf{F}}(\Delta^\circ)$ is a subset of the codimension k stratum $S_k(K_i)$ for some unique $k = 0, \dots, m$; in fact $n-k = \max\{j : p_j \text{ is a vertex of } \Delta\}$, so k is determined by Δ . Let $K_i^{\mathbf{F}, \Delta}$ be the unique $(n-k)$ -dimensional face of K_i containing $\sigma_{\mathbf{F}}(\Delta)$. Then $K_i^{\mathbf{F}, \Delta}$ is an $(n-k)$ -dimensional compact convex polyhedron in an affine subspace of $T_{p_i} X$ isomorphic to $\mathbb{R}^{n-k}$, and the barycentric subdivision of K_i restricts to the barycentric subdivision of $K_i^{\mathbf{F}, \Delta}$, with $\sigma_{\mathbf{F}}(\Delta)$ one of the simplices of the barycentric subdivision of $K_i^{\mathbf{F}, \Delta}$.

Now $\tilde{X}_i$ approximates K_i , and this approximation identifies the face structure of $\tilde{X}_i$ and K_i as manifolds with g-corners, the connected components of $S_k(\tilde{X}_i)$ and $S_k(K_i)$, and so on. As $(K_i^{\mathbf{F}, \Delta})^\circ$ is a connected component of $S_k(K_i)$, there is a unique corresponding connected component of $S_k(\tilde{X}_i)$. Let $\tilde{X}_i^{\mathbf{F}, \Delta}$ be its closure. Then $\tilde{X}_i^{\mathbf{F}, \Delta} \subseteq \tilde{X}_i$ is an $(n-k)$ -dimensional face of $\tilde{X}_i$, which approximates the convex polyhedron $K_i^{\mathbf{F}, \Delta}$. As $\tilde{X}_i \subset \partial Z_{X, \tilde{T}}$, the natural immersion $\iota : \partial Z_{X, \tilde{T}} \rightarrow Z_{X, \tilde{T}}$ restricts to $\iota : \tilde{X}_i \rightarrow Z_{X, \tilde{T}}$, and so to $\iota : \tilde{X}_i^{\mathbf{F}, \Delta} \rightarrow Z_{X, \tilde{T}}$. Then $\iota((\tilde{X}_i^{\mathbf{F}, \Delta})^\circ)$ is a connected component of $S_{k+1}(Z_{X, \tilde{T}})$, and $\iota(\tilde{X}_i^{\mathbf{F}, \Delta})$ is an $(n-k)$ -dimensional face of $Z_{X, \tilde{T}}$.

Let $(i, \mathbf{F}), (i', \mathbf{F}')$ lie in C . We use the same Δ and k for both. Then we get $(n-k)$ -dimensional faces $\iota(\tilde{X}_i^{\mathbf{F}, \Delta}), \iota(\tilde{X}_{i'}^{\mathbf{F}', \Delta})$ of $Z_{X, \tilde{T}}$. Suppose $\iota(\tilde{X}_i^{\mathbf{F}, \Delta}) = \iota(\tilde{X}_{i'}^{\mathbf{F}', \Delta})$. As $\iota(\tilde{X}_i^{\mathbf{F}, \Delta}), \iota(\tilde{X}_{i'}^{\mathbf{F}', \Delta})$ are approximately identified with convex polyhedra $K_i^{\mathbf{F}, \Delta}, K_{i'}^{\mathbf{F}', \Delta}$, there is an approximate identification between $K_i^{\mathbf{F}, \Delta}$ and $K_{i'}^{\mathbf{F}', \Delta}$, which identifies the faces of $K_i^{\mathbf{F}, \Delta}$ with the faces of $K_{i'}^{\mathbf{F}', \Delta}$, and the simplices of their barycentric subdivisions. Now $\sigma_{\mathbf{F}}(\Delta)$ and $\sigma_{\mathbf{F}'}(\Delta)$ are $(n-m)$ -simplices in the barycentric subdivisions of $K_i^{\mathbf{F}, \Delta}$ and $K_{i'}^{\mathbf{F}', \Delta}$.

Define a binary relation $\sim$ on $C \times S_n^m$ by $(i, \mathbf{F}, \Delta) \sim (i', \mathbf{F}', \Delta)$ if $\iota(\tilde{X}_i^{\mathbf{F}, \Delta}) = \iota(\tilde{X}_{i'}^{\mathbf{F}', \Delta})$, and $\sigma_{\mathbf{F}}(\Delta)$ and $\sigma_{\mathbf{F}'}(\Delta)$ correspond under the identification of simplices of the barycentric subdivisions of $K_i^{\mathbf{F}, \Delta}$ and $K_{i'}^{\mathbf{F}', \Delta}$. Define $(i, \mathbf{F}, \Delta) \not\sim (i', \mathbf{F}', \Delta')$ if $\Delta \neq \Delta'$. It is easy to see that $\sim$ is an *equivalence relation* on $C \times S_n^m$. Note that we allow $i = i'$ but $\mathbf{F} \neq \mathbf{F}'$, and then $\sigma_{\mathbf{F}}(\Delta_n), \sigma_{\mathbf{F}'}(\Delta_n)$ are

two n -simplices in the barycentric subdivision of K_i which share an $(n - m)$ -dimensional face $\sigma_{\mathbf{F}}(\Delta) = \sigma_{\mathbf{F}'}(\Delta)$.

By reverse induction on $m = n, n - 1, \dots, 0$, we will choose maps $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)} : \Delta \rightarrow \tilde{X}_i^{\mathbf{F}, \Delta} \subseteq \tilde{X}_i$ for all $(i, \mathbf{F}) \in C$ and $\Delta \in S_n^m$ satisfying the conditions:

- (i) Under the approximate identification of $\tilde{X}_i$ and K_i , $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ approximates $\sigma_{\mathbf{F}}|_{\Delta}$.
- (ii) If $\Delta \in S_n^m$ for $m < n$ then there is a natural identification $\partial\Delta \cong \Delta^0 \amalg \dots \amalg \Delta^{n-m}$, where $\Delta^0, \dots, \Delta^{n-m} \in S_n^{m+1}$. Under this identification we require that $\sigma_{(i, \mathbf{F}, \Delta)}|_{\Delta^j} \equiv \sigma_{(i, \mathbf{F}, \Delta^j)}$ for all $(i, \mathbf{F}) \in C$ and $j = 0, \dots, n - m$, where $\sigma_{(i, \mathbf{F}, \Delta^j)}$ was chosen in the previous inductive step.
- (iii) If $(i, \mathbf{F}), (i', \mathbf{F}') \in C$ and $\Delta \in S_n^m$ with $(i, \mathbf{F}, \Delta) \sim (i', \mathbf{F}', \Delta)$ then $\iota \circ \tilde{\sigma}_{(i, \mathbf{F}, \Delta)} \equiv \iota \circ \tilde{\sigma}_{(i', \mathbf{F}', \Delta)} : \Delta \rightarrow Z_{X, \tilde{T}}$.

The idea here is that $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)} = \tilde{\sigma}_{(i, \mathbf{F})}|_{\Delta}$. Then (i), (ii) are obvious, and (iii) is the boundary compatibilities that the $\tilde{\sigma}_{(i, \mathbf{F})}$ must satisfy over $\partial^m \Delta$. For the initial step $m = n$, each Δ in S_n^n is a point, one of the vertices $p_0, \dots, p_n$ of Δ_n , and $\tilde{X}_i^{\mathbf{F}, \Delta}$ is a point, one of the vertices of $\tilde{X}_i$. So the maps $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ are uniquely determined. Parts (i), (iii) are immediate, and (ii) is vacuous.

For the inductive step, suppose we have chosen $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ satisfying (i)–(iii) for all $(i, \mathbf{F}) \in C$ and $\Delta \in S_n^m$ for $m = n, n - 1, \dots, k + 1$, where $0 \leq k < n$. We shall choose $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ satisfying (i)–(iii) for all $(i, \mathbf{F}) \in C$ and $\Delta \in S_n^k$. Part (ii) determines $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}|_{\partial\Delta}$ uniquely. Part (ii) for $m = k + 1$ implies that these values for $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}|_{\partial\Delta}$, when restricted to $\partial^2 \Delta$, are invariant under the natural involution $\sigma : \partial^2 \Delta \rightarrow \partial^2 \Delta$ from §2.1. Since Δ is a manifold with corners (not g-corners), the Extension Principle, Principle 2.8(c) holds, and it is possible to smoothly extend the prescribed values for $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ on $\partial\Delta$ smoothly to Δ , at least near $\partial\Delta$. As in (i) we know approximately what $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ should be, and (i) when $m = k + 1$ implies that the prescribed values for $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}|_{\partial\Delta}$ are approximately right.

Thus there are no global topological problems in extending $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ from its prescribed boundary values to a global embedding satisfying (i), (ii), and for any given $(i, \mathbf{F}, \Delta)$ in $C \times S_n^k$ we can choose $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ satisfying (i), (ii). To make our choices satisfy (iii), we choose $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ as above for one representative in each equivalence class of $\sim$ in $C \times S_n^k$, and this choice and (iii) determine $\tilde{\sigma}_{(i', \mathbf{F}', \Delta)}$ uniquely for all $(i', \mathbf{F}', \Delta)$ in $C \times S_n^k$ with $(i', \mathbf{F}', \Delta) \sim (i, \mathbf{F}, \Delta)$. These choices then satisfy (i)–(iii) for all $(i, \mathbf{F}, \Delta)$ in $C \times S_n^k$, completing the inductive step. When $m = 0$ we have $S_n^0 = \{\Delta_n\}$. Define embeddings $\tilde{\sigma}_{(i, \mathbf{F})} = \tilde{\sigma}_{(i, \mathbf{F}, \Delta_n)} : \Delta_n \rightarrow \tilde{X}_i$ for all $(i, \mathbf{F}) \in C$.

Most of the rest of the proof follows the sketch above with few complications. Our choice of $\tilde{\sigma}_{(i, \mathbf{F})}$ determines $A_{(i, \mathbf{F})}$ on the simplex $\tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})}(\Delta_n)$ in $B_{2\epsilon}(p_i)$. Our definition of smooth maps between subsets of $\mathbb{R}^n$ in Definition 2.2 implies that $A_{(i, \mathbf{F})}$ extends smoothly to an open neighbourhood of $\tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})}(\Delta_n)$ in $B_{2\epsilon}(p_i)$, so there is no problem in extending $A_{(i, \mathbf{F})}$ from its prescribed values on $\tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})}(\Delta_n)$ to $B_{2\epsilon}(p_i)$, and in getting an approximately affine function. Then

provided $c_0, \dots, c_n$ and δ are chosen as above, we get a tent function T with the properties in the theorem, and unique diffeomorphisms $\sigma_{(i, \mathbf{F})} : \Delta_n \rightarrow X_{(i, \mathbf{F})}$ with $\pi \circ \sigma_{(i, \mathbf{F})} \equiv \tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})}$.

For the final part, writing $c = (i, \mathbf{F})$ and $c' = (i', \mathbf{F}')$, suppose that σ exchanges two $(n-1)$ -simplices $\sigma_{(i, \mathbf{F})} \circ F_j^n(\Delta_{n-1})$ and $\sigma_{(i', \mathbf{F}')} \circ F_{j'}^n(\Delta_{n-1})$ in (190). Applying $\iota : \partial^2 Z_{X, T} \rightarrow Z_{X, T}$ gives $\iota \circ \sigma_{(i, \mathbf{F})} \circ F_j^n(\Delta_{n-1}) = \iota \circ \sigma_{(i', \mathbf{F}')} \circ F_{j'}^n(\Delta_{n-1})$. Applying $\pi : Z_{X, T} \rightarrow X$ gives $\pi \circ \sigma_{(i, \mathbf{F})} \circ F_j^n(\Delta_{n-1}) = \pi \circ \sigma_{(i', \mathbf{F}')} \circ F_{j'}^n(\Delta_{n-1})$, since $\pi \circ \iota = \pi$. But $\tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})} = \pi \circ \sigma_{(i, \mathbf{F})}$ and $\tilde{\pi} \circ \tilde{\sigma}_{(i', \mathbf{F}')} = \pi \circ \sigma_{(i', \mathbf{F}')}$, so $\tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})} \circ F_j^n(\Delta_{n-1}) = \tilde{\pi} \circ \tilde{\sigma}_{(i', \mathbf{F}')} \circ F_{j'}^n(\Delta_{n-1})$. This then lifts to $\iota \circ \tilde{\sigma}_{(i, \mathbf{F})} \circ F_j^n(\Delta_{n-1}) = \iota \circ \tilde{\sigma}_{(i', \mathbf{F}')} \circ F_{j'}^n(\Delta_{n-1})$.

Now $F_j^n(\Delta_{n-1}), F_{j'}^n(\Delta_{n-1})$ lie in S_n^1 , so $\tilde{\sigma}_{(i, \mathbf{F})} = \tilde{\sigma}_{(i, \mathbf{F}, \Delta_n)}$ and (ii) above for $m = 0$ give $\tilde{\sigma}_{(i, \mathbf{F})}|_{F_j^n(\Delta_{n-1})} = \tilde{\sigma}_{(i, \mathbf{F}, F_j^n(\Delta_{n-1}))}$, and similarly $\tilde{\sigma}_{(i', \mathbf{F}')}|_{F_{j'}^n(\Delta_{n-1})} = \tilde{\sigma}_{(i', \mathbf{F}', F_{j'}^n(\Delta_{n-1}))}$. Therefore $\iota \circ \tilde{\sigma}_{(i, \mathbf{F}, F_j^n(\Delta_{n-1}))}(F_j^n(\Delta_{n-1})) = \iota \circ \tilde{\sigma}_{(i', \mathbf{F}', F_{j'}^n(\Delta_{n-1}))}(F_{j'}^n(\Delta_{n-1}))$, as in part (iii) above. This implies that $j = j'$, since the image of $\iota \circ \tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ in $Z_{X, \tilde{T}}$ determines Δ uniquely; this is why we put $(i, \mathbf{F}, \Delta) \not\sim (i', \mathbf{F}', \Delta')$ if $\Delta \neq \Delta'$.

By considering the faces of $K_i, K_{i'}$ corresponding to $\tilde{\sigma}_{(i, \mathbf{F})} \circ F_j^n(\Delta_{n-1})$ and $\tilde{\sigma}_{(i', \mathbf{F}')} \circ F_{j'}^n(\Delta_{n-1})$, we see from the definition of $\sim$ that $(i, \mathbf{F}, F_j^n(\Delta_{n-1})) \sim (i', \mathbf{F}', F_{j'}^n(\Delta_{n-1}))$, and so (iii) above with $m = 1$ gives $\iota \circ \tilde{\sigma}_{(i, \mathbf{F}, F_j^n(\Delta_{n-1}))} \equiv \iota \circ \tilde{\sigma}_{(i', \mathbf{F}', F_{j'}^n(\Delta_{n-1}))}$, yielding $\iota \circ \tilde{\sigma}_{(i, \mathbf{F})} \circ F_j^n \equiv \iota \circ \tilde{\sigma}_{(i', \mathbf{F}')} \circ F_{j'}^n$. Applying $\tilde{\pi}$ gives $\tilde{\pi} \circ \tilde{\sigma}_{(i, \mathbf{F})} \circ F_j^n \equiv \tilde{\pi} \circ \tilde{\sigma}_{(i', \mathbf{F}')} \circ F_{j'}^n$, as $\tilde{\pi} \circ \iota = \tilde{\pi}$, so $\pi \circ \sigma_{(i, \mathbf{F})} \circ F_j^n \equiv \pi \circ \sigma_{(i', \mathbf{F}')} \circ F_{j'}^n$. Since σ exchanges $\sigma_{(i, \mathbf{F})} \circ F_j^n(\Delta_{n-1})$ and $\sigma_{(i', \mathbf{F}')} \circ F_{j'}^n(\Delta_{n-1})$, and $\pi \circ \sigma = \pi$, this lifts to $\sigma \circ \sigma_{(i, \mathbf{F})} \circ F_j^n \equiv \sigma_{(i', \mathbf{F}')} \circ F_{j'}^n$, as we have to prove. $\square$

A.1.4 Tent functions and piecewise smooth extensions

As in §2.1, the smooth Extension Principle, Principle 2.8(c), fails for manifolds with g-corners. We will now introduce a notion of *piecewise-smooth* function associated to a tent function, and show that the Extension Principle does hold for such piecewise smooth functions.

Definition A.8. Let X be a compact manifold with g-corners, T a tent function on X , and $\pi : X_c \rightarrow X$ for $c \in C$ be as in Proposition A.2. A function $f : X \rightarrow \mathbb{R}$ is called *piecewise smooth subordinate to T* if f is continuous and $f|_{\pi(X_c)}$ is smooth for each $c \in C$, that is, $f \circ \pi : X_c \rightarrow \mathbb{R}$ is smooth for each $c \in C$.

Similarly, if $E \rightarrow X$ is a vector bundle then a section s of E is called *piecewise smooth subordinate to T* if s is continuous with $s|_{\pi(X_c)}$ smooth for each $c \in C$.

We shall define a tent function T on X for which a piecewise version of the Extension Principle holds.

Definition A.9. Let X be a compact n -manifold with g-corners. Choose a smooth vector field v on X such that v is nonzero and *inward pointing* at every point of ∂X . That is, we require that whenever $(p, B) \in \partial X$ and $(x_1, \dots, x_n)$ are coordinates on X near p such that $x_1 \geq 0$ on X and $x_1 = 0$ on B , then $dx_1(v|_p) > 0$. Then for all $t \geq 0$, the exponential map $\exp(tv) : X \rightarrow X$ is a

well-defined smooth embedding, which is the identity for $t = 0$, and maps X to a subset of X° for $t > 0$. Essentially, the flow $\exp(tv)$ for $t \geq 0$ *shrinks* X away from its boundary.

Write $X_1 = \exp(v)X$ and $X_2 = \exp(2v)X$. Then X_1, X_2 are embedded submanifolds of X° , and $\exp(v) : X \rightarrow X_1$, $\exp(2v) : X \rightarrow X_2$ are diffeomorphisms. The inclusion $\iota : \partial X_2 \rightarrow X$ is an immersion. We wish to extend ∂X_2 to a $(n-1)$ -manifold Y without boundary, not necessarily compact, with $\partial X_2 \subset Y$ an embedded $(n-1)$ -submanifold, and extend $\iota : \partial X_2 \rightarrow X$ to an immersion $\iota : Y \rightarrow X$ with the properties:

- (i) $\iota(Y \setminus \partial X_2) \subset X_1 \setminus X_2$.
- (ii) v is transverse to $\iota(Y)$ at every point of Y .
- (iii) there exists $M \geq 1$ such that for each $x \in X$ there are at most M points $(t, y) \in [0, 2] \times Y$ with $\exp(tv)x = \iota(y)$.
- (iv) $S_l = \{(y_1, \dots, y_l) \in Y^l : \iota(y_1) = \dots = \iota(y_l)\}$ is a disjoint union of embedded submanifolds of Y^l , for each $l \geq 1$.
- (v) In (iv), the map $i_l : S_l \rightarrow X$, $i_l : (y_1, \dots, y_l) \mapsto \iota(y_1)$ is an immersion, and $\text{di}_l(T_{(y_1, \dots, y_l)} S_l) = \bigcap_{i=1}^l d\iota(T_{y_i} Y)$ in $T_{\iota(y_1)} X$ for all $(y_1, \dots, y_l) \in S_l$.

The basic idea is that we are adding a small ‘collar’ to ∂X_2 near $\partial^2 X_2$. If $\partial^3 X_2 = \emptyset$ then we can think of Y as $\partial X_2 \amalg (0, \epsilon) \times \partial^2 X_2$ for $\epsilon > 0$ small. If we choose any Y, ι extending ∂X_2 , ι then $\iota(Y \setminus \partial X_2) \subset X_1 \setminus X_2$ near $\partial^2 X_2$, so making Y smaller if necessary (i) holds. Since v is $\exp(2v)$ -invariant and v is nonzero and inward-pointing at each point of ∂X , it follows that v is transverse to $\iota(\partial X_2)$ at every point of ∂X_2 . This is an open condition, so for any Y, ι extending ∂X_2 , v is transverse to $\iota(Y)$ on an open neighbourhood of ∂X_2 in Y . Thus making Y smaller if necessary, (ii) holds.

For (iii), using (ii) we find that the set of $(t, y) \in [0, 2] \times Y$ with $\exp(tv)x = \iota(y)$ is discrete, that is, it has no limit points. So if Z is any compact subset of Y , then there are only finitely many $(t, z) \in [0, 2] \times Z$ with $\exp(tv)x = \iota(z)$. The number of such (t, z) is an upper-semi-continuous function of $x \in X$, and so bounded by $M \geq 1$ as X is compact. Thus, given any Y, ι satisfying (ii), replacing Y by an open neighbourhood Y' of ∂X_2 in Y with $Z = \overline{Y'}$ compact in Y , there exists $M \geq 1$ with at most M pairs $(t, y) \in [0, 2] \times Y'$ with $\exp(tv)x = \iota(y)$ for any $x \in X$. So making Y smaller if necessary, (iii) holds.

To arrange for (iv) and (v) to hold, let $x \in \iota(\partial X_2) \subset X$, and choose a chart (U, ϕ) for X near x , so that $U \subset \mathbb{R}^n$ is open and $\phi : U \rightarrow X$ is a diffeomorphism with its image, with $x \in \phi(U)$. As X_2 is an embedded submanifold of X , $\phi^{-1}(X_2)$ is a *region with g -corners* in $\mathbb{R}^n$. Thus there exist smooth $f_1, \dots, f_N : U \rightarrow \mathbb{R}$ satisfying Definition 2.1(a),(b), such that $\phi^{-1}(X_2) = \{u \in U : f_i(u) \geq 0, i = 1, \dots, N\}$. Choosing such $f_1, \dots, f_N$ with N least, and making U smaller if necessary, we see that ∂X_2 is locally modelled on $\{(u, i) : u \in U, i = 1, \dots, N, f_i(u) = 0, f_j(u) \geq 0, j = 1, \dots, N\}$ over $\phi(U)$. We choose Y to be locally modelled on $\{(u, i) : u \in U, i = 1, \dots, N, f_i(u) = 0\}$ over $\phi(U)$. Then Definition 2.1(a),(b) for $f_1, \dots, f_N$ imply (iv),(v) above. Hence we can choose Y, ι satisfying

(i)–(v) near any point of $\iota(\partial X_2)$. Patching together such local choices, we can define Y, ι globally. Now define $T : X \rightarrow F([1, \infty))$ by

$$T(x) = \{2\} \cup \{3 - t : t \in (0, 2], \exp(tv)x \in \iota(Y)\}. \quad (192)$$

Proposition A.10. *In Definition A.9, T is a tent function on X . Furthermore, if $\pi : X_c \rightarrow X$ and C are as in Proposition A.2 for T , then there is a unique diffeomorphism $\Phi : X_1 \amalg [0, 1] \times \partial X \rightarrow \coprod_{c \in C} X_c$ such that $\pi \circ \Phi : X_1 \amalg [0, 1] \times \partial X \rightarrow X$ maps $x \mapsto x$ for $x \in X_1$ and $(t, (x, B)) \mapsto \exp(tv)x$ for $(t, (x, B)) \in [0, 1] \times \partial X$. If X is oriented then Φ is orientation-preserving on X_1 and orientation-reversing on $[0, 1] \times \partial X$, so we can regard Φ as an orientation-preserving diffeomorphism $\Phi : X_1 \amalg -[0, 1] \times \partial X \rightarrow \coprod_{c \in C} X_c$. Also $\min T|_{\partial X} \equiv 1$.*

Proof. Define $Z = \{(x, t, y) \in X \times (0, 3) \times Y : \exp(tv)x = \iota(y)\}$. Regarding Z as a fibre product $(X \times (0, 3)) \times_X Y$ and noting that $(x, t) \mapsto \exp(tv)x$ is a submersion $X \times (0, 3) \rightarrow X$, we see that Z is an n -manifold. Using Definition A.9(ii) and the fact that $(0, 3) \times Y$ has no boundary, we can show that the projection $\pi_X : Z \rightarrow X$, $\pi_X : (x, t, y) \mapsto x$ is a *local diffeomorphism*. Definition A.9(ii) implies that $\iota(Y) \cap X_2^\circ = \emptyset$. But if $x \in X$ and $t > 2$ then $\exp(tv)x \in X_2^\circ$, so there exists no $y \in Y$ with $\exp(tv)x = \iota(y)$. Hence $Z \subseteq X \times (0, 2] \times Y$. Therefore Definition A.9(iii) implies that $|\pi_X^{-1}(x)| \leq M$ for all $x \in X$, and π_X is a *globally finite* map. Note too that $T(x) = \{2\} \cup \{3 - t : (x, t, y) \in Z\}$.

Let $x \in X$. Since $\pi_X : Z \rightarrow X$ is a globally finite local diffeomorphism, if U is a sufficiently small open neighbourhood of x in X , then we may split $\pi_X^{-1}(U) = W_2 \amalg \dots \amalg W_N$ for some open subsets $W_2, \dots, W_N$ of Z for which $\pi_X|_{W_i} : W_i \rightarrow U$ is injective. Set $U_1 = U$ and $U_i = \pi_X(W_i)$ for $i = 2, \dots, N$. Then U_i are open subsets of U , and $\pi_X|_{W_i} : W_i \rightarrow U_i$ is a diffeomorphism for $i = 2, \dots, N$. Define $t_i : U_i \rightarrow [1, \infty)$ for $i = 1, \dots, N$ by $t_1 \equiv 2$ and $t_i(u) = 3 - t$ when $(u, t, y) \in W_i$ for $i = 2, \dots, N$. Since $\pi_X|_{W_i} : W_i \rightarrow U_i$ is a diffeomorphism there is a unique $(u, t, y) \in W_i$ for each $u \in U_i$, which depends smoothly on u , so t_i is well-defined and smooth.

We now see that $T(u) = \{t_i(u) : i = 1, \dots, N, u \in U_i\}$ for all $u \in U$, as in Definition A.1. We must verify Definition A.1(a)–(c). For subsets $\{i_1, \dots, i_l\}$ of $\{1, \dots, N\}$ with $1 \notin \{i_1, \dots, i_l\}$, Definition A.1(a), (b) follow directly from Definition A.9(iv), (v), with $S_{\{i_1, \dots, i_l\}} \subseteq \{x \in X : \exp(tv)x \in i_l(S_l), t \in (0, 3)\}$. When $1 \in \{i_1, \dots, i_l\}$, Definition A.1(a), (b) follow from Definition A.9(iv), (v), with $S_{\{i_1, \dots, i_l\}} \subseteq \{x \in X : \exp(v)x \in i_{l-1}(S_{l-1})\}$.

To prove Definition A.1(c), we shall show using (192) that

$$\min T(x) = \begin{cases} 2, & x \in X_1, \\ 1 + \inf\{t \in [0, 1] : x \in \exp(tv)X\}, & x \in X \setminus X_1. \end{cases} \quad (193)$$

For the first line of (193), note that if $x \in X_1$ and $t \in (1, 2]$ then $\exp(tv)x \in X_2^\circ$. But Definition A.9(i) implies that $\iota(Y) \cap X_2^\circ = \emptyset$. So $T(x)$ cannot contain $3 - t$

for $t \in (1, 2]$, implying that $\min T(x) = 2$. For the second line, let $x \in X \setminus X_1$, and set $s = \inf\{t \in [0, 1] : x \in \exp(tv)X\}$. (This set contains 1, so the infimum is well-defined.) By compactness of X we see that there exists $x' \in X$ with $\exp(sv)x' = x$. As s is smallest we must have $x' \in \iota(\partial X)$, so we can lift to $(x', B') \in \partial X$. We have a diffeomorphism $\exp(2v) : \partial X \rightarrow \partial X_2 \subset Y$. Define $y = \exp(2v)(x', B')$, and set $t = 2 - s$. Then $\iota(y) = \exp(2v)x' = \exp((t+s)v)x' = \exp(tv)\exp(sv)x' = \exp(tv)x$. Hence $3 - t = 1 + s \in T(x)$, and $\min T \leq 1 + s$. But using Definition A.9(i) we can show that $\min T \geq 1 + s$, proving (193).

Equation (193) implies that $\min T(x)$ occurs either at the extra branch 2, or at $3 - t$ when $\exp(tv)x = \iota(y)$ for $y \in \partial X_2$. The minimum can never occur at $3 - t$ when $\exp(tv)x = \iota(y)$ for $y \in Y \setminus \partial X_2$. Definition A.1(c) now follows using compactness of ∂X_2 . Therefore T is a *tent function* on X . The remainder of the proposition is a consequence of (193) and its proof. We showed that either $x \in X_1$ and $\min T(x) = 2$, or $\min T(x) = 1 + t$ for $t \in [0, 1]$ and there exists $(x', B') \in \partial X$ with $\exp(tv)x' = x$. These possibilities correspond to the fibres of $\pi : \coprod_{c \in C} X_c \rightarrow X$ over x , and show that we can write $\coprod_{c \in C} X_c \cong X_1 \amalg [0, 1] \times \partial X$ as in the proposition. $\square$

Here is our first version of Principle 2.8(c) for piecewise smooth functions. The proof is easy, we just have to check (194) is well-defined.

Proposition A.11. *Let X be a compact manifold with g -corners, and suppose $\zeta : \partial X \rightarrow \mathbb{R}$ is smooth, with $\zeta|_{\partial^2 X}$ invariant under the natural involution $\sigma : \partial^2 X \rightarrow \partial^2 X$ of §2.1. Let v, T, X_1 be as in Definition A.9, and Φ be as in Proposition A.10. Choose any smooth function $\eta_1 : X_1 \rightarrow \mathbb{R}$. Then there is a piecewise smooth function $\eta : X \rightarrow \mathbb{R}$ subordinate to the tent function T with $\eta|_{\partial X} \equiv \zeta$, given by*

$$\eta(x) = \begin{cases} \eta_1(x), & x \in X_1, \\ (1-t)\zeta(x', B') + t\eta_1(\exp(v)x'), & x = \pi_X \circ \Phi(t, (x', B')), \\ & (t, (x', B')) \in [0, 1] \times \partial X. \end{cases} \quad (194)$$

Similarly, if $E \rightarrow X$ is a vector bundle and f is a smooth section of $E|_{\partial X}$ with $f|_{\partial^2 X}$ σ -invariant, then by choosing a smooth section e_1 of $E|_{X_1}$ we can write down a piecewise smooth section e of E subordinate to T with $e|_{\partial X} \equiv f$.

Proposition A.11 is not adequate for our later applications as we cannot apply it iteratively, by induction on dimension, since the inputs ζ, f are smooth but the outputs η, e are piecewise smooth. We shall generalize it to a construction in which both input and output are piecewise smooth. First we generalize Definition A.9 to extend a tent function T' on ∂X to a tent function T on X .

Definition A.12. Let X be a compact n -manifold with g -corners, and let $T' : \partial X \rightarrow F([1, \infty))$ be a tent function on ∂X , such that $\min T'|_{\partial^2 X}$ is invariant under the natural involution $\sigma : \partial^2 X \rightarrow \partial^2 X$. We will construct a tent function $T : X \rightarrow F([1, \infty))$ with $T|_{\partial X} \equiv T'$, with useful properties we will explain in Propositions A.13 and A.14.

Let v, X_1, X_2 and $\iota : Y \rightarrow X$ be as in Definition A.9. Then $\exp(2v)$ induces a diffeomorphism $\exp(2v) : \partial X \rightarrow \partial X_2$, so that $T' \circ \exp(2v)$ is a tent function on ∂X_2 , and $\partial X_2 \subset Y$. Extend $T' \circ \exp(2v)$ to a tent function T'' on Y . Making Y smaller if necessary, this is possible, except that we may need to relax the condition that $\min T'' \geq 1$. So we take $T'' : Y \rightarrow F([\frac{1}{2}, \infty))$ to satisfy Definition A.1 replacing $[1, \infty)$ by $[\frac{1}{2}, \infty)$, with $T''|_{\partial X_2} \equiv T' \circ \exp(2v)$.

As $\min T' : \partial X \rightarrow [1, \infty)$ is continuous and ∂X is compact, $\min T'$ is bounded above. Choose $D > 1$ such that $\min T' < D$ on ∂X . So $\min T'' < D$ on ∂X_2 , as $T''|_{\partial X_2} \equiv T' \circ \exp(2v)$. Making Y smaller if necessary, we can also assume that $\min T'' < D$ on Y . Define $T : X \rightarrow F([1, \infty))$ by

$$T(x) = \{D\} \cup \{u + \theta(2-t) : t \in (0, 2], y \in Y, u \in T''(y), \exp(tv)x = \iota(y)\}, \quad (195)$$

where $\theta > D-1$ is chosen large enough to satisfy conditions in Proposition A.13.

Proposition A.13. *In Definition A.12, making Y smaller if necessary, if $\theta > D-1$ is sufficiently large then T is a tent function on X , and $\min T|_{\partial X} \equiv \min T'$, so that $Z_{\partial X, T|_{\partial X}} = Z_{\partial X, T'}$, and equation (187) becomes*

$$\partial Z_{X, T} = -Z_{\partial X, T'} \amalg -\{0\} \times X \amalg \coprod_{c \in C} X_c. \quad (196)$$

Let $X'_{c'}$ for $c' \in C'$ be as in (187) for $Z_{\partial X, T'}$, and X_c for $c \in C$ be as in (187) for $Z_{X, T}$. Define $\tilde{X} = \{x \in X : \min T(x) = D\}$. Then $\tilde{X}$ is a compact, embedded submanifold of X with g -corners, with $X_1 \subset \tilde{X} \subset X^\circ$. There are unique diffeomorphisms

$$\Phi : \tilde{X} \amalg ([0, 1] \times \coprod_{c' \in C'} X'_{c'}) \longrightarrow \coprod_{c \in C} X_c, \quad \Psi : \coprod_{c' \in C'} X'_{c'} \longrightarrow \partial \tilde{X} \quad (197)$$

which satisfy $\pi_X \circ \Phi(x) = x$ for $x \in \tilde{X}$, and for $t \in [0, 1]$, $x' \in \coprod_{c' \in C'} X'_{c'}$,

$$\pi_X \circ \Phi : (t, x') \longmapsto \exp\left(\frac{t}{\theta(D - \min T' \circ \pi_{\partial X}(x'))} v\right) \pi_X(x'), \quad (198)$$

$$\pi_X \circ \Psi : x' \longmapsto \exp\left(\frac{1}{\theta(D - \min T' \circ \pi_{\partial X}(x'))} v\right) \pi_X(x'), \quad (199)$$

$$\text{and} \quad \min T \circ \pi_X \circ \Phi(t, x') = (1-t) \min T' \circ \pi_{\partial X}(x') + tD, \quad (200)$$

writing $\pi_{\partial X} : \coprod_{c' \in C'} X'_{c'} \rightarrow \partial X$, $\pi_X : \coprod_{c' \in C'} X'_{c'}$ or $\coprod_{c \in C} X_c \rightarrow X$ for the natural projections. If X is oriented then Φ is orientation-preserving on $\tilde{X}$ and orientation-reversing on $[0, 1] \times \coprod_{c' \in C'} X'_{c'}$, and Ψ is orientation-preserving.

Proof. Our first goal is to prove that making Y smaller if necessary and taking $\theta > D-1$ sufficiently large, if $x \in X$ and $\min T(x) = u + \theta(2-t)$ for $t \in (0, 2]$, $y \in Y$ and $u \in T''(y)$ with $\exp(tv)x = \iota(y)$, then $y \in \partial X_2$. That is, the choice of ‘collar’ $Y \setminus \partial X_2$ and the values of T'' upon it do not affect $\min T$, since the branches of T coming from T'' on $Y \setminus \partial X_2$ are never minimal.

Let $y \in Y \setminus \partial X_2$. Then there exists unique $s \in (0, 1)$ with $\exp(sv)\iota(y) \in \iota(\partial X_2)$. Thus $\exp(sv)\iota(y) = \iota(z)$ for some (not necessarily unique) $z \in \partial X_2$. We claim that if Y is small enough and θ is large enough, then $\min T''(y) + \theta s > \min T''(z)$ for all such y, s, z . This implies what we want, for if $x \in X$ and $t \in (0, 2]$ with $\exp(tv)x = \iota(y)$, then $\exp((s+t)v)x = \iota(z)$, so $\min T''(z) + \theta(2-s-t) \in T(x)$, and $\min T''(z) + \theta(2-s-t) < \min T''(y) + \theta(2-t) \leq u + \theta(2-t)$ for all $u \in T''(y)$.

To prove the claim, choose a Riemannian metric g on X , let g_Y be the pullback metric on Y , and d_Y the induced metric on Y . As $\min T''$ is continuous and locally the minimum of differentiable functions, it is Lipschitz. Thus by compactness of ∂X_2 , making Y smaller if necessary, there exists $E \geq 0$ such that $|\min T''(y_1) - \min T''(y_2)| \leq E d_Y(y_1, y_2)$ for all $y_1, y_2 \in Y$. Now let y, s, z be as above. Then y, z are on different sheets of Y in X locally, but these sheets should intersect in X , so we can choose $y', z' \in \partial X_2$ such that y' is close to y in Y , and z' is close to z in Y , and $\iota(y') = \iota(z')$.

Since $\min T'|_{\partial^2 X}$ is σ -invariant, it follows that $\min T'$ pushes down to a function on $\iota(\partial X) \subset X$. Hence $\min T''|_{\partial X_2}$ pushes down to a function on $\iota(\partial X_2) \subset X$, and $\iota(y') = \iota(z')$ implies $\min T''(y') = \min T''(z')$. Think of $P = \iota(y)$, $Q = \iota(z)$ and $R = \iota(y') = \iota(z')$ as the vertices of a small triangle in X . The side PR has 'length' $d_Y(y, y')$. The side QR has 'length' $d_Y(z, z')$. The side PQ has 'length' at most $s\|v\|_{C^0}$, where $\|v\|_{C^0}$ is computed using g . The angle in PQR at R is the angle at $\iota(y')$ between the two sheets of $\iota(Y)$ containing $\iota(y), \iota(z)$. By compactness of $\partial^2 X_2$, and using Definition A.9(iv),(v), there is a global, positive lower bound for all angles of different sheets of Y meeting at $\iota(\partial^2 X_2)$. Hence there exists small $\epsilon > 0$ independent of y, s, z such that $|PQ| \geq \epsilon(|PR| + |QR|)$. So we have

$$\begin{aligned} s\|v\|_{C^0} &\geq \epsilon(d_Y(y, y') + d_Y(z, z')) \\ &\geq \epsilon E^{-1}(|\min T''(y) - \min T''(y')| + |\min T''(z) - \min T''(z')|) \\ &\geq \epsilon E^{-1}(|\min T''(y) - \min T''(z)|), \end{aligned}$$

since $\min T''(y') = \min T''(z')$. Hence if $\theta > \|v\|_{C^0} E \epsilon^{-1}$ then $\min T''(y) + \theta s > \min T''(z)$ for all such y, s, z , as we have to prove.

One can now show that T is a *tent function* by a modification of the proof of Proposition A.10, using the fact that T'' is a tent function. We can then give an expression for $\min T$: if $(x, B) \in \partial X$ and $s \in [0, 2]$ then

$$\min T(\exp(sv)x) = \min(\min T'(x, B) + \theta s, D). \quad (201)$$

This holds as putting $t = 2 - s$, we have $t \in (0, 2]$ and $\exp(tv)\exp(sv)x = \exp(2v)x = \iota(\exp(2v)(x, B))$, where $\exp(2v)(x, B) \in \partial X_2$, and $T''(\exp(2v)(x, B)) = T'(x, B)$. Hence by (195), $T(\exp(sv)x)$ is the union of $\{D\}$, and $\{u + \theta s : u \in T'(x, B)\}$, and contributions from $y \in Y \setminus \partial X_2$ which from above cannot be minimum in $T(\exp(sv)x)$. Equation (201) follows. When $s = 0$, it implies that $\min T|_{\partial X} \equiv \min T'$, as we want.

Since $\min T' \geq 1$ as T' is a tent function, and $\theta > D - 1$, we see from (201) that $\min T(\exp(sv)x) = D$ if $(x, B) \in \partial X$ and $s \in [1, 2)$. Hence $\min T \equiv D$ on $X_1 \setminus X_2$. But (195) gives $T(x) = \{D\}$ for $x \in X_2$, so $\min T \equiv D$ on X_2 , and so on X_1 . Thus $X_1 \subset \tilde{X}$. Since $\min T' < D$ on ∂X and $\min T|_{\partial X} \equiv \min T'$ we have $\min T < D$ on $\iota(\partial X_1)$. Therefore $X_1 \subset \tilde{X} \subset X^\circ$, as we have to prove.

We can now describe the pieces X_c for $c \in C$ in (187) for $Z_{X,T}$. Each X_c for $c \in C$ is the graph of a smooth branch of T on a connected component of the subset where it minimizes T . Therefore $\{D\} \times \tilde{X}$ is a union of some of the X_c in C , those associated to the constant branch $x \mapsto D$ of T . Hence $\tilde{X}$ is a compact, embedded submanifold of X with g-corners. We define Φ on $\tilde{X}$ by $\Phi : x \mapsto (D, x)$, and this is a diffeomorphism of $\tilde{X}$ with these X_c with $\pi_X \circ \Phi(x) = x$ for $x \in \tilde{X}$, as we want.

For the remaining X_c for $c \in C$ on which $\min T \neq D$, we can see from (201) that these correspond to $X'_{c'}$ for $c' \in C'$, since the smooth branches of T on their minimal subsets in X correspond to the smooth branches of T' on their minimal subsets in ∂X . In fact, if $c' \in C'$ then

$$X_c = \{(u + \theta s, \exp(sv)x) : (u, (x, B)) \in X'_{c'}, 0 \leq \theta s \leq D - u\} \quad (202)$$

is one of the X_c . We define a diffeomorphism $\Phi : [0, 1] \times X'_{c'} \rightarrow X_c$ by

$$\Phi : (t, (x, B)) \mapsto \left((1-t) \min T'(x, B) + tD, \exp\left(\frac{t}{\theta(D - \min T'(x, B))} v\right) x \right).$$

Equations (198) and (199) follow. For Ψ in (197) and (199), we note that $\Phi|_{\{1\} \times \coprod_{c' \in C'} X'_{c'}}$ maps $\coprod_{c' \in C'} X'_{c'}$ to the faces of X_c in (202) on which $\min T \equiv D$, and these correspond to faces of $\tilde{X}$. The remainder of the proposition follows as for Proposition A.10. $\square$

Here is a generalization of Proposition A.11. The proof is easy.

Proposition A.14. *Suppose X is a compact manifold with g-corners, and $T' : \partial X \rightarrow F([1, \infty))$ is a tent function, and $\zeta : \partial X \rightarrow \mathbb{R}$ is a piecewise smooth function subordinate to T' , such that $\min T'|_{\partial^2 X}$ and $\zeta|_{\partial^2 X}$ are invariant under the natural involution $\sigma : \partial^2 X \rightarrow \partial^2 X$.*

Let T be as in Definition A.12, and $\tilde{X}, \Phi, \Psi$ be as in Proposition A.13. Choose any smooth function $\tilde{\eta} : \tilde{X} \rightarrow \mathbb{R}$. Then there is a piecewise smooth function $\eta : X \rightarrow \mathbb{R}$ subordinate to the tent function T with $\eta|_{\partial X} \equiv \zeta$, given by

$$\eta(x) = \begin{cases} \tilde{\eta}(x), & x \in \tilde{X}, \\ (1-t)\zeta \circ \pi'(x') + t\tilde{\eta}|_{\partial \tilde{X}} \circ \Psi(x'), & \begin{matrix} x = \pi_X \circ \Phi(t, x'), \\ (t, x') \in [0, 1] \times (\coprod_{c' \in C'} X'_{c'}). \end{matrix} \end{cases} \quad (203)$$

Similarly, if $E \rightarrow X$ is a vector bundle and f is a piecewise smooth section of $E|_{\partial X}$ subordinate to T' with $\min T'|_{\partial^2 X}$ and $f|_{\partial^2 X}$ σ -invariant, then by choosing a smooth section $\tilde{e}$ of $E|_{\tilde{X}}$ we can write down a piecewise smooth section e of E subordinate to T with $e|_{\partial X} \equiv f$.

A.1.5 Cutting a manifold into small pieces with boundary conditions

We now combine the ideas of §A.1.2 and §A.1.4. Given a compact manifold X and a tent function T' on ∂X with $T'|_{\partial^2 X}$ σ -invariant, we construct a tent function T on X with $\min T|_{\partial X} \equiv \min T'$, which cuts X into ‘arbitrarily small pieces’ X_c for $c \in C$, as far as this is possible given the boundary conditions.

Definition A.15. Let X be a compact n -manifold with g-corners and $T' : \partial X \rightarrow F([1, \infty))$ a tent function such that $\min T'|_{\partial^2 X}$ is invariant under the involution $\sigma : \partial^2 X \rightarrow \partial^2 X$. Let $v, \iota : Y \rightarrow X, T'', D, \theta$ be as in Definition A.12.

Choose a Riemannian metric g on X . Note that g is not required to take any particular values on ∂X — it is unrelated to any metric used to construct T' . Choose $\epsilon > 0$ with $2\epsilon < \delta(g)$, the injectivity radius of g . Write $d(\cdot, \cdot)$ for the metric on X determined by g . For $p \in X$ and $r > 0$ write $B_r(p)$ for the open ball of radius r about p . Then $x \mapsto d(p, x)^2$ is smooth on $B_{2\epsilon}(p)$, as $2\epsilon < \delta(g)$.

The open sets $B_\epsilon(p)$ for $p \in X^\circ$ cover X , so as X is compact we can choose a finite subset $\{p_1, \dots, p_N\}$ of X° with $X = \bigcup_{i=1}^N B_\epsilon(p_i)$. As in Definition A.3, choose $p_1, \dots, p_N$ to be *generic* amongst all such N -tuples $p'_1, \dots, p'_N$. Since $p_i \in X^\circ$ and ∂X is compact, there exists $\zeta \in (0, \epsilon)$ such that $d(p_i, x) \geq \zeta$ for all $i = 1, \dots, N$ and $(x, B) \in \partial X$. Make θ larger if necessary so that $\theta > D + \epsilon^2 - 1$ and $1 + \theta\zeta\|v\|_{C^0} > D$, where $\|v\|_{C^0}$ is computed using g . Define $T : X \rightarrow F([1, \infty))$ by

$$T(x) = \{D + d(p_i, x)^2 : i = 1, \dots, N, \ d(p_i, x) < 2\epsilon\} \cup \{u + \theta(2 - t) : t \in (0, 2], \ y \in Y, \ u \in T''(y), \ \exp(tv)x = \iota(y)\}. \quad (204)$$

Proposition A.16. In Definition A.15, T is a tent function on X , and $\min T' \equiv \min T|_{\partial X}$. Let $X'_{c'}$ for $c' \in C'$ be as in (187) for $Z_{\partial X, T'}$, and X_c for $c \in C$ be as in (187) for $Z_{X, T}$. Then we can take $C = C' \amalg \{1, \dots, N\}$, where for $c' \in C'$ there is a natural identification between $X'_{c'}$ and a component of $\partial X_{c'}$, and for $i = 1, \dots, N$ we have $p_i \in \pi(X_i^\circ) \subseteq \pi(X_i) \subseteq B_\epsilon(p_i)$.

We can choose T so that the pieces X_c are ‘arbitrarily small’, given the $X'_{c'}$ which are already fixed. That is, if $\{V_i : i \in I\}$ is any open cover for X such that for all $c' \in C'$ there exists $i \in I$ with $\pi'(X'_{c'}) \subseteq \partial V_i$, then we can choose T so that for all $c \in C$ there exists $i \in I$ with $\pi(X_c) \subseteq V_i$.

Proof. The proof is a straightforward generalization of those of Propositions A.4 and A.13. The first and second lines of (204) are analogues of (188) and (195) respectively. Since $X = \bigcup_{i=1}^N B_\epsilon(p_i)$, the minimum of the terms $D + d(p_i, x)^2$ in the first line of (204) is a continuous function on X with values between D and $D + \epsilon^2$. The condition $\theta > D + \epsilon^2 - 1$ strengthening $\theta > D - 1$ in Definition A.12 ensures that in the second line of (204), $u + \theta(2 - t)$ is never minimal in $T(x)$ for $t \in [1, 2)$, which helps to ensure Definition A.1(c) holds and $\min T$ is continuous. It follows as in Propositions A.4 and A.13 that T is a *tent function*, and $\min T' \equiv \min T|_{\partial X}$.

For $i = 1, \dots, N$, we claim that $\min T(p_i) = D$, and $D + d(p_i, x)^2$ is the only branch of T achieves this minimum. For $j \neq i$ we have $d(p_j, p_i) > 0$, so $D + d(p_i, x)^2 > D$ at $x = p_i$. Suppose $t \in (0, 2]$, $y \in Y$, $u \in T''(y)$ and $\exp(tv)x = \iota(y) = p_i$ such that $u + \theta(2 - t)$ is minimal amongst the branches of the second line of (204) at p_i . Then as in the proof of Proposition A.13, we have $y \in \partial X_2$, so $y = \exp(2v)(x', B')$ for $(x', B') \in \partial X$. Hence $p_i = \exp((2 - t)v)x'$. Therefore $(2 - t)\|v\|_{C^0} \geq \zeta$, since $d(p_i, x') \geq \zeta$. Also $u = \min T'(x', B') \geq 1$. The condition $1 + \theta\zeta\|v\|_{C^0} > D$ thus gives $u + \theta(2 - t) > D$, proving the claim.

The pieces X_c for $c \in C$ are thus of two kinds: the branches $D + d(p_i, x)^2$ of T for $i = 1, \dots, N$ from the first line of (204) are minimal near p_i and so yield components X_i with $p_i \in \pi(X_i^\circ) \subseteq \pi(X_i) \subseteq B_\epsilon(p_i)$, as in Proposition A.4. The remaining X_c come from minimal branches $u + \theta(2 - t)$ from the second line of (204), and these correspond to minimal branches of T' on ∂X and so to components $X_{c'}$, with $\pi(X_{c'}) \subset X$ the union of segments of flow lines of v starting in $\pi(X_{c'}) \subset \iota(\partial X)$. This proves the next part of the proposition. The final part holds provided $\epsilon > 0$ is taken sufficiently small with fixed g and θ is taken sufficiently large, since then the $\pi(X_{c'})$ for $c' \in C'$ become very thin neighbourhoods of the $\pi(X_{c'})$, and the $\pi(X_i)$ for $i = 1, \dots, N$ are contained in small balls $B_\epsilon(p_i)$. $\square$

A.1.6 Cutting a manifold into simplices with boundary conditions

We would also like to extend Lemma A.5 and Theorem A.7 to the case of prescribed boundary data. However, we cannot do this with arbitrary $T' : \partial X \rightarrow F([1, \infty))$, since Lemma A.5 relies on taking ϵ very small so that the pieces $\pi(X_c)$ approximate small convex polyhedra in $\mathbb{R}^n$, but as in Proposition A.16 the $X_{c'}$ for $c' \in C'$ appear as components of ∂X_c , and the $X_{c'}$ depend on T' and are of a fixed size, so this prevents us making the X_c arbitrarily small.

To get round this, we assume that T' has been constructed using Lemma A.5 or Theorem A.7 (or, recursively, using Lemma A.17 or Theorem A.18 below with ∂X in place of X), using a sufficiently small constant $\epsilon' > 0$, where ‘sufficiently small’ depends not just on ∂X and choices made during the construction of T' , but also on X and choices to be made during the construction of T . Thus, T' depends not only on ∂X , it also has a minimal dependence on X as well. The argument here appears circular: we are trying to choose T' depending on T , and T depending on T' . However, it is not circular, as the only data on X that ϵ' must depend on are the choices of g, v, D in Definition A.15, and these are independent of T' , that is, they are chosen without reference to T' .

Here is our generalization of Lemma A.5. There is one subtle point: at the junction between a region $\pi(X_i)$ in X coming from a branch $D + d(p_i, x)^2$ of T in the first line of (204), and a region $\pi(X_{c'})$ in X coming from a branch $u + \theta(2 - t)$ in the second line of (204), the common boundary may not approximate a hyperplane $\mathbb{R}^{n-1}$ in $\mathbb{R}^n$, but rather a piece of a quadric in $\mathbb{R}^n$. The condition $\theta \inf_{\partial X} |v| \gg \epsilon^2$ ensures this piece is approximately flat and parallel to ∂X , so we can still approximate $\pi(X_i)$ and $\pi(X_{c'})$ by compact convex polyhedra.

Lemma A.17. *Suppose X is a compact manifold with g -corners. Let $T' :$*

$\partial X \rightarrow F([1, \infty))$ be constructed in Definition A.3 (or Definition A.15) using some metric g' on ∂X and constant $\epsilon' > 0$, and suppose $\min T'|_{\partial^2 X}$ is invariant under the natural involution $\sigma : \partial^2 X \rightarrow \partial^2 X$. Let $T : X \rightarrow F([1, \infty))$ be constructed in Definition A.15 for this T' , using some metric g on X , vector field v and constants D, θ, ϵ . We regard g', g, v, D as chosen in advance and fixed, and $\epsilon', \epsilon, \theta$ as satisfying inequalities which may depend on g', g, v, D . In particular, ϵ' can depend on g, v, D .

Suppose that ϵ is small compared to the natural length-scales of (X, g) and v , that ϵ' is small compared to the natural length-scales of $(X, g), v, (\partial X, g')$ and constants comparing $g|_{\partial X}$ and g' , and that θ is chosen large enough that $\theta \inf_{\partial X} |v| \gg \epsilon^2$. Then geodesic normal coordinates in X approximately identify the $\pi(X_c) \subset X$ for $c \in C$ with compact convex polyhedra in $\mathbb{R}^n$.

For $c' \in C'$ Lemma A.5 approximately identifies $\pi'(X'_{c'}) \subset \partial X$ with a compact convex polyhedron in $\mathbb{R}^{n-1}$. Also, Proposition A.16 identifies $X'_{c'}$ with a component of ∂X_c , and we have approximately identified X_c with a compact convex polyhedron in $\mathbb{R}^n$. Combining these three gives an approximate identification of a compact convex polyhedron in $\mathbb{R}^{n-1}$ with a codimension one face of a compact convex polyhedron in $\mathbb{R}^n$. This is approximately affine.

Here is our generalization of Theorem A.7. The proof is an easy modification, as we have already done the hard work in Definition A.15 and Lemma A.17. The main points are to take T' and $\sigma'_{(i', \mathbf{F}')}$ to be built in Theorem A.7 using data on ∂X in Lemma A.5 chosen with ϵ' sufficiently small as in Lemma A.17. Then when we choose the $\tilde{\sigma}_{(i, \mathbf{F}, \Delta)}$ inductively in Theorem A.7, those lying over ∂X rather than X° are equal to $\sigma'_{(i', \mathbf{F}')}|_{\Delta'}$ for some natural choice of $i', \mathbf{F}', \Delta'$.

In the last part of Lemma A.17 we comment that identifications of $X'_{c'}$ and a component of ∂X_c with convex polyhedra in $\mathbb{R}^{n-1}$ and $\mathbb{R}^n$ are related by an approximately affine transformation. The point here is that we have two metrics g' and $g|_{\partial X}$ on ∂X , with geodesic normal coordinates on $X'_{c'}$ are defined using g' , and on the component of ∂X_c using $g|_{\partial X}$ (approximately). Since ϵ' is chosen small compared to the length scales of both g' and $g|_{\partial X}$ and any comparison between them, the geodesic normal coordinates on ∂X using g' and $g|_{\partial X}$ differ by approximately affine transformations at length scale ϵ' . This is important in Theorem A.17, since the barycentric subdivision of a compact convex polyhedron is invariant under affine transformations, so the barycentric subdivision steps in Theorem A.7 for T' and Theorem A.18 for T are compatible.

In (ii) below, only the last sentence is an assumption, the rest follows from Theorem A.7 for T' and from (i).

Theorem A.18. *Suppose X is a compact manifold with g -corners. Let $T' : \partial X \rightarrow F([1, \infty))$, $X'_{c'}$, $c' \in C'$ and diffeomorphisms $\sigma'_{c'} : \Delta_{n-1} \rightarrow X'_{c'}$ for $c' \in C'$ be constructed in Theorem A.7 (or Theorem A.18) for ∂X . Suppose:*

- (i) *Assume $\min T'|_{\partial^2 X}$ is invariant under the involution $\sigma : \partial^2 X \rightarrow \partial^2 X$.*
- (ii) *The maps $\pi' \circ \sigma'_{c'} : \Delta_{n-1} \rightarrow \partial X$ for $c' \in C'$ are a triangulation of ∂X by $(n-1)$ -simplices. They induce a triangulation of $\partial^2 X$ by $(n-2)$ -simplices, of the following form: let $F_{n-1}^{n-1} : \Delta_{n-2} \rightarrow \Delta_{n-1}$ be as in §4.1.*

Let $\check{C}'$ be the subset of $c' \in C'$ for which $\pi' \circ \sigma'_{c'} \circ F_{n-1}^{n-1} : \Delta_{n-2} \rightarrow \partial X$ maps Δ_{n-2} to $\iota(\partial^2 X)$. For each $c' \in \check{C}'$ there is a unique embedding $\check{\sigma}'_{c'} : \Delta_{n-2} \rightarrow \partial^2 X$ with $\iota \circ \check{\sigma}'_{c'} \equiv \pi' \circ \sigma'_{c'} \circ F_{n-1}^{n-1}$, where $\iota : \partial^2 X \rightarrow \partial X$ is the natural immersion. Then $\check{\sigma}'_{c'}$ for $c' \in \check{C}'$ are a triangulation of $\partial^2 X$ by $(n-2)$ -simplices. Part (i) implies that the set of images $\check{\sigma}'_{c'}(\Delta_{n-2})$ in $\partial^2 X$ for $c' \in \check{C}'$ is σ -invariant, since $\check{\sigma}'_{c'}(\Delta_{n-2})$ are the $\pi(X_c)$ for $T'|_{\partial^2 X}$.

Assume the $\check{\sigma}'_{c'}$ for $c' \in \check{C}'$ are invariant under $\sigma : \partial^2 X \rightarrow \partial^2 X$, in the sense that if $\sigma(\check{\sigma}'_{c'}(\Delta_{n-2})) = \check{\sigma}'_{\tilde{c}'}(\Delta_{n-2})$ for $c', \tilde{c}' \in \check{C}'$ then $\sigma \circ \check{\sigma}'_{c'} \equiv \check{\sigma}'_{\tilde{c}'}$.

- (iii) Let the initial application of Definition A.3 in the construction of T' use some metric g' on ∂X and constant $\epsilon' > 0$. Assume ϵ' is chosen small enough, compared to geometry on X , that Lemma A.17 applies.

Then we can construct a tent function $T : X \rightarrow F([1, \infty))$, with $\min T|_{\partial X} \equiv T'$, such that the components X_c , $c \in C$ of $\partial Z_{X,T}$ in (187) are all diffeomorphic to the n -simplex Δ_n , with diffeomorphisms $\sigma_c : \Delta_n \rightarrow X_c$ for $c \in C$, and for each $c' \in C'$ there exists a unique $c \in C$ such that $\iota \circ \pi' \circ \sigma'_{c'} \equiv \pi \circ \sigma_c \circ F_n^n$ as maps $\Delta_{n-1} \rightarrow X$, where $\iota : \partial X \rightarrow X$ is the natural immersion. We can also choose the σ_c to have the boundary compatibility described in Theorem A.7 under $\sigma : \partial^2 Z_{X,T} \rightarrow \partial^2 Z_{X,T}$.

The reason why we use $F_{n-1}^{n-1} : \Delta_{n-2} \rightarrow \Delta_{n-1}$ in (ii), and $F_n^n : \Delta_{n-1} \rightarrow \Delta_n$ in the last part, rather than just referring to some codimension 1 faces of Δ_{n-1} and Δ_n , is that the barycentric subdivision of a convex n -polyhedron K in Definition A.6 automatically triangulates ∂K by $(n-1)$ -simplices $\sigma_F \circ F_n^n(\Delta_{n-1})$, so we know that the $(n-2)$ -simplices in the triangulation of $\partial^2 X$ induced by $\sigma_{c'}$, $c' \in C'$ are all of the form $\pi' \circ \sigma'_{c'} \circ F_{n-1}^{n-1}(\Delta_{n-2})$ for some $c' \in C'$, and similarly the $(n-1)$ -simplices in the triangulation of ∂X induced by σ_c , $c \in C$ are all of the form $\pi \circ \sigma_c \circ F_n^n(\Delta_{n-1})$ for some $c \in C$.

A.2 Tent functions on orbifolds

Next we extend §A.1 to orbifolds X . Here is the analogue of Definition A.1.

Definition A.19. Let X be an n -orbifold with g-corners. Then as in §2.2, X is covered by orbifold charts (U, Γ, ϕ) , where Γ acts linearly on $\mathbb{R}^n$, $U \subseteq \mathbb{R}^n$ is a Γ -invariant region with g-corners, and $\phi : U/\Gamma \rightarrow X$ is a homeomorphism with an open set $\phi(U/\Gamma) \subseteq X$. Write $\pi : U \rightarrow U/\Gamma$ for the natural projection. Then $\phi \circ \pi$ maps $U \rightarrow X$. A function $T : X \rightarrow F([1, \infty))$ is called a *tent function* if for all orbifold charts (U, Γ, ϕ) on X , the map $T \circ \phi \circ \pi : U \rightarrow F([1, \infty))$ is a tent function on the n -manifold U , in the sense of Definition A.1.

Equivalently, T is a tent function if for each $x \in X$ there exists a small orbifold chart (U, Γ, ϕ) with $x \in \phi(U/\Gamma)$, open subsets $U_1, \dots, U_N \subseteq U$ and smooth functions $t_i : U_i \rightarrow [1, \infty)$ for $i = 1, \dots, N$ satisfying Definition A.1(a)–(c), such that $T \circ \phi \circ \pi(u) = \{t_i(u) : i = 1, \dots, N, u \in U_i\}$ for all $u \in U$. We can take the t_i to be locally distinct, that is, $t_i \neq t_j$ in any nonempty subset of

$U_i \cap U_j$, and the U_j to be connected. Then U and $T \circ \phi \circ \pi$ determine the U_i, t_i uniquely up to permutations of $\{1, \dots, N\}$.

Since $T \circ \phi \circ \pi$ is invariant under the action of Γ on U , it follows that Γ acts on $\{1, \dots, N\}$ such that $\gamma(U_i) = U_{\gamma \cdot i}$ and $t_{\gamma \cdot i} \circ \gamma \equiv t_i$ for all $\gamma \in \Gamma$ and $i = 1, \dots, N$. Note that we do *not* require this action of Γ on $\{1, \dots, N\}$ to be trivial. This means that tent functions on an orbifold X in general *cannot* be written locally as the union of finitely many smooth, single-valued functions on X near orbifold strata of X , it really is necessary to lift to orbifold charts.

Much of §A.1 extends from manifolds to orbifolds with almost no change; this holds for §A.1.4, for instance. We will comment only on new issues in the orbifold case. There is one significant idea in Definition A.3 and Proposition A.4 which does not work for orbifolds, and needs revision. It is this: in Definition A.3, with X a compact manifold, g a Riemannian metric on X , and $d(\cdot, \cdot)$ the metric induced by g , we used the notion of *injectivity radius* $\delta(g) > 0$, and the fact that for any $p \in X$ and $0 < r < \delta(g)$ the map $B_r(p) \rightarrow [0, r^2)$ taking $q \mapsto d(p, q)^2$ is smooth.

For orbifolds this is no longer true: if X is an orbifold and g, d are as above, then for any $r > 0$, if $p \in X$ is sufficiently close to an orbifold stratum of X , then the map $B_r(p) \rightarrow [0, r^2)$ taking $q \mapsto d(p, q)^2$ is only piecewise smooth, not smooth. We get round this by replacing $q \mapsto d(p, q)^2$ by a smooth, *multivalued* function measuring the squared lengths of all geodesic segments of length less than r joining p and q .

Here are our generalizations of Definition A.3 and Proposition A.4.

Definition A.20. Let X be a compact n -orbifold with g-corners. Choose a Riemannian metric g on X , and write $d(\cdot, \cdot)$ for the metric on X determined by g . As X is compact we can choose a finite system of orbifold charts (U^j, Γ^j, ϕ^j) for $j \in J$ with $X = \bigcup_{j \in J} \phi^j(U^j/\Gamma^j)$. Write $\pi^j : U^j \rightarrow U^j/\Gamma^j$, so that $\phi^j \circ \pi^j$ maps $U^j \rightarrow X$. Define $g^j = (\phi^j \circ \pi^j)^*(g)$, a Riemannian metric on U^j . Write $d^j(\cdot, \cdot)$ for the metric on U^j induced by g^j . Even though U^j may not be compact, one can use the compactness of X to show that g^j has a positive injectivity radius $\delta(g^j) > 0$ on U^j , so that $u \mapsto d^j(p, u)^2$ is a smooth map $B_r^j(p) \rightarrow [0, r^2)$ for all $p \in U$ and $0 < r \leq \delta(g^j)$, where $B_r^j(p)$ is the open ball of radius r about p in U defined using d^j .

Choose $\epsilon > 0$ satisfying two conditions: (a) $2\epsilon < \delta(g^j)$ for all $j \in J$, and (b) for all $p \in X$, there exists $j \in J$ such that $B_{3\epsilon}(p) \subseteq \phi^j(U^j/\Gamma^j)$. This holds for all sufficiently small ϵ as J is finite, $\delta(g^j) > 0$ and X is compact. The open sets $B_\epsilon(p)$ for $p \in X$ cover X , so as X is compact we can choose a finite subset $\{p_1, \dots, p_N\}$ of X with $X = \bigcup_{c=1}^N B_\epsilon(p_c)$. Choose $p_1, \dots, p_N$ to be *generic* amongst all such N -tuples $p'_1, \dots, p'_N$, as in Definition A.3. Define $T : X \rightarrow F([1, \infty))$ by

$$T(x) = \{1 + l^2 : l \in [0, 2\epsilon), \text{ there exists a geodesic segment of length } l \text{ in } (X, g) \text{ with end points } x \text{ and } p_c \text{ for some } c = 1, \dots, N\}. \quad (205)$$

Note that T does not depend on the choice of (U^j, Γ^j, ϕ^j) , $j \in J$, except through

the smallness conditions (a),(b) on ϵ . These are a substitute for the condition $2\epsilon < \delta(g)$ in Definition A.3, and are needed because the notion of injectivity radius does not behave well on orbifolds.

Proposition A.21. *In Definition A.20, T is a tent function on X . In (187), we can take the indexing set C to be $\{1, \dots, N\}$ with $p_c \in \pi(X_c^\circ) \subseteq \pi(X_c) \subseteq B_\epsilon(p_c)$ for $c = 1, \dots, N$. We can choose T so that the pieces X_c are ‘arbitrarily small’, in the sense that if $\{V_i : i \in I\}$ is any open cover for X then we can choose T so that for all $c = 1, \dots, N$ there exists $i \in I$ with $\pi(X_c) \subseteq V_i$.*

*If X is an **effective** orbifold then the X_c for $c \in C$ in (187) are compact **manifolds** with g -corners.*

Proof. Let $x \in X$. Then by choice of $\epsilon > 0$ in Definition A.20, there exists $j \in J$ such that $B_{3\epsilon}(p) \subseteq \phi^j(U^j/\Gamma^j)$. Choose $y \in U^j$ with $\phi \circ \pi(y) = x$. Let $\Gamma = \{\gamma \in \Gamma^j : \gamma \cdot y = y\}$, the stabilizer group of y in Γ^j . Then if $\gamma \in \Gamma^j \setminus \Gamma$ we have $y \neq \gamma \cdot y$, so $d^j(y, \gamma \cdot y) > 0$. Choose $\delta \in (0, \epsilon]$ such that $d^j(y, \gamma \cdot y) \geq 2\delta$ for all $\gamma \in \Gamma^j \setminus \Gamma$, and define $U = B_\delta^j(y)$, an open neighbourhood of y in U^j . Then U is a Γ -invariant subset of U^j , and $U \cap \gamma \cdot U = \emptyset$ for all $\gamma \in \Gamma^j \setminus \Gamma$. Hence the projection $U/\Gamma \rightarrow U^j/\Gamma^j$ given by $\Gamma u \mapsto \Gamma^j u$ is a homeomorphism with its image. Define $\phi : U/\Gamma \rightarrow X$ by $\phi : \Gamma u \mapsto \phi^j(\Gamma^j u)$. Then (U, Γ, ϕ) is an orbifold chart on X with $x \in \phi(U/\Gamma)$.

Write $q_1, \dots, q_M$ for the points of $B_{2\epsilon+\delta}^j(y) \cap (\phi^j \circ \pi^j)^{-1}(\{p_1, \dots, p_N\})$. That is, $q_1, \dots, q_M$ are those preimages of $p_1, \dots, p_N$ in U^j with distance less than $2\epsilon + \delta$ from y . Since $|(\phi^j \circ \pi^j)^{-1}(\{p_1, \dots, p_N\})| \leq N|\Gamma^j|$ this is a finite set, and $M \leq |\Gamma^j|N$. The action of Γ on $B_{2\epsilon+\delta}^j(y) \subseteq U^j$ permutes $q_1, \dots, q_M$, and $\gamma \in \Gamma^j \setminus \Gamma$ may also identify distinct q_a, q_b . For $a = 1, \dots, M$, define an open set $U_a \subseteq U$ by $U_a = U \cap B_{2\epsilon}(q_j)$, and define a smooth function $t_a : U_a \rightarrow [1, \infty)$ by $t_a(u) = 1 + d^j(q_a, u)^2$. It is now easy to check from (205) that

$$T \circ \phi \circ \pi : u \mapsto \{t_a(u) : a = 1, \dots, M, u \in U_a\}. \quad (206)$$

We claim that Definition A.1(a)–(c) hold for these $U_1, \dots, U_M, t_1, \dots, t_M$. The proof is a little different to that in the manifold case in Proposition A.4, and the extra property $\dim S_{\{i_1, \dots, i_l\}} = n - l + 1$ for all $\{i_1, \dots, i_l\} \subseteq \{1, \dots, N\}$ with $l \geq 2$ in Proposition A.4 in general will not hold here. The point is that although $p_1, \dots, p_N$ are generic in X , as each p_i in X lifts to finitely many q_a in U , the M -tuple $q_1, \dots, q_M$ may *not* be generic in U .

However, $(\phi^j \circ \pi^j)^{-1}(\{p_1, \dots, p_N\})$ is generic amongst Γ^j -invariant finite subsets of U^j , and this is enough to imply Definition A.1(a),(b). The conditions $d^j(q_{i_1}, u) = \dots = d^j(q_{i_l}, u)$ may intersect non-transversely, so that $\dim S_{\{i_1, \dots, i_l\}} > n - l + 1$, but this can happen only if a nontrivial subset of $\{q_{i_1}, \dots, q_{i_l}\}$ is preserved by a nontrivial subgroup G of Γ^j , and $S_{\{i_1, \dots, i_l\}}$ is contained in the fixed-point set $\text{Fix}(G)$ of G in U^j . Since $\text{Fix}(G)$ is a submanifold of U^j , one can show $S_{\{i_1, \dots, i_l\}}$ is a submanifold of U . Therefore $T \circ \phi \circ \pi$ in (206) is a tent function on the manifold U . As every $x \in X$ has an orbifold chart (U, Γ, ϕ) with $x \in \phi(U/\Gamma)$ and $T \circ \phi \circ \pi$ a tent function on U , equation (205) defines a tent function on the orbifold X , as we have to prove.

The second and third parts of the proposition, that we can take $C = \{1, \dots, N\}$ and choose the X_c arbitrarily small, follow as in Proposition A.4. For the final part, suppose X is an *effective* orbifold. Then Γ^j acts effectively on U^j for each $j \in J$. Let $x \in X$, and define $(U, \Gamma, \phi), U_1, \dots, U_M, t_1, \dots, t_M$ as above. Write $\pi : Z_{X,T} \rightarrow X$ for the projection. Then there is a diffeomorphism

$$Z_{X,T} \supseteq \pi^{-1}(\phi(U/\Gamma)) \cong \{(t, u) : u \in U, 0 \leq t \leq \min T \circ \phi \circ \pi(u)\} / \Gamma = Z_{U, T \circ \phi \circ \pi} / \Gamma.$$

Points of $\coprod_{c \in C} X_c \subset \partial Z_{X,T}$ are of the form $((t, x'), B)$ for $(t, x') \in Z_{X,T}$ with $t = \min T(x')$, and B a local boundary component of $Z_{X,T}$ containing (t, x') . Lifting from $Z_{X,T}$ up to $Z_{U, T \circ \phi \circ \pi}$, choices of B correspond to choices of branch t_b of $T \circ \phi \circ \pi$ inducing the value $\min T(x')$, for $b = 1, \dots, M$. Therefore

$$\coprod_{c \in C} X_c \supseteq (\pi \circ \iota)^{-1}(\phi(U/\Gamma)) \cong \{(t, u, b) : u \in U, t = \min T \circ \phi \circ \pi(u), b = 1, \dots, M, \min T \circ \phi \circ \pi(u) = t_b(u)\} / \Gamma, \quad (207)$$

where $\iota : \coprod_{c \in C} X_c \rightarrow Z_{X,T}$ is the natural immersion.

Here in (207), Γ acts trivially on $t \in [0, \infty)$, in the usual way on $u \in U$, and on $b \in \{1, \dots, M\}$, the action is induced from the action of Γ on $\{q_1, \dots, q_M\} \subset U$. Now Γ acts effectively on U , and $p_1, \dots, p_N$ are generic in X , so that each q_b is generic in U . Together these imply that the stabilizer group of each q_b in Γ is trivial, so Γ acts freely on $\{1, \dots, M\}$. Thus the action of Γ in (207) is free, so the quotient $(\pi \circ \iota)^{-1}(\phi(U/\Gamma))$ is a manifold. Since $\coprod_{c \in C} X_c$ is covered by such open sets $(\pi \circ \iota)^{-1}(\phi(U/\Gamma))$, we see that $\coprod_{c \in C} X_c$ and each X_c for $c \in C$ are manifolds. This completes the proof. $\square$

Remark A.22. In Remark 2.10(b), we explained that for orbifolds X with corners or g-corners, restricting to *orbifold strata* $X^{\Gamma, \rho}$ in the sense of §5.6 does not commute with taking boundaries, and therefore no information from orbifold strata $X^{\Gamma, \rho}$ survives in Kuranishi homology. Proposition A.21 is an illustration of this. Given some Kuranishi cycle K defined using effective orbifolds X , Propositions A.2 and A.21 enable us to construct a homologous Kuranishi cycle K' defined using manifolds X_c for $c \in C$, where the homology between K and K' is defined using the orbifolds $Z_{X,T}$. This works because $\partial(Z_{X,T}^{\Gamma, \rho}) \not\cong (\partial Z_{X,T})^{\Gamma, \rho}$. So, a homology theory defined using effective orbifolds with (g-)corners as chains is equivalent to a homology theory defined using manifolds as chains. This will be important in making the X_{ac} have trivial stabilizers in the proof of Step 1 in §B.1.

The rest of §A.1 extends to orbifolds X without any further significant problems or changes, except those already explained in Definitions A.19–A.20 and Proposition A.21. In fact we will not need analogues of §A.1.3 and §A.1.6 for orbifolds, but we will need versions of §A.1.4–§A.1.5. Section A.1.4 holds for orbifolds with essentially no change. The choice of vector field v and flows $\exp(tv)$ work for orbifolds X as well as manifolds; note that v is automatically tangent to each orbifold stratum $X^{\Gamma, \rho}$ of X , for Γ a finite group, and $\exp(tv)$ takes $X^{\Gamma, \rho}$ to $X^{\Gamma, \rho}$. In §A.1.5 we restrict to *effective* orbifolds, so that we can

use the last part of Proposition A.21, and we must modify the first line of (204) as in (205), and the proof of Proposition A.16 as in Proposition A.21.

Theorem A.23. *The results of §A.1.4–§A.1.5 all hold when X is a compact n -orbifold with g -corners, restricting to effective orbifolds in §A.1.5. In Proposition A.16, if X is an effective orbifold and the $X'_{c'}$ are manifolds for all $c' \in C'$, then the X_c are manifolds for all $c \in C$.*

A.3 Tent functions on Kuranishi chains $[X, \mathbf{f}, \mathbf{G}], [X, \mathbf{f}, \underline{\mathbf{G}}]$

In §A.3.1 we discuss some of the new problems in generalizing §A.1–§A.2 from manifolds and orbifolds to Kuranishi chains $[X, \mathbf{f}, \mathbf{G}], [X, \mathbf{f}, \underline{\mathbf{G}}]$, and how we solve them. Section A.3.2 defines tent functions on Kuranishi chains, and §A.3.3–§A.3.5 give analogues of sections A.1.2, A.1.4 and A.1.5 respectively. We will not need Kuranishi chain analogues of §A.1.3 and §A.1.6.

A.3.1 Preliminary discussion

Let X be a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ be strongly smooth, and $\mathbf{G}$ (or $\underline{\mathbf{G}}$) be (effective) gauge-fixing data for $(X, \mathbf{f})$. Then $\mathbf{G}, \underline{\mathbf{G}}$ include an excellent coordinate system $(\mathbf{I}, \boldsymbol{\eta})$ for $(X, \mathbf{f})$, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i) : i \in I, \dots)$. Roughly speaking, a *tent function* for $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \underline{\mathbf{G}})$ is $\mathbf{T} = (T^i : i \in I)$, where $T^i : V^i \rightarrow F([1, \infty))$ is a tent function on V^i for each $i \in I$, such that if $j \leq i$ in I then $\min T^j|_{V^{ij}} \equiv \min T^i \circ \phi^{ij}$.

However, we need to modify this definition a little, to make it easier to construct tent functions $\mathbf{T}$. Here are the two main new problems we will meet in trying to extend the results of §A.1–§A.2 to Kuranishi chains:

- (a) One thing we will need to do very often below is to choose some smooth data δ^i on V^i for each $i \in I$, (such as a Riemannian metric g , or a transverse perturbation $\tilde{s}^i$ of the Kuranishi map s^i), with compatibility conditions $\delta^j|_{V^{ij}} \equiv (\phi^{ij})^*(\delta^i)$ when $j \leq i$ in I . Our basic method for doing this is to choose δ^i by induction on increasing $i \in I$, where $\delta^i|_{\phi^{ij}(V^{ij})}$ for all $j < i$ in I is (partially) prescribed by $\delta^j|_{V^{ij}} \equiv (\phi^{ij})^*(\delta^i)$ and the choice of δ^j in a previous inductive step.

Since $\phi^{ij}(V^{ij})$ need not be closed in V^i , we may not be able to extend smooth prescribed values for δ^i on $\phi^{ij}(V^{ij})$ smoothly to V^i . In fact, if $\phi^{ij}(V^{ij})$ is badly placed in V^i , then smooth prescribed values for δ^i on $\phi^{ij}(V^{ij})$ may not even extend continuously to the closure $\overline{\phi^{ij}(V^{ij})}$ of $\phi^{ij}(V^{ij})$ in V^i . Thus, the inductive step may not be possible.

- (b) All our results in §A.1–§A.2 involve *compact* manifolds and orbifolds. In particular, compactness is used to choose finite covers by balls $B_\epsilon(p_c)$. However, the V^i are in general noncompact orbifolds.

Here is an example of problem (a).

Example A.24. Let X be the compact topological space $\{(x, y) \in [-1, 1]^2 : xy = 0\}$. Define Kuranishi neighbourhoods (V^1, E^1, s^1, ψ^1) , (V^2, E^2, s^2, ψ^2) on X by $V^1 = \{(x, 0) : 0 < |x| \leq 1\} \cup \{(0, y) : 0 < |y| \leq 1\}$, and $E^1 = V^1$, regarded as the zero vector bundle over V^1 , $s^1 \equiv 0$ and $\psi^1 = \text{id}_{V^1}$, and $V^2 = [-1, 1]^2$, $E^2 = \mathbb{R} \times V^2$, $s^2 : V^2 \rightarrow \mathbb{R}$ is $s^2(x, y) = xy$, and $\psi^2 = \text{id}_X$. Then X has a unique Kuranishi structure whose Kuranishi neighbourhoods $(V_p, \dots, \psi_p)$ are equivalent to $(V^1, \dots, \psi^1)$ for $p \neq (0, 0)$ and to $(V^2, \dots, \psi^2)$ for $p = (0, 0)$.

We can complete $(V^1, \dots, \psi^1), (V^2, \dots, \psi^2)$ to a very good coordinate system for X , with indexing set $I = \{1, 2\}$, and $V^{21} = V^1$. Suppose we want to choose some continuous or smooth data for each $(V^i, \dots, \psi^i)$ by induction over i in $I = \{1, 2\}$ with compatibilities on V^{ij} . Having chosen data for $(V^1, \dots, \psi^1)$, we have to choose data for $(V^2, \dots, \psi^2)$ taking prescribed values over the subset V^1 of V^2 . In general this is *not possible*, even continuously, near $(0, 0)$ in $\bar{V}^1 \setminus V^1$, since the four different ends of V^1 at $(0, 0)$ can prescribe four different limits at $(0, 0)$, if the limits exist at all.

This example also illustrates a different problem with choosing *transverse perturbations*, which appears to have been overlooked in the proofs of Fukaya and Ono [25, Th. 6.4], [24, Th. A1.23]. To carry out Step 2 in §B.2, we would like to choose small, smooth, transverse perturbations $\tilde{s}^1, \tilde{s}^2$ of the Kuranishi maps s^1, s^2 , satisfying $\hat{\phi}^{21} \circ \tilde{s}^1 \equiv \tilde{s}^2 \circ \phi^{21}$ on V^{21} . This is not possible near $(0, 0)$, not even using multisections, and not because bad choices of $\tilde{s}^1$ may not extend smoothly, but for a different reason. As E^1 is the zero vector bundle, the only choice for $\tilde{s}^1$ is $\tilde{s}^1 \equiv 0$, and thus we must have $\tilde{s}^2 \equiv 0$ on V^1 , and hence on $\bar{V}^1$ in V^2 . But this implies $\tilde{s}^2|_{(0,0)} = d\tilde{s}^2|_{(0,0)} = 0$, and $\tilde{s}^2$ is not transverse at $(0, 0)$.

Our solution to problem (a) above, which we used in the proof of Proposition 3.15 above, is to replace the V^i by smaller open subsets $\dot{V}^i$. We now define these $\dot{V}^i$ using the partition of unity data $\boldsymbol{\eta}$ in $(\mathbf{I}, \boldsymbol{\eta})$, and complete the $\dot{V}^i$ to a modified really good coordinate system $(\dot{\mathbf{I}}, \dot{\boldsymbol{\eta}})$. The advantage of doing it this way, rather than choosing some arbitrary smaller subsets $\dot{V}^i \subset V^i$, is that passing from $(\mathbf{I}, \boldsymbol{\eta})$ to $(\dot{\mathbf{I}}, \dot{\boldsymbol{\eta}})$ is functorial in chains $[X, \mathbf{f}, \mathbf{G}]$ or $[X, \mathbf{f}, \underline{\mathbf{G}}]$, and compatible with relations in $KC_*, KC_*^{\text{ef}}(Y; R)$ that we need to preserve.

Definition A.25. Let X be a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ be strongly smooth, and $(\mathbf{I}, \boldsymbol{\eta})$ an excellent coordinate system for $(X, \mathbf{f})$, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i) : i \in I, \dots)$ and $\boldsymbol{\eta} = (\eta_i : i \in I, \eta_i^j : i, j \in I)$.

Let $\vartheta > 0$ be small. For each $j \in I$, define $\dot{\eta}_i^j : V^j \rightarrow [0, 1]$ for $i \in I$ by

$$\dot{\eta}_i^j(v) = \text{mid}(0, (1 + \vartheta)\eta_i^j(v) + c(v), 1), \quad (208)$$

where $\text{mid}(x, y, z)$ is the middle one of x, y, z , that is, $\text{mid}(x, y, z) = y$ when $x \leq y \leq z$ or $z \leq y \leq x$, and so on, and $c(v) \in \mathbb{R}$ is chosen so that $\sum_{i \in I} \dot{\eta}_i^j(v) = 1$. From (208) we see that $\sum_{i \in I} \dot{\eta}_i^j(v)$ is a continuous, monotone increasing function of $c(v)$ with minimum 0 and maximum $|I| \geq 1$, so there exists $c(v)$ with $\sum_{i \in I} \dot{\eta}_i^j(v) = 1$ by the Intermediate Value Theorem, and this defines the $\dot{\eta}_i^j(v)$ (though not necessarily $c(v)$) uniquely. It is easy to see that the $\dot{\eta}_i^j$ are continuous. Define continuous $\dot{\eta}_i : X \rightarrow [0, 1]$ from the η_i in the same way.

By construction $\sum_{i \in I} \dot{\eta}_i^j \equiv 1$ and $\sum_{i \in I} \dot{\eta}_i \equiv 1$, as in Definition 3.4(i),(ii), Definition 3.4(iii) implies that $\dot{\eta}_i^j|_{(s^j)^{-1}(0)} \equiv \dot{\eta}_i \circ \psi^j$ for all $i, j \in I$, and Definition 3.4(iv) implies that if $i, j, k \in I$ with $k \leq j$ then $\dot{\eta}_i^k|_{V^{jk}} \equiv \dot{\eta}_i^j \circ \phi^{jk}$. Define open subsets $\dot{V}^i \subset V^i$ and $\dot{V}^{ij} \subset V^{ij}$ by $\dot{V}^i = \{v \in V^i : \dot{\eta}_i^i(v) > 0\}$ and $\dot{V}^{ij} = \{v \in V^{ij} : \dot{\eta}_i^j(v) > 0\}$. Let $\dot{E}^i, \dot{s}^i, \dot{\psi}^i$ be the restrictions of E^i, s^i, ψ^i to $\dot{V}^i$ for all $i \in I$, and $\dot{\phi}^{ij}, \dot{\phi}^{ij}$ the restrictions of ϕ^{ij}, ϕ^{ij} to $\dot{V}^{ij}$. Set $\dot{I} = \{i \in I : \dot{V}^i \neq \emptyset\}$.

When $\vartheta > 0$, the effect of (208) is to move the points $\eta_i^j(v)$ in $[0, 1]$ away from the middle and towards the end points 0,1. Thus, if $\eta_i^j(v) > 0$ is small then $\dot{\eta}_i^j(v)$ is smaller, and may be zero. So we have $\text{supp}(\dot{\eta}_i^j) \subseteq \text{supp}(\eta_i^j)$, and $\dot{V}^i \subseteq V^i$. When $\vartheta = 0$ we have $\dot{\eta}_i^j = \eta_i^j$ and $\dot{V}^i = V^i$. Also $\dot{\eta}_i^j$ varies continuously with ϑ , and $\text{supp}(\eta_i^j)$ is a monotone decreasing, lower semicontinuous function of ϑ , so $\dot{V}^i$ and $\text{Im } \dot{\psi}^i$ are also monotone decreasing and lower semicontinuous in ϑ . Hence, when $\vartheta > 0$, is small, $\dot{V}^i, \text{Im } \dot{\psi}^i$ are just a little smaller than $V^i, \text{Im } \psi^i$.

We have $X = \bigcup_{i \in I} \text{Im } \psi^i$, where X is compact, and $\{\text{Im } \psi^i : i \in I\}$ is a finite open cover for X . Being a finite open cover of a compact space is not changed by making all the sets a little smaller. Therefore, for sufficiently small $\vartheta > 0$, we have $X = \bigcup_{i \in I} \text{Im } \dot{\psi}^i$. Let $\vartheta > 0$ be chosen small enough to achieve this.

It is not difficult to show that apart from the condition $X = \bigcup_{i \in I} \text{Im } \dot{\psi}^i$, all the conditions for this data $\dot{I}, (\dot{V}^i, \dots, \dot{\psi}^i), \dots$ to define a really good coordinate system follow from the corresponding conditions on $I, (V^i, \dots, \psi^i), \dots$. Thus we have defined a *really good coordinate system* $(\dot{I}, \dot{\eta})$ for $(X, \mathbf{f})$. We can apply Algorithm 3.16 to $(\dot{I}, \dot{\eta})$ to get an excellent coordinate system $(\check{I}, \check{\eta})$ if we wish.

We shall show that if $j \leq i$ in I then $\phi^{ij}(\dot{V}^{ij}) \subseteq \phi^{ij}(V^{ij})$, where $\phi^{ij}(\dot{V}^{ij})$ is the closure of $\phi^{ij}(\dot{V}^{ij})$ in V^i . This property will be important in problems involving choosing smooth data on the $\dot{V}^i$ with compatibilities over the $\dot{V}^{ij}$. Set $\epsilon = \frac{\vartheta}{(1+\vartheta)|I|} > 0$. Suppose $\eta_j^j(v) \leq \epsilon$ for some $v \in V^j$. We will prove that $\dot{\eta}_j^j(v) = 0$. Since $\sum_{i' \in I} \eta_{i'}^j \equiv 1$, the average value of the $\eta_{i'}^j(v)$ for $i' \in |I|$ is $\frac{1}{|I|} > \epsilon$, so $\eta_j^j(v)$ is not the maximum of the $\eta_{i'}^j(v)$. If any $\eta_{i''}^j(v) = 1$ then $\eta_{i''}^j(v)$ must be maximum amongst the $\eta_{i'}^j(v)$, so $i'' \neq i$, and thus $\dot{\eta}_j^j(v) = 0$. So suppose $\eta_{i'}^j(v) \neq 1$ for any $i' \in I$. Then (208) gives $\dot{\eta}_{i'}^j(v) \geq (1+\vartheta)\eta_{i'}^j(v) + c(v)$, so summing over $i' \in |I|$ and using $\sum_{i' \in I} \eta_{i'}^j \equiv \sum_{i' \in I} \dot{\eta}_{i'}^j \equiv 1$ implies that $c(v) \leq -\frac{\vartheta}{|I|}$. But then $(1+\vartheta)\eta_j^j(v) + c(v) \leq 0$, so (208) gives $\dot{\eta}_j^j(v) = 0$, as we want. Thus $\dot{\eta}_j^j = 0$ wherever $\eta_j^j \leq \epsilon$.

Since $\dot{\eta}_j^j > 0$ on $\dot{V}^{ij}$ we have $\eta_j^j > \epsilon$ on $\dot{V}^{ij}$, so $\eta_j^i > \epsilon$ on $\phi^{ij}(\dot{V}^{ij})$ by Definition 3.4(iv), and thus $\eta_j^i \geq \epsilon > 0$ on $\phi^{ij}(\dot{V}^{ij})$. But $\phi^{ij}(V^{ij})$ is closed in $\{v \in V^i : \eta_j^i(v) > 0\}$ by Definition 3.4(ii), and $\phi^{ij}(\dot{V}^{ij})$ lies in the intersection of the closure of $\phi^{ij}(V^{ij})$ and $\{v \in V^i : \eta_j^i(v) > 0\}$, so $\phi^{ij}(\dot{V}^{ij}) \subseteq \phi^{ij}(V^{ij})$.

We can now explain our solutions to problems (a) and (b) above.

Remark A.26. Here is how to use Definition A.25 to avoid problem (a) above. Suppose we want to choose some smooth data δ^i on $\dot{V}^i$ for $i \in I$ with $\delta^j|_{\dot{V}^{ij}} \equiv$

$\delta^i \circ \dot{\phi}^{ij}$ when $j \leq i$ in I . Then, what we actually do is to choose δ^i on $\dot{V}^i$ by induction on increasing $i \in I$, with $\delta^j|_{\dot{V}^{ij}} \equiv \delta^i \circ \dot{\phi}^{ij}$ for all $j < i$ in I , such that δ^i extends smoothly to an open neighbourhood U^i of the closure $\overline{\dot{V}^i}$ of $\dot{V}^i$ in V^i . Thus, in the inductive step where we choose δ^i , we have prescribed values for δ^i on the subsets $\dot{\phi}^{ij}(\dot{V}^{ij})$ of V^i for $j < i$ in I . The prescribed values are consistent on the overlaps $\dot{\phi}^{ij}(\dot{V}^{ij}) \cap \dot{\phi}^{ik}(\dot{V}^{ik})$ for $k < j < i$ because $\delta^k|_{\dot{V}^{jk}} \equiv \delta^j \circ \dot{\phi}^{jk}$.

Since $\overline{\dot{\phi}^{ij}(\dot{V}^{ij})} \subseteq \dot{\phi}^{ij}(V^{ij})$ as in Definition A.25, we see that $\overline{\dot{\phi}^{ij}(\dot{V}^{ij})} = \dot{\phi}^{ij}(\overline{\dot{V}^{ij}}) \subseteq \dot{\phi}^{ij}(V^{ij} \cap U^j)$. As δ^j extends smoothly to U^j , it follows that the prescribed values for δ^i on $\dot{\phi}^{ij}(\dot{V}^{ij})$ extend smoothly to $\dot{\phi}^{ij}(\overline{\dot{V}^{ij}})$ in V^i , and thus, to a neighbourhood of $\dot{\phi}^{ij}(\dot{V}^{ij})$ in V^i . This overcomes problem (a).

Here is how we deal with problem (b) above. In choosing data such as a tent function T^i on $\dot{V}^i$, the values of T^i only really matter near $\dot{V}^i \cap (s^i)^{-1}(0)$, since this is the part of $\dot{V}^i$ identified with X . So we can use compactness of X to control the T^i near the $\dot{V}^i \cap (s^i)^{-1}(0)$, and then take T^i to be constant away from $\dot{V}^i \cap (s^i)^{-1}(0)$, for instance.

Note that when we apply Definition A.25 we must choose some $\vartheta > 0$, which must be chosen small enough that $X = \bigcup_{i \in I} \text{Im } \psi^i$. When we do this in Appendices B and C, we will generally be working not with one $X, \mathbf{f}, (\mathbf{I}, \boldsymbol{\eta})$ but with a finite collection of $X_a, \mathbf{f}_a, (\mathbf{I}_a, \boldsymbol{\eta}_a)$ for $a \in A$. We choose the same ϑ for all $a \in A$, and make it small enough that $X_a = \bigcup_{i \in I_a} \text{Im } \psi_a^i$ for all $a \in A$, which is possible as A is finite.

A.3.2 The definition of tent functions on Kuranishi chains

Definition A.27. Let X be a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ be strongly smooth, and either $\mathbf{G}$ be gauge-fixing data for $(X, \mathbf{f})$, or $\underline{\mathbf{G}}$ be effective gauge-fixing data for $(X, \mathbf{f})$. Then $\mathbf{G}$ or $\underline{\mathbf{G}}$ includes an excellent coordinate system $(\mathbf{I}, \boldsymbol{\eta})$ for $(X, \mathbf{f})$, where $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i) : i \in I, \dots)$. Let $(\dot{\mathbf{I}}, \dot{\boldsymbol{\eta}})$ be as in Definition A.25, where $\dot{\mathbf{I}} = (I, (\dot{V}^i, \dot{E}^i, \dot{s}^i, \dot{\psi}^i) : i \in I, \dots)$, depending on the choice of some small $\vartheta > 0$.

A *tent function* for $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \underline{\mathbf{G}})$ is $\mathbf{T} = (T^i : i \in I)$, where $T^i : \dot{V}^i \rightarrow F([1, \infty))$ is a tent function on the orbifold $\dot{V}^i$ for each $i \in I$, such that T^i extends to a tent function on an open neighbourhood of the closure $\overline{\dot{V}^i}$ of $\dot{V}^i$ in V^i , and if $j \leq i$ in I then $\min T^j|_{\dot{V}^{ij}} \equiv \min T^i \circ \dot{\phi}^{ij}$, and the submanifolds $S_{\{i_1, \dots, i_l\}}$ of Definition A.1 for T^i intersect $\dot{\phi}^{ij}(\dot{V}^{ij})$ transversely in $\dot{V}^i$ wherever $t_{i_1}(u) = \min T^i(u)$. When $\dot{V}^i$ is an orbifold, this last condition must be expressed in orbifold charts, as follows. Suppose (U, Γ, ϕ) is an orbifold chart for $\dot{V}^i$. Since $\dot{\phi}^{ij}(\dot{V}^{ij})$ is an embedded suborbifold of $\dot{V}^i$, $V = (\phi \circ \pi)^{-1}(\dot{\phi}^{ij}(\dot{V}^{ij}))$ is a submanifold of U . By Definition A.19, $T^i \circ \phi \circ \pi : U \rightarrow F([1, \infty))$ is a tent function on the manifold U . We require that each of the submanifolds $S_{\{i_1, \dots, i_l\}}$ of U given in Definition A.1 for $T = T^i \circ \phi \circ \pi$ should intersect V transversely wherever $t_{i_1}(u) = \min T^i \circ \phi \circ \pi(u)$.

If $\mathbf{T}, \tilde{\mathbf{T}}$ are tent functions for $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \underline{\mathbf{G}})$ then we write $\min \mathbf{T} = \min \tilde{\mathbf{T}}$ if $\min T^i \equiv \min \tilde{T}^i$ on $\dot{V}^i$ for all $i \in I$. If $\mathbf{T} = (T^i : i \in I)$ is a tent function

for $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \underline{\mathbf{G}})$ then $\mathbf{T}|_{\partial X} = (T_{\partial}^{i-1} = T^i|_{\partial \dot{V}^i} : i \in I, \partial V^i \neq \emptyset)$ is a tent function for $(\partial X, \mathbf{f}|_{\partial X}, \mathbf{G}|_{\partial X})$ or $(\partial X, \mathbf{f}|_{\partial X}, \underline{\mathbf{G}}|_{\partial X})$. Often, given some tent function $\mathbf{T}'$ for $(\partial X, \mathbf{f}|_{\partial X}, \mathbf{G}|_{\partial X})$, we will wish to construct a tent function $\mathbf{T}$ for $(X, \mathbf{f}, \mathbf{G})$ with $\min(\mathbf{T}|_{\partial X}) = \min \mathbf{T}'$.

For tent functions T on an oriented manifold or orbifold X , we defined an oriented manifold or orbifold $Z_{X,T}$ in (186), with $\partial Z_{X,T}$ given by (187). Here are analogues of these for tent functions on (effective) Kuranishi chains.

Definition A.28. Suppose X is a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ strongly smooth, and $\mathbf{G}$ (or $\underline{\mathbf{G}}$) is (effective) gauge-fixing data for $(X, \mathbf{f})$. Let $\mathbf{T}$ be a tent function for $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \underline{\mathbf{G}})$. Since the continuous functions $\min T^i : \dot{V}^i \rightarrow [1, \infty)$ for $i \in I$ satisfy $\min T^i \circ \dot{\phi}^{ij} \equiv \min T^j|_{\dot{V}^{ij}}$, they induce a unique continuous function $\min T : X \rightarrow [1, \infty)$ with $\min T \circ \dot{\psi}^i \equiv \min T^i|_{(\dot{s}^i)^{-1}(0)}$ for $i \in I$. As in (186), define $Z_{X,T}$ as a compact topological space by

$$Z_{X,T} = \{(t, x) \in [0, \infty) \times X : t \leq \min T(x)\}.$$

We will define an oriented Kuranishi structure on $Z_{X,T}$ and (effective) gauge-fixing data $\mathbf{H}_{X,T}$ or $\underline{\mathbf{H}}_{X,T}$ for $(Z_{X,T}, \mathbf{f} \circ \pi)$, where $\pi : Z_{X,T} \rightarrow X$ is the projection. Set $J = \{i+1 : i \in I\}$. For each $i \in I$, define

$$W^{i+1} = [0, \frac{1}{2}] \times V^i \amalg \{(t, v) \in [\frac{1}{2}, \infty) \times \dot{V}^i : t \leq \min T^i(v)\} \subset V^i \times [0, \infty). \quad (209)$$

Then W^{i+1} is an *orbifold with g-corners*, as $T^i : \dot{V}^i \rightarrow F([1, \infty))$ is a tent function, by the orbifold version of Proposition A.2. Write $\pi_{V^i} : W^{i+1} \rightarrow V^i$ for the projection. Define an orbifold vector bundle $F^{i+1} \rightarrow W^{i+1}$ by $F^{i+1} = \pi_{V^i}^*(E^i)$, a section t^{i+1} of F^{i+1} by $t^{i+1} = s^i \circ \pi_{V^i}$, and a continuous map $\xi^{i+1} : (t^{i+1})^{-1}(0) \rightarrow Z_{X,T}$ by $\xi^{i+1}(t, v) = (t, \psi^i(v))$. Define a smooth map $g^{i+1} : W^{i+1} \rightarrow Y$ by $g^{i+1}(t, v) = f^i(v)$. Then (W^j, F^j, t^j, ξ^j) for $j \in J$ are *Kuranishi neighbourhoods* on $Z_{X,T}$.

If $j \leq i$ in I , define an open subset $W^{(i+1)(j+1)}$ in W^{j+1} by $W^{(i+1)(j+1)} = W^{j+1} \cap ([0, \infty) \times V^{ij})$, and define a *coordinate change* $(\psi^{(i+1)(j+1)}, \hat{\psi}^{(i+1)(j+1)})$ from $(W^{(i+1)(j+1)}, \dots, \xi^{j+1}|_{W^{(i+1)(j+1)}})$ to $(W^{i+1}, \dots, \xi^{i+1})$ by $\psi^{(i+1)(j+1)}(t, v) = (t, \phi^{ij}(v))$ and $\hat{\psi}^{(i+1)(j+1)}(t, e) = (t, \hat{\phi}^{ij}(e))$. The condition in Definition A.27 that the $S_{\{i_1, \dots, i_l\}}$ for T^i should intersect $\dot{\phi}^{ij}(\dot{V}^{ij})$ transversely in $\dot{V}^i$ ensures that $\psi^{(i+1)(j+1)}$ is compatible with boundaries and g-corners, as in Definition 2.18(a).

Since X is an oriented Kuranishi space, $[0, \infty) \times X$ is an oriented Kuranishi space. It is now easy to check that there is a unique oriented Kuranishi structure on $Z_{X,T}$ such that $Z_{X,T}$ is a Kuranishi subspace of $[0, \infty) \times X$, with natural strongly smooth map $\pi : Z_{X,T} \rightarrow X$ induced by the projection $[0, \infty) \times X \rightarrow X$, and $\mathbf{J} = (J, (W^j, F^j, t^j, \xi^j), g^j : j \in J, (W^{jk}, \psi^{jk}, \hat{\psi}^{jk}) : k \leq j \in J)$ is a *very good coordinate system* for $(Z_{X,T}, \mathbf{f} \circ \pi)$.

For $i, j \in I$, define continuous $\zeta_{i+1} : Z_{X,T} \rightarrow [0, 1]$, $\zeta_{i+1}^{j+1} : W^{j+1} \rightarrow [0, 1]$ by

$$\begin{aligned}\zeta_{i+1}(t, x) &= \begin{cases} (1 - 2t)\eta_i(x) + 2t\dot{\eta}_i(x), & t \in [0, \frac{1}{2}), \\ \dot{\eta}_i(x), & t \geq \frac{1}{2}, \end{cases} \\ \zeta_{i+1}^{j+1}(t, v) &= \begin{cases} (1 - 2t)\eta_i^j(v) + 2t\dot{\eta}_i^j(v), & t \in [0, \frac{1}{2}), \\ \dot{\eta}_i^j(v), & t \geq \frac{1}{2}. \end{cases}\end{aligned}\quad (210)$$

Write $\zeta = (\zeta_j : j \in J, \zeta_j^k : j, k \in J)$. Since $(\mathbf{I}, \boldsymbol{\eta})$ and $(\dot{\mathbf{I}}, \dot{\boldsymbol{\eta}})$ are both really good coordinate systems for $(X, \mathbf{f})$, one can show that $(\mathbf{J}, \zeta)$ is a *really good coordinate system* for $(Z_{X,T}, \mathbf{f} \circ \pi)$.

Let $(\check{\mathbf{J}}, \check{\zeta})$ be the excellent coordinate system for $(Z_{X,T}, \mathbf{f} \circ \pi)$ constructed from $(\mathbf{J}, \zeta)$ by Algorithm 3.16. Then $\check{W}^j$ is an open subset of W^j for each $j \in J$. As $(\mathbf{I}, \boldsymbol{\eta})$ is excellent, no pieces of $[0, \frac{1}{2}) \times V^i \subseteq W^{i+1}$ are discarded, so $[0, \frac{1}{2}) \times V^i \subseteq \check{W}^{i+1}$ for all $i \in I$. This implies $\check{J} = J = \{i + 1 : i \in I\}$.

If we have gauge-fixing data $\mathbf{G}$ for $(X, \mathbf{f})$, define $H^{i+1} : \check{W}^{i+1} \rightarrow P$ for each $i \in I$ by $H^{i+1}(t, v) = G^i(v)$ if $t = 0$, and $H^{i+1}(t, v) = S_{n+1}(t, x_1, \dots, x_n)$ if $t > 0$ and $G^i(v) = S_n(x_1, \dots, x_n)$, and write $\mathbf{H}_{Z,T} = ((\check{\mathbf{J}}, \check{\zeta}), H^j : j \in \check{J})$. If we have effective gauge-fixing data $\underline{\mathbf{G}}$ for $(X, \mathbf{f})$, define $\underline{H}^{i+1} : \check{W}^{i+1} \rightarrow \underline{P}$ for each $i \in I$ by $\underline{H}^{i+1}(t, v) = \underline{G}^i(v)$ if $t = 0$, and $\underline{H}^{i+1}(t, v) = (t, x_1, \dots, x_n)$ if $t > 0$ and $\underline{G}^i(v) = (x_1, \dots, x_n)$, and write $\underline{\mathbf{H}}_{Z,T} = ((\check{\mathbf{J}}, \check{\zeta}), \underline{H}^j : j \in \check{J})$. Then $\mathbf{H}_{Z,T}$ or $\underline{\mathbf{H}}_{Z,T}$ is (effective) gauge-fixing data for $(Z_{X,T}, \mathbf{f} \circ \pi)$.

If $\mathbf{T}, \tilde{\mathbf{T}}$ are tent functions for $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \underline{\mathbf{G}})$ and $\min \mathbf{T} = \min \tilde{\mathbf{T}}$ then $Z_{X,T}, \pi$ and $\mathbf{H}_{X,T}$ or $\underline{\mathbf{H}}_{X,T}$ coincide with $Z_{X,\tilde{T}}, \pi, \mathbf{H}_{X,\tilde{T}}$ or $\underline{\mathbf{H}}_{X,\tilde{T}}$.

Here is the analogue of Proposition A.2. The proof is straightforward.

Proposition A.29. *In Definition A.28 we have*

$$\begin{aligned}\partial[Z_{X,T}, \mathbf{f} \circ \pi, \mathbf{H}_{X,T}] &= -[Z_{\partial X, T|_{\partial X}}, \mathbf{f}|_{\partial X} \circ \pi, \mathbf{H}_{\partial X, T|_{\partial X}}] \\ &\quad - [X, \mathbf{f}, \mathbf{G}] + \sum_{c \in C} [X_c, \mathbf{f}_c, \mathbf{G}_c],\end{aligned}\quad (211)$$

$$\begin{aligned}\partial[Z_{X,T}, \mathbf{f} \circ \pi, \underline{\mathbf{H}}_{X,T}] &= -[Z_{\partial X, T|_{\partial X}}, \mathbf{f}|_{\partial X} \circ \pi, \underline{\mathbf{H}}_{\partial X, T|_{\partial X}}] \\ &\quad - [X, \mathbf{f}, \underline{\mathbf{G}}] + \sum_{c \in C} [X_c, \mathbf{f}_c, \underline{\mathbf{G}}_c],\end{aligned}\quad (212)$$

in $KC_*(Y; R)$ and $KC_*^{\text{ef}}(Y; R)$. Here C is a finite indexing set and $(X_c, \mathbf{f}_c, \mathbf{G}_c)$ for c in C in (211) are the connected components (in the sense of Lemma 3.7) of $(\partial Z_{X,T}, \mathbf{f} \circ \pi|_{\partial Z_{X,T}}, \mathbf{H}_{X,T}|_{\partial Z_{X,T}})$ which meet $(0, \infty) \times X^\circ$, and similarly for $(X_c, \mathbf{f}_c, \underline{\mathbf{G}}_c)$ in (212). For each $c \in C$, $\pi|_{X_c} : X_c \rightarrow X$ is an orientation-preserving immersion of Kuranishi spaces which is an embedding on X_c° , with $X = \bigcup_{c \in C} \pi(X_c)$, and $\pi(X_c^\circ) \cap \pi(X_{c'}^\circ) = \emptyset$ for all $c \neq c'$ in C .

We think of this as cutting X into finitely many pieces $\pi(X_c)$ for $c \in C$.

A.3.3 Cutting a Kuranishi chain into arbitrarily small pieces

We now generalize §A.1.2 to Kuranishi chains. Here are analogues of Definitions A.3 and A.20 and Propositions A.4 and A.21.

Definition A.30. Suppose X is a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ strongly smooth, and $\mathbf{G}$ (or $\underline{\mathbf{G}}$) is (effective) gauge-fixing data for $(X, \mathbf{f})$. Let $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i) : i \in I, \dots)$ and $\dot{\mathbf{I}} = (I, (\dot{V}^i, \dot{E}^i, \dot{s}^i, \dot{\psi}^i) : i \in I, \dots)$ be as in Definition A.27.

In the following, when we choose some data on $\dot{V}^i$ and require that it should extend to an open neighbourhood U^i of $\bar{V}^i$ in V^i , we do not mean a fixed neighbourhood U^i , but a different U^i each time. We should think of U^i as getting smaller each time it is used in an inductive proof, so that in effect we are choosing a series of neighbourhoods $U_1^i, U_2^i, U_3^i, \dots$ with

$$V^i \supseteq \bar{U}_1^i \supseteq U_1^i \supseteq \bar{U}_2^i \supseteq U_2^i \supseteq \bar{U}_3^i \supseteq U_3^i \supseteq \dots \supseteq \bar{V}^i \supseteq \dot{V}^i.$$

This convention eliminates problems about extending data from U^i to $\bar{U}^i$.

By induction on increasing $i \in I$, choose a Riemannian metric g^i on $\dot{V}^i$ which extends to a neighbourhood U^i of $\bar{V}^i$ in V^i , such that whenever $j \leq i$ in I we have $(\phi^{ij})^*(g^i) \equiv g^j$ on $V^{ij} \cap U^j \cap (\phi^{ij})^{-1}(U^i)$, and $\phi^{ij}(V^{ij})$ is geodesically closed in (U^i, g^i) . This is possible as in Remark A.26. These conditions imply that the metrics d^i, d^j on $\dot{V}^i, \dot{V}^j$ induced by g^i, g^j satisfy $(\phi^{ij})^*(d^i) \equiv d^j$ on $\dot{V}^{ij}$, for small distances, and ϕ^{ij} maps g^j -geodesic segments in $\dot{V}^{ij}$ to g^i -geodesics in $\dot{V}^i$.

In Definition A.20, to construct tent functions T on an orbifold X , we introduced a finite system of orbifold charts on X , but the only way in which we used them was to determine injectivity-radius-type smallness conditions on $\epsilon > 0$. We need to do the same here, but for simplicity we will not go into detail. In brief, for each $i \in I$ we choose a finite system of orbifold charts on the orbifold $U^i \subseteq V^i$ which cover the compact subset $\bar{V}^i \cap (s^i)^{-1}(0)$, but need not cover the possibly noncompact orbifold U^i . Then we choose $\epsilon > 0$ sufficiently small such that the conditions of Definition A.20 hold for these charts on U^i with metric g^i , for all $i \in I$.

Now choose finite subsets $\{p_1^i, \dots, p_{N^i}^i\}$ in $\dot{V}^i$ for each $i \in I$ satisfying:

- (a) $\bar{V}^i \cap (s^i)^{-1}(0) \subseteq \bigcup_{b=1}^{N^i} B_\epsilon^i(p_b^i)$, where $B_r^i(p)$ is the open ball of radius r about p in (U^i, g^i) .
- (b) $\phi^{ij}(\{p_1^j, \dots, p_{N^j}^j\} \cap \dot{V}^{ij}) = \{p_1^i, \dots, p_{N^i}^i\} \cap \phi^{ij}(\dot{V}^{ij})$ for all $j < i$ in I .
- (c) $\{p_1^i, \dots, p_{N^i}^i\} \setminus \bigcup_{j \in I: j < i} \phi^{ij}(\dot{V}^{ij})$ is generic amongst finite subsets of $\dot{V}^i$, for all $i \in I$.
- (d) if $j < i$ in I and $v \in \dot{V}^{ij} \subseteq \dot{V}^j$, then either $d^j(p_b^j, v) > \epsilon$ for all $b = 1, \dots, N^j$ with $p_b^j \notin \dot{V}^{ij}$, or there exists $p_c^j \in \dot{V}^{ij}$ for some $c = 1, \dots, N^j$ such that $d^j(p_c^j, v) < d^j(p_b^j, v)$ for all $b = 1, \dots, N^j$ with $p_b^j \notin \dot{V}^{ij}$.
- (e) if $j < i$ in I and $v \in \phi^{ij}(\dot{V}^{ij}) \subseteq \dot{V}^i$, then either $d^i(p_b^i, v) > \epsilon$ for all $b = 1, \dots, N^i$ with $p_b^i \notin \phi^{ij}(\dot{V}^{ij})$, or there exists $p_c^i \in \phi^{ij}(\dot{V}^{ij})$ for some $c = 1, \dots, N^i$ such that $d^i(p_c^i, v) < d^i(p_b^i, v)$ for all $b = 1, \dots, N^i$ with $p_b^i \notin \phi^{ij}(\dot{V}^{ij})$.

Here is one way to do this. For each subset $\emptyset \neq J \subseteq I$, let $j = \min J$, and choose a finite subset

$$S_J \subset \bigcap_{i \in J} \dot{V}^{ij} \setminus \bigcup_{k \in I: k < j} \phi^{jk}(\dot{V}^{jk}) \subseteq \dot{V}^j \quad (213)$$

such that

$$(s^j)^{-1}(0) \cap \bigcap_{i \in J} \overline{\dot{V}^{ij}} \subseteq \bigcup_{p \in S_J} B_\epsilon^j(p), \quad (214)$$

where in (214), the closure is taken in U^j , and balls $B_\epsilon^j(p)$ are in (U^j, g^j) . It is possible to choose such a finite set as the left hand side of (214) is compact, and is also contained in the closure of the right hand side of (213), since the deletions $\phi^{jk}(\dot{V}^{jk})$ have strictly smaller dimension than $\dot{V}^j$, so they do not change the closure. Then for each $i \in I$, define

$$\{p_1^i, \dots, p_{N^i}^i\} = \bigcup_{\substack{\text{sets } J \text{ with } i \in J \subseteq I: \\ \text{set } j = \min J}} \phi^{ij}(S_J).$$

Then conditions (a),(b) above hold automatically. For (c), it is enough to take each S_J to be generic amongst such subsets. For (d),(e) we have to work a bit harder. Basically we should choose the S_J by induction on $|J|$, and when we choose S_K with $k = \min K$, we should ensure that for all $\emptyset \neq J \subset K$ with $J \neq K$ and $j = \min J$, each point of $\phi^{jk}(\bigcap_{i \in K} \dot{V}^{ik})$ is either distance at least ϵ from each point of S_J (which was chosen in a previous inductive step), or is closer to some point of $\phi^{jk}(S_K)$ than to any point of S_J . This is possible if the points in S_K are sufficiently close together. But to deal with some noncompactness issues we may need to make ϵ smaller, and to require the points in S_J to be close to $(s^j)^{-1}(0)$, so that we are really working in a compact subset of U^j .

Following (205), define $T^i : \dot{V}^i \rightarrow F([1, \infty))$ for each $i \in I$ by

$$T^i(v) = \{1 + l^2 : l \in [0, 2\epsilon), \text{ there exists a geodesic segment of length } l \text{ in } (\dot{V}^i, g^i) \text{ with end points } v \text{ and } p_b^i, \text{ some } b = 1, \dots, N^i\} \cup \{1 + \epsilon^2\}. \quad (215)$$

Then by a very similar proof to Proposition A.21, T^i is a *tent function* on $\dot{V}^i$. We added the extra branch $1 + \epsilon^2$ in (215) because the balls $B_\epsilon^i(p_b^i)$ for $b = 1, \dots, N^i$ need not cover $\dot{V}^i$, but only $\dot{V}^i \cap (s^i)^{-1}(0)$, so without including $1 + \epsilon^2$ we might have $T^i(v) = \emptyset$, which is not allowed.

Let $j \leq i$ in I . Then (d) above implies that $\min T^j|_{\dot{V}^{ij}}$ is the same whether we compute it using all of $p_1^j, \dots, p_{N^j}^j$, or only those p_b^j lying in $\dot{V}^{ij}$. Part (e) implies $\min T^i|_{\phi^{ij}(\dot{V}^{ij})}$ is the same whether we compute it using all of $p_1^i, \dots, p_{N^i}^i$, or only those p_b^i lying in $\phi^{ij}(\dot{V}^{ij})$. Part (a), $(\phi^{ij})^*(g^i) \equiv g^j$ on $\dot{V}^{ij}$, and $\phi^{ij}(\dot{V}^{ij})$ geodesically closed in $(\dot{V}^i, g^i)$, imply that $\min T^j|_{\dot{V}^{ij}}$ computed using only p_b^i in $\phi^{ij}(\dot{V}^{ij})$ agrees with $\min T^i|_{\phi^{ij}(\dot{V}^{ij})}$ computed using only p_b^i in $\phi^{ij}(\dot{V}^{ij})$ under ϕ^{ij} . Therefore $\min T^i \circ \phi^{ij} \equiv \min T^j|_{\dot{V}^{ij}}$, as we have to prove.

It remains only to verify the condition in Definition A.27 that the submanifolds $S_{\{i_1, \dots, i_l\}}$ of Definition A.1 for T^i intersect $\phi^{ij}(\dot{V}^{ij})$ transversely in $\dot{V}^i$

wherever $t_{i_1}(u) = \min T^i(u)$. Part (e) above implies that any branch of t_k of T^i on $\dot{\phi}^{ij}(\dot{V}^{ij})$ with $t_k(u) = \min T^i(u)$ comes either from a point p_b^i in $\dot{\phi}^{ij}(\dot{V}^{ij})$, or the extra branch $1 + \epsilon^2$ in (215); we cannot get such branches t_k from p_c^i in $\dot{V}^i \setminus \dot{\phi}^{ij}(\dot{V}^{ij})$. Using this and the genericness condition (c), one can show that the $S_{\{i_1, \dots, i_l\}}$ intersect $\dot{\phi}^{ij}(\dot{V}^{ij})$ transversely. This proves that $\mathbf{T} = (T^i : i \in I)$ is a *tent function* for $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \mathbf{\underline{G}})$.

Proposition A.31. *In Definition A.30, if $\mathbf{I}_c = (I_c, (V_c^i, \dots, \psi_c^i) : i \in I_c, \dots)$ is the very good coordinate system for $(X_c, \mathbf{f}_c)$ in $\mathbf{G}_c$ or $\mathbf{\underline{G}}_c$ in Proposition A.29, then by construction $I_c \subseteq I$ and $\pi : X_c \rightarrow X$ lifts to natural embeddings $\pi : V_c^i \rightarrow \dot{V}^i$ for $i \in I_c$. For each $i \in I_c$ there is a unique $b^{i,c} = 1, \dots, N^i$ with $p_{b^{i,c}}^i \in \pi(V_c^i)$, and $\pi(V_c^i) \subseteq B_\epsilon^i(p_{b^{i,c}}^i)$. If $j = \min I_c$ then for each $i \in I_c$ we have $p_{b^{j,c}}^i \in \dot{V}^{ij}$ and $p_{b^{i,c}}^i = \phi^{ij}(p_{b^{j,c}}^i)$.*

By taking $\epsilon > 0$ sufficiently small we can choose $\mathbf{T}$ to make the X_c and V_c^i ‘arbitrarily small’, *in* the sense that given open covers $\{U_j : j \in J\}$ for X and $\{U_j^i : j \in J^i\}$ for $\dot{V}^i \subseteq V^i$ for each $i \in I$, we can choose $\mathbf{T}$ such that for all $c \in C$ there exists $j \in J$ with $\pi(X_c) \subseteq U_j$, and for all $i \in I_c$ there exists $j \in J^i$ with $\pi(V_c^i) \subseteq U_j^i$.

In the case of **effective** gauge-fixing data $\mathbf{\underline{G}}, \mathbf{\underline{H}}, \mathbf{\underline{G}}_c$ the Kuranishi space X_c has trivial stabilizers for all $c \in C$, and the V_c^i are manifolds for all $i \in I_c$.

Proof. The first part follows quickly from the construction. To make sense of it, note that W^{i+1} in (209) has boundary composed of $\{0\} \times V^i$, and a portion lying over ∂V^i , and a connected component for each p_b^i , $b = 1, \dots, N^i$, and possibly also a piece from the branch $1 + \epsilon^2$ in (215). However, when we apply Algorithm 3.16 and pass to $\tilde{W}^{i+1}$, we discard boundary components not intersecting $(t^{i+1})^{-1}(0)$. This includes the whole of the piece from the branch $1 + \epsilon^2$, since Definition A.30(a) implies that $\min T^i < 1 + \epsilon^2$ on $(s^i)^{-1}(0) \cap \dot{V}^i$. It may also include some components from the p_b^i . Hence $\partial \tilde{W}^{i+1}$ is the union of $\{0\} \times V^i$, a portion lying over ∂V^i , and a component associated to p_b^i for a subset of $b \in \{1, \dots, N^i\}$. Definition A.30(b) then implies that the X_c for $c \in C$ must correspond to points $p_{b^{i,c}}^i$ as in the proposition. In particular, no X_c corresponds to the branch $1 + \epsilon^2$ in (215).

For the second part, note that each $\pi(V_c^i)$ is contained in $B_\epsilon^i(p_{b^{i,c}}^i)$ and intersects $(s^i)^{-1}(0) \cap \dot{V}^i$, and $(s^i)^{-1}(0) \cap \dot{V}^i$ is compact. Given any open cover $\{U_j^i : j \in J^i\}$ for $\dot{V}^i$, as $(s^i)^{-1}(0) \cap \dot{V}^i$ is compact, if $\epsilon > 0$ is sufficiently small then every ball of radius ϵ intersecting $(s^i)^{-1}(0) \cap \dot{V}^i$ must lie in U_j^i for some $j \in J^i$. Hence, if $\epsilon > 0$ is sufficiently small then each $\pi(V_c^i)$ lies in some U_j^i . For X we use a similar argument, noting that $\pi(X_c)$ has diameter at most 2ϵ in the metric on X obtained by gluing together the metrics $\psi_*^i(d^i)$ on $\text{Im } \psi^i \subseteq X$ for $i \in I$. For the last part, as $\mathbf{\underline{G}}$ is effective gauge-fixing data the V^i are effective orbifolds, so the V_c^i are manifolds as in Proposition A.21, and thus X_c has trivial stabilizers as $\text{Stab}_{X_c}(p) = \text{Stab}_{V_c^i}(v) = \{1\}$ for p in X_c and v in $(s_c^i)^{-1}(0) \subseteq V_c^i$ with $p = \psi_c^i(v)$. $\square$

A.3.4 Piecewise smooth extensions on Kuranishi chains

We extend §A.1.4 to Kuranishi chains. Here is our analogue of Definition A.9.

Definition A.32. Suppose X is a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ strongly smooth, and $\mathbf{G}$ (or $\underline{\mathbf{G}}$) is (effective) gauge-fixing data for $(X, \mathbf{f})$. Let $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i) : i \in I, \dots)$ and $\dot{\mathbf{I}} = (I, (\dot{V}^i, \dot{E}^i, \dot{s}^i, \dot{\psi}^i) : i \in I, \dots)$ be as in Definition A.27.

We will choose a smooth vector field v^i on V^i for each $i \in I$ such that:

- (a) v^i is nonzero and *inward pointing* at every point of ∂V^i .
- (b) For any $x_0 \in V^i$, there exists a smooth path $x : [0, \infty) \rightarrow V^i$ such that $x(0) = x_0$ and $\frac{dx}{dt}(t) = v^i|_{x(t)}$ for all $t \in [0, \infty)$. That is, v^i is a *complete* vector field on V^i . This implies that $\exp(tv^i) : V^i \rightarrow V^i$ is a well-defined smooth embedding for all $t \geq 0$.
- (c) There exist open neighbourhoods U^i of $\overline{V^i}$ in V^i for each $i \in I$, such that whenever $j \leq i$ in I we have $\phi_*^{ij}(v^j) \equiv v^i$ on $\phi^{ij}(V^{ij} \cap U^j) \cap U^i$.
- (d) The U^i in (c) have the following property: if $x \in V^i$ and $\exp(tv^i) \in \overline{V^i}$ for some $t \in [0, 2]$, then $x \in U^i$.

Before doing this, we pause to explain the issues involved. One of our problems is ensuring that the exponential maps $\exp(tv^i) : V^i \rightarrow V^i$ are well-defined, at least where we need them. If X is a compact orbifold without boundary and v is a smooth vector field on X then $\exp(tv) : X \rightarrow X$ is a well-defined diffeomorphism for all $t \in \mathbb{R}$. If X is a compact orbifold with boundary or (g-)corners and v is a vector field on X then $\exp(tv) : X \rightarrow X$ is a well-defined embedding for all $t \geq 0$ provided v is inward-pointing along ∂X .

However, if X is a *noncompact* orbifold and v a vector field on X then conditions for $\exp(tv) : X \rightarrow X$ to exist for $t \geq 0$ are more complicated. In effect, X has two kinds of boundary: the usual boundary ∂X , along which v must be inward-pointing for $\exp(tv)$ to exist, plus some noncompact ends, which we think of as being ‘at infinity’ in X . If some flow-line of v in X goes to infinity in finite time, then $\exp(tv)$ does not exist for $t > 0$. For instance, if $X = \mathbb{R}$ then $v = \frac{\partial}{\partial x}$ is complete, with $\exp(tv) : x \mapsto x + t$, but $v = x^2 \frac{\partial}{\partial x}$ is not complete, as the flow-line $t \mapsto (1 - t)^{-1}$ of v for $t \in [0, 1)$ goes to infinity in finite time. To avoid this kind of behaviour we have to ensure that $|v|$ does not grow too quickly near infinity in X , so that flow-lines of v can only go to infinity in infinite time.

Unfortunately, this issue of completeness of vector fields causes difficulties with compatibility of v^i, v^j under ϕ^{ij} . In general, if v^i, v^j are to be compatible on U^i, U^j as in (c), we cannot also choose v^i, v^j to be complete on U^i, U^j , because as v^j must grow slowly near the noncompact boundary of U^j , we find that the limit of $\phi_*^{ij}(v^j)$ at any point of $\phi^{ij}(\overline{V^{ij} \cap U^j}) \setminus \phi^{ij}(V^{ij} \cap U^j)$ must be zero, so v^i must be zero there by continuity, but this could contradict v^i being nonzero at boundary points in (a). Therefore we make v^i complete on V^i but not on U^i . Part (d) will be used later to ensure that the tent functions T^i on $\dot{V}^i$ we construct using v^i satisfy $\min T^j|_{\dot{V}^{ij}} \equiv \min T^i \circ \dot{\phi}^{ij}$, as in Definition A.27.

Next we explain how to choose v^i, U^i satisfying (a)–(d). Firstly, for each fixed $i \in I$, here is a way of constructing v^i satisfying (a),(b). Choose some Riemannian metric g^i on V^i . For each $x \in V^i$, define $f^i(x) = \sup\{\epsilon \in (0, 1] : \overline{B}_\epsilon^i(x) \text{ is a compact subset of } V^i\}$, where $\overline{B}_\epsilon^i(x)$ is the closed ball of radius ϵ about x in (V^i, g^i) . Then $f^i(x)$ is the minimum of 1 and the ‘distance $d(x, \infty)$ from x to infinity in V^i ’, measured using g^i , and $f^i : V^i \rightarrow (0, 1]$ is continuous.

Let v^i be a vector field on V^i which is nonzero and inward pointing at every point of ∂V^i . Then a sufficient condition for v^i to satisfy (b) is that $|v^i(x)| \leq C f^i(x)$ for some $C > 0$ and all $x \in V^i$, where $|v^i(x)|$ is computed using g^i . This is because if $d(x, \infty) = \infty$ then as flow of v moves at speed at most C then it must take infinite time to reach infinity, and if $d(x, \infty)$ is finite then $\log d(x, \infty)$ decreases with speed at most C , so $\log d(x, \infty) \rightarrow -\infty$ in infinite time, and the flow of v takes infinite time to reach infinity in V^i .

Thus, given any v^i satisfying (a), which are easy to make by joining local choices together by a partition of unity, we can make it satisfy (b) as well by multiplying it by a smooth function $F : V^i \rightarrow (0, 1]$ which approaches zero sufficiently fast at infinity that $|F(x)v^i(x)| \leq C f^i(x)$ for all $x \in V^i$ and some $C > 0$. Note too that if v^i satisfies (a),(b) and $F : V^i \rightarrow (0, 1]$ is smooth then Fv^i also satisfies (a),(b), since the flow-lines of Fv^i are just the flow-lines of v^i reparametrized to move more slowly, so they still must take infinite time to reach infinity in V^i .

Secondly, by induction on increasing $i \in I$, we choose v^i, U^i for each $i \in I$ satisfying (a)–(c). In the inductive step, having chosen v^j, U^j for all $j < i$, choosing smooth v^i, U^i satisfying (c) is possible as in Remark A.26, making the U^j for $j < i$ smaller if necessary as in Definition A.30. To choose v^i to satisfy (a) is also possible, since the prescribed values $\phi_*^{ij}(v^j)$ for $\phi^{ij}(V^{ij} \cap U^j) \cap U^i$ satisfy (a) by (a) for v^j , and (a) is an open convex condition. To make v^i satisfy (b) as well, we make some initial choice not necessarily satisfying (b), and then multiply it by some smooth function $F : V^i \rightarrow (0, 1]$ which is 1 on $\phi^{ij}(V^{ij} \cap U^j) \cap U^i$ for each $j < i$ in I and decreases sufficiently fast near infinity in V^i . For portions of $\phi^{ij}(V^{ij} \cap U^j) \cap U^i$ which go to infinity in V^i , the prescribed values for v^i are already complete by (b) for v^j , so it is not necessary to make v^i smaller there.

Finally, having chosen v^i, U^i for $i \in I$ satisfying (a)–(c) but not necessarily (d), we fix these U^i , and choose some smooth functions $F^i : V^i \rightarrow (0, 1]$ for $i \in I$ with $F^i \circ \phi^{ij} \equiv F^j$ on $V^{ij} \cap U^j \cap (\phi^{ij})^{-1}(U^i)$, such that $\tilde{v}^i = F^i v^i$ for $i \in I$ satisfy (d) with the given U^i . This holds provided F^i is sufficiently small; what we need is that the flow of $-\tilde{v}^i = -F^i v^i$ cannot start in $\overline{V}^i$ and reach $V^i \setminus U^i$ within time 2. Hence we can choose v^i, U^i for $i \in I$ satisfying (a)–(d) above.

We now continue the argument of Definition A.9. Write $V_1^i = \exp(v^i)V^i$ and $V_2^i = \exp(2v^i)V^i$. Then V_1^i, V_2^i are embedded submanifolds of $(V^i)^\circ$, and $\exp(v^i) : V^i \rightarrow V_1^i$, $\exp(2v^i) : V^i \rightarrow V_2^i$ are diffeomorphisms. The inclusion $\iota^i : \partial V_2^i \rightarrow V^i$ is an immersion, with closed image. As for Y in Definition A.9, for each $i \in I$ we extend ∂V_2^i to an $(i-1)$ -orbifold Y^i without boundary, with $\partial V_2^i \subset Y$ a closed embedded $(i-1)$ -submanifold, and extend $\iota^i : \partial V_2^i \rightarrow V^i$ to

an immersion $\iota^i : Y^i \rightarrow V^i$ satisfying Definition A.9(i)–(v) with v^i, V^i, V_1^i, V_2^i in place of v, X, X_1, X_2 , and where we only need (iii) to hold locally in V^i , that is, such a bound M should exist for each compact subset of V^i . In addition, we require the Y^i, ι^i to be compatible with coordinate changes ϕ^{ij} , in the sense that if $j < i$ in I then defining $Y^{ij} = (\iota^j)^{-1}(V^{ij} \cap U^j \cap (\phi^{ij})^{-1}(U^i))$, and open set in Y^j , there should exist an embedding $\psi^{ij} : Y^{ij} \rightarrow Y^i$ such that $\iota^i \circ \psi^{ij} \equiv \phi^{ij} \circ \iota^j$ on Y^{ij} . By Remark A.26 and the proof in Definition A.9, we can choose such Y^i, ι^i for all $i \in I$. As in (192), define $T^i : \dot{V}^i \rightarrow F([1, \infty))$ by

$$T^i(x) = \{2\} \cup \{3 - t : t \in (0, 2], \exp(tv^i)x \in \iota^i(Y^i)\} \quad (216)$$

for each $i \in I$. Write $\mathbf{T} = (T^i : i \in I)$.

The proof of Proposition A.10 shows that T^i is a *tent function* on $\dot{V}^i$ for each $i \in I$. It extends to a tent function on V^i also given by (216), so T^i certainly extends to a tent function on an open neighbourhood of $\dot{V}^i$ in V^i . The construction implies that if $j \leq i$ in I then $T^j|_{\dot{V}^{ij}} \equiv T^i \circ \phi^{ij}$, and thus $\min T^j|_{\dot{V}^{ij}} \equiv \min T^i \circ \phi^{ij}$. Note that we use (d) above to prove this: (216) and (d) imply that for each $x \in \dot{V}^{ij}$, $T^j(x)$ depends only on the values of v^j on U^j , rather than on the whole of V^j , and then (c) shows that computing $T^j(x)$ in V^j is equivalent to computing $T^i \circ \phi^{ij}(x)$ in V^i .

The suborbifolds $S_{\{i_1, \dots, i_l\}}$ of Definition A.1 for T^i are all locally either the images of some codimension k corner of V^i under the flow of v^i , or the intersection of such an image with $\iota^i(Y^i)$. Since $\phi^{ij} : \dot{V}^{ij} \rightarrow \dot{V}^i$ is an embedding which takes v^j to v^i , it follows that the $S_{\{i_1, \dots, i_l\}}$ for T^i intersect $\phi^{ij}(\dot{V}^{ij})$ transversely. Hence $\mathbf{T}$ is a *tent function* for $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \underline{\mathbf{G}})$ by Definition A.27.

We can also combine Definitions A.12 and A.32 to extend a tent function for $(\partial X, \mathbf{f}|_{\partial X}, \mathbf{G}|_{\partial X})$ to one for $(X, \mathbf{f}, \mathbf{G})$.

Definition A.33. Let X be a compact Kuranishi space, Y an orbifold, $\mathbf{f} : X \rightarrow Y$ strongly smooth, and $\mathbf{G}$ or $\underline{\mathbf{G}}$ (effective) gauge-fixing data for $(X, \mathbf{f})$. Suppose $\mathbf{T}'$ is a tent function for $(\partial X, \mathbf{f}|_{\partial X}, \mathbf{G}|_{\partial X})$ or $(\partial X, \mathbf{f}|_{\partial X}, \underline{\mathbf{G}}|_{\partial X})$ such that $\min(\mathbf{T}')|_{\partial^2 X}$ is invariant under the natural involution $\sigma : \partial^2 X \rightarrow \partial^2 X$. That is, for $\mathbf{I} = (I, (V^i, E^i, s^i, \psi^i) : i \in I, \dots)$ in $\mathbf{G}$ or $\underline{\mathbf{G}}$, for each $i \in I$ the tent function $\mathbf{T}'$ gives a tent function $T'^{(i-1)}$ on $\partial \dot{V}^i$, and we require that $\min T'^{(i-1)}|_{\partial^2 \dot{V}^i}$ should be invariant under the involution $\sigma : \partial^2 \dot{V}^i \rightarrow \partial^2 \dot{V}^i$. For simplicity, suppose also that $\min T'^{(i-1)}$ is bounded above on $\partial \dot{V}^i$ for all $i \in I$. This holds automatically in all our constructions.

We will construct a tent function $\mathbf{T}$ for $(X, \mathbf{f}, \mathbf{G}$ or $\underline{\mathbf{G}})$ with $\min(\mathbf{T}|_{\partial X}) = \min \mathbf{T}'$. Choose v^i, U^i, Y^i, ι^i for $i \in I$ as in Definition A.32. Then we have a diffeomorphism $\exp(2v^i) : \partial V^i \rightarrow \partial V_2^i$, so that $T'^{(i-1)} \circ \exp(2v^i)$ is a tent function on $\exp(2v^i)(\partial \dot{V}^i) \subseteq \partial V_2^i \subseteq Y^i$. Extend $T'^{(i-1)} \circ \exp(2v^i)$ to a tent function $T''^{(i-1)}$ on an open neighbourhood Y'^i of $\exp(2v^i)(\partial \dot{V}^i)$ in Y^i , relaxing the condition $\min T''^{(i-1)} \geq 1$. We require too that $\min T''^{(i-1)}$ should be bounded above on Y'^i . This is possible, as $\min T'^{(i-1)}$ is bounded above.

Choose $D > 1$ such that $\min T''^{(i-1)} < D$ on Y^i for all $i \in I$. As in (195), define $T^i : \dot{V}^i \rightarrow F([1, \infty))$ by

$$T^i(x) = \{D\} \cup \{u + \theta(2 - t) : t \in (0, 2], y \in Y^i, u \in T''^{(i-1)}(y), \exp(tv^i)x = \iota^i(y)\}, \quad (217)$$

where $\theta > D - 1$ is chosen large enough to satisfy conditions in Proposition A.13 for each $i \in I$. Actually, because of the noncompactness of $\dot{V}^i$ it may not be possible to choose θ sufficiently large everywhere. We can deal with this by replacing θ by smooth functions $\theta^i : \dot{V}^i \rightarrow (D - 1, \infty)$ for $i \in I$ with $\theta^i \circ \dot{\phi}^{ij} \equiv \theta^j|_{\dot{V}^{ij}}$ for $j < i$ in I , which are sufficiently large locally in $\dot{V}^i$.

The proof of Proposition A.13 now shows that T^i is a tent function on $\dot{V}^i$ for each $i \in I$, with $\min(T^i|_{\partial\dot{V}^i}) \equiv \min T'^{(i-1)}$. It extends to a tent function on an open neighbourhood of $\dot{V}^i$ in V^i , given by the same formula (217). For $j < i$ in I we find as in Definition A.32 that $\min T^j|_{\dot{V}^{ij}} \equiv \min T^i \circ \dot{\phi}^{ij}$, and the $S_{\{i_1, \dots, i_l\}}$ for T^i intersect $\dot{\phi}^{ij}(\dot{V}^{ij})$ transversely. Therefore $\mathbf{T} = (T^i : i \in I)$ is a *tent function* for $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \underline{\mathbf{G}})$, with $\min(\mathbf{T}|_{\partial X}) = \min \mathbf{T}'$.

Analogues of Definition A.8 and Propositions A.10, A.11, A.13 and A.14 hold for $\mathbf{T}$ in Definitions A.32 and A.33, so that we have a well-defined notion of *piecewise strongly smooth functions subordinate to a tent function $\mathbf{T}$* on $(X, \mathbf{f}, \mathbf{G})$, for which an Extension Principle holds. We will not write these out, as they are cumbersome to state. However, Step 2 of the proof of Theorem 4.8 in §B.2 depends on this piecewise smooth Extension Principle for effective Kuranishi chains, so we will discuss some of the ideas there.

A.3.5 Cutting into small pieces with boundary conditions

Section A.1.5 combined §A.1.2 and §A.1.4 to define tent functions T cutting a compact manifold X into arbitrarily small pieces, with boundary conditions $\min T|_{\partial X} \equiv \min T'$ over ∂X . We now combine §A.3.3 and §A.3.4 to do the same thing for Kuranishi chains. Here is our analogue of Definition A.15.

Definition A.34. Let $X, Y, \mathbf{f}, \mathbf{G}$ or $\underline{\mathbf{G}}$ and $\mathbf{T}'$ be as in Definition A.33. We will construct a tent function $\mathbf{T}$ for $(X, \mathbf{f}, \mathbf{G}$ or $\underline{\mathbf{G}})$ with $\min(\mathbf{T}|_{\partial X}) = \min \mathbf{T}'$, such that the pieces X_c in (211)–(212) can be made ‘arbitrarily small’, subject to the boundary conditions $\mathbf{T}'$.

Choose metrics g^i on $U^i \supset \dot{V}^i$ for $i \in I$, $\epsilon > 0$, finite subsets $\{p_1^i, \dots, p_{N^i}^i\}$ in $\dot{V}^i$ for $i \in I$, as in Definition A.30, with the extra condition that $p_c^i \in (\dot{V}^i)^\circ$ for all i, c . Choose v^i, U^i, Y^i, ι^i for $i \in I$ as in Definition A.32. Choose $Y^i, T''^{(i-1)}$ for $i \in I$ and $D > 1$, $\theta > D - 1$ or $\theta^i : \dot{V}^i \rightarrow (D - 1, \infty)$ as in Definition A.33. Make θ , or the functions θ^i for $i \in I$, larger if necessary so that $\theta > D + \epsilon^2 - 1$, and an analogue of the condition $1 + \theta\zeta\|v\|_{C^0} > D$ in Definition A.15 holds

locally in $\dot{V}^i$. Combining (204), (215) and (217), define $T^i : \dot{V}^i \rightarrow F([1, \infty))$ by

$$\begin{aligned} T^i(v) = & \{D + l^2 : l \in [0, 2\epsilon), \text{ there exists a geodesic segment of length } l \\ & \text{in } (\dot{V}^i, g^i) \text{ with end points } v \text{ and } p_b^i, \text{ some } b = 1, \dots, N^i\} \cup \{D + \epsilon^2\} \cup \\ & \{u + \theta(2 - t) : t \in (0, 2], y \in Y'^i, u \in T''^{(i-1)}(y), \exp(tv^i)x = \iota^i(y)\}, \end{aligned} \quad (218)$$

for each $i \in I$. Write $\mathbf{T} = (T^i : i \in I)$.

The proofs of Propositions A.16, A.31 and Definitions A.30, A.33 then yield:

Theorem A.35. *In Definition A.34, $\mathbf{T}$ is a tent function for $(X, \mathbf{f}, \mathbf{G})$ or $(X, \mathbf{f}, \underline{\mathbf{G}})$, with $\min(\mathbf{T}|_{\partial X}) = \min \mathbf{T}'$.*

Let $X_c, \mathbf{f}_c, \mathbf{G}_c$ for $c \in C$ be as in Proposition A.29 for $Z_{X, \mathbf{T}}$, and $X'_{c'}, \mathbf{f}'_{c'}, \mathbf{G}'_{c'}$ for $c' \in C'$ be as in Proposition A.29 for $Z_{\partial X, \mathbf{T}'}$. Then we can take $C' \subseteq C$, such that for each $c' \in C'$ there is a natural identification between $(X'_{c'}, \mathbf{f}'_{c'}, \mathbf{G}'_{c'})$ and a component of $(\partial X_c, \mathbf{f}_c|_{\partial X_{c'}}, \mathbf{G}_c|_{\partial X_{c'}})$. For $c \in C \setminus C'$, the X_c correspond to points $p_{b^i, c}^i$ as in Proposition A.31. The same holds for $\underline{\mathbf{G}}_c, \underline{\mathbf{G}}'_{c'}$.

By taking $\epsilon > 0$ sufficiently small and θ^i for $i \in I$ sufficiently large, we can choose $\mathbf{T}$ to make the X_c and V_c^i ‘arbitrarily small’, given the $X'_{c'}, V_{c'}^{i'}$ which are already fixed. That is, given an open cover $\{U_j : j \in J\}$ for X such that for all $c' \in C'$ there exists $j \in J$ with $\pi'(X'_{c'}) \subseteq U_j$, then we can choose $\mathbf{T}$ so that for all $c \in C$ there exists $j \in J$ with $\pi(X_c) \subseteq U_j$, and similarly for the $V_c^i, V_{c'}^{i'}$.

In the case of **effective** gauge-fixing data $\underline{\mathbf{G}}, \underline{\mathbf{H}}, \underline{\mathbf{G}}_c$, if the $X'_{c'}$ have trivial stabilizers and the $V_{c'}^{i'}$ are manifolds for all $c' \in C'$ and $i \in I'_{c'}$, then the X_c have trivial stabilizers and the V_c^i are manifolds for all $c \in C$ and $i \in I_c$.

The tent functions $\mathbf{T}$ of Definition A.34 are also suitable for piecewise smooth extensions, as in §A.3.4, and we will use them for this in Step 2 of the proof of Theorem 4.8 in §B.2.

Remark A.36. It is instructive to compare our notion of tent functions on Kuranishi chains $(X, \mathbf{f}, \mathbf{G})$ with Fukaya and Ono’s notion of *multisection* on a Kuranishi space X [25, §3, §6], [24, §A1.1]. There are many similarities, as in writing this book I started with Fukaya and Ono’s definitions of Kuranishi spaces and virtual cycles and tried to improve them. Here are some:

- In both cases we fix a good coordinate system $\mathbf{I} = (I, (V^i, \dots, \psi^i) : i \in I, \dots)$ on X , and then consider a *smooth, multivalued* function T^i on $\dot{V}^i \subset V^i$, or section $\mathbf{s}^i$ of $E^i \rightarrow V^i$ for each $i \in I$, which must be compatible with coordinate changes ϕ^{ij} for $j < i$ in I .
- Fukaya and Ono use their multisections to perturb Kuranishi spaces to (non-Hausdorff, $\mathbb{Q}$ -weighted) manifolds, and then to triangulate these manifolds by simplices. We use tent functions for both these purposes.

However, there are also important differences:

- For a multisection with n -branches, each branch has equal weight $1/n$. Therefore by its nature the multisection method produces singular chains over $\mathbb{Q}$, not over $\mathbb{Z}$. In contrast, though a tent function T may have many smooth branches, only the minimal branch $\min T$ really counts, and it has weight 1. Because of this, we are able to make effective Kuranishi (co)homology work over $\mathbb{Z}$ as well as $\mathbb{Q}$.
- Fukaya and Ono use *multivalued* sections of *orbifold vector bundles* so that they can perturb to transverse sections, which is generally not possible for single-valued sections of orbifold vector bundles.

For *effective* Kuranishi neighbourhoods (V, E, s, ψ) , single-valued sections of E can always be perturbed to transverse, so this problem does not arise in Appendix B, where we actually do our perturbations.

However, in Appendix C we use a tent function to cut a general Kuranishi space X into small pieces X_c for $c \in C$ which are quotients X'_c/Γ_c for X'_c an effective Kuranishi space and Γ_c a finite group, and then we replace X_c by X'_c with weight $1/|\Gamma_c|$ in $\mathbb{Q}$. So we are forced to work over $\mathbb{Q}$ in Appendix C, for essentially the same reason as Fukaya and Ono.

B Proof that $H_*^{\text{si}}(Y; R) \cong KH_*^{\text{ef}}(Y; R)$

We now prove Theorem 4.8. Let Y be an orbifold and R a commutative ring. We must show that $\Pi_{\text{si}}^{\text{ef}} : H_k^{\text{si}}(Y; R) \rightarrow KH_k^{\text{ef}}(Y; R)$ in (76) is an isomorphism. We do this in four steps, which we describe briefly below and then cover in detail in §B.1–§B.4. In Steps 1–3 we start with an arbitrary cycle $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a]$ in $KC_k^{\text{ef}}(Y; R)$ representing some $\alpha \in KH_k^{\text{ef}}(Y; R)$, and progressively modify it to get other homologous cycles representing α with better and better properties, until we represent α by the image under $\Pi_{\text{si}}^{\text{ef}}$ in (73) of a cycle in $C_k^{\text{si}}(Y; R)$, proving that $\Pi_{\text{si}}^{\text{ef}}$ in (76) is *surjective*. Step 4 proves $\Pi_{\text{si}}^{\text{ef}}$ is *injective*, using the same method but for chains with boundary rather than cycles. We allow $k < 0$ in Steps 1 and 2, and prove the last part of Theorem 4.8 at the end of Step 2.

Step 1. Let $k \in \mathbb{Z}$, $\alpha \in KH_k^{\text{ef}}(Y; R)$, and $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a] \in KC_k^{\text{ef}}(Y; R)$ represent α . Then each X_a is an *effective Kuranishi space*, and in the excellent coordinate system $\mathbf{I}_a = (I_a, (V_a^i, \dots, \psi_a^i) : i \in I_a, \dots)$ for $(X_a, \mathbf{f}_a)$ in $\underline{\mathbf{G}}_a$, V_a^i is an *effective orbifold*. As in §A.3, we use a *tent function* $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a)$ to ‘cut’ each X_a into finitely many smaller pieces X_{ac} for $c \in C_a$, in triples $(X_{ac}, \mathbf{f}_{ac}, \underline{\mathbf{G}}_{ac})$, where X_{ac} is a Kuranishi space with *trivial stabilizers*, and in the excellent coordinate system $\mathbf{I}_{ac} = (I_{ac}, (V_{ac}^i, \dots, \psi_{ac}^i) : i \in I_{ac}, \dots)$ for $(X_{ac}, \mathbf{f}_{ac})$ in $\underline{\mathbf{G}}_{ac}$, each V_{ac}^i is a *manifold*. We show $\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \underline{\mathbf{G}}_{ac}]$ is homologous to $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a]$ in $KC_k^{\text{ef}}(Y; R)$, and so represents α .

As $\partial(\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a]) = 0$, the $[\partial X_a, \mathbf{f}_a|_{\partial X_a}, \underline{\mathbf{G}}_a|_{\partial X_a}]$ must satisfy relations in $KC_{k-1}^{\text{ef}}(Y; R)$. The main problem is to choose the $\mathbf{T}_a$ for $a \in A$ in a way compatible with these relations, so that $\partial(\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \underline{\mathbf{G}}_{ac}]) = 0$. That is, the choices of $\mathbf{T}_a, C_a$ and $X_{ac}, \mathbf{f}_{ac}, \underline{\mathbf{G}}_{ac}$ for $c \in C_a$ cannot be

made independently for each $a \in A$, but rather, they must satisfy complicated boundary conditions over ∂X_a relating the choices for different $a \in A$.

To make these choices in a way compatible with the boundary conditions, we introduce some more notation. By Lemma 3.7, for each $a \in A$ and $m \geq 0$ the triple $(\partial^m X_a, \mathbf{f}_a|_{\partial^m X_a}, \mathbf{G}_a|_{\partial^m X_a})$ has a splitting $\partial^m X_a = X_{a1}^m \amalg \cdots \amalg X_{an_a^m}^m$ into *connected* triples $(X_{ab}^m, \mathbf{f}_a|_{X_{ab}^m}, \mathbf{G}_a|_{X_{ab}^m})$, which is unique up to order of the X_{ab}^m for $b = 1, \dots, n_a^m$. Write $\mathbf{f}_{ab}^m = \mathbf{f}_a|_{X_{ab}^m}$ and $\mathbf{G}_{ab}^m = \mathbf{G}_a|_{X_{ab}^m}$. Then $\text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) = \{1\}$ by Theorem 3.39(b).

Define $P^m = \{(a, b) : a \in A, b = 1, \dots, n_a^m\}$. Define an equivalence relation $\sim$ on P^m by $(a, b) \sim (a', b')$ if there exists an isomorphism $(\mathbf{a}, \mathbf{b}) : (X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) \rightarrow (X_{a'b'}^m, \mathbf{f}_{a'b'}^m, \mathbf{G}_{a'b'}^m)$ as in Definition 3.36, where $\mathbf{a}$ need not identify orientations. Note that $(\mathbf{a}, \mathbf{b})$ is unique as $\text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) = \{1\}$. Let $Q^m = P^m / \sim$ be the set of equivalence classes of (a, b) in P^m . We shall treat Q^m as an *indexing set*. For each $q \in Q^m$, choose $(a^q, b^q) \in P^m$ representing the equivalence class q . Write $R^m = \{(a^q, b^q) : q \in Q^m\}$. Define $\phi^m : P^m \rightarrow R^m$ by $\phi^m(a, b) = (a^q, b^q)$ when the $\sim$ -equivalence class of (a, b) in Q^m is q . Write $(\mathbf{a}, \mathbf{b})_{ab}^m$ for the unique isomorphism $(\mathbf{a}, \mathbf{b}) : (X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) \rightarrow (X_{a^q b^q}^m, \mathbf{f}_{a^q b^q}^m, \mathbf{G}_{a^q b^q}^m)$.

Since A is finite and $\partial^m X_a = \emptyset$ for $m \gg 0$, P^m, Q^m, R^m are finite with $P^m = Q^m = R^m = \emptyset$ for $m \gg 0$. Let $M \geq 0$ be largest with $P^M, Q^M, R^M \neq \emptyset$. Write $\epsilon_{ab}^m = 1$ if $\mathbf{a}$ in $(\mathbf{a}, \mathbf{b})_{ab}^m$ is orientation-preserving, and $\epsilon_{ab}^m = -1$ if $\mathbf{a}$ is orientation-reversing. As $X_{ab}^m \subseteq \partial^m X_a$, we have $\partial X_{ab}^m \subseteq \partial^{m+1} X_a = \coprod_{b'=1}^{n_a^{m+1}} X_{ab'}^{m+1}$. Clearly, ∂X_{ab}^m is a disjoint union of some subset of the $X_{ab'}^{m+1}$. Define B_{ab}^m to be the set of $b' = 1, \dots, n_a^{m+1}$ with $X_{ab'}^{m+1} \subseteq \partial X_{ab}^m$. Then $\partial X_{ab}^m = \coprod_{b' \in B_{ab}^m} X_{ab'}^{m+1}$.

By induction on decreasing $m = M, M-1, \dots, 1, 0$, we use Definition A.34 and Theorem A.35 to choose a tent function $\mathbf{T}_{ab}^m$ for $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ for all (a, b) in R^m , satisfying the following boundary condition: we have $\partial X_{ab}^m = \coprod_{b' \in B_{ab}^m} X_{ab'}^{m+1}$, and for each such $X_{ab'}^{m+1}$ we have $(\bar{a}, \bar{b}) = \phi^{m+1}(a, b') \in R^{m+1}$, an isomorphism $(\mathbf{a}, \mathbf{b})_{ab'}^{m+1} : (X_{ab'}^{m+1}, \mathbf{f}_{ab'}^{m+1}, \mathbf{G}_{ab'}^{m+1}) \rightarrow (X_{\bar{a}\bar{b}}^{m+1}, \mathbf{f}_{\bar{a}\bar{b}}^{m+1}, \mathbf{G}_{\bar{a}\bar{b}}^{m+1})$, and a tent function $\mathbf{T}_{\bar{a}\bar{b}}^{m+1}$ for $(X_{\bar{a}\bar{b}}^{m+1}, \mathbf{f}_{\bar{a}\bar{b}}^{m+1}, \mathbf{G}_{\bar{a}\bar{b}}^{m+1})$ chosen in the previous inductive step. We need $\min(\mathbf{T}_{ab}^m|_{X_{ab'}^{m+1}})$ to agree with $\min \mathbf{T}_{\bar{a}\bar{b}}^{m+1}$ under the isomorphism $(\mathbf{a}, \mathbf{b})_{ab'}^{m+1}$, for all b' in B_{ab}^m . This is possible as in §A.3.5.

After completing the induction, for $a \in A$ we have $X_a = \coprod_{b=1}^{n_a^0} X_{ab}^0$, and for each such (a, b) we have $(\bar{a}, \bar{b}) = \phi^0(a, b) \in R^0$, an isomorphism $(\mathbf{a}, \mathbf{b})_{ab}^0 : (X_{ab}^0, \mathbf{f}_{ab}^0, \mathbf{G}_{ab}^0) \rightarrow (X_{\bar{a}\bar{b}}^0, \mathbf{f}_{\bar{a}\bar{b}}^0, \mathbf{G}_{\bar{a}\bar{b}}^0)$, and a tent function $\mathbf{T}_{\bar{a}\bar{b}}^0$ for $(X_{\bar{a}\bar{b}}^0, \mathbf{f}_{\bar{a}\bar{b}}^0, \mathbf{G}_{\bar{a}\bar{b}}^0)$. We define the tent function $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ to be the pullback of $\mathbf{T}_{\bar{a}\bar{b}}^0$ under $(\mathbf{a}, \mathbf{b})_{ab}^0$ on X_{ab}^0 , for each $b = 1, \dots, n_a^0$. Then by construction, the $\mathbf{T}_a|_{\partial X_a}$ for $a \in A$ satisfy relations corresponding to $\partial(\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]) = 0$.

Let $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ for $c \in C_a$ be the decomposition of $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ into pieces defined using $\mathbf{T}_a$ in §A.3.2. Theorem A.35 and induction on m implies that the X_{ac} have trivial stabilizers, and the V_{ac}^i are manifolds. We show using Proposition A.29 that $\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$ is homologous to $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]$ in $KC_k^{\text{ef}}(Y; R)$.

Step 2. We shall make a small deformation of X_{ac} into a *compact manifold* $\tilde{X}_{ac}$ with *g-corners*, for each $a \in A$ and $c \in C_a$. This yields a new cycle $\sum_{a \in A} \sum_{c \in C_a} \rho_a[\tilde{X}_{ac}, \tilde{f}_{ac}, \tilde{\underline{G}}_{ac}] \in KC_k^{\text{ef}}(Y; R)$ which is homologous to $\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \underline{G}_{ac}]$, and so represents α . We write $\tilde{f}_{ac} : \tilde{X}_{ac} \rightarrow Y$ rather than $\tilde{\mathbf{f}}_{ac} : \tilde{X}_{ac} \rightarrow Y$ since $\tilde{f}_{ac}$ is a smooth map of orbifolds, rather than a strongly smooth map of Kuranishi spaces.

The equation $\partial(\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \underline{G}_{ac}]) = 0$ yields relations on the $[\partial X_{ac}, \mathbf{f}_{ac}|_{\partial X_{ac}}, \underline{G}_{ac}|_{\partial X_{ac}}]$ in $KC_{k-1}^{\text{ef}}(Y; R)$. The main problem is to choose the deformations $\tilde{X}_{ac}$ in a way compatible with these relations, so that $\partial(\sum_{a \in A} \sum_{c \in C_a} \rho_a[\tilde{X}_{ac}, \tilde{f}_{ac}, \tilde{\underline{G}}_{ac}]) = 0$. That is, the perturbations $\tilde{X}_{ac}$ must satisfy complicated boundary conditions. Before explaining how we do this, we discuss how to perturb a single $(X_{ac}, \mathbf{f}_{ac}, \underline{G}_{ac})$ without boundary conditions.

Let $(\mathbf{I}_{ac}, \boldsymbol{\eta}_{ac})$ be the excellent coordinate system for $(X_{ac}, \mathbf{f}_{ac})$ in $\underline{G}_{ac}$, where $\mathbf{I}_{ac} = (I_{ac}, (V_{ac}^i, E_{ac}^i, s_{ac}^i, \psi_{ac}^i), f_{ac}^i : i \in I_{ac}, \dots)$, the V_{ac}^i are manifolds, and each $E_{ac}^i \rightarrow V_{ac}^i$ is a vector bundle.

Working by induction on $i \in I_{ac}$, we will choose C^1 small perturbations $\tilde{s}_{ac}^i$ of s_{ac}^i on V_{ac}^i for $i \in I_{ac}$ such that $\tilde{s}_{ac}^i$ is *transverse* along $(\tilde{s}_{ac}^i)^{-1}(0)$, that is, $d\tilde{s}_{ac}^i : TV_{ac}^i \rightarrow E_{ac}^i$ is surjective on $(\tilde{s}_{ac}^i)^{-1}(0)$, and $\hat{\phi}_{ac}^{ij} \circ \tilde{s}_{ac}^j \equiv \tilde{s}_{ac}^i \circ \phi_{ac}^{ij}$ on V_{ac}^{ij} whenever $j \leq i$ in I_{ac} .

By gluing together the sets $(\tilde{s}_{ac}^i)^{-1}(0)$ we make a *compact, oriented k-manifold with g-corners* $\tilde{X}_{ac}$ which is a small transverse perturbation of X_{ac} , with a smooth map $\tilde{f}_{ac} : \tilde{X}_{ac} \rightarrow Y$, and an excellent coordinate system $(\tilde{\mathbf{I}}_{ac}, \tilde{\boldsymbol{\eta}}_{ac})$ for $(\tilde{X}_{ac}, \tilde{f}_{ac})$ with indexing set $\tilde{I}_{ac} = \{k\}$, one Kuranishi neighbourhood $(\tilde{V}_{ac}^k, \tilde{E}_{ac}^k, \tilde{s}_{ac}^k, \tilde{\psi}_{ac}^k) = (\tilde{X}_{ac}, \tilde{X}_{ac}, 0, \text{id}_{\tilde{X}_{ac}})$ and $\tilde{\eta}_{ac,k} \equiv 1 \equiv \tilde{\eta}_{ac,k}^k$. We choose $\tilde{\underline{G}}_{ac}^k : \tilde{X}_{ac} \rightarrow \underline{P}$ such that $\tilde{\underline{G}}_{ac} = (\tilde{\mathbf{I}}_{ac}, \tilde{\boldsymbol{\eta}}_{ac}, \tilde{\underline{G}}_{ac}^k)$ is effective gauge-fixing data for $(\tilde{X}_{ac}, \tilde{f}_{ac})$. We also construct a homology $[Z_{ac}, \mathbf{g}_{ac}, \underline{H}_{ac}]$ between $[X_{ac}, \mathbf{f}_{ac}, \underline{G}_{ac}]$ and $[\tilde{X}_{ac}, \tilde{f}_{ac}, \tilde{\underline{G}}_{ac}]$ in $KC_{k+1}^{\text{ef}}(Y; R)$, modulo terms over ∂X_{ac} .

Next we explain how to make these choices of perturbations $\tilde{s}_{ac}^i$ for $a \in A$ and $c \in C_a$ compatible with the boundary conditions. Given how we chose the $\mathbf{T}_a$ compatible with boundary conditions in Step 1, one obvious approach would be to use the same method, and to work by induction on decreasing $m = M, M-1, \dots, 0$ choosing the $\tilde{s}_{ac}^i|_{\partial^m V_{ac}^i}$, by splitting $\partial^m[X_{ac}, \mathbf{f}_{ac}, \underline{G}_{ac}]$ into connected components, and ensuring our choices depend only on the isomorphism class of the component.

However, there is a problem with this: as the V_{ac}^i are manifolds *with g-corners*, not corners, the Extension Principle for smooth sections, Principle 2.8(c), does not hold. Thus, given prescribed values for $\tilde{s}_{ac}^i$ over ∂V_{ac}^i which are σ -invariant over $\partial^2 V_{ac}^i$, we may *not* be able to extend these prescribed values smoothly over V_{ac}^i to choose $\tilde{s}_{ac}^i$ satisfying the necessary boundary conditions.

Our solution to this was explained in §A.1.4. There are notions of *continuous, piecewise smooth functions and sections subordinate to a tent function*, and an Extension Principle holds for such functions and sections on manifolds and orbifolds with g-corners. So, we could solve the problem by using continuous, piecewise smooth sections $\tilde{s}_{ac}^i$ on V_{ac}^i . That is, we could choose tent functions

$\mathbf{T}_{ac}$ for $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ which cut X_{ac} into still smaller pieces X_{acd} for $d \in D_{ac}$, and continuous, piecewise smooth sections $\tilde{s}_{ac}^i$ over V_{ac}^i subordinate to the tent function T_{ac}^i in $\mathbf{T}_{ac}^i$, satisfying the necessary boundary conditions. The restriction of $\tilde{s}_{ac}^i$ to V_{acd}^i would then be smooth for each $d \in D_{ac}$.

In fact there is a simpler method. We do not need to choose more tent functions $\mathbf{T}_{ac}$ and divide the X_{ac} into smaller pieces X_{acd} . Instead, we treat the smooth sections $\tilde{s}_{ac}^i$ on V_{ac}^i for $c \in C_a$ as coming from a piecewise smooth section $\tilde{s}_a^i$ on $\dot{V}_a^i$ subordinate to T_a^i . So, we can re-use the tent functions $\mathbf{T}_a$ chosen in Step 1 to inductively construct piecewise smooth sections over X_a and $\dot{V}_a^i$ satisfying boundary conditions, which restrict to smooth sections over X_{ac} and V_{ac}^i satisfying the necessary boundary conditions.

The $\mathbf{T}_a$ are suitable for this purpose, as we were careful to define them this way in §A.3.5. Also, despite the discussion after Example A.24 in §A.3, in this case we do not need to define the $\tilde{s}_{ac}^i$ on smaller subsets $\dot{V}_{ac}^i \subset V_{ac}^i$, since the V_{ac}^i are constructed from subsets $\dot{V}_a^i$ in V_a^i , and by choosing the $\tilde{s}_a^i$ to extend to open neighbourhoods of $\bar{\dot{V}}_a^i$ in V_a^i for $i \in I_a$, we can define $\tilde{s}_a^i$ on all of $\dot{V}_a^i$ and so define $\tilde{s}_{ac}^i$ on all of V_{ac}^i .

Here, then, is how we actually choose the $\tilde{s}_{ac}^i$. Use the notation of Step 1, and let the effective gauge-fixing data $\mathbf{G}_{ab}^m$ for $(X_{ab}^m, \mathbf{f}_{ab}^m)$ contain $\mathbf{I}_{ab}^m = ((V_{ab}^{i,m}, E_{ab}^{i,m}, s_{ab}^{i,m}, \psi_{ab}^{i,m}) : i \in I_{ab}^m, \dots)$, and $\mathbf{T}_{ab}^m = (T_{ab}^{i,m} : i \in I_{ab}^m)$, where $T_{ab}^{i,m}$ is a tent function on $\dot{V}_{ab}^{i,m} \subset V_{ab}^{i,m}$. By induction on decreasing $m = M, M-1, \dots, 1, 0$, we choose a piecewise smooth section $\tilde{s}_{ab}^{i,m}$ of $E_{ab}^{i,m}$ over $\dot{V}_{ab}^{i,m}$ subordinate to $T_{ab}^{i,m}$, which is a piecewise C^1 small perturbation of $s_{ab}^{i,m}$, with $\tilde{s}_{ab}^{i,m}$ prescribed over $\partial \dot{V}_{ab}^{i,m}$ in the same way as for the boundary conditions for $\min T_{ab}^{i,m}$ in Step 1. This is possible, in a similar way to §A.1.4.

After completing the induction, we define piecewise smooth sections $\tilde{s}_a^i$ of E_a^i over $\dot{V}_a^i$ from the $\tilde{s}_{ab}^{i,0}$ subordinate to T_a^i , as for the construction of $\mathbf{T}_a$ from $\mathbf{T}_{ab}^0$ in Step 1. These $\tilde{s}_a^i$ pull back to smooth sections $\tilde{s}_{ac}^i$ on each V_{ac}^i , under the natural embeddings $\pi_a : V_{ac}^i \rightarrow \dot{V}_a^i$. The $\tilde{s}_{ac}^i$ then satisfy all the necessary boundary conditions.

We then show that $\sum_{a \in A} \sum_{c \in C_a} \rho_a[\tilde{X}_{ac}, \tilde{f}_{ac}, \tilde{\mathbf{G}}_{ac}] \in KC_k^{\text{ef}}(Y; R)$ is homologous to $\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$, and so represents α . If $k < 0$ then $\tilde{X}_a = \emptyset$ as $\dim \tilde{X}_a < 0$, so $\alpha = 0$. Thus $KH_k^{\text{ef}}(Y; R) = 0$ for $k < 0$, proving the last part of Theorem 4.8. Therefore we suppose $k \geq 0$ from now on.

Step 3. For simplicity we now *change notation* from $\tilde{X}_{ac}$ for $a \in A$ and $c \in C_a$ to $\tilde{X}_a$ for $a \in A$. That is, Steps 1 and 2 have shown that we can represent α by a cycle $\sum_{a \in A} \rho_a[\tilde{X}_a, \tilde{f}_a, \tilde{\mathbf{G}}_a] \in KC_k^{\text{ef}}(Y; R)$, with each $\tilde{X}_a$ a compact, oriented k -manifold with g-corners, $\tilde{f}_a : \tilde{X}_a \rightarrow Y$ a smooth map, and $\tilde{\mathbf{G}}_a = (\tilde{\mathbf{I}}_a, \tilde{\eta}_a, \tilde{\mathbf{G}}_a^k)$ effective gauge-fixing data for $(\tilde{X}_a, \tilde{f}_a)$, where $\tilde{\mathbf{I}}_a$ has indexing set $\tilde{I}_a = \{k\}$, one Kuranishi neighbourhood $(\tilde{V}_a^k, \tilde{E}_a^k, \tilde{s}_a^k, \tilde{\psi}_a^k) = (\tilde{X}_a, \tilde{X}_a, 0, \text{id}_{\tilde{X}_a})$, and $\tilde{\eta}_a = (\tilde{\eta}_{a,k}, \tilde{\eta}_{a,k}^k)$ with $\tilde{\eta}_{a,k} \equiv 1 \equiv \tilde{\eta}_{a,k}^k$, and $\tilde{\mathbf{G}}_a^k : \tilde{X}_a \rightarrow \underline{P}$ is some map satisfying injectivity conditions as in §3.9.

As in §A.1.3 and §A.1.6, we can choose a tent function $\tilde{T}_a : \tilde{X}_a \rightarrow F([1, \infty))$ for each $a \in A$ such that the components $\tilde{X}_{ac}$, $c \in C_a$ of $\partial Z_{\tilde{X}_a, \tilde{T}_a}$ in (187) are all

diffeomorphic to the k -simplex Δ_k , with diffeomorphisms $\sigma_{ac} : \Delta_k \rightarrow \tilde{X}_{ac}$ for $c \in C_a$. In order to make these choices in a way compatible with the boundary relations involved in $\partial(\sum_{a \in A} \rho_a [\tilde{X}_a, \tilde{f}_a, \tilde{\mathbf{G}}_a]) = 0$, we follow the strategy and use the notation of Step 1, and by reverse induction on $m = k, k-1, \dots, 1, 0$ we use Theorem A.18 to choose tent functions $\tilde{T}_{ab}^m$ on $\tilde{X}_{ab}^m$ for $(a, b) \in R^m$ and diffeomorphisms $\sigma_{abc}^m : \Delta_{k-m} \rightarrow \tilde{X}_{abc}^m$ for $(a, b) \in R^m$ and $c \in C_{ab}^m$, such that $\min \tilde{T}_{ab}^m|_{\partial \tilde{X}_{ab}^m}$ and $\sigma_{abc}^m|_{\partial \tilde{X}_{ab}^m}$ are identified on $X_{ab'}^{m+1}$ for each $b' \in B_{ab}^m$ with $\min \tilde{T}_{\bar{a}\bar{b}}^{m+1}$ and $\sigma_{\bar{a}\bar{b}\bar{c}}^{m+1}$ under $(\mathbf{a}, \mathbf{b})_{ab'}^{m+1}$, for $(\bar{a}, \bar{b}) = \phi^{m+1}(a, b')$ in R^{m+1} , and $\bar{c} \in C_{\bar{a}\bar{b}}^{m+1}$. At the end of the induction we define $\tilde{T}_a, C_a$ and σ_{ac} using the $\tilde{T}_{ab}^0, C_{ab}^0$ and σ_{abc} for $b = 1, \dots, n_a^0$ and $c \in C_{ab}^0$.

For $a \in A$ and $c \in C_a$, define $\epsilon_{ac} = 1$ if the diffeomorphism $\sigma_{ac} : \Delta_k \rightarrow \tilde{X}_{ac}$ is orientation-preserving, and $\epsilon_{ac} = -1$ otherwise. We then show that $\sum_{a \in A} \rho_a [\tilde{X}_a, \tilde{f}_a, \tilde{\mathbf{G}}_a]$ is homologous in $KC_k^{\text{ef}}(Y; R)$ to the cycle

$$\begin{aligned} & \sum_{a \in A} \sum_{c \in C_a} \rho_a \epsilon_{ac} [\Delta_k, \tilde{f}_a \circ \pi \circ \sigma_{ac}, \mathbf{G}_{\Delta_k}] \\ &= \Pi_{\text{si}}^{\text{ef}} \left(\sum_{a \in A} \sum_{c \in C_a} (\rho_a \epsilon_{ac}) (\tilde{f}_a \circ \pi \circ \sigma_{ac}) \right). \end{aligned} \quad (219)$$

Since $\Pi_{\text{si}}^{\text{ef}} : C_*^{\text{si}}(Y; R) \rightarrow KC_*^{\text{ef}}(Y; R)$ is injective at the chain level, with $\Pi_{\text{si}}^{\text{ef}} \circ \partial = \partial \circ \Pi_{\text{si}}^{\text{ef}}$, as (219) is a cycle in $KC_k^{\text{ef}}(Y; R)$ we have

$$\partial \left(\sum_{a \in A} \sum_{c \in C_a} (\rho_a \epsilon_{ac}) (\tilde{f}_a \circ \pi \circ \sigma_{ac}) \right) = 0 \quad \text{in } C_{k-1}^{\text{si}}(Y; R). \quad (220)$$

Thus $\beta = [\sum_{a \in A} \sum_{c \in C_a} (\rho_a \epsilon_{ac}) (\tilde{f}_a \circ \pi \circ \sigma_{ac})]$ is well-defined in $H_k^{\text{si}}(Y; R)$, and (219) implies that $\Pi_{\text{si}}^{\text{ef}}(\beta) = \alpha$. Hence $\Pi_{\text{si}}^{\text{ef}}$ in (76) is *surjective*.

Step 4. Suppose $\beta \in H_k^{\text{si}}(Y; R)$ with $\Pi_{\text{si}}^{\text{ef}}(\beta) = 0$. Represent β by $\sum_{d \in D} \eta_d \tau_d$ in $C_k^{\text{si}}(Y; R)$, for D a finite indexing set, $\eta_d \in R$ and $\tau_d : \Delta_k \rightarrow Y$ smooth. Then $\Pi_{\text{si}}^{\text{ef}}(\sum_{d \in D} \eta_d \tau_d)$ is exact in $KC_k^{\text{ef}}(Y; R)$, so there exists $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a] \in KC_{k+1}^{\text{ef}}(Y; R)$ with

$$\partial \left(\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a] \right) = \sum_{d \in D} \eta_d [\Delta_k, \tau_d, \mathbf{G}_{\Delta_k}]. \quad (221)$$

We now apply Steps 1–3 to this $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$, replacing k -cycles by $(k+1)$ -chains. This eventually yields a singular $(k+1)$ -chain $\sum_{a \in A} \sum_{c \in C_a} (\rho_a \epsilon_{ac}) (\tilde{f}_a \circ \pi \circ \sigma_{ac})$. We make choices in Steps 1–3 so that

$$\partial \left(\sum_{a \in A} \sum_{c \in C_a} (\rho_a \epsilon_{ac}) (\tilde{f}_a \circ \pi \circ \sigma_{ac}) \right) = \sum_{d \in D} \sum_{e \in E} (\eta_d \zeta_e) (\tau_d \circ v_e), \quad (222)$$

where $\{v_e : e \in E\}$ are the simplices in the N^{th} barycentric subdivision of Δ_k for some $N \gg 0$, so that $v_e : \Delta_k \rightarrow \Delta_k$ is an affine embedding for each $e \in E$, with $\zeta_e = 1$ if v_e is orientation-preserving, and $\zeta_e = -1$ otherwise. But $\sum_{d \in D} \sum_{e \in E} (\eta_d \zeta_e) (\tau_d \circ v_e)$ is homologous to $\sum_{d \in D} \eta_d \tau_d$ in $C_k^{\text{si}}(Y; R)$ by Bredon [10, §IV.17], so $\sum_{d \in D} \eta_d \tau_d$ is also a boundary in $C_k^{\text{si}}(Y; R)$. Thus $\beta = 0$, and $\Pi_{\text{si}}^{\text{ef}}$ in (76) is *injective*.

Remark B.1. (a) Steps 1–3 above are related to Fukaya and Ono’s construction [25, §6], [24, §A1.1] of virtual moduli chains for Kuranishi spaces by using *transverse multisections*, small multi-valued perturbations $\mathfrak{s}_{ac}^i$ of the obstruction maps s_{ac}^i . In our approach, we first pass from Kuranishi spaces and $KH_*(Y; R)$ to *effective* Kuranishi spaces and $KH_*^{\text{ef}}(Y; R)$, as in Theorem 4.9 and its proof in Appendix C. Then, Steps 1–3 above in effect construct virtual moduli chains for effective Kuranishi spaces using *single-valued* transverse perturbations $\tilde{s}_{ac}^i$. Since we use single-valued perturbations rather than multisections, we are able to work over $\mathbb{Z}$, or an arbitrary commutative ring R , rather than $\mathbb{Q}$.

(b) We require our perturbations $\tilde{s}_{ac}^i$ in Step 2 to be close to the Kuranishi maps s_{ac}^i in C^1 , whereas Fukaya and Ono only require their multisections $\mathfrak{s}_{ac}^i$ to be close to s_{ac}^i in C^0 . The difference comes from our new Definition 2.18(e), which replaces Fukaya and Ono’s notion of ‘having a tangent bundle’, and alters the notion of compatibility of perturbations $\tilde{s}_{ac}^i$ under coordinate changes. Under a coordinate change $(\phi_{ac}^{ij}, \hat{\phi}_{ac}^{ij})$ from $(V_{ac}^j, \dots, \psi_{ac}^j)$ to $(V_{ac}^i, \dots, \psi_{ac}^i)$, in our set-up $\tilde{s}_{ac}^j$ determines $\tilde{s}_{ac}^i$ only on $\phi_{ac}^{ij}(\tilde{V}_{ac}^{ij})$, but for Fukaya and Ono $\mathfrak{s}_{ac}^j$ determines $\mathfrak{s}_{ac}^i$ on a neighbourhood of $\phi_{ac}^{ij}(V_{ac}^{ij})$. We need C^1 closeness to ensure Definition 2.18(e) holds for the coordinate changes (233) below. Remark 2.19 discussed ways of doing without Definition 2.18(e).

(c) In fact the notion that ‘ $\tilde{s}_{ac}^i$ is C^1 close to s_{ac}^i ’ in Step 2 does not really make sense, as we have not said how close $\tilde{s}_{ac}^i, s_{ac}^i$ need to be. As in Fukaya and Ono [25, Th. 6.4], [24, Th A1.23], what we should probably have done throughout Step 2 is not to choose a single perturbation $\tilde{s}_{ac}^i$ of s_{ac}^i , but a series $\tilde{s}_{ac,n}^i$ for $n = 1, 2, \dots$, such that $\tilde{s}_{ac,n}^i \rightarrow s_{ac}^i$ in C^1 as $n \rightarrow \infty$, compatibilities between $\tilde{s}_{ac}^i, \tilde{s}_{ac}^j$ are replaced by compatibilities between $\tilde{s}_{ac,n}^i, \tilde{s}_{ac,n}^j$ for each $n = 1, 2, \dots$, and so on. Then we take $n \gg 0$ to ensure C^1 -closeness wherever we need it. But we suppress this point, as there are too many indices in the proof already.

B.1 Step 1: cutting X_a into X_{ac} with trivial stabilizers

We work in the situation and with the notation of Step 1 above, so that $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a] \in KC_k^{\text{ef}}(Y; R)$ is a cycle representing $\alpha \in KH_k^{\text{ef}}(Y; R)$. We will use Theorem A.35 to choose a *tent function* $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a)$ in the sense of §A.3.2, for each $a \in A$. Then Definition A.28 defines $[Z_{X_a, \mathbf{T}_a}, \mathbf{f}_a \circ \pi_a, \underline{\mathbf{H}}_{X_a, \mathbf{T}_a}]$ in $KC_{k+1}^{\text{ef}}(Y; R)$, and equation (212) of Proposition A.29 yields

$$\begin{aligned} \partial[Z_{X_a, \mathbf{T}_a}, \mathbf{f}_a \circ \pi_a, \underline{\mathbf{H}}_{X_a, \mathbf{T}_a}] &= \sum_{c \in C_a} [X_{ac}, \mathbf{f}_{ac}, \underline{\mathbf{G}}_{ac}] - [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a] \\ &\quad - [Z_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}, \mathbf{f}_a|_{\partial X_a} \circ \pi_a, \underline{\mathbf{H}}_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}], \end{aligned} \quad (223)$$

where C_a is some finite indexing set, and Theorem A.35 says that each X_{ac} has *trivial stabilizers*, and the V_{ac}^i in $\underline{\mathbf{G}}_{ac}$ are *manifolds*. As $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a]$ is a cycle we have

$$\sum_{a \in A} \rho_a [\partial X_a, \mathbf{f}_a|_{\partial X_a}, \underline{\mathbf{G}}_a|_{\partial X_a}] = 0. \quad (224)$$

We want to choose the $\mathbf{T}_a$ for $a \in A$ such that

$$\sum_{a \in A} \rho_a [Z_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}, \mathbf{f}_a|_{\partial X_a} \circ \pi_a, \underline{\mathbf{H}}_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}] = 0. \quad (225)$$

Note that if the $\mathbf{T}_a$ depend functorially on $[X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a]$ in some sense compatible with boundaries, then (225) would follow from (224).

Assuming (225), multiplying (223) by ρ_a and summing over $a \in A$ yields

$$\begin{aligned} & \partial \left(\sum_{a \in A} \rho_a [Z_{X_a, \mathbf{T}_a}, \mathbf{f}_a \circ \pi_a, \underline{\mathbf{H}}_{X_a, \mathbf{T}_a}] \right) \\ &= \sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \underline{\mathbf{G}}_{ac}] - \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a]. \end{aligned} \quad (226)$$

Thus $\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \underline{\mathbf{G}}_{ac}]$ is a cycle homologous to $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a]$, and so represents α , as we have to prove.

The obvious way to do this is to first choose tent functions $\mathbf{T}_a|_{\partial X_a}$ for $(\partial X_a, \mathbf{f}_a|_{\partial X_a}, \underline{\mathbf{G}}_a|_{\partial X_a})$ satisfying the necessary compatibilities that will ensure (225) holds, and then extend these choices from ∂X_a to X_a . However, there is a problem. Let $\sigma_{X_a} : \partial^2 X_a \rightarrow \partial^2 X_a$ be the orientation-reversing involution from Definition 2.27. It is a general principle, as in Principle 2.8 and Property 3.1(e) for instance, that a choice of data over ∂X_a should extend to X_a only if the restriction of the choice to $\partial^2 X_a$ is invariant under σ_{X_a} . So our choice of $\mathbf{T}_a$ over ∂X_a should satisfy extra conditions over $\partial^2 X_a$.

We can achieve this by first choosing $\mathbf{T}_a$ over $\partial^2 X_a$ satisfying the compatibilities and these extra conditions, and then extending the choice from $\partial^2 X_a$ to ∂X_a . But this extension is only possible provided the restriction of the choice to $\partial^3 X_a$ is invariant under $\sigma_{\partial X_a} : \partial^2(\partial X_a) \rightarrow \partial^2(\partial X_a)$. Continuing this argument indefinitely, we see that the right approach is to choose data $\mathbf{T}_a$ over $\partial^m X_a$ for $a \in A$ by *reverse induction on m* , where in the inductive step we choose data over $\partial^m X_a$ which restricts on $\partial^{m+1} X_a$ to the data already chosen, and has the compatibilities necessary to prove (225), and is invariant under the strong diffeomorphisms $\sigma_l : \partial^m X_a \rightarrow \partial^m X_a$ induced by $\sigma_{\partial^l X_a} : \partial^2(\partial^l X_a) \rightarrow \partial^2(\partial^l X_a)$ for all $0 \leq l \leq m-2$.

Use the notation $n_a^m, (X_{ab}^m, \mathbf{f}_{ab}^m, \underline{\mathbf{G}}_{ab}^m), P^m, Q^m, R^m, M, (a^q, b^q), \phi^m, (\mathbf{a}, \mathbf{b})_{ab}^m, \epsilon_{ab}^m, B_{ab}^m$ of Step 1 above. By induction on decreasing $m = M, M-1, \dots, 1, 0$, for all (a, b) in R^m , we choose tent functions $\mathbf{T}_{ab}^m$ for $(X_{ab}^m, \mathbf{f}_{ab}^m, \underline{\mathbf{G}}_{ab}^m)$ using Definition A.34 and Theorem A.35, satisfying the inductive hypothesis:

(*)_{ab}^m By definition of B_{ab}^m we have $\partial X_{ab}^m = \coprod_{b' \in B_{ab}^m} X_{ab'}^{m+1}$, and if $\phi^{m+1}(a, b') = (\bar{a}, \bar{b})$ in R^{m+1} , we have an isomorphism $(\mathbf{a}, \mathbf{b})_{ab'}^{m+1} : (X_{ab'}^{m+1}, \mathbf{f}_{ab'}^{m+1}, \underline{\mathbf{G}}_{ab'}^{m+1}) \rightarrow (X_{\bar{a}\bar{b}}^{m+1}, \mathbf{f}_{\bar{a}\bar{b}}^{m+1}, \underline{\mathbf{G}}_{\bar{a}\bar{b}}^{m+1})$. So there is an isomorphism

$$\coprod_{b' \in B_{ab}^m} (\mathbf{a}, \mathbf{b})_{ab'}^{m+1} : \partial(X_{ab}^m, \mathbf{f}_{ab}^m, \underline{\mathbf{G}}_{ab}^m) \longrightarrow \coprod_{b' \in B_{ab}^m} (X_{\bar{a}\bar{b}}^{m+1}, \mathbf{f}_{\bar{a}\bar{b}}^{m+1}, \underline{\mathbf{G}}_{\bar{a}\bar{b}}^{m+1}). \quad (227)$$

$b' \in B_{ab}^m$: define $(\bar{a}, \bar{b}) = \phi^{m+1}(a, b')$

On the left hand side $\partial(X_{ab}^m, \mathbf{f}_{ab}^m, \underline{\mathbf{G}}_{ab}^m)$ of (227) we have a tent function $\mathbf{T}_{ab}^m|_{\partial X_{ab}^m}$. On the right hand side we have a tent function $\coprod_{b'} \mathbf{T}_{\bar{a}\bar{b}}^{m+1}$. We require that (227) should identify $\min(\mathbf{T}_{ab}^m|_{\partial X_{ab}^m})$ with $\min(\coprod_{b'} \mathbf{T}_{\bar{a}\bar{b}}^{m+1})$.

For the first step, when $m = M$, as $P^{M+1} = \emptyset$, if $(a, b) \in R^M$ then $\partial X_{ab}^M = \emptyset$, so the boundary conditions in $(*)_{ab}^M$ are trivial. Thus for all $(a, b) \in R^M$ we choose $\mathbf{T}_{ab}^M$ for $(X_{ab}^M, \mathbf{f}_{ab}^M, \underline{\mathbf{G}}_{ab}^M)$ using Definition A.30 and Proposition A.31.

For the inductive step, suppose that for some $l = 0, \dots, M-1$ we have chosen $\mathbf{T}_{ab}^m$ satisfying $(*)_{ab}^m$ for all $l < m \leq M$ and $(a, b) \in R^m$. We will choose $\mathbf{T}_{ab}^l$ satisfying $(*)_{ab}^l$ for all $(a, b) \in R^l$. The important point is this. Let $\mathbf{T}_{ab}^l$ be the tent function for $\partial(X_{ab}^l, \mathbf{f}_{ab}^l, \underline{\mathbf{G}}_{ab}^l)$ obtained by pulling back $\coprod_{b'} \mathbf{T}_{ab'}^{l+1}$. By $(*)_{ab}^l$, we have to choose a tent function $\mathbf{T}_{ab}^l$ for $(X_{ab}^l, \mathbf{f}_{ab}^l, \underline{\mathbf{G}}_{ab}^l)$ such that $\min(\mathbf{T}_{ab}^l|_{\partial X_{ab}^l}) = \min \mathbf{T}_{ab}^l$.

As in Definition A.34, we can do this provided $\min \mathbf{T}_{ab}^l|_{\partial^2 X_{ab}^l}$ is invariant under the natural involution $\sigma : \partial^2 X_{ab}^l \rightarrow \partial^2 X_{ab}^l$. To see this, note that $\partial X_{ab}^l = \coprod_{b' \in B_{ab}^l} X_{ab'}^{l+1}$, and so $\partial^2 X_{ab}^l = \coprod_{b' \in B_{ab}^l} \coprod_{b'' \in B_{ab'}^{l+1}} X_{ab''}^{l+2}$. Thus, the splitting of $(\partial^2 X_{ab}^l, \mathbf{f}_{ab}^l|_{\partial^2 X_{ab}^l}, \underline{\mathbf{G}}_{ab}^l|_{\partial^2 X_{ab}^l})$ into connected components in Lemma 3.7 is

$$(\partial^2 X_{ab}^l, \mathbf{f}_{ab}^l|_{\partial^2 X_{ab}^l}, \underline{\mathbf{G}}_{ab}^l|_{\partial^2 X_{ab}^l}) \cong \coprod_{b' \in B_{ab}^l} \coprod_{b'' \in B_{ab'}^{l+1}} (X_{ab''}^{l+2}, \mathbf{f}_{ab''}^{l+2}, \underline{\mathbf{G}}_{ab''}^{l+2}). \quad (228)$$

The natural orientation-reversing involution $\sigma : \partial^2 X_{ab}^l \rightarrow \partial^2 X_{ab}^l$ extends to an involution (σ, τ) of $(\partial^2 X_{ab}^l, \mathbf{f}_{ab}^l|_{\partial^2 X_{ab}^l}, \underline{\mathbf{G}}_{ab}^l|_{\partial^2 X_{ab}^l})$. Thus, σ must permute the components $X_{ab''}^{l+2}$ in (228). Now $\text{Aut}(X_{ab''}^{l+2}, \mathbf{f}_{ab''}^{l+2}, \underline{\mathbf{G}}_{ab''}^{l+2}) = \{1\}$ by Theorem 3.39(b), as $\underline{\mathbf{G}}_{ab''}^{l+2}$ is effective. Since σ is orientation-reversing, it cannot act as the identity. Therefore σ must act *freely* on the set of $X_{ab''}^{l+2}$ appearing in (228), as if σ fixed any component, that component would have a nontrivial automorphism.

Let $X_{ab''}^{l+2}$ be a component on the r.h.s. of (228). Then σ takes $X_{ab''}^{l+2}$ to some unique $X_{ab''' }^{l+2}$ with $b'' \neq b'''$. The isomorphism (σ, τ) between these two components implies that (a, b'') and (a, b''') lie in the same equivalence class in P^{l+2} , so that $\phi^{l+2}(a, b'') = \phi^{l+2}(a, b''') = (\bar{a}, \bar{b}) \in R^{l+2}$. The associated isomorphisms $(\mathbf{a}, \mathbf{b})_{ab''}^{l+2} : (X_{ab''}^{l+2}, \mathbf{f}_{ab''}^{l+2}, \underline{\mathbf{G}}_{ab''}^{l+2}) \rightarrow (X_{\bar{a}\bar{b}}^{l+2}, \mathbf{f}_{\bar{a}\bar{b}}^{l+2}, \underline{\mathbf{G}}_{\bar{a}\bar{b}}^{l+2})$ and $(\mathbf{a}, \mathbf{b})_{ab'''}^{l+2} : (X_{ab'''}^{l+2}, \mathbf{f}_{ab'''}^{l+2}, \underline{\mathbf{G}}_{ab'''}^{l+2}) \rightarrow (X_{\bar{a}\bar{b}}^{l+2}, \mathbf{f}_{\bar{a}\bar{b}}^{l+2}, \underline{\mathbf{G}}_{\bar{a}\bar{b}}^{l+2})$ satisfy $(\mathbf{a}, \mathbf{b})_{ab''}^{l+2} = (\mathbf{a}, \mathbf{b})_{ab'''}^{l+2} \circ (\sigma, \tau)$.

The restrictions of $\min \mathbf{T}_{ab}^l$ to $X_{ab''}^{l+2}$ and $X_{ab'''}^{l+2}$ are the pullbacks of $\min \mathbf{T}_{\bar{a}\bar{b}}^{l+2}$ by $(\mathbf{a}, \mathbf{b})_{ab''}^{l+2}$ and $(\mathbf{a}, \mathbf{b})_{ab'''}^{l+2}$ respectively, and so $(\mathbf{a}, \mathbf{b})_{ab''}^{l+2} = (\mathbf{a}, \mathbf{b})_{ab'''}^{l+2} \circ (\sigma, \tau)$ implies that the restrictions of $\min \mathbf{T}_{ab}^l$ to $X_{ab''}^{l+2}$ and to $X_{ab'''}^{l+2}$ are identified by σ . Hence $\min \mathbf{T}_{ab}^l|_{\partial^2 X_{ab}^l}$ is σ -invariant, and Definition A.34 and Theorem A.35 apply. This completes the inductive step.

Hence by induction we can choose $\mathbf{T}_{ab}^m$ satisfying $(*)_{ab}^m$ for all m, a, b . Now $X_a = \coprod_{b=1}^{n_a^0} X_{ab}^0$, and for each $b = 1, \dots, n_a^0$, if $\phi^0(a, b) = (\bar{a}, \bar{b})$ in R^0 , we have an isomorphism $(\mathbf{a}, \mathbf{b})_{ab}^0 : (X_{ab}^0, \mathbf{f}_{ab}^0, \underline{\mathbf{G}}_{ab}^0) \rightarrow (X_{\bar{a}\bar{b}}^0, \mathbf{f}_{\bar{a}\bar{b}}^0, \underline{\mathbf{G}}_{\bar{a}\bar{b}}^0)$, and a tent function $\mathbf{T}_{\bar{a}\bar{b}}^0$ for $(X_{\bar{a}\bar{b}}^0, \mathbf{f}_{\bar{a}\bar{b}}^0, \underline{\mathbf{G}}_{\bar{a}\bar{b}}^0)$. Define a tent function $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a)$ by

$T_a|_{X_{ab}^0} = ((a, b)_{ab}^0)^*(T_{ab}^0)$ for $b = 1, \dots, n_a^0$, for each $a \in A$. We have

$$\begin{aligned} 0 &= \partial \left[\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a] \right] = \sum_{a \in A} \rho_a \sum_{b=1}^{n_a^1} [X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1] \\ &= \sum_{\substack{a \in A, b=1, \dots, n_a^1 \\ \text{Set } (\bar{a}, \bar{b}) = \phi^1(a, b)}} \rho_a \epsilon_{ab}^1 [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1] = \sum_{(\bar{a}, \bar{b}) \in R^1} \left\{ \sum_{\substack{a \in A, b=1, \dots, n_a^1 \\ \phi^1(a, b) = (\bar{a}, \bar{b})}} \rho_a \epsilon_{ab}^1 \right\} [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1]. \end{aligned} \quad (229)$$

By construction the triples $(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)$ for $(\bar{a}, \bar{b}) \in R^1$ are connected, have no non-trivial automorphisms, and are mutually non-isomorphic, even through orientation-reversing isomorphisms. It follows that $[X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1]$ for $(\bar{a}, \bar{b}) \in R^1$ are linearly independent over R in $KC_{k-1}^{\text{ef}}(Y; R)$, as Definition 4.3(i)–(iii) impose no nontrivial relations between them. Thus (229) implies that the coefficient $\{\dots\}$ on the last line is zero, for all $(\bar{a}, \bar{b}) \in R^1$. The same argument as for (229) shows that

$$\begin{aligned} &\sum_{a \in A} \rho_a [Z_{\partial X_a, T_a|_{\partial X_a}}, \mathbf{f}_a|_{\partial X_a} \circ \pi_a, \mathbf{H}_{\partial X_a, T_a|_{\partial X_a}}] = \\ &\sum_{(\bar{a}, \bar{b}) \in R^1} \left\{ \sum_{\substack{a \in A, b=1, \dots, n_a^1 \\ \phi^1(a, b) = (\bar{a}, \bar{b})}} \rho_a \epsilon_{ab}^1 \right\} [Z_{X_{\bar{a}\bar{b}}^1, T_{\bar{a}\bar{b}}^1}, \mathbf{f}_{\bar{a}\bar{b}}^1 \circ \pi_{\bar{a}\bar{b}}^1, \mathbf{H}_{X_{\bar{a}\bar{b}}^1, T_{\bar{a}\bar{b}}^1}] = 0, \end{aligned} \quad (230)$$

since the coefficient $\{\dots\}$ on the second line is zero. This proves (225), and (226) follows, completing Step 1.

B.2 Step 2: deforming X_{ac} to compact manifolds $\tilde{X}_{ac}$

We work in the situation and with the notation of Steps 1 and 2 above, so that $\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$ is a cycle representing $\alpha \in KH_k^{\text{ef}}(Y; R)$, where X_{ac} has trivial stabilizers and the V_{ac}^i are manifolds, and our goal is to construct a homologous cycle $\sum_{a \in A} \sum_{c \in C_a} \rho_a [\tilde{X}_{ac}, \tilde{\mathbf{f}}_{ac}, \tilde{\mathbf{G}}_{ac}]$ with $\tilde{X}_{ac}$ a compact k -manifold with g-corners, constructed from X_{ac} by choosing C^1 small, smooth, transverse perturbations $\tilde{s}_{ac}^i$ of the Kuranishi maps s_{ac}^i of the Kuranishi neighbourhoods $(V_{ac}^i, E_{ac}^i, s_{ac}^i, \psi_{ac}^i)$ on X_{ac} in $\mathbf{I}_{ac}$ in $\tilde{\mathbf{G}}_{ac}$. Our argument is based on Fukaya and Ono [25, Th. 6.4], [24, Th. A1.23], but using single-valued sections rather than multisections.

To prove that $\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$ and $\sum_{a \in A} \sum_{c \in C_a} \rho_a [\tilde{X}_{ac}, \tilde{\mathbf{f}}_{ac}, \tilde{\mathbf{G}}_{ac}]$ are homologous, we will also construct a homology $[Z_{ac}, \mathbf{g}_{ac}, \mathbf{H}_{ac}]$ between $[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$ and $[\tilde{X}_{ac}, \tilde{\mathbf{f}}_{ac}, \tilde{\mathbf{G}}_{ac}]$, modulo terms over ∂X_{ac} . We first explain how to choose the $\tilde{s}_{ac}^i$ and define $\tilde{X}_{ac}, \tilde{\mathbf{f}}_{ac}, \tilde{\mathbf{G}}_{ac}, Z_{ac}, \mathbf{g}_{ac}, \mathbf{H}_{ac}$ for a single pair a, c , ignoring compatibility conditions between other pairs a', c' . Then later we will explain how to choose the $\tilde{s}_{ac}^i$ for all $a \in A$ and $c \in C_a$ so that the necessary compatibility conditions hold. For simplicity, in this first part we omit the subscripts ‘ ac ’, so we begin with a triple $(X, \mathbf{f}, \mathbf{G})$, choose perturbations $\tilde{s}^i$, and define $\tilde{X}, \tilde{\mathbf{f}}, \tilde{\mathbf{G}}, Z, \mathbf{g}, \mathbf{H}$.

Let $(X, \mathbf{f}, \mathbf{G})$ be a triple with $\mathbf{G} = (I, \boldsymbol{\eta}, \mathbf{G}^i : i \in I)$ and $I = (I, (V^i, \dots, \psi^i), f^i : i \in I, \dots)$, where X has trivial stabilizers and the V^i are manifolds. Define $\dot{V}^i, \dot{V}^{ij}, (\dot{I}, \dot{\boldsymbol{\eta}})$ as in Definition A.25, so that $\dot{V}^i \subset V^i$, $\dot{V}^{ij} \subset V^{ij}$ are open sets for $j \leq i$ with $\phi^{ij}(\dot{V}^{ij}) \subseteq \phi^{ij}(V^{ij})$. We shall choose perturbations $\tilde{s}^i$ of $s^i|_{\dot{V}^i}$ for $i \in I$. The necessity to use smaller open subsets $\dot{V}^i \subset V^i$ on which to define the perturbations $\tilde{s}^i$ is explained after Example A.24: without shrinking the domains V^i, V^{ij} we may be unable to choose the $\tilde{s}^i$ to be transverse over points in V^i lying in the boundary of $\phi^{ij}(V^{ij})$ for $j \leq i$ in I .

We will choose C^1 small perturbations $\tilde{s}^i$ of $s^i|_{\dot{V}^i}$ with $\tilde{s}^i$ *transverse* along $(\tilde{s}^i)^{-1}(0)$, such that if $j < i$ in I then $\hat{\phi}^{ij} \circ \tilde{s}^j \equiv \tilde{s}^i \circ \phi^{ij}$ on $\dot{V}^{ij}$, by induction on $i \in I$. Here is our inductive hypothesis. Suppose that $i \in I$, and that for $j < i$ in I we have already chosen a smooth section $\tilde{s}^j$ of $E^j|_{\dot{V}^j}$, such that $\tilde{s}^j$ extends smoothly and transversely from $\dot{V}^j$ to some open neighbourhood U^j of the closure $\overline{\dot{V}^j}$ of $\dot{V}^j$ in V^j , and $\tilde{s}^j$ is C^1 close to s^j , and $\tilde{s}^j$ is *transverse* along $(\tilde{s}^j)^{-1}(0)$, and if $k \leq j < i$ in I then $\hat{\phi}^{jk} \circ \tilde{s}^k \equiv \tilde{s}^j \circ \phi^{jk}$ on $\dot{V}^{jk}$. The first step $i = \min I$ is vacuous.

To do the inductive step we have to choose a smooth transverse section $\tilde{s}^i$ of $E^i|_{\dot{V}^i}$ which extends smoothly from $\dot{V}^i$ to some open neighbourhood U^i of $\overline{\dot{V}^i}$ in V^i , such that if $j < i$ in I then $\hat{\phi}^{ij} \circ \tilde{s}^j \equiv \tilde{s}^i \circ \phi^{ij}$ on $\dot{V}^{ij}$. That is, $\tilde{s}^i$ is prescribed on the subset $\bigcup_{j < i \text{ in } I} \phi^{ij}(\dot{V}^{ij})$ of $\dot{V}^i$, and we must extend it smoothly and transversely to $\dot{V}^i$.

As in the proof of Theorem 3.18, the prescribed values for $\tilde{s}^i$ from different $j, k < i$ are consistent, and the subset $\bigcup_{j < i \text{ in } I} \phi^{ij}(\dot{V}^{ij})$ is a locally closed submanifold of $\dot{V}^i$ except possibly at points of the *boundaries* $\partial(\phi^{ij}(\dot{V}^{ij})) = \phi^{ij}(\dot{V}^{ij}) \setminus \phi^{ij}(\dot{V}^{ij})$. Thus, except possibly at these boundaries, locally we can extend $\tilde{s}^i$ smoothly to $\dot{V}^i$, and since the $\tilde{s}^j$ for $j < i$ are *transverse*, locally we can choose $\tilde{s}^i$ to be transverse too. Note that as $V^i, \dot{V}^i$ are manifolds and E^i is a vector bundle, there are no problems with orbifold groups, and generic sections are *transverse*.

We have $\phi^{ij}(\dot{V}^{ij}) \subseteq \phi^{ij}(V^{ij})$ as in Definition A.25. But we already know that $\tilde{s}^j$ extends smoothly and transversely to some open neighbourhood U^j of $\overline{\dot{V}^j}$ in V^j , and therefore the prescribed values of $\tilde{s}^i$ on $\phi^{ij}(\dot{V}^{ij})$ extend smoothly and transversely to $\phi^{ij}(U^j \cap V^{ij})$, which contains $\partial(\phi^{ij}(\dot{V}^{ij}))$. Thus $\tilde{s}^i$ can locally be extended smoothly and transversely over the boundaries $\partial(\phi^{ij}(\dot{V}^{ij}))$, so the extension is locally possible everywhere in $\dot{V}^i$. Patching the local choices together with a partition of unity, we can choose $\tilde{s}^i$ with all the properties we need. This completes the inductive step.

We shall now construct $\tilde{X}, \tilde{\mathbf{f}}, \tilde{\mathbf{G}}$. As in Remark B.1(a), this is similar to parts of Fukaya and Ono's construction of *virtual cycles* [25, §6], [24, §A1]. Define

$$\tilde{X} = [\coprod_{i \in I} (\tilde{s}^i)^{-1}(0)] / \sim, \quad (231)$$

where $\sim$ is the equivalence relation induced by the maps $\phi^{ij} : (\tilde{s}^j)^{-1}(0) \cap \dot{V}^{ij} \rightarrow (\tilde{s}^i)^{-1}(0)$ for all $i, j \in I$ with $j \leq i$. Here ϕ^{ij} maps $(\tilde{s}^j)^{-1}(0)$ to $(\tilde{s}^i)^{-1}(0)$ since

$\hat{\phi}^{ij} \circ \tilde{s}^j \equiv \tilde{s}^i \circ \phi^{ij}$, and $\hat{\phi}^{ij}$ is an embedding and so injective.

Since the $\tilde{V}^i$ are manifolds and the $\tilde{s}^i$ are transverse, each subset $(\tilde{s}^i)^{-1}(0)$ is actually a manifold with g-corners, of dimension k . As the gluing maps ϕ^{ij} are embeddings $V^{ij} \rightarrow V^i$ they restrict to local diffeomorphisms from $(\tilde{s}^j)^{-1}(0)$ to $(\tilde{s}^i)^{-1}(0)$. Therefore $\tilde{X}$ has the structure of a k -manifold with g-corners, which is *Hausdorff* by the argument of Remark 3.14, and *compact* provided the $\tilde{s}^i$ are sufficiently close to s^i in C^0 , so that $(\tilde{s}^i)^{-1}(0)$ is close to $(s^i)^{-1}(0)$.

Define $\tilde{f} : \tilde{X} \rightarrow Y$ by $\tilde{f}|_{(\tilde{s}^i)^{-1}(0)} = f^i|_{(\tilde{s}^i)^{-1}(0)}$ for $i \in I$. Since $f^i \circ \phi^{ij} \equiv f^j$ on V^{ij} , this is compatible with the gluing maps ϕ^{ij} defining $\sim$, so $\tilde{f}$ is well-defined, and is clearly a smooth map. Define an *excellent coordinate system* $(\tilde{I}, \tilde{\eta})$ for $(\tilde{X}, \tilde{f})$ to have indexing set $\tilde{I} = \{k\}$ and Kuranishi neighbourhood $(\tilde{V}^k, \tilde{E}^k, \tilde{s}^k, \tilde{\psi}^k) = (\tilde{X}, \tilde{X}, 0, \text{id}_{\tilde{X}})$, where $\tilde{E}^k \rightarrow \tilde{V}^k = \tilde{X}$ is the zero vector bundle, so that $\tilde{E}^k = \tilde{X}$ as a manifold, and define $\tilde{f}^k : \tilde{V}^k \rightarrow Y$ by $\tilde{f}^k = \tilde{f}$, so that $\tilde{f}^k$ represents $\tilde{f}$. Define $\tilde{\eta}_k : \tilde{X} \rightarrow [0, 1]$ and $\tilde{\eta}_k^k : \tilde{V}^k \rightarrow [0, 1]$ by $\tilde{\eta}_k \equiv 1 \equiv \tilde{\eta}_k^k$.

Choose some map $\tilde{\underline{G}}^k : \tilde{X} \rightarrow \underline{P}$ such that $\tilde{\underline{G}} = (\tilde{I}, \tilde{\eta}, \tilde{\underline{G}}^k)$ is *effective gauge-fixing data* for $(\tilde{X}, \tilde{f})$. Ignoring boundary conditions, it is enough to take $\tilde{\underline{G}}^k : \tilde{X} \rightarrow \mathbb{R}^n \subset \underline{P}$ injective for some $n \gg 0$, but when we discuss how to choose $\tilde{s}_{ac}^i, \tilde{X}_{ac}, \tilde{f}_{ac}, \tilde{\underline{G}}_{ac}, Z_{ac}, \underline{g}_{ac}, \underline{H}_{ac}$ with compatibility between different $a \in A, c \in C_a$, we will explain in more detail how to choose these maps $\tilde{\underline{G}}^k$. Then $[\tilde{X}, \tilde{f}, \tilde{\underline{G}}]$ is well-defined in $KC_k^{\text{ef}}(Y; R)$.

Next we define $(Z, \underline{g}, \underline{H})$ such that Z is a perturbation of $[0, 1] \times X$, and $[Z, \underline{g}, \underline{H}]$ is a homology between $[X, \underline{f}, \underline{G}]$ and $[\tilde{X}, \tilde{f}, \tilde{\underline{G}}]$, modulo terms over a perturbation of $[0, 1] \times \partial X$. In $\underline{H}$ we will have a very good coordinate system $\underline{J}$ with indexing set $\underline{J} = \{i+1 : i \in I\} \cup \{k+1\}$ and Kuranishi neighbourhoods (W^j, F^j, t^j, ξ^j) for $j \in \underline{J}$, and we define these first. For $i \in I \setminus \{k\}$, define $W^{i+1} = ([0, \frac{1}{3}] \times V^i) \cup ([\frac{1}{3}, \frac{3}{4}] \times \tilde{V}^i)$. This is an open set in $[0, \frac{3}{4}] \times V^i$, and so is a manifold with g-corners. Define $F^{i+1} = \pi_{V^i}^*(E^i)$, so that $F^{i+1} \rightarrow W^{i+1}$ is a vector bundle.

If $k \notin I$, define $W^{k+1} = (\frac{2}{3}, 1] \times \tilde{X}$. If $k \in I$, define $W^{k+1} = ([0, \frac{1}{3}] \times V^k) \cup ([\frac{1}{3}, \frac{3}{4}] \times \tilde{V}^k) \cup (\frac{2}{3}, 1] \times \tilde{X}$. Here as $\text{vdim } X = k$ we have $\text{rank } E^k = 0$, so $(\tilde{s}^k)^{-1}(0) = \tilde{V}^k$, and we regard $\tilde{V}^k$ as an open subset of $\tilde{X}$, so that $[\frac{1}{3}, \frac{3}{4}] \times \tilde{V}^k$ and $(\frac{2}{3}, 1] \times \tilde{X}$ intersect in $(\frac{2}{3}, \frac{3}{4}] \times \tilde{V}^k$. In both cases define $F^{k+1} \rightarrow W^{k+1}$ to be the zero vector bundle, so that $F^{k+1} = W^{k+1}$ as a manifold.

Fix a smooth map $\eta : [0, 1] \rightarrow [0, 1]$ independent of $a \in A$ such that $\eta(x) = 0$ for $x \in [0, \frac{1}{3}]$ and $\eta(x) = 1$ for $x \in [\frac{2}{3}, 1]$. For $i \in I \setminus \{k\}$ define a smooth section t^{i+1} of F^{i+1} by $t^{i+1}(u, v) = s^i(v)$ for $u \in [0, \frac{1}{3}]$ and $t^{i+1}(u, v) = (1 - \eta(u))s^i(v) + \eta(u)\tilde{s}^i(v)$ for $u \in [\frac{1}{3}, \frac{3}{4}]$. Define $t^{k+1} = 0$. Following (231), define

$$Z = [\coprod_{j \in \underline{J}} (t^j)^{-1}(0)] / \approx, \quad (232)$$

where $\approx$ is the equivalence relation induced by $(u, v^i) \approx (u, v^j)$ if $u \in [0, \frac{3}{4}]$, $j \leq i$ in I , $(u, v^i) \in (t^{i+1})^{-1}(0)$, $(u, v^j) \in (t^{j+1})^{-1}(0)$, and $v^j \in V^{ij}$ with $\phi^{ij}(v^j) = v^i$, and $(u, v^i) \approx (u, v^k)$ if $u \in (\frac{2}{3}, \frac{3}{4}]$, $i \in I$, $(u, v^i) \in (t^{i+1})^{-1}(0)$, $(u, v^k) \in (\frac{2}{3}, 1] \times \tilde{X}$, and $v^i = v^k$ in $\tilde{X}$, where $\eta(u) = 1$ and $(u, v^i) \in (t^i)^{-1}(0)$ imply that $v^i \in (\tilde{s}^i)^{-1}(0)$, so we can regard v^i as an element of $\tilde{X}$. As the

gluing maps are local homeomorphisms, Z has the structure of a topological space, which is compact and Hausdorff provided $\tilde{s}^i$ is close to s^i in C^0 .

Define $\xi^j : (t^j)^{-1}(0) \rightarrow Z$ by $\xi^j = \text{id}_{(t^j)^{-1}(0)}$. Then (W^j, F^j, t^j, ξ^j) is a *Kuranishi neighbourhood* on Z for $j \in J$. When $k < j \leq i$ in I , define an open subset $W^{(i+1)(j+1)}$ in W^{j+1} to be $([0, \frac{1}{3}] \times V^{ij}) \cup ([\frac{1}{3}, \frac{3}{4}] \times \tilde{V}^{ij})$. Define

$$(\psi^{(i+1)(j+1)}, \hat{\psi}^{(i+1)(j+1)}) : (W^{(i+1)(j+1)}, F^{j+1}|_{W^{(i+1)(j+1)}}, t^{j+1}|_{W^{(i+1)(j+1)}}, \xi^{j+1}|_{W^{(i+1)(j+1)}}) \longrightarrow (W^{i+1}, F^{i+1}, t^{i+1}, \xi^{i+1}) \quad (233)$$

by $\psi^{(i+1)(j+1)} = (\text{id}_{[0, \frac{3}{4}]} \times \phi^{ij})|_{W^{(i+1)(j+1)}}$, $\hat{\psi}^{(i+1)(j+1)} = (\text{id}_{[0, \frac{3}{4}]} \times \hat{\phi}^{ij})|_{\pi^*(W^{(i+1)(j+1)})}$. We must check (233) satisfies Definition 2.18. Parts (a),(b) follow as $(\phi^{ij}, \hat{\phi}^{ij})$ is a coordinate change, (c) as $\hat{\phi}^{ij} \circ s^j \equiv s^i \circ \phi^{ij}$ on V^{ij} and $\hat{\phi}^{ij} \circ \tilde{s}^j \equiv \tilde{s}^i \circ \phi^{ij}$ on $\tilde{V}^{ij}$, and (d) by the definition of Z using gluing maps $\text{id}_{[0, \frac{3}{4}]} \times \phi^{ij}$ in $\approx$. The only subtle part is (e). This holds as $\tilde{s}^i, \tilde{s}^j$ are C^1 close to s^i, s^j , so t^{i+1}, t^{j+1} are C^1 -close to $s^i \circ \pi_{V^i}, s^i \circ \pi_{V^i}$. We need $d\hat{t}^{i+1}$ to be an isomorphism over $(t^{j+1})^{-1}(0)$. This would hold if t^{i+1}, t^{j+1} were replaced by $s^i \circ \pi_{V^i}, s^j \circ \pi_{V^j}$, and being an isomorphism is an open condition, so $d\hat{t}^{i+1}$ is an isomorphism over $(t^{j+1})^{-1}(0)$ provided $\tilde{s}^i, \tilde{s}^j$ are sufficiently close to s^i, s^j in C^1 .

If $k \notin I$ and $k < i \in I$, define $W^{(i+1)(k+1)}$ in $W^{k+1} = (\frac{2}{3}, 1] \times \tilde{X}$ by $W^{(i+1)(k+1)} = (\frac{2}{3}, \frac{3}{4}] \times (\tilde{s}^i)^{-1}(0)$. Define $\psi^{(i+1)(k+1)} : W^{(i+1)(k+1)} \rightarrow W^{i+1}$ to be inclusion, regarding $W^{(i+1)(k+1)}$ as an open subset of $(t^{i+1})^{-1}(0) \subseteq W^{i+1}$. Define $\hat{\psi}^{(i+1)(k+1)} : F^{k+1}|_{W^{(i+1)(k+1)}} \rightarrow F^{i+1}$ by $\hat{\psi}^{(i+1)(k+1)} = z \circ \psi^{(i+1)(k+1)}$, where $z : W^{i+1} \rightarrow F^{i+1}$ is the zero section. This is valid as $F^{k+1} = W^{k+1}$, so $F^{k+1}|_{W^{(i+1)(k+1)}} = W^{(i+1)(k+1)}$. If $k \in I$ and $k < i \in I$, we combine the previous two definitions for $W^{(i+1)(k+1)}, \psi^{(i+1)(k+1)}, \hat{\psi}^{(i+1)(k+1)}$ in the obvious way. We also define $W^{(k+1)(k+1)} = W^{k+1}$, and $\psi^{(k+1)(k+1)}, \hat{\psi}^{(k+1)(k+1)}$ to be the identities. Then (233) is a coordinate change also for $j = k$ or $i = j = k$.

Define a Kuranishi structure on Z as follows. There is a natural projection $\pi_{[0,1]} : Z \rightarrow [0, 1]$, with homeomorphisms $\pi_{[0,1]}^{-1}([0, \frac{1}{3}]) \cong [0, \frac{1}{3}] \times X$ and $\pi_{[0,1]}^{-1}([\frac{2}{3}, 1]) \cong [\frac{2}{3}, 1] \times \tilde{X}$. On $\pi_{[0,1]}^{-1}([0, \frac{1}{3}])$, let Z have the Kuranishi structure of $[0, \frac{1}{3}] \times X$. On $\pi_{[0,1]}^{-1}([\frac{2}{3}, 1])$, let Z have the Kuranishi structure (manifold structure) of $(\frac{2}{3}, 1] \times \tilde{X}$. For each $p \in \pi_{[0,1]}^{-1}([\frac{1}{3}, \frac{2}{3}])$, let $j \in J$ be least such that $p \in (t^j)^{-1}(0)$, regarding $(t^j)^{-1}(0)$ as a subset of Z by (232). Then $(W^j, \dots, \xi^j)$ is a Kuranishi neighbourhood of p . Define the germ of Kuranishi neighbourhoods on Z at p to be the equivalence class of $(W^j, \dots, \xi^j)$.

Since we use the Kuranishi structures from $[0, 1] \times X$ and $[0, 1] \times \tilde{X}$ on open subsets of Z , and choose $j \in J$ least for each $p \in \pi_{[0,1]}^{-1}([\frac{1}{3}, \frac{2}{3}])$ so that the function $p \mapsto j$ is upper semicontinuous, to define the germ of coordinate changes in the Kuranishi structure on Z , we need coordinate changes as follows:

- (i) from $(V_q, \dots, \psi_q)$ to $(W^j, \dots, \xi^j)$, where $(V_q, \dots, \psi_q)$ lies in the germ of Kuranishi neighbourhoods on $[0, 1] \times X$ at $q \in [0, \frac{1}{3}] \times \tilde{X}$, and $(W^j, \dots, \xi^j)$ is in the germ of Kuranishi neighbourhoods of $p \in \pi_{[0,1]}^{-1}(\{\frac{1}{3}\})$;

- (ii) from $(V_q, V_q, 0, \text{id}_{V_q})$ to $(W^j, \dots, \xi^j)$, where V_q is a small open neighbourhood of $q \in (\frac{2}{3}, 1] \times \tilde{X}$, so that $(V_q, V_q, 0, \text{id}_{V_q})$ lies in the germ of Kuranishi neighbourhoods on $[0, 1] \times \tilde{X}$, and $(W^j, \dots, \xi^j)$ is in the germ of Kuranishi neighbourhoods of $p \in \pi_{[0,1]}^{-1}(\{\frac{2}{3}\})$; and
- (iii) If $p, q \in \pi_{[0,1]}^{-1}([\frac{1}{3}, \frac{2}{3}])$ with $p \in (t^j)^{-1}(0)$, $j \in J$ least, and $q \in (t^j)^{-1}(0) \cap (t^{j'})^{-1}(0)$ for $j' < j \in J$ least, a coordinate change from $(W^{jj'}, \dots, \xi^{jj'}|_{W^{jj'}})$ to $(W^j, \dots, \xi^j)$ for $j' < j$ in J , where $(W^{jj'}, \dots, \xi^{jj'}|_{W^{jj'}})$ is equivalent to $(W^{j'}, F^{j'}, t^{j'}, \xi^{j'})$ as a Kuranishi neighbourhood of q in Z , and so lies in the germ at q .

For (i) the coordinate changes exist as $(W^j, \dots, \xi^j)$ is a compatible Kuranishi neighbourhood on $[0, 1] \times X$ over $[0, \frac{1}{3}] \times X$, since it is constructed from a compatible Kuranishi neighbourhood $(V^{j-1}, \dots, \psi^{j-1})$ on X . For (ii) the coordinate changes are easily constructed from the inclusions $V_q \subseteq (t^j)^{-1}(0) \subseteq W^j$. For (iii), the coordinate change is $(\psi^{jj'}, \hat{\psi}^{jj'})$ in (233). It is now not difficult to show this defines a *Kuranishi structure* on Z , for which the $(W^j, \dots, \xi^j)$ for $j \in J$ are compatible Kuranishi neighbourhoods.

Then Z is a compact Kuranishi space with $\text{vdim } Z = k + 1$. The product orientation on $[0, 1] \times X$ induces orientations on the $(W^i, \dots, \xi^i)$, yielding an orientation on Z . Then

$$\partial Z \cong (\{1\} \times \tilde{X}) \amalg -(\{0\} \times X) \amalg -Z^\partial \quad (234)$$

in oriented Kuranishi spaces, where Z^∂ is a perturbation of $[0, 1] \times \partial X$, obtained by applying the construction above to ∂X rather than X .

Define $g^{i+1} : W^{i+1} \rightarrow Y$ by $g^i = f^i \circ \pi_{V^i}$ for $i \in I \setminus \{k\}$, and $g^{k+1} : W^{k+1} \rightarrow Y$ by $g^{k+1} = \tilde{f} \circ \pi_{\tilde{X}}$ when $k \notin I$ or on $(\frac{2}{3}, 1] \times \tilde{X}$, and $g^{k+1} = \tilde{f} \circ \pi_{V^i}$ when $k \in I$ on $([0, \frac{1}{3}] \times V^k) \cup ([\frac{1}{3}, \frac{3}{4}] \times \tilde{V}^k)$. These represent a strongly smooth map $\mathbf{g} : Z \rightarrow Y$, which restricts to $\mathbf{f}$ on $\{0\} \times X$ and $\tilde{\mathbf{f}}$ on $\{1\} \times \tilde{X}$, under the identification (234). Then $\mathbf{J} = (J, (W^j, F^j, t^j, \xi^j), g^j : j \in J, \dots)$ is a *very good coordinate system* for $(Z, \mathbf{g})$.

Choose continuous functions $\alpha_1, \alpha_2, \alpha_3 : [0, 1] \rightarrow [0, 1]$ with $\alpha_1 + \alpha_2 + \alpha_3 \equiv 1$ such that $\alpha_1 \equiv 1$ on $[0, \frac{1}{4}]$, $\alpha_1 \equiv 0$ on $[\frac{1}{3}, 1]$, $\alpha_2 \equiv 0$ on $[0, \frac{1}{4}]$, $\alpha_2 \equiv 1$ on $[\frac{1}{3}, \frac{2}{3}]$, $\alpha_2 \equiv 0$ on $[\frac{3}{4}, 1]$, $\alpha_3 \equiv 0$ on $[0, \frac{2}{3}]$, and $\alpha_3 \equiv 1$ on $[\frac{3}{4}, 1]$. We shall define functions $\zeta_j^{\tilde{j}} : W^{\tilde{j}} \rightarrow [0, 1]$ for $j, \tilde{j} \in J$. For $i, i' \in I \setminus \{k\}$, set

$$\zeta_{i+1}^{i'+1}(w) = \alpha_1 \circ \pi_{[0,1]}(w) \cdot \eta_i^{i'} \circ \pi_{V^{i'}}(w) + \alpha_2 \circ \pi_{[0,1]}(w) \cdot \dot{\eta}_i^{i'} \circ \pi_{V^{i'}}(w), \quad (235)$$

$$\begin{aligned} \zeta_{k+1}^{i'+1}(w) &= \alpha_1 \circ \pi_{[0,1]}(w) \cdot \eta_k^{i'} \circ \pi_{V^{i'}}(w) + \alpha_2 \circ \pi_{[0,1]}(w) \cdot \dot{\eta}_k^{i'} \circ \pi_{V^{i'}}(w) \\ &\quad + \alpha_3 \circ \pi_{[0,1]}(w). \end{aligned} \quad (236)$$

When $k \in I$, define ζ_{i+1}^{k+1} on $([0, \frac{1}{3}] \times V^k) \cup ([\frac{1}{3}, \frac{3}{4}] \times \tilde{V}^k)$ by (235) and (236) with $i' = k$. Regarding $\tilde{X}$ as a union of subsets $(\tilde{s}^{i'})^{-1}(0)$, define ζ_{i+1}^{k+1} on

$(\frac{2}{3}, 1] \times (\tilde{s}^{i'})^{-1}(0)$ for each $i' \in I$ by

$$\zeta_{i+1}^{k+1}(w) = \alpha_2 \circ \pi_{[0,1]}(w) \cdot \dot{\eta}_i^{i'} \circ \pi_{(\tilde{s}^{i'})^{-1}(0)}, \quad i \neq k, \quad (237)$$

$$\zeta_{k+1}^{k+1}(w) = \alpha_2 \circ \pi_{[0,1]}(w) \cdot \dot{\eta}_k^{i'} \circ \pi_{(\tilde{s}^{i'})^{-1}(0)} + \alpha_3 \circ \pi_{[0,1]}(w). \quad (238)$$

One can show that (235)–(238) are consistent on their overlaps, so the $\zeta_j^{\tilde{j}}$ are well-defined, and that the $\zeta_j^{\tilde{j}}$ are continuous and satisfy $\sum_{j \in J} \zeta_j^{\tilde{j}} \equiv 1$ on $W^{\tilde{j}}$ and $\zeta_j^{\tilde{j}} \circ \psi^{\tilde{j}\tilde{j}'} \equiv \zeta_j^{j'}|_{W^{\tilde{j}\tilde{j}'}}$ for $j, \tilde{j}, j' \in J$ with $j' \leq \tilde{j}$. Define $\zeta_j : Z \rightarrow [0, 1]$ for $j \in J$ by $\zeta_j|_{(t^{\tilde{j}})^{-1}(0)} \equiv \zeta_j^{\tilde{j}}|_{(t^{\tilde{j}})^{-1}(0)}$ for all $\tilde{j} \in J$. Then the ζ_j are well-defined and continuous with $\sum_{j \in J} \zeta_j \equiv 1$. Write $\zeta = (\zeta_j, \zeta_j^{\tilde{j}} : j, \tilde{j} \in J)$. Then (J, ζ) is a *really good coordinate system* for (Z, g) .

Furthermore, since (I, η) is an excellent coordinate system, $\partial^l V^i$ has no unnecessary connected components for all $i \in I$ and $l \geq 0$, so $\partial^l W^{i+1}$ has no unnecessary connected components. (To see this, note that for $i \neq k$, $\partial^l W^{i+1}$ is the disjoint union of l copies of $\partial^{l-1} V^i$, and a piece which retracts onto $\partial^l V^i$. When $i = k$, the whole of W^{i+1} is necessary as $t^{k+1} \equiv 0$.) Hence (J, ζ) is an *excellent good coordinate system* for (Z, g) .

Choose some maps $\underline{H}^j : F^j \rightarrow \underline{P}$ for $j \in J$ such that $\underline{H} = (J, \zeta, \underline{H}^j : j \in J)$ is *effective gauge-fixing data* for (Z, g) , satisfying the boundary conditions

$$\underline{H}^{i+1}(0, e) = \underline{G}^i(e), \quad i \in I, \quad e \in E^i, \quad \text{and} \quad \underline{H}^{k+1}(0, e) = \tilde{G}^k(e), \quad e \in \tilde{X}. \quad (239)$$

Ignoring boundary conditions, it is enough to define $\underline{H}^{i+1}(u, e)$ for $u = 0$ or 1 by (239), and to define $\underline{H}^{i+1}(u, e)$ for $u \in (0, 1)$ by some injective map $F^j \cap \pi_{[0,1]}^{-1}((0, 1)) \rightarrow \mathbb{R}^n \subset \underline{P}$ for some $n \gg 0$. When we discuss how to choose $Z_{ac}, g_{ac}, \underline{H}_{ac}$ with compatibility between different $a \in A$, $c \in C_a$ below, we will explain in more detail how to choose these maps $\underline{H}^j$. Then $[Z, g, \underline{H}]$ is well-defined in $KC_{k+1}^{\text{ef}}(Y; R)$. Using (234) we find that in $KC_k^{\text{ef}}(Y; R)$ we have

$$\partial[Z, g, \underline{H}] = [\tilde{X}, \tilde{f}, \tilde{\underline{G}}] - [X, f, \underline{G}] - [Z^\partial, g|_{Z^\partial}, \underline{H}|_{Z^\partial}]. \quad (240)$$

This concludes our discussion of perturbing a single $(X, f, \underline{G})$.

Next we explain how to choose such $(\tilde{X}_{ac}, \tilde{f}_{ac}, \tilde{\underline{G}}_{ac})$ and $(Z_{ac}, g_{ac}, \underline{H}_{ac})$ for all $a \in A$ and $c \in C_a$, with boundary compatibilities. The equation we need to hold, similar to (225), is

$$\sum_{a \in A} \sum_{c \in C_a} \rho_a [Z_{ac}^\partial, g_{ac}|_{Z_{ac}^\partial}, \underline{H}_{ac}|_{Z_{ac}^\partial}] = 0. \quad (241)$$

We would like (241) to follow from $\sum_{a \in A} \sum_{c \in C_a} \rho_a [\partial X_{ac}, f_{ac}|_{\partial X_{ac}}, \underline{G}_{ac}|_{\partial X_{ac}}] = 0$, in the same way as (225) follows from (224). Assuming (241), inserting subscripts ‘ ac ’ in (240), multiplying it by ρ_a , and summing over all $a \in A$ and $c \in C_a$ yields

$$\begin{aligned} & \partial \left(\sum_{a \in A} \sum_{c \in C_a} \rho_a [Z_{ac}, g_{ac}, \underline{H}_{ac}] \right) \\ &= \sum_{a \in A} \sum_{c \in C_a} \rho_{ac} [\tilde{X}_{ac}, \tilde{f}_{ac}, \tilde{\underline{G}}_{ac}] - \sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, f_{ac}, \underline{G}_{ac}]. \end{aligned} \quad (242)$$

Thus $\sum_{a \in A} \sum_{c \in C_a} \rho_a [\tilde{X}_{ac}, \tilde{f}_{ac}, \tilde{\mathbf{G}}_{ac}]$ is a cycle homologous to $\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$, and so represents α , by Step 1.

The only arbitrary choices made in the construction of $\tilde{X}, \tilde{f}, \tilde{\mathbf{G}}, Z, \mathbf{g}, \mathbf{H}$ above were the sections $\tilde{s}^i$ of $E^i|_{\tilde{V}^i}$ for $i \in I$, and the maps $\tilde{\mathbf{G}}^k : \tilde{X} \rightarrow \underline{P}$ and $\tilde{\mathbf{H}}^j : \tilde{F}^j \rightarrow \underline{P}$ for $j \in J$. Inserting subscripts ‘ ac ’, we will give a method for choosing $\tilde{s}_{ac}^i$ for $i \in I_{ac}$, $\tilde{\mathbf{G}}_{ac}^k$ and $\tilde{\mathbf{H}}_{ac}^j$ for $j \in J_{ac}$ for all $a \in A$ and $c \in C_a$ such that (241) holds. As explained in Step 2, we will actually construct the smooth $\tilde{s}_{ac}^i$ for all $c \in C_a$ from piecewise smooth $\tilde{s}_a^i$ on X_a , and similarly, we will define $\tilde{\mathbf{G}}_{ac}^k, \tilde{\mathbf{H}}_{ac}^j$ for $c \in C_a$ from maps $\tilde{\mathbf{G}}_a^k$ and $\tilde{\mathbf{H}}_a^j$ over X_a .

We choose the $\tilde{s}_a^i$ by an inductive procedure on decreasing codimension like that in Step 1. Use the notation of Step 1 above. For $m = 0, \dots, M$ and (a, b) in R^m , here is our inductive hypothesis:

- ($\star$) $_{ab}^m$ In the triple $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ of Step 1, write $(\mathbf{I}_{ab}^m, \boldsymbol{\eta}_{ab}^m)$ for the excellent coordinate system in $\mathbf{G}_{ab}^m$, with $\mathbf{I}_{ab}^m = (I_{ab}^m, (V_{ab}^{m,i}, \dots, \psi_{ab}^{m,i}), f_{ab}^{mi} : i \in I_{ab}^m, \dots)$, and construct $(\dot{\mathbf{I}}_{ab}^m, \dot{\boldsymbol{\eta}}_{ab}^m), \dot{V}_{ab}^{m,i}, \dot{V}_{ab}^{m,ij}, \dots$ as in Definition A.25. Let $\mathbf{T}_{ab}^m = (T_{ab}^{m,i} : i \in I_{ab}^m)$ be the tent function for $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ in ($\star$) $_{ab}^m$ in Step 1, so that $T_{ab}^{m,i}$ is a tent function on $\dot{V}_{ab}^{m,i}$.

Then $\tilde{s}_{ab}^{m,i}$ is a continuous, piecewise smooth section of $E_{ab}^{m,i}|_{\dot{V}_{ab}^{m,i}}$ subordinate to the tent function $T_{ab}^{m,i}$ for each $i \in I_{ab}^m$, satisfying:

- (a) $\tilde{s}_{ab}^{m,i}$ is C^1 close to $s_{ab}^{m,i}$, and transverse (both in a piecewise smooth sense).
- (b) If $j < i$ in I_{ab}^m then $\hat{\phi}_{ab}^{m,ij} \circ \tilde{s}_{ab}^{m,j} \equiv \tilde{s}_{ab}^{m,i} \circ \phi_{ab}^{m,ij}$ on $\dot{V}_{ab}^{m,ij}$.
- (c) As in ($\star$) $_{ab}^m$ we have an isomorphism (227) between $\partial(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ and $\coprod_{b' \in B_{ab}^m} (X_{ab'}^{m+1}, \mathbf{f}_{ab'}^{m+1}, \mathbf{G}_{ab'}^{m+1})$. We require that this isomorphism should identify $\tilde{s}_{ab}^{m,i}|_{\partial \dot{V}_{ab}^{m,i}}$ with $\coprod_{b' \in B_{ab}^m} \tilde{s}_{ab'}^{m+1,i-1}$ for all $i \in I_{ab}^m$.

We now choose $\tilde{s}_{ab}^{m,i}$ satisfying ($\star$) $_{ab}^m$ by induction on decreasing $m = M, M-1, \dots, 1, 0$, for all (a, b) in R^m and $i \in I_{ab}^m$. For the first step, when $m = M$, as $P^{M+1} = \emptyset$, if $(a, b) \in R^M$ then $\partial X_{ab}^M = \emptyset$, so the boundary conditions (c) in ($\star$) $_{ab}^M$ are trivial. By induction on increasing $i \in I_{ab}^M$, we choose smooth (*not* piecewise smooth) transverse sections $\tilde{s}_{ab}^{M,i}$ of $E_{ab}^{M,i}|_{\dot{V}_{ab}^{M,i}}$ satisfying (a),(b) above.

This is possible as in the construction of $\tilde{X}$ without boundary conditions above, with one extra remark about orbifolds.

As $\mathbf{G}_{ab}^M$ is effective gauge-fixing data, $(V_{ab}^{M,i}, E_{ab}^{M,i}, s_{ab}^{M,i}, \psi_{ab}^{M,i})$ is an *effective* Kuranishi neighbourhood in the sense of Definition 3.35. Thus, even though $\dot{V}_{ab}^{M,i}$ may be an orbifold, its orbifold groups act trivially on the fibres of $E_{ab}^{M,i}$. This implies that generic smooth sections of $E_{ab}^{M,i}$ are transverse, so we can indeed choose $\tilde{s}_{ab}^{M,i}$ transverse as in (a). However, if the orbifold groups of $\dot{V}_{ab}^{M,i}$ were allowed to act nontrivially on the fibres of $E_{ab}^{M,i}$, it could happen that $E_{ab}^{M,i}$ admits no transverse sections $\tilde{s}_{ab}^{M,i}$ at all, since being invariant under the

orbifold groups might force $\tilde{s}_{ab}^{M,i}$ to be zero on an orbifold stratum of $\dot{V}_{ab}^{M,i}$ in a non-transverse way. This finishes the first step.

For the inductive step, suppose that for some $l = 0, \dots, M-1$ we have chosen data $\tilde{s}_{ab}^{m,i}$ satisfying $(\star)_{ab}^m$ for all $l < m \leq M$, $(a, b) \in R^m$ and $i \in I_{ab}^m$. We will construct $\tilde{s}_{ab}^{l,i}$ satisfying $(\star)_{ab}^l$ for all $(a, b) \in R^l$ and $i \in I_{ab}^l$. First, by induction on increasing $i \in I_{ab}^l$, choose generic smooth (*not* piecewise smooth) sections $\hat{s}_{ab}^{l,i}$ of $E_{ab}^{l,i}|_{\dot{V}_{ab}^{l,i}}$ satisfying (a),(b) above with $\hat{s}_{ab}^{l,i}$ in place of $\tilde{s}_{ab}^{m,i}$, but *without* boundary conditions such as (c). This is possible as in the first step.

To define $\tilde{s}_{ab}^{l,i}$ we use a version of the piecewise smooth extension idea of §A.1.4, in Definition A.12 and Proposition A.14. That is, $(\star)_{ab}^l$ (c) prescribes piecewise smooth values for $\tilde{s}_{ab}^{l,i}$ over $\partial\dot{V}_{ab}^{l,i}$, and we want to extend these to piecewise smooth values over $\dot{V}_{ab}^{l,i}$. We do this by choosing an arbitrary smooth section $\hat{s}_{ab}^{l,i}$ on the interior of $\dot{V}_{ab}^{l,i}$, corresponding to $\tilde{\eta}$ in Proposition A.14, and then interpolating between $\hat{s}_{ab}^{l,i}$ and the boundary prescribed values for $\tilde{s}_{ab}^{l,i}$ near $\partial\dot{V}_{ab}^{l,i}$ using the flow of $v_{ab}^{l,i}$. However, we have to modify the construction of Proposition A.14 slightly, as $T_{ab}^{l,i}$ was made not using Definition A.12, but with Definition A.32, a noncompact, orbifold version of Definition A.15.

Define subsets W_1, W_2 of $\dot{V}_{ab}^{l,i}$ by

$$\begin{aligned} W_1 &= \{w \in \dot{V}_{ab}^{l,i} : \min T_{ab}^{l,i}(w) \geq D\}, \\ W_2 &= \{w \in \dot{V}_{ab}^{l,i} : w = \exp(tv_{ab}^{l,i})x, t \in [0, 2), (x, B) \in \partial U_{ab}^{l,i}\}. \end{aligned}$$

Then W_1 is closed in $\dot{V}_{ab}^{l,i}$ as $\min T_{ab}^{l,i}$ is continuous, and W_2 is open in $\dot{V}_{ab}^{l,i}$, and $W_1 \cup W_2 = \dot{V}_{ab}^{l,i}$, since the branches $u + \theta(2-t)$ in (218) are supported on W_2 , so for $v \notin W_2$ we see that $\min T_{ab}^{l,i}(v)$ is of the form $D + l^2$ or $D + \epsilon^2$, so $\min T_{ab}^{l,i}(v) \geq D$ and $v \in W_1$.

Write $\tilde{s}_{ab}^{l,i}$ for the piecewise smooth section of $E_{ab}^{l,i}|_{\partial\dot{V}_{ab}^{l,i}}$ prescribed as boundary conditions for $\tilde{s}_{ab}^{l,i}$ in $(\star)_{ab}^l$ (c). Then as in §A.3.4–§A.3.5, $\tilde{s}_{ab}^{l,i}$ extends to $\partial U_{ab}^{l,i}$, for some open neighbourhood $U_{ab}^{l,i}$ of $\overline{V_{ab}^{l,i}}$ in $V_{ab}^{l,i}$, such that Definition A.32(a)–(d) hold for $U_{ab}^{l,i}$ and the vector field $v_{ab}^{l,i}$ on $V_{ab}^{l,i}$ used to construct $T_{ab}^{l,i}$. From $(\star)_{ab}^{l+1}$ (c) for $(\bar{a}, \bar{b}) = \phi^{l+1}(a, b')$, all $b' \in B_{ab}^l$, we can deduce that $\tilde{s}_{ab}^{l,i}|_{\partial^2 U_{ab}^{l,i}}$ is invariant under the natural involution $\sigma : \partial^2 U_{ab}^{l,i} \rightarrow \partial^2 \dot{V}_{ab}^{l,i}$, in the same way as we showed that $\min T_{ab}^{l,i}|_{\partial^2 X_{ab}^l}$ is invariant under the natural involution $\sigma : \partial^2 X_{ab}^l \rightarrow \partial^2 X_{ab}^l$ in the proof of $(\star)_{ab}^m$ in §B.1.

We will extend $\tilde{s}_{ab}^{l,i}$ to a continuous, piecewise smooth section $\tilde{s}_{ab}^{l,i}$ of $E_{ab}^{l,i}$ over W_2 subordinate to the tent function $T_{ab}^{l,i}|_{W_2}$. Let $\nabla_{ab}^{l,i}$ be a connection on the orbifold vector bundle $E_{ab}^{l,i} \rightarrow V_{ab}^{l,i}$. If $w \in W_2$ then $w = \exp(tv_{ab}^{l,i})x$ for some $t \in [0, 2)$ and $(x, B) \in \partial U_{ab}^{l,i}$. Here t, x are uniquely determined by w , as the flow-line of $-v_{ab}^{l,i}$ first hits $\iota(\partial U_{ab}^{l,i})$ at x in time t . The choice of local boundary component B of $U_{ab}^{l,i}$ at x is not uniquely determined.

However, since $\tilde{s}_{ab}^{l,i}|_{\partial^2 U_{ab}^{l,i}}$ is invariant under the natural involution $\sigma : \partial^2 U_{ab}^{l,i}$

$\rightarrow \partial^2 \dot{V}_{ab}^{l,i}$, it follows that $\tilde{s}_{ab}^{l,i}(x, B) \in E_{ab}^{l,i}|_x$ depends only on x , not on the choice of B . Thus $\tilde{s}_{ab}^{l,i}(x, B)$ is also uniquely determined by w . Define

$$\tilde{s}_{ab}^{l,i}(w) = s_{ab}^{l,i}(w) + P_{\nabla_{ab}^{l,i}}(\tilde{s}_{ab}^{l,i}(x, B) - s_{ab}^{l,i}(x)), \quad (243)$$

where $P_{\nabla_{ab}^{l,i}} : E_{ab}^{l,i}|_x \rightarrow E_{ab}^{l,i}|_w$ is the *parallel transport map* along the flow-line of $v_{ab}^{l,i}$ from x to w using the connection $\nabla_{ab}^{l,i}$. Considering the construction of $T_{ab}^{l,i}$, we see that $\tilde{s}_{ab}^{l,i}$ is continuous and piecewise smooth subordinate to $T_{ab}^{l,i}|_{W_2}$, with $\tilde{s}_{ab}^{l,i}|_{\partial \dot{V}_{ab}^{l,i}} \equiv \tilde{s}_{ab}^{l,i}$.

By the construction in §A.3.4–§A.3.5, $\min T_{ab}^{l,i}|_{\partial \dot{V}_{ab}^{l,i}}$ is bounded above, and D is strictly greater than this bound. Thus there exists $D' < D$ with $\min T_{ab}^{l,i}|_{\partial \dot{V}_{ab}^{l,i}} \leq D'$. Choose a smooth function $\eta : \mathbb{R} \rightarrow [0, 1]$ with $\eta(t) = 0$ for $t \leq D'$ and $\eta(t) = 1$ for $t \geq D$. Define a section $\tilde{s}_{ab}^{l,i}$ of $E_{ab}^{l,i}$ over $\dot{V}_{ab}^{l,i}$ by

$$\tilde{s}_{ab}^{l,i}(w) = \begin{cases} \hat{s}_{ab}^{l,i}(w), & w \in W_1, \\ (\eta \circ \min T_{ab}^{l,i})(w) \tilde{s}_{ab}^{l,i}(w) + (1 - \eta \circ \min T_{ab}^{l,i})(w) \hat{s}_{ab}^{l,i}(w), & w \in W_2. \end{cases} \quad (244)$$

On the overlap $W_1 \cap W_2$ we have $\min T_{ab}^{l,i}(w) \geq D$ and $\eta \circ \min T_{ab}^{l,i}(w) = 1$, so $\tilde{s}_{ab}^{l,i}$ is well-defined. As $\hat{s}_{ab}^{l,i}, \eta$ are smooth and $\min T_{ab}^{l,i}, \tilde{s}_{ab}^{l,i}$ are continuous and piecewise smooth subordinate to $T_{ab}^{l,i}$, we see that $\tilde{s}_{ab}^{l,i}$ is *continuous* and *piecewise smooth subordinate to $T_{ab}^{l,i}$* .

On $\partial \dot{V}_{ab}^{l,i}$ we have $\min T_{ab}^{l,i} \leq D'$, $\eta \circ \min T_{ab}^{l,i} \equiv 0$ and $\tilde{s}_{ab}^{l,i}|_{\partial \dot{V}_{ab}^{l,i}} \equiv \hat{s}_{ab}^{l,i}$, so $\tilde{s}_{ab}^{l,i}|_{\partial \dot{V}_{ab}^{l,i}} \equiv \hat{s}_{ab}^{l,i}$. Hence $\tilde{s}_{ab}^{l,i}$ satisfies $(\star)_{ab}^{l,i}(c)$. For part (a), from $(\star)_{ab}^{l,i+1}(a)$ for $(\bar{a}, \bar{b}) = \phi^{l+1}(a, b')$, all $b' \in B_{ab}^l$, it follows that $\tilde{s}_{ab}^{l,i}$ is C^1 -close to $\hat{s}_{ab}^{l,i}|_{\partial \dot{V}_{ab}^{l,i}}$ and transverse, in a piecewise smooth sense. Thus in (243), $\tilde{s}_{ab}^{l,i}(x, B) - s_{ab}^{l,i}(x)$ is C^1 -small as a function of $(x, B) \in \partial \dot{V}_{ab}^{l,i}$, which implies that $\tilde{s}_{ab}^{l,i}(w) - s_{ab}^{l,i}(w)$ is C^1 -small as a function of $w \in W_2$. So $\tilde{s}_{ab}^{l,i}$ is C^1 -close to $s_{ab}^{l,i}$. Also $\tilde{s}_{ab}^{l,i}$ was chosen above to be C^1 -close to $s_{ab}^{l,i}$. Therefore $\tilde{s}_{ab}^{l,i}$ in (244) is also C^1 -close to $s_{ab}^{l,i}$.

As $\tilde{s}_{ab}^{l,i}$ and $\hat{s}_{ab}^{l,i}$ were chosen generic and transverse, we see that $\tilde{s}_{ab}^{l,i}$ is transverse, and $(\star)_{ab}^{l,i}(a)$ holds. For part (b), the $\hat{s}_{ab}^{l,i}$ for $i \in I_{ab}^l$ were chosen to satisfy $\hat{\phi}_{ab}^{l,ij} \circ \hat{s}_{ab}^{l,j} \equiv \hat{s}_{ab}^{l,i} \circ \hat{\phi}_{ab}^{l,ij}$ if $j < i$ in I_{ab}^l . Also $(\star)_{ab}^{l,i+1}(b)$ for $(\bar{a}, \bar{b}) = \phi^{l+1}(a, b')$, all $b' \in B_{ab}^l$, imply that the $\tilde{s}_{ab}^{l,i}$ satisfy $\hat{\phi}_{ab}^{l,ij} \circ \tilde{s}_{ab}^{l,j} \equiv \tilde{s}_{ab}^{l,i} \circ \hat{\phi}_{ab}^{l,ij}$ if $j < i$ in I_{ab}^l . Now $\tilde{s}_{ab}^{l,i}$ was constructed using $\tilde{s}_{ab}^{l,i}$, the vector field $v_{ab}^{l,i}$, and the connection $\nabla_{ab}^{l,i}$ on $E_{ab}^{l,i}$. We have $(\phi_{ab}^{l,ij})^*(v_{ab}^{l,j}) \equiv v_{ab}^{l,i}$ if $j < i$ in I_{ab}^l by Definition A.32(c). Thus, if the connections $\nabla_{ab}^{l,i}, \nabla_{ab}^{l,j}$ on $E_{ab}^{l,i}, E_{ab}^{l,j}$ are chosen compatible under $(\phi_{ab}^{l,ij}, \hat{\phi}_{ab}^{l,ij})$, which we can do, then $\hat{\phi}_{ab}^{l,ij} \circ \tilde{s}_{ab}^{l,j} \equiv \tilde{s}_{ab}^{l,i} \circ \hat{\phi}_{ab}^{l,ij}$ if $j < i$ in I_{ab}^l . So $(\star)_{ab}^{l,i}(b)$ follows, by (244). This completes the inductive step. Hence by induction we can choose $\tilde{s}_{ab}^{m,i}$ satisfying $(\star)_{ab}^{m,i}$ for all m, a, b .

Now $X_a = \coprod_{b=1}^{n_a^0} X_{ab}^0$, and for each $b = 1, \dots, n_a^0$, if $\phi^0(a, b) = (\bar{a}, \bar{b})$ in R^0 , we have an isomorphism $(a, b)_{ab}^0 : (X_{ab}^0, \mathbf{f}_{ab}^0, \underline{\mathbf{G}}_{ab}^0) \rightarrow (X_{\bar{a}\bar{b}}^0, \mathbf{f}_{\bar{a}\bar{b}}^0, \underline{\mathbf{G}}_{\bar{a}\bar{b}}^0)$, and

piecewise smooth sections $\tilde{s}_{\bar{a}\bar{b}}^{0,i}$ of $E_{\bar{a}\bar{b}}^{0,i} \rightarrow \dot{V}_{\bar{a}\bar{b}}^{0,i}$ for all $i \in I_{\bar{a}\bar{b}}^{0,i}$. Define sections $\tilde{s}_a^i$ of $E_a^i \rightarrow \dot{V}_a^i$ by $\hat{b}_{ab}^{0,i} \circ \tilde{s}_a^i|_{\dot{V}_{ab}^{0,i}} = \tilde{s}_{\bar{a}\bar{b}}^{0,i} \circ b_{ab}^{0,i}$ for $b = 1, \dots, n_a^0$, where $(b_{ab}^{0,i}, \hat{b}_{ab}^{0,i}) : (V_{ab}^{0,i}, \dots, \psi_{ab}^{0,i}) \rightarrow (V_{\bar{a}\bar{b}}^{0,i}, \dots, \psi_{\bar{a}\bar{b}}^{0,i})$ is the coordinate change in $(\mathbf{a}, \mathbf{b})_{ab}^0$. Then $\tilde{s}_a^i$ is continuous, and piecewise smooth subordinate to T_a^i .

For each $c \in C_a$ we have projections $\pi_a : V_{ac}^i \rightarrow \dot{V}_a^i$ for all $i \in I_{ac}$, with $E_{ac}^i = \pi_a^*(E_a^i)$. Define a section $\tilde{s}_{ac}^i$ of $E_{ac}^i \rightarrow V_{ac}^i$ by $\tilde{s}_{ac}^i = \tilde{s}_a^i \circ \pi_a$. Since $\tilde{s}_a^i$ is piecewise smooth subordinate to T_a^i and V_{ac}^i for $i \in I_{ac}$ are the pieces in the decomposition of $\dot{V}_a^i$ induced by T_a^i , it follows that $\tilde{s}_{ac}^i$ is a *smooth* section, not just piecewise smooth. From $(\star)_{\bar{a}\bar{b}}^0(a), (b)$ we see that $\tilde{s}_{ac}^i$ is C^1 close to s_{ac}^i and transverse, and $j < i$ in I_{ac} then $\phi_{ac}^{ij} \circ \tilde{s}_{ac}^j \equiv \tilde{s}_{ac}^i \circ \phi_{ac}^{ij}$ on V_{ac}^{ij} .

We have now constructed perturbations $\tilde{s}_{ac}^i$ of s_{ac}^i for all $a \in A$, $c \in C_a$ and $i \in I_{ac}$ satisfying all the conditions we need. Thus we can apply the construction of $\tilde{X}, \tilde{f}, \tilde{\underline{G}}, Z, \underline{g}, \underline{H}$ from $(X, \mathbf{f}, \underline{G})$ above to $(X_{ac}, \mathbf{f}_{ac}, \underline{G}_{ac})$ for $a \in A$ and $c \in C_a$ using these $\tilde{s}_{ac}^i$. This gives a compact, oriented k -manifold $\tilde{X}_{ac}$, a smooth map $\tilde{f}_{ac} : \tilde{X}_{ac} \rightarrow Y$, a compact oriented Kuranishi space Z_{ac} and a strongly smooth map $\underline{g}_{ac} : Z_{ac} \rightarrow Y$.

To define the effective gauge-fixing data $\tilde{\underline{G}}_{ac}, \underline{H}_{ac}$ for $(\tilde{X}_{ac}, \tilde{f}_{ac}), (Z_{ac}, \underline{g}_{ac})$ we need in addition to choose maps $\tilde{\underline{G}}_{ac}^k : \tilde{X}_{ac} \rightarrow \underline{P}$ and $\underline{H}_{ac}^j : F_{ac}^j \rightarrow \underline{P}$ for $j \in J_{ac}$. To ensure the necessary boundary compatibilities hold, these must also be chosen by reverse induction on codimension by a procedure similar to $(\star)_{ab}^m$ above. In effect, from the piecewise smooth perturbations $\tilde{s}_{ab}^{m,i}$ for $i \in I_{ab}^m$ in $(\star)_{ab}^m$ above, we construct piecewise smooth $\tilde{X}_{ab}^m, \tilde{f}_{ab}^m, Z_{ab}^m, \underline{g}_{ab}^m$ for $m = M, M-1, \dots, 0$ and $(a, b) \in R^m$, and then in an analogue of $(\star)_{ab}^m$ we choose $\tilde{\underline{G}}_{ab}^{m,k-m} : \tilde{X}_{ab}^m \rightarrow \underline{P}$ and $\underline{H}_{ab}^{m,j} : F_{ab}^{m,j} \rightarrow \underline{P}$ for $j \in J_{ab}^m$ for all $m = M, M-1, \dots, 0$, $(a, b) \in R^m$ and $j \in J_{ab}^m$ satisfying boundary conditions similar to $(\star)_{ab}^m(c)$. This is much simpler than the inductive proof above — at each step, the values of $\tilde{\underline{G}}_{ab}^{m,k-m}|_{\partial \tilde{X}_{ab}^m}$ and $\underline{H}_{ab}^{m,j}|_{\partial F_{ab}^{m,j}}$ are prescribed, and we choose arbitrary injective extensions $\tilde{\underline{G}}_{ab}^{m,k-m} : (\tilde{X}_{ab}^m)^\circ \rightarrow \mathbb{R}^n \subset \underline{P}$ and $\underline{H}_{ab}^{m,j} : (F_{ab}^{m,j})^\circ \rightarrow \mathbb{R}^{n'} \subset \underline{P}$ for $n, n' \gg 0$ and all $j \in J_{ab}^m$, satisfying no compatibility conditions at all.

This completes the construction of $(\tilde{X}_{ac}, \tilde{f}_{ac}, \tilde{\underline{G}}_{ac})$ and $(Z_{ac}, \underline{g}_{ac}, \underline{H}_{ac})$ for all $a \in A$ and $c \in C_a$. We claim that the boundary compatibilities we imposed on the data $\tilde{s}_{ac}^i, \tilde{\underline{G}}_{ac}^k, \underline{H}_{ac}^j$ during the inductive construction $(\star)_{ab}^m$ for the $\tilde{s}_{ac}^i$ and the analogues for $\tilde{\underline{G}}_{ac}^k, \underline{H}_{ac}^j$ imply that (241) holds. To see this, note that Z_{ac}^∂ is a perturbation of $[0, 1] \times \partial X_{ac}$. Now the X_{ac} for $c \in C_a$ are the result of cutting X_a into pieces using T_a . Therefore we can split $\partial X_{ac} = (\partial X_{ac})^{\text{int}} \amalg (\partial X_{ac})^{\text{ext}}$, where $(\partial X_{ac})^{\text{int}}$ is the *internal* boundary, the new boundaries creating in the interior of X_a by cutting X_a into pieces, and $(\partial X_{ac})^{\text{ext}}$ is the *external* boundary, part of ∂X_a . Thus there is a corresponding splitting $Z_{ac}^\partial = (Z_{ac}^\partial)^{\text{int}} \amalg (Z_{ac}^\partial)^{\text{ext}}$.

Each component of $[(\partial X_{ac})^{\text{int}}, \mathbf{f}_{ac}|_{(\partial X_{ac})^{\text{int}}}, \underline{G}_{ac}|_{(\partial X_{ac})^{\text{int}}}]$ is cancelled by a component of $[(\partial X_{ac'})^{\text{int}}, \mathbf{f}_{ac'}|_{(\partial X_{ac'})^{\text{int}}}, \underline{G}_{ac'}|_{(\partial X_{ac'})^{\text{int}}}]$ for some other $c' \in C_a$, the ‘other side of the cut’. Therefore for each $a \in A$ we have $\sum_{c \in C_a} [(\partial X_{ac})^{\text{int}}, \mathbf{f}_{ac}|_{(\partial X_{ac})^{\text{int}}}, \underline{G}_{ac}|_{(\partial X_{ac})^{\text{int}}}] = 0$. Because the perturbations $\tilde{s}_a^i$ for $i \in I_a$ were chosen *continuous* and piecewise smooth, and the $\tilde{\underline{G}}_{ac}^k, \underline{H}_{ac}^j$ are also restrictions

of functions independent of $c \in C_a$, the construction of $Z_{ac}^\partial, \mathbf{g}_{ac}|_{Z_{ac}^\partial}, \underline{\mathbf{H}}_{ac}|_{Z_{ac}^\partial}$ over $[0, 1] \times \partial X_{ac}$ is compatible with this cancellation of internal boundary components. Hence for each $a \in A$ we have

$$\sum_{c \in C_a} [(Z_{ac}^\partial)^{\text{int}}, \mathbf{g}_{ac}|_{(Z_{ac}^\partial)^{\text{int}}}, \underline{\mathbf{H}}_{ac}|_{(Z_{ac}^\partial)^{\text{int}}}] = 0. \quad (245)$$

For the external boundaries, using the argument of (229) and (230) we have

$$\begin{aligned} & \sum_{a \in A} \rho_a [(Z_{ac}^\partial)^{\text{ext}}, \mathbf{g}_{ac}|_{(Z_{ac}^\partial)^{\text{ext}}}, \underline{\mathbf{H}}_{ac}|_{(Z_{ac}^\partial)^{\text{ext}}}] = \\ & \sum_{(\bar{a}, \bar{b}) \in R^1} \left\{ \sum_{\substack{a \in A, b=1, \dots, n_a^1: \\ \phi^1(a, b) = (\bar{a}, \bar{b})}} \rho_a \epsilon_{ab}^1 \right\} \sum_{\bar{c} \in C_{\bar{a}\bar{b}}^1} [Z_{\bar{a}\bar{b}\bar{c}}^1, \mathbf{g}_{\bar{a}\bar{b}\bar{c}}^1, \underline{\mathbf{H}}_{\bar{a}\bar{b}\bar{c}}^1] = 0, \end{aligned} \quad (246)$$

where $C_{\bar{a}\bar{b}}^1$ is an indexing set for the decomposition induced by the tent function $\mathbf{T}_{\bar{a}\bar{b}}^1$ for $(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \underline{\mathbf{G}}_{\bar{a}\bar{b}}^1)$ chosen in $(*)_{\bar{a}\bar{b}}^1$ in §B.1, and $(Z_{\bar{a}\bar{b}\bar{c}}^1, \mathbf{g}_{\bar{a}\bar{b}\bar{c}}^1, \underline{\mathbf{H}}_{\bar{a}\bar{b}\bar{c}}^1)$ for $C_{\bar{a}\bar{b}}^1$ are the corresponding chains constructed above using the piecewise smooth sections $\tilde{s}_{\bar{a}\bar{b}}^1$ in $(*)_{\bar{a}\bar{b}}^1$. The coefficient $\{\dots\}$ on the second line of (246) is zero as in the proof of (230) in §B.1. Combining (245), (246) and $Z_{ac}^\partial = (Z_{ac}^\partial)^{\text{int}} \amalg (Z_{ac}^\partial)^{\text{ext}}$ proves (241), and completes Step 2.

B.3 Step 3: triangulating the $\tilde{X}_a$ with simplices $\sigma_{ac}(\Delta_k)$

As in Step 3 above we now *change notation* from $\tilde{X}_{ac}$ for $a \in A$ and $c \in C_a$ to $\tilde{X}_a$ for $a \in A$. So, we represent α by a cycle $\sum_{a \in A} \rho_a [\tilde{X}_a, \tilde{f}_a, \tilde{\mathbf{G}}_a] \in KC_k^{\text{ef}}(Y; R)$, with each $\tilde{X}_a$ a compact, oriented k -manifold with g-corners, $\tilde{f}_a : \tilde{X}_a \rightarrow Y$ a smooth map, and $\tilde{\mathbf{G}}_a = (\tilde{\mathbf{I}}_a, \tilde{\eta}_a, \tilde{\mathbf{G}}_a^k)$ effective gauge-fixing data for $(\tilde{X}_a, \tilde{f}_a)$, where $\tilde{\mathbf{I}}_a$ has indexing set $\tilde{I}_a = \{k\}$, one Kuranishi neighbourhood $(\tilde{V}_a^k, \tilde{E}_a^k, \tilde{s}_a^k, \tilde{\psi}_a^k) = (\tilde{X}_a, \tilde{X}_a, 0, \text{id}_{\tilde{X}_a})$, and $\tilde{\eta}_a = (\tilde{\eta}_{a,k}, \tilde{\eta}_{a,k}^k)$ with $\tilde{\eta}_{a,k} \equiv 1 \equiv \tilde{\eta}_{a,k}^k$.

We will use Theorem A.18 to choose a tent function $\tilde{T}_a : \tilde{X}_a \rightarrow F([1, \infty))$ for each $a \in A$ such that the components $\tilde{X}_{ac}$, $c \in C_a$ of $\partial Z_{\tilde{X}_a, \tilde{T}_a}$ in (187) are all diffeomorphic to the k -simplex Δ_k , with chosen diffeomorphisms $\sigma_{ac} : \Delta_k \rightarrow \tilde{X}_{ac}$ for $c \in C_a$. (Note that these C_a and $\tilde{X}_{ac}$ are different to those in Steps 1 and 2.) To make these choices of $\tilde{T}_a, \sigma_{ac}$ compatible with the boundary relations in $\partial(\sum_{a \in A} \rho_a [\tilde{X}_a, \tilde{f}_a, \tilde{\mathbf{G}}_a]) = 0$, we use reverse induction on codimension in a similar way to Steps 1 and 2.

Define notation $n_a^m, (\tilde{X}_{ab}^m, \tilde{f}_{ab}^m, \tilde{\mathbf{G}}_{ab}^m), P^m, Q^m, R^m, (a^q, b^q), \phi^m, (\mathbf{a}, \mathbf{b})_{ab}^m, \epsilon_{ab}^m, B_{ab}^m$ as in Step 1 above, but using $(\tilde{X}_a, \tilde{f}_a, \tilde{\mathbf{G}}_a)$ for $a \in A$ rather than $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ for $a \in A$, so that $(\partial^m \tilde{X}_a, \tilde{f}_a|_{\partial^m \tilde{X}_a}, \tilde{\mathbf{G}}_a|_{\partial^m \tilde{X}_a}) = \coprod_{b=1, \dots, n_a^m} (\tilde{X}_{ab}^m, \tilde{f}_{ab}^m, \tilde{\mathbf{G}}_{ab}^m)$, and so on. As $\tilde{X}_{ab}^m$ is a manifold of dimension $k-m$ we have $\tilde{X}_{ab}^m = \emptyset$ for $m > k$, so we fix $M = k$. By induction on decreasing $m = k, k-1, \dots, 1, 0$, for all (a, b) in R^m , we choose tent functions $\tilde{T}_{ab}^m$ on $\tilde{X}_{ab}^m$ using Theorem A.18, such that the expression (187) for $\partial Z_{\tilde{X}_{ab}^m, \tilde{T}_{ab}^m}$ involves components $\tilde{X}_{abc}^m$ for $c \in C_{ab}^m$, with C_{ab}^m a finite indexing set, and for all (a, b) in R^m and $c \in C_{ab}^m$ we choose diffeomorphisms $\sigma_{abc}^m : \Delta_{k-m} \rightarrow \tilde{X}_{abc}^m$, satisfying the inductive hypothesis:

(†)_{ab}^m By definition of B_{ab}^m we have $\partial\tilde{X}_{ab}^m = \coprod_{b' \in B_{ab}^m} \tilde{X}_{ab'}^{m+1}$, and if $\phi^{m+1}(a, b') = (\bar{a}, \bar{b})$ in R^{m+1} , we have an isomorphism $(\mathbf{a}, \mathbf{b})_{ab'}^{m+1} : (\tilde{X}_{ab'}^{m+1}, \tilde{f}_{ab'}^{m+1}, \tilde{\mathbf{g}}_{ab'}^{m+1}) \rightarrow (\tilde{X}_{\bar{a}\bar{b}}^{m+1}, \tilde{f}_{\bar{a}\bar{b}}^{m+1}, \tilde{\mathbf{g}}_{\bar{a}\bar{b}}^{m+1})$. So as for (227) there is a diffeomorphism

$$\coprod_{b' \in B_{ab}^m} \mathbf{a}_{ab'}^{m+1} : \partial\tilde{X}_{ab}^m \longrightarrow \coprod_{b' \in B_{ab}^m} \text{define } (\bar{a}, \bar{b}) = \phi^{m+1}(a, b') \tilde{X}_{\bar{a}\bar{b}}^{m+1}. \quad (247)$$

On the left hand side $\partial\tilde{X}_{ab}^m$ of (247) we have a tent function $\tilde{T}_{ab}^m|_{\partial\tilde{X}_{ab}^m}$. On the right hand side we have a tent function $\coprod_{b'} \tilde{T}_{\bar{a}\bar{b}}^{m+1}$. We require:

- (a) Equation (247) must identify $\min(\tilde{T}_{ab}^m|_{\partial\tilde{X}_{ab}^m})$ with $\min(\coprod_{b'} \tilde{T}_{\bar{a}\bar{b}}^{m+1})$.
- (b) Suppose the natural involution $\sigma : \partial^2 Z_{\tilde{X}_{ab}^m, \tilde{T}_{ab}^m} \rightarrow \partial^2 Z_{\tilde{X}_{ab}^m, \tilde{T}_{ab}^m}$ exchanges two $(k-m-1)$ -simplices $\sigma_{abc}^m \circ F_j^{k-m}(\Delta_{k-m-1})$ and $\sigma_{abc'}^m \circ F_j^{k-m}(\Delta_{k-m-1})$ for $c, c' \in C_{ab}^m$. Then $j = j'$, $c \neq c'$ and $\sigma \circ \sigma_{abc}^m \circ F_j^{k-m} \equiv \sigma_{abc'}^m \circ F_j^{k-m}$ as maps $\Delta_{k-m-1} \rightarrow \partial^2 Z_{\tilde{X}_{ab}^m, \tilde{T}_{ab}^m}$.
- (c) If $b' \in B_{ab}^m$, $(\bar{a}, \bar{b}) = \phi^{m+1}(a, b')$ and $\bar{c} \in C_{\bar{a}\bar{b}}^{m+1}$ then $\tilde{X}_{\bar{a}\bar{b}\bar{c}}^{m+1} = \sigma_{\bar{a}\bar{b}\bar{c}}^{m+1}(\Delta_{k-m-1})$ is a component of $\partial Z_{\tilde{X}_{\bar{a}\bar{b}}^{m+1}, \tilde{T}_{\bar{a}\bar{b}}^{m+1}}$. Applying the projection $\pi : \partial Z_{\tilde{X}_{\bar{a}\bar{b}}^{m+1}, \tilde{T}_{\bar{a}\bar{b}}^{m+1}} \rightarrow \tilde{X}_{\bar{a}\bar{b}}^{m+1}$ gives an embedding $\pi \circ \sigma_{\bar{a}\bar{b}\bar{c}}^{m+1} : \Delta_{k-m-1} \rightarrow \tilde{X}_{\bar{a}\bar{b}}^{m+1}$, so $((\mathbf{a}, \mathbf{b})_{ab'}^{m+1})^{-1} \circ \pi \circ \sigma_{\bar{a}\bar{b}\bar{c}}^{m+1}$ is an embedding $\Delta_{k-m-1} \rightarrow \tilde{X}_{ab'}^{m+1} \subset \partial\tilde{X}_{ab}^m$. Composing with $\iota : \partial\tilde{X}_{ab}^m \rightarrow \tilde{X}_{ab}^m$ yields an embedding $\iota \circ ((\mathbf{a}, \mathbf{b})_{ab'}^{m+1})^{-1} \circ \pi \circ \sigma_{\bar{a}\bar{b}\bar{c}}^{m+1} : \Delta_{k-m-1} \rightarrow \tilde{X}_{ab}^m$. There should exist some unique $c \in C_{ab}^m$ such that $\pi \circ \sigma_{abc}^m \circ F_j^{k-m} \equiv \iota \circ ((\mathbf{a}, \mathbf{b})_{ab'}^{m+1})^{-1} \circ \pi \circ \sigma_{\bar{a}\bar{b}\bar{c}}^{m+1}$ as embeddings $\Delta_{k-m-1} \rightarrow \tilde{X}_{ab}^m$.

Here part (a) is almost the same as $(*)_{ab}^m$. Part (b) gives compatibility between the ‘internal’ codimension one faces of $\sigma_{abc}^m(\Delta_{k-m})$ within a fixed X_{ab}^m , and corresponds to the boundary compatibility between the σ_c in Theorem A.7. Part (c) gives compatibility of the ‘external’ codimension one faces of $\sigma_{abc}^m(\Delta_{k-m})$ between two different $X_{ab}^m, X_{a'b'}^m$, and corresponds to the boundary compatibility between $\sigma_{c'}, \sigma_c$ in Theorem A.18.

The first step, when $m = k$, is trivial. Each $\tilde{X}_{ab}^k$ is a compact, connected 0-manifold, that is, a single point. We define $\tilde{T}_{ab}^k \equiv \{1\}$, $C_{ab}^k = \{1\}$ for all $(a, b) \in R^k$, and take $\sigma_{ab1} : \Delta_0 \rightarrow \tilde{X}_{ab1}^k = \{1\} \times \tilde{X}_{ab}^k$ to be the unique diffeomorphism between the two points. Parts (a)–(c) are vacuous as $\partial\tilde{X}_{ab}^k = \emptyset$.

For the inductive step, suppose that for some $l = 0, \dots, k-1$ we have chosen $\tilde{T}_{ab}^l, C_{ab}^l$ and σ_{abc}^l satisfying $(\dagger)_{ab}^m$ for all $l < m \leq k$, $(a, b) \in R^m$ and $c \in C_{ab}^m$. For each $(a, b) \in R^l$, we will apply Theorem A.18 to choose $\tilde{T}_{ab}^l, C_{ab}^l$ and σ_{abc}^l for $c \in C_{ab}^l$ satisfying $(\dagger)_{ab}^l$. We have a tent function $T' = \coprod_{b' \in B_{ab}^m} \tilde{T}_{\bar{a}\bar{b}}^{m+1} \circ \mathbf{a}_{ab'}^{m+1}$ on $\partial\tilde{T}_{ab}^m$, and part (a) of $(\dagger)_{ab}^m$ says that $\min \tilde{T}_{ab}^m|_{\partial X_{ab}^m} \equiv \min T'$, which corresponds to $\min T|_{\partial X} \equiv \min T'$ in Theorem A.18.

We must verify the hypotheses Theorem A.18(i)–(iii). Parts (i), (ii) hold by the argument showing that $\min \mathbf{T}_{ab}^l|_{\partial^2 X_{ab}^l}$ is invariant under the natural involution $\sigma : \partial^2 X_{ab}^l \rightarrow \partial^2 X_{ab}^l$ in the proof of the inductive step of $(*)_{ab}^m$ in §B.1.

Part (iii) is a little more subtle. To satisfy it, before starting the induction, we first choose Riemannian metrics g_{ab}^m on $\tilde{X}_{ab}^m$ for all $m = 0, \dots, k$ and $(a, b) \in R^m$, and then we choose $\epsilon > 0$ very small compared to length scales in $(\tilde{X}_{ab}^m, g_{ab}^m)$ and constants comparing $g_{ab}^m|_{\tilde{X}_{ab'}^{m+1}}$ and $g_{\bar{a}\bar{b}}^{m+1}$ under the inclusion $\tilde{X}_{ab'}^{m+1} \subseteq \partial\tilde{X}_{ab}^m$ and diffeomorphism $\mathbf{a}_{ab'}^{m+1} : \tilde{X}_{ab'}^{m+1} \rightarrow \tilde{X}_{\bar{a}\bar{b}}^{m+1}$, for all $m, (a, b), b', (\bar{a}, \bar{b})$. Then we use these g_{ab}^m and ϵ to construct $\tilde{T}_{ab}^l$ in Theorem A.18.

The point is that as the ϵ used to construct T' and $\sigma_{c'}$ on $\partial\tilde{X}_{ab}^m$ has been chosen small compared not just to geometry on $\partial\tilde{X}_{ab}^m$, but also to geometry on $\tilde{X}_{ab}^m$ and constants comparing the two, Theorem A.18(iii) holds. So Theorem A.18 gives $\tilde{T}_{ab}^m, C_{ab}^m$ and σ_{abc}^m for $c \in C_{ab}^m$. Part (a) of $(\dagger)_{ab}^m$ follows from $\min T|_{\partial X} \equiv \min T'$ in Theorem A.18, part (b) from the boundary compatibility between the σ_c in Theorem A.7 referred to in Theorem A.18, and (c) from the compatibility between the $\sigma_{c'}$ and σ_c in Theorem A.18. This proves the inductive step. Hence by induction $(\dagger)_{ab}^m$ holds for all m, a, b .

Now $\tilde{X}_a = \coprod_{b=1}^{n_a^0} \tilde{X}_{ab}^0$, and if $b = 1, \dots, n_a^0$ and $\phi^0(a, b) = (\bar{a}, \bar{b})$ in R^0 , we have a diffeomorphism $\mathbf{a}_{ab}^0 : \tilde{X}_{ab}^0 \rightarrow \tilde{X}_{\bar{a}\bar{b}}^0$ and a tent function $\tilde{T}_{ab}^0$ on $\tilde{X}_{ab}^0$. Define a tent function $\tilde{T}_a$ on $\tilde{X}_a$ by $\tilde{T}_a|_{\tilde{X}_{ab}^0} = \tilde{T}_{ab}^0 \circ \mathbf{a}_{ab}^0$ for $b = 1, \dots, n_a^0$, for each $a \in A$. Then we may form a compact $(k+1)$ -manifold $Z_{\tilde{X}_a, \tilde{T}_a}$ as in (186). Let $\tilde{X}_{ac}$ for $c \in C_a$ be the extra pieces in the decomposition of $\partial Z_{\tilde{X}_a, \tilde{T}_a}$ in (187). Then from above we may take $C_a = \coprod_{b=1}^{n_a^0} C_{\bar{a}\bar{b}}^0$, where $(\bar{a}, \bar{b}) = \phi^0(a, b)$, and the diffeomorphisms $\sigma_{\bar{a}\bar{b}\bar{c}}^0 : \Delta_k \rightarrow \tilde{X}_{\bar{a}\bar{b}\bar{c}}^0$ for $b = 1, \dots, n_a^0$, $(\bar{a}, \bar{b}) = \phi^0(a, b)$ and $\bar{c} \in C_{\bar{a}\bar{b}}^0$ induce diffeomorphisms $\sigma_{ac} : \Delta_k \rightarrow \tilde{X}_{ac}$ for all $c \in C_a$.

Write $\pi : Z_{\tilde{X}_a, \tilde{T}_a} \rightarrow \tilde{X}_a$ for the natural projection. We will define effective gauge-fixing data $\tilde{\mathbf{H}}_a = (\tilde{\mathbf{J}}_a, \tilde{\zeta}_a, \tilde{\mathbf{H}}_a^{k+1})$ for $(Z_{\tilde{X}_a, \tilde{T}_a}, \tilde{f}_a \circ \pi)$. Let $\tilde{\mathbf{J}}_a$ have indexing set $\tilde{J}_a = \{k+1\}$, one Kuranishi neighbourhood $(\tilde{W}_a^{k+1}, \tilde{F}_a^{k+1}, \tilde{t}_a^{k+1}, \tilde{\chi}_a^{k+1}) = (Z_{\tilde{X}_a, \tilde{T}_a}, Z_{\tilde{X}_a, \tilde{T}_a}, 0, \text{id}_{Z_{\tilde{X}_a, \tilde{T}_a}})$, map $\tilde{f}_a \circ \pi : \tilde{W}_a^{k+1} \rightarrow Y$ representing $\tilde{f}_a \circ \pi$, and let $\tilde{\zeta}_a = (\tilde{\zeta}_{a, k+1}, \tilde{\zeta}_{a, k+1}^{k+1})$ with $\tilde{\zeta}_{a, k+1} \equiv 1 \equiv \tilde{\zeta}_{a, k+1}^{k+1}$. Then $(\tilde{\mathbf{J}}_a, \tilde{\zeta}_a)$ is an excellent coordinate system for $(Z_{\tilde{X}_a, \tilde{T}_a}, \tilde{f}_a \circ \pi)$. Define $\tilde{\mathbf{H}}_a^{k+1} : Z_{\tilde{X}_a, \tilde{T}_a} \rightarrow \underline{P}$ by

$$\begin{aligned} \tilde{\mathbf{H}}_a^{k+1}(u, x) = & \\ \begin{cases} \tilde{\mathbf{G}}_a^k(x), & u = 0, \\ (u, x_1, \dots, x_n), & 0 < u < \min \tilde{T}_a(x), \tilde{\mathbf{G}}_a^k(x) = (x_1, \dots, x_n), \\ \mathbf{G}_{\Delta_k}^k((y_0, \dots, y_k)), & u = \min \tilde{T}_a(x), (u, x) = \iota \circ \sigma_{ac}(y_0, \dots, y_k), \\ & c \in C_a, (y_0, \dots, y_k) \in \Delta_k, \end{cases} \end{aligned} \quad (248)$$

where points of $Z_{\tilde{X}_a, \tilde{T}_a}$ are (u, x) for $x \in \tilde{X}_a$ and $0 \leq u \leq \min \tilde{T}_a(x)$, and $\mathbf{G}_{\Delta_k}^k$ is as in (70), and $\iota : \partial Z_{\tilde{X}_a, \tilde{T}_a} \rightarrow Z_{\tilde{X}_a, \tilde{T}_a}$ is the natural immersion, restricted to $\tilde{X}_{ac} \subset \partial Z_{\tilde{X}_a, \tilde{T}_a}$.

Note that if $(u, x) \in Z_{\tilde{X}_a, \tilde{T}_a}$ with $u = \min \tilde{T}_a(x)$ then $(u, x) \in \iota(\tilde{X}_{ac})$ for some $c \in C_a$, so that $(u, x) = \iota \circ \sigma_{ac}(y_0, \dots, y_k)$ for some $(y_0, \dots, y_k) \in \Delta_k$ as $\sigma_{ac} : \Delta_k \rightarrow \tilde{X}_{ac}$ is a diffeomorphism, and the third line of (248) makes sense.

However, $c \in C_a$ may not be unique, and we may also have $(u, x) \in \iota(\tilde{X}_{ac'})$ for some other $c' \in C_a$, with $(u, x) = \iota \circ \sigma_{ac'}(y'_0, \dots, y'_k)$ for $(y'_0, \dots, y'_k) \in \Delta_k$. Thus, to show (248) is well-defined we must prove that $\underline{G}_{\Delta_k}^k((y_0, \dots, y_k)) = \underline{G}_{\Delta_k}^k((y'_0, \dots, y'_k))$. Let $(z_1, \dots, z_l)$ and $(z'_1, \dots, z'_{l'})$ be the result of deleting all zero entries in $(y_0, \dots, y_k)$ and $(y'_0, \dots, y'_k)$ respectively. Then using $(\dagger)_{ab}^m(b)$ one can show that $l = l'$ and $(z_1, \dots, z_l) = (z'_1, \dots, z'_{l'})$, and (70) then implies that $\underline{G}_{\Delta_k}^k((y_0, \dots, y_k)) = \underline{G}_{\Delta_k}^k((y'_0, \dots, y'_k))$.

Hence $\tilde{\underline{H}}_a^{k+1}$ is well-defined, and it is easy to check that $\tilde{\underline{H}}_a = (\tilde{\underline{J}}_a, \tilde{\underline{\zeta}}_a, \tilde{\underline{H}}_a^{k+1})$ is *effective gauge-fixing data* for $(Z_{\tilde{X}_a, \tilde{T}_a}, \tilde{f}_a \circ \pi)$. As in (212) we see that

$$\begin{aligned} \partial[Z_{\tilde{X}_a, \tilde{T}_a}, \tilde{f}_a \circ \pi, \tilde{\underline{H}}_a] &= \sum_{c \in C_a} [\tilde{X}_{ac}, \tilde{f}_a \circ \pi|_{\tilde{X}_{ac}}, \tilde{\underline{H}}_a|_{\tilde{X}_{ac}}] \\ &\quad - [\tilde{X}_a, \tilde{f}_a, \tilde{\underline{G}}_a] - [Z_{\partial\tilde{X}_a, \tilde{T}_a|_{\partial\tilde{X}_a}}, \tilde{f}_a|_{\partial\tilde{X}_a} \circ \pi, \tilde{\underline{H}}_a|_{Z_{\partial\tilde{X}_a, \tilde{T}_a|_{\partial\tilde{X}_a}}}] \end{aligned} \quad (249)$$

For $a \in A$ and $c \in C_a$, define $\epsilon_{ac} = 1$ if the diffeomorphism $\sigma_{ac} : \Delta_k \rightarrow \tilde{X}_{ac}$ is orientation-preserving, and $\epsilon_{ac} = -1$ otherwise. Then from (73) and the last line of (248) we see that

$$\begin{aligned} [\tilde{X}_{ac}, \tilde{f}_a \circ \pi|_{\tilde{X}_{ac}}, \tilde{\underline{H}}_a|_{\tilde{X}_{ac}}] &= \epsilon_{ac} [\Delta_k, \tilde{f}_a \circ \pi \circ \sigma_{ac}, \underline{f}_{ac}, \underline{G}_{\Delta_k}] \\ &= \epsilon_{ac} \Pi_{\text{si}}^{\text{ef}}(\tilde{f}_a \circ \pi \circ \sigma_{ac}). \end{aligned} \quad (250)$$

A similar proof to (229)–(230) and (246) shows that

$$\sum_{a \in A} \rho_a [Z_{\partial\tilde{X}_a, \tilde{T}_a|_{\partial\tilde{X}_a}}, \tilde{f}_a|_{\partial\tilde{X}_a} \circ \pi, \tilde{\underline{H}}_a|_{Z_{\partial\tilde{X}_a, \tilde{T}_a|_{\partial\tilde{X}_a}}}] = 0. \quad (251)$$

Multiplying (249) by ρ_a , summing over $a \in A$, and using (250) and (251) yields

$$\begin{aligned} &\partial\left(\sum_{a \in A} \rho_a [Z_{\tilde{X}_a, \tilde{T}_a}, \tilde{f}_a \circ \pi, \tilde{\underline{H}}_a]\right) \\ &= \Pi_{\text{si}}^{\text{ef}}\left(\sum_{a \in A} \sum_{c \in C_a} (\rho_a \epsilon_{ac}) (\tilde{f}_a \circ \pi \circ \sigma_{ac})\right) - \sum_{a \in A} \rho_a [\tilde{X}_a, \tilde{f}_a, \tilde{\underline{G}}_a]. \end{aligned} \quad (252)$$

Thus $\Pi_{\text{si}}^{\text{ef}}(\sum_{a \in A} \sum_{c \in C_a} (\rho_a \epsilon_{ac}) (\tilde{f}_a \circ \pi \circ \sigma_{ac}))$ is homologous to $\sum_{a \in A} \rho_a [\tilde{X}_a, \tilde{f}_a, \tilde{\underline{G}}_a]$, and is a cycle representing α .

Now Definition 4.3(i)–(iii) impose no nontrivial relations upon chains of the form $[\Delta_k, \sigma, \underline{G}_{\Delta_k}]$ for smooth $\sigma : \Delta_k \rightarrow Y$, and thus $\Pi_{\text{si}}^{\text{ef}}$ in (73) is injective at the chain level. As $\Pi_{\text{si}}^{\text{ef}} \circ \partial = \partial \circ \Pi_{\text{si}}^{\text{ef}}$ and $\Pi_{\text{si}}^{\text{ef}}(\sum_{a \in A} \sum_{c \in C_a} (\rho_a \epsilon_{ac}) (\tilde{f}_a \circ \pi \circ \sigma_{ac}))$ is a cycle in $KC_k^{\text{ef}}(Y; R)$, (220) holds, and $\beta = [\sum_{a \in A} \sum_{c \in C_a} (\rho_a \epsilon_{ac}) (\tilde{f}_a \circ \pi \circ \sigma_{ac})]$ is well-defined in $H_k^{\text{si}}(Y; R)$, with $\Pi_{\text{si}}^{\text{ef}}(\beta) = \alpha$. This completes Step 3.

B.4 Step 4: $\Pi_{\text{si}}^{\text{ef}} : H_k^{\text{si}}(Y; R) \rightarrow KH_k^{\text{ef}}(Y; R)$ is injective

Let β in $H_k^{\text{si}}(Y; R)$ with $\Pi_{\text{si}}^{\text{ef}}(\beta) = 0$, the singular k -chain $\sum_{d \in D} \eta_d \tau_d$, and $\sum_{a \in A} \rho_a [X_a, \underline{f}_a, \underline{G}_a]$ in $KC_{k+1}^{\text{ef}}(Y; R)$ be as in Step 4, so that (221) holds. We now explain how to apply Steps 1–3 to this $\sum_{a \in A} \rho_a [X_a, \underline{f}_a, \underline{G}_a]$, replacing k -chains by $(k+1)$ -chains, to construct a chain in $C_{k+1}^{\text{si}}(Y; R)$ with boundary

$\sum_{d \in D} \eta_d \tau_d$, proving that $\beta = 0$. Write $\partial X_a = X_{a1}^1 \amalg \cdots \amalg X_{an_a^1}^1$ for the decomposition of $[\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a}]$ into connected pieces given by Lemma 3.7.

From (221) and the proof that $\Pi_{\text{si}}^{\text{ef}} : C_k^{\text{si}}(Y; R) \rightarrow KC_k^{\text{ef}}(Y; R)$ is injective, we see that we can divide the pieces X_{ab}^1 for $a \in A$ and $b = 1, \dots, n_a^1$ into two kinds: (i) those with $[X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1] = \pm[\Delta_k, \tau_d, \mathbf{G}_{\Delta_k}]$ for some $d \in D$, and (ii) the rest. Moreover, the sum of $\rho_a[X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1]$ over all (a, b) of type (i) is $\sum_{d \in D} \eta_d [\Delta_k, \tau_d, \mathbf{G}_{\Delta_k}]$, and the sum over all (a, b) of type (ii) is zero. Let us rewrite the chain $\sum_{d \in D} \eta_d \tau_d$ if necessary, allowing repeats among the τ_d , so that there is a 1-1 correspondence between $d \in D$ and those (a, b) of type (i), so that if d corresponds to (a, b) then $[X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1] = \pm[\Delta_k, \tau_d, \mathbf{G}_{\Delta_k}]$ and $\rho_a = \pm \eta_d$. Clearly, this is possible.

If X_{ab}^1 is a boundary component of type (i) then as X_{ab}^1 is diffeomorphic to the smooth k -manifold Δ_k , it follows that X_a is near X_{ab}^1 a smooth $(k+1)$ -manifold. So in Step 1 there is no need to cut X_a into pieces near X_{ab}^1 to get trivial stabilizers, and in Step 2 there is no need to deform X_a near X_{ab}^1 to make it a manifold; we can make choices in Steps 1 and 2 so that the type (i) boundary components X_{ab}^1 are not changed at all.

In Step 1 we do this by choosing the tent functions $\mathbf{T}_a$ for $a \in A$ so that $\min \mathbf{T}_a|_{X_{ab}^1} \equiv 1$ for each type (i) boundary component X_{ab}^1 . That is, in $(*)_{ab}^m$ we impose the extra condition that $T_{ab}^{m, k-m} \equiv \{1\}$ for any triple $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ which is isomorphic to a component of $\partial^{m-1}(X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1)$ for some type (i) boundary component X_{ab}^1 and all $m = k, k-1, \dots, 1$. Then at the end of Step 1, to each $d \in D$ there corresponds a unique type (i) boundary component $X_{ab}^1 \cong \Delta_k$ with $[X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1] = \pm[\Delta_k, \tau_d, \mathbf{G}_{\Delta_k}]$, and X_{ab}^1 is naturally diffeomorphic to a unique component of ∂X_{ac} for some unique $c \in C_a$.

In Step 2, on X_{ac} near a component of ∂X_{ac} naturally diffeomorphic to a type (i) boundary component X_{ab}^1 , since the excellent coordinate system $(\mathbf{I}_{ab}^1, \boldsymbol{\eta}_{ab}^1)$ for $(X_{ab}^1, \mathbf{f}_{ab}^1)$ in $\mathbf{G}_{ab}^1$ has indexing set $I_{ab}^1 = \{k\}$ and Kuranishi neighbourhood $(V_{ab}^{1,k}, E_{ab}^{1,k}, s_{ab}^{1,k}, \psi_{ab}^{1,k}) = (X_{ab}^1, X_{ab}^1, 0, \text{id}_{X_{ab}^1})$, the excellent coordinate system $(\mathbf{I}_{ac}, \boldsymbol{\eta}_{ac})$ for $(X_{ac}, \mathbf{f}_{ac})$ in $\mathbf{G}_{ac}$ includes a Kuranishi neighbourhood $(V_{ac}^{k+1}, E_{ac}^{k+1}, s_{ac}^{k+1}, \psi_{ac}^{k+1})$ covering the component of ∂X_{ac} diffeomorphic to X_{ab}^1 .

But $E_{ac}^{k+1} \equiv 0$, so $\tilde{s}_{ac}^{k+1} \equiv 0$, the $\tilde{s}_{ac}^i$ for $i \in I_{ac}$ are zero near this component of ∂X_{ac} . Thus $\tilde{X}_{ac}$ coincides with X_{ac} near this component of ∂X_{ac} . We can also choose the map $\tilde{\mathbf{G}}_{ac}^{k+1} : \tilde{X}_{ac} \rightarrow \underline{P}$ to agree with $\mathbf{G}_{ab}^1 : X_{ab}^1 \rightarrow \underline{P}$, or equivalently with $\mathbf{G}_{\Delta_k}^k : \Delta_k \rightarrow \underline{P}$, on the component of $\partial \tilde{X}_{ac}$ diffeomorphic to $X_{ab}^1 \cong \Delta_k$. Therefore at the end of Step 2, to each $d \in D$ there corresponds a type (i) boundary component $X_{ab}^1 \cong \Delta_k$ which is naturally diffeomorphic to a unique component of $\partial \tilde{X}_{ac}$ for unique $c \in C_a$.

As in Step 3, we change notation from $\tilde{X}_{ac}$ to $\tilde{X}_a$. Thus, at the beginning of Step 3 we have a chain $\sum_{a \in A} \rho_a [\tilde{X}_a, \tilde{\mathbf{f}}_a, \tilde{\mathbf{G}}_a]$ in $KC_{k+1}^{\text{ef}}(Y; R)$ with

$$\partial(\sum_{a \in A} \rho_a [\tilde{X}_a, \tilde{\mathbf{f}}_a, \tilde{\mathbf{G}}_a]) = \sum_{d \in D} \eta_d [\Delta_k, \tau_d, \mathbf{G}_{\Delta_k}], \quad (253)$$

where for each $d \in D$ we are given $a \in A$ and a component $\tilde{X}_{ab}^1$ of $\partial(\tilde{X}_a, \tilde{\mathbf{f}}_a, \tilde{\mathbf{G}}_a)$ with a diffeomorphism $\tilde{X}_{ab}^1 \cong \Delta_k$ such that $[\tilde{X}_{ab}^1, \tilde{\mathbf{f}}_{ab}^1, \tilde{\mathbf{G}}_{ab}^1] = \pm[\Delta_k, \tau_d, \mathbf{G}_{\Delta_k}]$

and $\eta_d = \pm \rho_a$. Step 3 then chooses tent functions $\tilde{T}_a$ on $\tilde{X}_a$ for $a \in A$ and diffeomorphisms $\sigma_{ac} : \Delta_{k+1} \rightarrow \tilde{X}_{ac}$. If $d \in D$ and $\tilde{X}_{ab}^1 \cong \Delta_k$ is the given diffeomorphism, then $\tilde{T}_a|_{\tilde{X}_{ab}^1}$ is a tent function on Δ_k , which cuts Δ_k up into pieces, each of which has a diffeomorphism with Δ_k of the form $\pi \circ \sigma_{ac} \circ F_{k+1}^{k+1} : \Delta_k \rightarrow \tilde{X}_a$ for some $c \in C_a$.

Unlike in Steps 1 and 2, we cannot choose the $\tilde{T}_a$ in Step 3 so that the type (i) boundary components $\tilde{X}_{ab}^1$, that is, the chains $[\Delta_k, \tau_d, \underline{\mathbf{G}}_{\Delta_k}]$ for $d \in D$, are unchanged, because to apply the results of §A.1.6 during Step 3 we need to cut the $\tilde{X}_a$ into *small pieces*, which includes the boundary components $\tilde{X}_{ab}^1 \cong \Delta_k$. This is necessary in the proof of the inductive step in $(\dagger)_{ab}^m$, when we have to ensure that the smallness condition Theorem A.18(iii) holds, and means that we must cut the $\tilde{X}_{ab}^1 \cong \Delta_k$ into pieces small compared to the length scales involved in $\tilde{X}_{ab}^m$ for all $m = k, k-1, \dots, 0$ and $(a, b) \in R^m$.

However, we need not cut the boundary components $\tilde{X}_{ab}^1 \cong \Delta_k$ into small pieces in an arbitrary way. In Definition A.6 we defined the *barycentric subdivision* of a polyhedron, and showed how to realize it using a tent function. The barycentric subdivision of Δ_k divides it into $(k+1)!$ simplices Δ_k , each of which is smaller than the original Δ_k by a factor of at most $\frac{k}{k+1}$ in every direction, for $k \geq 1$. Barycentric subdivision can be iterated, so that for $N = 1, 2, \dots$ the N^{th} barycentric subdivision of Δ_k divides it up into $((k+1)!)^N$ smaller simplices Δ_k , each of which is smaller than the original Δ_k by a factor of at most $(\frac{k}{k+1})^N$ in every direction. Since $(\frac{k}{k+1})^N \rightarrow 0$ as $N \rightarrow \infty$, by taking N arbitrarily large we make the simplices of the N^{th} barycentric subdivision of Δ_k arbitrarily small. The N^{th} barycentric subdivision of Δ_k can also be realized using a tent function, combining the constructions of Definition A.6 and Theorem A.7.

Therefore we choose $N \gg 0$ such that the N^{th} barycentric subdivision of each boundary component $\tilde{X}_{ab}^1 \cong \Delta_k$ corresponding to some $d \in D$ cuts $\tilde{X}_{ab}^1$, and all its boundary faces of dimension $k-1, k-2, \dots, 1$, into sufficiently small pieces that the smallness condition Theorem A.18(iii) holds for these faces if we use the N^{th} barycentric subdivision and its associated tent function for $\tilde{X}_{ab}^1 \cong \Delta_k$, rather than just some tent function chosen arbitrarily by iteration using Theorem A.18. Then in $(\dagger)_{ab}^m$ we impose the extra condition that for any triple $(\tilde{X}_{ab}^m, \tilde{f}_{ab}^m, \tilde{\mathbf{G}}_{ab}^m)$ which is isomorphic to a component of $\partial^{m-1}(\tilde{X}_{ab}^1, \tilde{f}_{ab}^1, \tilde{\mathbf{G}}_{ab}^1)$ for some $\tilde{X}_{ab}^1 \cong \Delta_k$ with $[\tilde{X}_{ab}^1, \tilde{f}_{ab}^1, \tilde{\mathbf{G}}_{ab}^1] = \pm [\Delta_k, \tau_d, \underline{\mathbf{G}}_{\Delta_k}]$ for some $d \in D$, so that we have a natural isomorphism $\tilde{X}_{ab}^m \cong \Delta_{k-m}$, then $\tilde{T}_{ab}^m$ and $\sigma_{ab\bar{c}}^m$ for $c \in C_{ab}^m$ should come from the N^{th} barycentric subdivision of Δ_{k-m} .

Then writing $\{v_e : e \in E\}$ for the simplices in the N^{th} barycentric subdivision of Δ_k , and setting $\zeta_e = 1$ if v_e is orientation-preserving and $\zeta_e = -1$ otherwise, the proof of (220) in Step 3 generalizes to give (222). Now barycentric subdivision is used by Bredon [10, §IV.17] to prove the Excision Axiom for singular homology. Bredon defines $\Upsilon : C_*^{\text{si}}(Y; R) \rightarrow C_*^{\text{si}}(Y; R)$ which replaces each singular simplex by its barycentric subdivision, and $T : C_*^{\text{si}}(Y; R) \rightarrow C_{*+1}^{\text{si}}(Y; R)$ with $\partial \circ T + T \circ \partial = \Upsilon - \text{id}$. Thus Υ induces the identity on $H_*^{\text{si}}(Y; R)$. In this

notation, $\sum_{d \in D} \sum_{e \in E} (\eta_d \zeta_e) (\tau_d \circ v_e) = \Upsilon^N (\sum_{d \in D} \eta_d \tau_d)$, so

$$\beta = [\sum_{d \in D} \eta_d \tau_d] = [\sum_{d \in D} \sum_{e \in E} (\eta_d \zeta_e) (\tau_d \circ v_e)] = 0 \quad \text{in } H_k^{\text{si}}(Y; R),$$

as Υ is the identity on $H_k^{\text{si}}(Y; R)$, and using (222). Therefore $\Pi_{\text{si}}^{\text{ef}}$ in (76) is *injective*, as we want. This completes Step 4, and the proof of Theorem 4.8.

C Proof that $KH_*(Y; R) \cong KH_*^{\text{ef}}(Y; R)$

We now prove Theorem 4.9. Let Y be an orbifold, R a $\mathbb{Q}$ -algebra and $k \in \mathbb{Z}$. We must show that $\Pi_{\text{ef}}^{\text{Kh}} : KH_k^{\text{ef}}(Y; R) \rightarrow KH_k(Y; R)$ in (68) is an isomorphism. In a similar way to the proof of Theorem 4.8 in Appendix B, we do this in five steps, which we describe briefly below and then cover in detail in §C.1–§C.5. Steps 1–4 show that $\Pi_{\text{ef}}^{\text{Kh}}$ in (68) is surjective, and Step 5 that it is injective.

Step 1. Let $k \in \mathbb{Z}$, $\alpha \in KH_k(Y; R)$, and $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a] \in KC_k(Y; R)$ represent α . The goal of this step is to choose a *tent function* $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ for each $a \in A$ to ‘cut’ each X_a into finitely many *arbitrarily small pieces* X_{ac} for $c \in C_a$, in triples $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$, using the results of §A.3.5. This construction will be used for two different purposes in Steps 2 and 3, and each time we apply it, we will explain what we mean by ‘arbitrarily small’.

As in Step 1 of Appendix B, we wish to choose the $\mathbf{T}_a$ such that $\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$ is homologous to $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]$ in $KC_k(Y; R)$. Since $\partial(\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]) = 0$, the $[\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a}]$ must satisfy relations in $KC_{k-1}(Y; R)$. The issue is to choose the $\mathbf{T}_a$ for $a \in A$ in a way compatible with these boundary relations, so that $\partial(\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]) = 0$. In Step 1 of Appendix B we solved a similar problem by introducing notation $n_a^m, (X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m), P^m, Q^m, R^m, \dots$, and then choosing data on X_{ab}^m for $(a, b) \in R^m$ with boundary conditions by reverse induction on codimension $m = M, M-1, \dots, 0$. Here we adopt a similar strategy, but the problem is more difficult because of the extra relation Definition 4.3(iv).

Define $\widetilde{KC}_k(Y; R)$ to be the R -module of finite R -linear combinations of isomorphism classes $[X, \mathbf{f}, \mathbf{G}]$ as in Definition 4.3 for which $\text{vdim } X = k$, with relations Definition 4.3(i)–(iii), but *not* (iv). Write $\Pi : \widetilde{KC}_k(Y; R) \rightarrow KC_k(Y; R)$ for the natural, surjective projection $\Pi : \sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a] \mapsto \sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]$, whose kernel is the R -submodule of $\widetilde{KC}_k(Y; R)$ set to zero by equation (65) in Definition 4.3(iv). That is, $\text{Ker } \Pi$ is the subspace of elements of $\widetilde{KC}_k(Y; R)$ of the form

$$\sum_{d \in D} \eta_d ([X_d / \Gamma_d, \pi_*(\mathbf{f}_d), \pi_*(\mathbf{G}_d)] - \frac{1}{|\Gamma_d|} [X_d, \mathbf{f}_d, \mathbf{G}_d]), \quad (254)$$

where D is a finite indexing set, and for each $d \in D$ we have $\eta_d \in R$ and $X_d, \mathbf{f}_d, \mathbf{G}_d, \Gamma_d$ are as in Definition 4.3(iv). Using Definition 4.3(i)–(iii) we can also suppose that each $(X_d, \mathbf{f}_d, \mathbf{G}_d)$ is connected.

The proofs in Appendix B relied on the fact that triples $[X, \mathbf{f}, \mathbf{G}]$ with $(X, \mathbf{f}, \mathbf{G})$ connected form a basis for $KC_k^{\text{ef}}(Y; R)$ over R , modulo only the relation $[-X, \mathbf{f}, \mathbf{G}] = -[X, \mathbf{f}, \mathbf{G}]$; this was how we showed the coefficient $\{\dots\}$ on

the last line of (229) is zero. Since $\widetilde{KC}_k(Y; R)$ is defined using the same relations as $KC_k^{\text{ef}}(Y; R)$, it is easy to see that triples $[X, \mathbf{f}, \mathbf{G}]$ such that $(X, \mathbf{f}, \mathbf{G})$ is connected and does not admit an orientation-reversing isomorphism form a basis for $\widetilde{KC}_k(Y; R)$ over R , modulo only the relation $[-X, \mathbf{f}, \mathbf{G}] = -[X, \mathbf{f}, \mathbf{G}]$. However, we do *not* have a good basis for $KC_k(Y; R)$. This makes choosing the $\mathbf{T}_a$ for $a \in A$ compatible with relations in $KC_{k-1}(Y; R)$ more complicated.

Our strategy is to lift from $KC_*(Y; R)$ to $\widetilde{KC}_*(Y; R)$, where we have a good basis, and deal with the extra relation Definition 4.3(iv) ‘by hand’. As $\partial(\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]) = 0$ in $KC_{k-1}(Y; R)$, it lifts to $\text{Ker } \Pi$ in $\widetilde{KC}_{k-1}(Y; R)$, and so is of the form (254). Thus we may write

$$\begin{aligned} \sum_{a \in A} \rho_a [\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a}] = \\ \sum_{d \in D} \eta_d ([\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d)] - \frac{1}{|\Gamma_d|} [\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d]) \text{ in } \widetilde{KC}_{k-1}(Y; R), \end{aligned} \quad (255)$$

with each $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ connected.

We shall choose tent functions $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ for $a \in A$, and $\tilde{\mathbf{T}}_d$ for $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ for $d \in D$, satisfying the conditions:

- (a) $\mathbf{T}_a|_{\partial X_a}$ is invariant under $\text{Aut}(\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a})$ for each $a \in A$, and $\tilde{\mathbf{T}}_d$ is invariant under $\text{Aut}(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$, and $\pi_*(\tilde{\mathbf{T}}_d)$ is invariant under $\text{Aut}(\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d))$ for each $d \in D$.
- (b) If two connected components of any two of $(\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a})$ for $a \in A$ and $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$, $(\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d))$ for $d \in D$ are isomorphic, the restrictions of $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ to these components are identified by the isomorphism.
- (c) The $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ cut $(X_a, \mathbf{f}_a, \mathbf{G}_a), (\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ into ‘arbitrarily small pieces’, where we will explain how small the pieces need be whenever we apply the result.

To do this consistently, we must actually work by induction on decreasing codimension m in a similar way to Appendix B, and choose data on $\partial^m(X_a, \mathbf{f}_a, \mathbf{G}_a)$ and $\partial^{m-1}(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ for $m = M, M-1, \dots, 1, 0$ satisfying analogues of (a)–(c), with boundary values chosen in the previous inductive step. The details will be explained in §C.1.

Haven chosen $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ for all a, d satisfying (a)–(c), equation (211) of §A.3 gives, in both $KC_k(Y; R)$ and $\widetilde{KC}_k(Y; R)$,

$$\begin{aligned} \partial[Z_{X_a, \mathbf{T}_a}, \mathbf{f}_a \circ \pi, \mathbf{H}_{X_a, \mathbf{T}_a}] = -[Z_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}, \mathbf{f}_a|_{\partial X_a} \circ \pi, \mathbf{H}_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}] \\ - [X_a, \mathbf{f}_a, \mathbf{G}_a] + \sum_{c \in C_a} [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]. \end{aligned} \quad (256)$$

From equation (255) and (a),(b) above we deduce that

$$\begin{aligned} \sum_{a \in A} \rho_a [Z_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}, \mathbf{f}_a|_{\partial X_a} \circ \pi, \mathbf{H}_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}] = \\ \sum_{d \in D} \eta_d ([Z_{\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{T}}_d)}, \pi_*(\tilde{\mathbf{f}}_d) \circ \pi, \mathbf{H}_{\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{T}}_d)}] \text{ in } \widetilde{KC}_k(Y; R). \\ - \frac{1}{|\Gamma_d|} [Z_{\tilde{X}_d, \tilde{\mathbf{T}}_d}, \tilde{\mathbf{f}}_d \circ \pi, \mathbf{H}_{\tilde{X}_d, \tilde{\mathbf{T}}_d}]) \end{aligned} \quad (257)$$

The action of Γ_d on $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ preserves $\tilde{\mathbf{T}}_d$ and so lifts to an action of Γ_d on $(Z_{\tilde{X}_d, \tilde{\mathbf{T}}_d}, \tilde{\mathbf{f}}_d \circ \pi, \mathbf{H}_{\tilde{X}_d, \tilde{\mathbf{T}}_d})$, and the quotient is isomorphic to $(Z_{\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{T}}_d)}, \pi_*(\tilde{\mathbf{f}}_d) \circ \pi, \mathbf{H}_{\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{T}}_d)})$. Thus the right hand side of (257) lies in $\text{Ker } \Pi$, and projecting to $KC_k(Y; R)$ yields

$$\sum_{a \in A} \rho_a [Z_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}, \mathbf{f}_a|_{\partial X_a} \circ \pi, \mathbf{H}_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}] = 0 \text{ in } KC_k(Y; R). \quad (258)$$

Multiplying (256) by ρ_a , summing over $a \in A$ and using (258) yields

$$\begin{aligned} & \partial \left(\sum_{a \in A} \rho_a [Z_{X_a, \mathbf{T}_a}, \mathbf{f}_a \circ \pi, \mathbf{H}_{X_a, \mathbf{T}_a}] \right) \\ &= \sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}] - \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a] \end{aligned} \quad (259)$$

in $KC_k(Y; R)$, as for (226). Thus $\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$ is a cycle homologous to $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$, as we want.

Step 2. Let $k \in \mathbb{Z}$, $\alpha \in KH_k(Y; R)$, and $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a] \in KC_k(Y; R)$ represent α . Using Step 1, we choose a tent function $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ for all $a \in A$ which ‘cuts’ each X_a into small pieces X_{ac} for $c \in C_a$, in triples $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$, such that for each $a \in A$ and $c \in C_a$ there exists a triple $(\dot{X}_{ac}, \dot{\mathbf{f}}_{ac}, \dot{\mathbf{G}}_{ac})$ with $\dot{X}_{ac}$ an oriented Kuranishi space with *trivial stabilizers*, and writing $\dot{\mathbf{G}}_{ac} = (\dot{\mathbf{I}}_{ac}, \dot{\eta}_{ac}, G_{ac}^i : i \in I_{ac})$ with $\dot{\mathbf{I}}_{ac} = (\dot{I}_{ac}, (\dot{V}_{ac}^i, \dots, \dot{\psi}_{ac}^i) : i \in \dot{I}_{ac}, \dots)$, each $\dot{V}_{ac}^i$ is a *manifold*, and there exists a finite group Γ_{ac} of automorphisms $(\mathbf{a}, \mathbf{b}) : (\dot{X}_{ac}, \dot{\mathbf{f}}_{ac}, \dot{\mathbf{G}}_{ac}) \rightarrow (\dot{X}_{ac}, \dot{\mathbf{f}}_{ac}, \dot{\mathbf{G}}_{ac})$ with $\mathbf{a} : \dot{X}_{ac} \rightarrow \dot{X}_{ac}$ orientation-preserving, such that the triple $(\dot{X}_{ac}/\Gamma_{ac}, \pi_*(\dot{\mathbf{f}}_{ac}), \pi_*(\dot{\mathbf{G}}_{ac}))$ is isomorphic to $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$. Relation Definition 4.3(iv) thus gives

$$[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}] = \frac{1}{|\Gamma_{ac}|} [\dot{X}_{ac}, \dot{\mathbf{f}}_{ac}, \dot{\mathbf{G}}_{ac}] \quad \text{in } KC_k(Y; R). \quad (260)$$

Such $(\dot{X}_{ac}, \dot{\mathbf{f}}_{ac}, \dot{\mathbf{G}}_{ac})$ exist provided the X_{ac} and V_{ac}^i in $\mathbf{I}_{ac}$ in $\mathbf{G}_{ac}$ are ‘sufficiently small’, just as in an orbifold V , any sufficiently small open set in V may be written as U/Γ for U a manifold and Γ a finite group.

Combining equation (260) and Step 1 shows that

$$\sum_{a \in A} \sum_{c \in C_a} \rho_a |\Gamma_{ac}|^{-1} [\dot{X}_{ac}, \dot{\mathbf{f}}_{ac}, \dot{\mathbf{G}}_{ac}] = \sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}] \quad (261)$$

is homologous to $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$ in $KC_k(Y; R)$, and so is a cycle representing α . Thus we can represent α by a cycle (261) in which the Kuranishi spaces $\dot{X}_{ac}$ have *trivial stabilizers*, and in the Kuranishi neighbourhoods $(\dot{V}_{ac}^i, \dots, \dot{\psi}_{ac}^i)$ in $\dot{\mathbf{G}}_{ac}$, the $\dot{V}_{ac}^i$ are *manifolds*. Hence $\dot{X}_{ac}$ is an *effective* Kuranishi space, and the $(\dot{V}_{ac}^i, \dots, \dot{\psi}_{ac}^i)$ are *effective* Kuranishi neighbourhoods. We will need this when we lift to effective Kuranishi chains in Step 4.

Step 3. For simplicity we now *change notation* from $\dot{X}_{ac}$ for $a \in A$ and $c \in C_a$ back to X_a for $a \in A$, and absorb the factors $|\Gamma_{ac}|^{-1}$ in (261) into new constants $\rho_a \in R$. That is, Steps 1 and 2 have shown that we can represent α by a cycle

$\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a] \in KC_k(Y; R)$, where each X_a is a compact, oriented Kuranishi space with *trivial stabilizers*, and writing $\mathbf{G}_a = (\mathbf{I}_a, \boldsymbol{\eta}_a, G_a^i : i \in I_a)$ with $\mathbf{I}_a = (I_a, (V_a^i, \dots, \psi_a^i) : i \in I_a, \dots)$, each V_a^i is a *manifold*. Equation (255) holds for connected $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ for $d \in D$. We stress that these $A, X_a, \mathbf{f}_a, \mathbf{G}_a, \rho_a, D, \tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d, \Gamma_d, \eta_d$ are different from those in Steps 1 and 2.

We now repeat Step 1, choosing tent functions $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ for $a \in A$, and $\tilde{\mathbf{T}}_d$ for $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ for $d \in D$, satisfying conditions (a)–(c), such that $\mathbf{T}_a$ cuts $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ into finitely many small pieces $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ for $c \in C_a$, and $\tilde{\mathbf{T}}_d$ cuts $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ into finitely many small pieces $(\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df})$ for $f \in F_d$.

As in Step 1, equations (255)–(257) hold in $\widetilde{KC}_*(Y; R)$. Taking ∂ of (256) and using $\partial^2 = 0$ gives

$$\begin{aligned} \partial[Z_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}, \mathbf{f}_a|_{\partial X_a} \circ \boldsymbol{\pi}, \mathbf{H}_{\partial X_a, \mathbf{T}_a|_{\partial X_a}}] = & \quad \text{in } \widetilde{KC}_{k-1}(Y; R). \\ - \partial[X_a, \mathbf{f}_a, \mathbf{G}_a] + \sum_{c \in C_a} \partial[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}] \end{aligned} \quad (262)$$

The analogue of (256) for $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ in $\widetilde{KC}_{k-1}(Y; R)$ is

$$\begin{aligned} \partial[Z_{\tilde{X}_d, \tilde{\mathbf{T}}_d}, \tilde{\mathbf{f}}_d \circ \boldsymbol{\pi}, \mathbf{H}_{\tilde{X}_d, \tilde{\mathbf{T}}_d}] = & -[Z_{\partial \tilde{X}_d, \tilde{\mathbf{T}}_d|_{\partial \tilde{X}_d}}, \tilde{\mathbf{f}}_d|_{\partial \tilde{X}_d} \circ \boldsymbol{\pi}, \mathbf{H}_{\partial \tilde{X}_d, \tilde{\mathbf{T}}_d|_{\partial \tilde{X}_d}}] \\ & - [\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d] + \sum_{f \in F_d} [\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df}]. \end{aligned} \quad (263)$$

The action of Γ_d on $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ lifts to $\coprod_{f \in F_d} (\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df})$, and

$$\left(\coprod_{f \in F_d} (\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df}) \right) / \Gamma_d \cong \coprod_{f \Gamma_d \in F_d / \Gamma_d} (\tilde{X}_{df} / \text{Stab}_{\Gamma_d}(f), \boldsymbol{\pi}_*(\tilde{\mathbf{f}}_{df}), \boldsymbol{\pi}_*(\tilde{\mathbf{G}}_{df})),$$

where to interpret the right hand side, for each orbit $f \Gamma_d$ of Γ_d in F_d we pick a representative f , and then divide $(\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df})$ by the stabilizer group $\text{Stab}_{\Gamma_d}(f)$ of f in Γ_d . From this we deduce an analogue of (263) for $(\tilde{X}_d / \Gamma_d, \boldsymbol{\pi}_*(\tilde{\mathbf{f}}_d), \boldsymbol{\pi}_*(\tilde{\mathbf{G}}_d))$ in $\widetilde{KC}_{k-1}(Y; R)$:

$$\begin{aligned} \partial[Z_{\tilde{X}_d / \Gamma_d, \boldsymbol{\pi}_*(\tilde{\mathbf{T}}_d)}, \boldsymbol{\pi}_*(\tilde{\mathbf{f}}_d) \circ \boldsymbol{\pi}, \mathbf{H}_{\tilde{X}_d / \Gamma_d, \boldsymbol{\pi}_*(\tilde{\mathbf{T}}_d)}] = & \\ - [Z_{\partial(\tilde{X}_d / \Gamma_d), \boldsymbol{\pi}_*(\tilde{\mathbf{T}}_d)|_{\partial(\tilde{X}_d / \Gamma_d)}}, \boldsymbol{\pi}_*(\tilde{\mathbf{f}}_d)|_{\partial(\tilde{X}_d / \Gamma_d)} \circ \boldsymbol{\pi}, & \\ \mathbf{H}_{\partial(\tilde{X}_d / \Gamma_d), \boldsymbol{\pi}_*(\tilde{\mathbf{T}}_d)|_{\partial(\tilde{X}_d / \Gamma_d)}}] - [\tilde{X}_d / \Gamma_d, \boldsymbol{\pi}_*(\tilde{\mathbf{f}}_d), \boldsymbol{\pi}_*(\tilde{\mathbf{G}}_d)] & \\ + \sum_{f \Gamma_d \in F_d / \Gamma_d} [\tilde{X}_{df} / \text{Stab}_{\Gamma_d}(f), \boldsymbol{\pi}_*(\tilde{\mathbf{f}}_{df}), \boldsymbol{\pi}_*(\tilde{\mathbf{G}}_{df})]. & \end{aligned} \quad (264)$$

Taking the boundary of (255) and using $\partial^2 = 0$ gives

$$\begin{aligned} 0 = \sum_{d \in D} \eta_d ([\partial(\tilde{X}_d / \Gamma_d), \boldsymbol{\pi}_*(\tilde{\mathbf{f}}_d)|_{\partial(\tilde{X}_d / \Gamma_d)}, \boldsymbol{\pi}_*(\tilde{\mathbf{G}}_d)|_{\partial(\tilde{X}_d / \Gamma_d)}] & \\ - |\Gamma_d|^{-1} [\partial \tilde{X}_d, \tilde{\mathbf{f}}_d|_{\partial \tilde{X}_d}, \tilde{\mathbf{G}}_d|_{\partial \tilde{X}_d}]) & \quad \text{in } \widetilde{KC}_{k-2}(Y; R). \end{aligned} \quad (265)$$

As the choice of tent functions $\tilde{\mathbf{T}}_d, \boldsymbol{\pi}_*(\tilde{\mathbf{T}}_d)$ in Step 1 actually depends only on the isomorphism class of each connected component of $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$, $(\tilde{X}_d / \Gamma_d,$

$\pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d))$ in $\widetilde{KC}_{k-1}(Y; R)$, and by the inductive construction this also holds for their boundaries in $\widetilde{KC}_{k-2}(Y; R)$, we can lift (265) to an equation in the $Z_{\dots}$ chains in $\widetilde{KC}_{k-1}(Y; R)$:

$$0 = \sum_{d \in D} \eta_d \left(\left[Z_{\partial(\tilde{X}_d/\Gamma_d), \pi_*(\tilde{\mathbf{T}}_d)|_{\partial(\tilde{X}_d/\Gamma_d)}, \pi_*(\tilde{\mathbf{f}}_d)|_{\partial(\tilde{X}_d/\Gamma_d)} \circ \pi, \right. \right. \\ \left. \left. \mathbf{H}_{\partial(\tilde{X}_d/\Gamma_d), \pi_*(\tilde{\mathbf{T}}_d)|_{\partial(\tilde{X}_d/\Gamma_d)}} \right] \right. \\ \left. - |\Gamma_d|^{-1} [Z_{\partial\tilde{X}_d, \tilde{\mathbf{T}}_d|_{\partial\tilde{X}_d}, \tilde{\mathbf{f}}_d|_{\partial\tilde{X}_d} \circ \pi, \mathbf{H}_{\partial\tilde{X}_d, \tilde{\mathbf{T}}_d|_{\partial\tilde{X}_d}}] \right). \quad (266)$$

Take the boundary ∂ of (257) to get an equation in $\widetilde{KC}_{k-1}(Y; R)$. Use (262), (264) and (263) to substitute for the boundaries of the first, second and third lines of (257) respectively. Add on (255), which cancels three sets of terms, and then add on (266), which cancels two more. This yields an equation in $\widetilde{KC}_{k-1}(Y; R)$:

$$\partial \left(\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}] \right) = \\ \sum_{d \in D} \eta_d \left(\sum_{f \in F_d/\Gamma_d} [\tilde{X}_{df}/\text{Stab}_{\Gamma_d}(f), \pi_*(\tilde{\mathbf{f}}_{df}), \pi_*(\tilde{\mathbf{G}}_{df})] \right. \\ \left. - |\Gamma_d|^{-1} \sum_{f \in F_d} [\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df}] \right). \quad (267)$$

We will prove that provided the $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ are chosen to make the $X_{ac}, \tilde{X}_{df}$ sufficiently small, the terms on the right hand side of (267) all cancel, that is, the right hand side is zero. To get a rough idea of why, note that since the X_a have trivial stabilizers, if $\tilde{X}_d/\Gamma_d$ is needed to cancel part of any ∂X_a in (255), then $\tilde{X}_d/\Gamma_d$ must have trivial stabilizers, and so Γ_d acts freely on $\tilde{X}_d$. But if Γ_d acts freely on $\tilde{X}_d$, and the pieces $\tilde{X}_{df}$ for $f \in F_d$ are sufficiently small, then Γ_d must also act freely on F_d . And if Γ_d acts freely on F_d then $\sum_{f \in F_d/\Gamma_d} [\tilde{X}_{df}/\text{Stab}_{\Gamma_d}(f), \dots] = |\Gamma_d|^{-1} \sum_{f \in F_d} [\tilde{X}_{df}, \dots]$, so the second and third lines of (267) cancel.

We have now constructed $\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$, where X_{ac} has trivial stabilizers and the V_{ac}^i are manifolds, and shown that regarded as an element of $KC_k(Y; R)$ this is a cycle representing α , and regarded as an element of $\widetilde{KC}_k(Y; R)$ satisfies

$$\partial \left(\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}] \right) = 0 \quad \text{in } \widetilde{KC}_{k-1}(Y; R). \quad (268)$$

Thus, we have lifted from a cycle in $(KC_*(Y; R), \partial)$ to one in $(\widetilde{KC}_*(Y; R), \partial)$. For the rest of the proof we will work in the complex $(\widetilde{KC}_*(Y; R), \partial)$, which as in Step 1 has an easily understood basis, and we no longer need to worry about the relation Definition 4.3(iv).

In this step we also prove we can take the $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ to have an additional property that we will need later. Fix $a \in A$ and $c \in C_a$, and for $m \geq 0$ write $(X_{acd}^m, \mathbf{f}_{acd}^m, \mathbf{G}_{acd}^m)$ for $d = 1, \dots, n_{ac}^m$ for the connected components of $(\partial^m X_{ac}, \mathbf{f}_{ac}|_{\partial^m X_{ac}}, \mathbf{G}_{ac}|_{\partial^m X_{ac}})$. Then in the usual notation the Kuranishi

neighbourhoods $(V_{acd}^{m,i-m}, \dots, \psi_{acd}^{m,i-m})$ for $i-m \in I_{ac}^m$ in $\mathbf{G}_{acd}^m$ have $V_{acd}^{m,i-m}$ an open subset of $\partial^m V_{ac}^i$, where V_{ac}^i is a *manifold*.

If V is a manifold with corners and $m \geq 2$ then points of $\partial^m V$ may be written as $(v, B_1, \dots, B_m)$, where $B_1, \dots, B_m$ are distinct local boundary components of V . Hence $(v, B_1), \dots, (v, B_m)$ lie in ∂V . Thus, a point of $V_{acd}^{m,i-m}$ is of the form $(v, B_1, \dots, B_m)$, where $(v, B_l) \in \partial V_{ac}^i$ for $l = 1, \dots, m$. It is necessary that $(v, B_l) \in V_{acd_l}^{1,i-1}$ for some unique $d_l = 1, \dots, n_{ac}^1$. That is, (v, B_l) cannot lie in the part of ∂V_{ac}^i discarded by Algorithm 3.16, since then $(v, B_1, \dots, B_m)$ would also be discarded, so (v, B_l) lies in $\coprod_{d'=1}^{n_{ac}^1} V_{acd'}^{1,i-1}$, and so in $V_{acd_l}^{1,i-1}$ for unique d_l .

Hence, each $i \in I_{acd}^m$ and $(v, B_1, \dots, B_m)$ in $V_{acd}^{m,i-m}$ determine an l -tuple $(d_1, \dots, d_m)$ with $d_l = 1, \dots, n_{ac}^1$. We claim that this $(d_1, \dots, d_m)$ is independent of i and $(v, B_1, \dots, B_m)$, and depends only on $d = 1, \dots, n_{ac}^m$. To see this, note that $(d_1, \dots, d_m)$ is locally constant on $V_{acd}^{m,i-m}$, and is also preserved under coordinate changes $(\phi_{acd}^{m,(i-m)(j-m)}, \hat{\phi}_{acd}^{m,(i-m)(j-m)})$ in $\mathbf{G}_{acd}^m$, so the part of $(\partial^m X_{ac}, \mathbf{f}_{ac}|_{\partial^m X_{ac}}, \mathbf{G}_{ac}|_{\partial^m X_{ac}})$ with fixed $(d_1, \dots, d_m)$ must be a union of connected components. Thus, each component of $(\partial^m X_{ac}, \mathbf{f}_{ac}|_{\partial^m X_{ac}}, \mathbf{G}_{ac}|_{\partial^m X_{ac}})$ determines an ordered m -tuple of components of $(\partial X_{ac}, \mathbf{f}_{ac}|_{\partial X_{ac}}, \mathbf{G}_{ac}|_{\partial X_{ac}})$.

We will prove that provided the X_{ac} are chosen sufficiently small, for all $a \in A$, $c \in C_a$, $m \geq 2$ and $d = 1, \dots, n_{ac}^m$, the numbers $d_1, \dots, d_m$ above are *distinct*. That is, we show that each component of $(\partial^m X_{ac}, \mathbf{f}_{ac}|_{\partial^m X_{ac}}, \mathbf{G}_{ac}|_{\partial^m X_{ac}})$ occurs as the intersection of m *distinct* components of $(\partial X_{ac}, \mathbf{f}_{ac}|_{\partial X_{ac}}, \mathbf{G}_{ac}|_{\partial X_{ac}})$.

Here is why we need this. Suppose $(X, \mathbf{f}, \mathbf{G})$ is a triple in effective Kuranishi homology. Then there is a natural action of the symmetric group S_m on $(\partial^m X, \mathbf{f}|_{\partial^m X}, \mathbf{G}|_{\partial^m X})$ by isomorphisms, which acts on points $(v, B_1, \dots, B_m)$ in $V_{acd}^{m,i-m}$ in $\mathbf{G}|_{\partial^m X}$ by permutation of $B_1, \dots, B_m$. As connected components of $(\partial^m X, \mathbf{f}|_{\partial^m X}, \mathbf{G}|_{\partial^m X})$ have no automorphisms by Theorem 3.39(b), S_m must act *freely* on the set of connected components of $(\partial^m X, \mathbf{f}|_{\partial^m X}, \mathbf{G}|_{\partial^m X})$.

In Step 4 we will modify $\mathbf{G}_{ac}$ to effective gauge-fixing data. A necessary condition to be able to do this is that S_m acts freely on the connected components of $(\partial^m X_{ac}, \mathbf{f}_{ac}|_{\partial^m X_{ac}}, \mathbf{G}_{ac}|_{\partial^m X_{ac}})$. Since each component determines some $(d_1, \dots, d_m)$ and S_m permutes $d_1, \dots, d_m$, if $d_1, \dots, d_m$ are *distinct* then the action of S_m on the set of components is automatically free, as we want.

Step 4. For simplicity we again *change notation* from X_{ac} for $a \in A$ and $c \in C_a$ back to X_a for $a \in A$. That is, Step 3 has shown that we can represent α by a cycle $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]$ in $KC_k(Y; R)$ or $\widetilde{KC}_k(Y; R)$, where each X_a is a compact, oriented Kuranishi space with *trivial stabilizers*, and writing $\mathbf{G}_a = (\mathbf{I}_a, \boldsymbol{\eta}_a, G_a^i : i \in I_a)$ with $\mathbf{I}_a = (I_a, (V_a^i, \dots, \psi_a^i) : i \in I_a, \dots)$, each V_a^i is a *manifold*, each $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ is *connected*, and

$$\partial(\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]) = 0 \quad \text{in } \widetilde{KC}_{k-1}(Y; R). \quad (269)$$

Moreover, for $a \in A$ and $m \geq 2$ each component of $(\partial^m X_a, \mathbf{f}_a|_{\partial^m X_a}, \mathbf{G}_a|_{\partial^m X_a})$ occurs as the intersection of m *distinct* components of $(\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a})$.

Define notation $n_a^m, (X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m), P^m, \sim, Q^m, R^m, \phi^m, (\mathbf{a}, \mathbf{b})_{ab}^m, M, \epsilon_{ab}^m, B_{ab}^m$ exactly as in Step 1 of Appendix B, but with $\mathbf{G}_a, \mathbf{G}_{ab}^m$ in place of $\underline{\mathbf{G}}_a, \underline{\mathbf{G}}_{ab}^m$. Note

one significant difference: in Appendix B we have $\text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) = \{1\}$ by Theorem 3.39(b), so $(\mathbf{a}, \mathbf{b})_{ab}^m$ is unique. However, in the noneffective case we know only that $\text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ is finite by Theorem 3.20, so when $\phi^m(a, b) = (\bar{a}, \bar{b})$ the isomorphism $(\mathbf{a}, \mathbf{b})_{ab}^m : (X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) \rightarrow (X_{\bar{a}\bar{b}}^m, \mathbf{f}_{\bar{a}\bar{b}}^m, \mathbf{G}_{\bar{a}\bar{b}}^m)$ may not be unique, we have to make an arbitrary choice of $(\mathbf{a}, \mathbf{b})_{ab}^m$ out of finitely many possibilities.

Then as for equation (229), from (269) in $\widetilde{KC}_{k-1}(Y; R)$ we have

$$\begin{aligned} 0 &= \partial \left[\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a] \right] = \sum_{a \in A} \rho_a \sum_{b=1}^{n_a^1} [X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1] \\ &= \sum_{\substack{a \in A, b=1, \dots, n_a^1 \\ \text{Set } (\bar{a}, \bar{b}) = \phi^1(a, b)}} \rho_a \epsilon_{ab}^1 [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1] = \sum_{(\bar{a}, \bar{b}) \in R^1} \left\{ \sum_{\substack{a \in A, b=1, \dots, n_a^1 \\ \phi^1(a, b) = (\bar{a}, \bar{b})}} \rho_a \epsilon_{ab}^1 \right\} [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1]. \end{aligned} \quad (270)$$

By construction the triples $(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)$ for $(\bar{a}, \bar{b}) \in R^1$ are connected and mutually non-isomorphic, even through orientation-reversing isomorphisms. As in Step 1, triples $[X, \mathbf{f}, \mathbf{G}]$ such that $(X, \mathbf{f}, \mathbf{G})$ is connected and does not admit an orientation-reversing automorphism form a basis for $\widetilde{KC}_{k-1}(Y; R)$ over R , modulo $[-X, \mathbf{f}, \mathbf{G}] = -[X, \mathbf{f}, \mathbf{G}]$.

We separate $(\bar{a}, \bar{b}) \in R^1$ into two cases: (A) $(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)$ admits an orientation-reversing automorphism, and (B) it does not. In case (A) we have $[X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1] = 0$ by Definition 4.3(ii). The $[X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1]$ in case (B) form part of a basis for $\widetilde{KC}_{k-1}(Y; R)$ over R , and so are linearly independent over R . Thus, in the same way that we argued in §B.1 that the coefficients $\{\dots\}$ in the last line of (229) are zero, from (270) we see that for each $(\bar{a}, \bar{b}) \in R^1$, either

- (A) $(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)$ admits an orientation-reversing automorphism, so that $[X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1] = 0$ in $KC_{k-1}(Y; R)$ and $\widetilde{KC}_{k-1}(Y; R)$; or
- (B) $\sum_{a \in A, b=1, \dots, n_a^1: \phi^1(a, b) = (\bar{a}, \bar{b})} \rho_a \epsilon_{ab}^1 = 0$ in R .

The rough idea of this step is to choose maps $\underline{G}_a^i : E_a^i \rightarrow \underline{P}$ for all $a \in A$ and $i \in I_a$ such that $\underline{G}_a = (\mathbf{I}_a, \boldsymbol{\eta}_a, \underline{G}_a^i : i \in I_a)$ is effective gauge-fixing data for $(X_a, \mathbf{f}_a)$. (Note that as X_a has trivial stabilizers, it is an *effective* Kuranishi space, and as V_a^i is a manifold, $(V_a^i, \dots, \psi_a^i)$ is an *effective* Kuranishi neighbourhood, so we have already satisfied some of the conditions in Definition 3.36 for $\underline{G}_a$.) Thus we can form a chain $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{G}_a] \in KC_k^{\text{ef}}(Y; R)$, and project to $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \Pi(\underline{G}_a)] = \Pi_{\text{ef}}^{\text{Kh}}(\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{G}_a])$ in $KC_k(Y; R)$.

If we could choose the $\underline{G}_a^i$ for all a, i compatible with the relations between the $[\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a}]$ implied by (269)–(270), we would expect to prove that $\partial(\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{G}_a]) = 0$ in $KC_{k-1}^{\text{ef}}(Y; R)$, giving a homology class $\beta = [\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \underline{G}_a]]$ in $KH_k^{\text{ef}}(Y; R)$, and to show $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \Pi(\underline{G}_a)]$ is homologous to $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$ in $KC_k(Y; R)$, so that $\Pi_{\text{ef}}^{\text{Kh}}(\beta) = \alpha$. As $\alpha \in KH_k(Y; R)$ was arbitrary, this would show that $\Pi_{\text{ef}}^{\text{Kh}} : KH_k^{\text{ef}}(Y; R) \rightarrow KH_k(Y; R)$ is *surjective*.

However, there is a problem. In the $\text{Aut}(X_a, \mathbf{f}_a, \mathbf{G}_a)$ are finite by Theorem 3.20, but for connected $(X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a)$ the $\text{Aut}(X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a)$ are trivial by Theorem 3.39(b). Thus, when we lift from $\mathbf{G}_a$ to $\underline{\mathbf{G}}_a$ we must *break the symmetries* of $(X_a, \mathbf{f}_a, \mathbf{G}_a)$. But breaking these symmetries is unnatural, and may cause $\partial(\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \underline{\mathbf{G}}_a])$ to be nonzero. To see this, note that (270) implicitly uses an isomorphism

$$(\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a}) \cong \coprod_{b=1, \dots, n_a^1} (X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1). \quad (271)$$

$\bar{a} = 1, \dots, n_a^1$. Set $(\bar{a}, \bar{b}) = \phi^1(a, b)$

The set of isomorphisms (271) has a free, transitive action of $\prod_b \text{Aut}(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)$, and in general there is no natural choice. To choose $\underline{\mathbf{G}}_a$ for $a \in A$ satisfying boundary relations, we expect (as part of an induction on reverse codimension) to choose $\underline{\mathbf{G}}_{\bar{a}\bar{b}}^1$ for all $(\bar{a}, \bar{b}) \in R^1$, and to use these and (271) to determine boundary values $\underline{\mathbf{G}}_a|_{\partial X_a}$ for $\underline{\mathbf{G}}_a$, and to extend these over X_a to get $\underline{\mathbf{G}}_a$. But to do this we must make an arbitrary choice of isomorphism (271), and similarly in previous inductive steps. There is no reason for these arbitrary choices to be compatible with involutions $\sigma : \partial^2 X_a \rightarrow \partial^2 X_a$. Thus the prescribed values for $\underline{\mathbf{G}}_a|_{\partial X_a}$, when restricted to $\partial^2 X_a$, may not be σ -invariant, and so by Principle 2.8, we cannot extend them over ∂X_a .

To get round this, we choose not one set of effective gauge-fixing data $\underline{\mathbf{G}}_a$ for $(X_a, \mathbf{f}_a)$ for each $a \in A$, but a finite family $\underline{\mathbf{G}}_{ao}$ for $o \in O_a$, such that $\text{Aut}(X_a, \mathbf{f}_a, \mathbf{G}_a)$ acts freely on O_a , and if $(\mathbf{a}, \mathbf{b}) \in \text{Aut}(X_a, \mathbf{f}_a, \mathbf{G}_a)$ and $d \in O_a$ with $o' = (\mathbf{a}, \mathbf{b}) \cdot o$ then $(\mathbf{a}, \mathbf{b})$ lifts to an isomorphism $(\mathbf{a}', \mathbf{b}') : (X_a, \mathbf{f}_a, \underline{\mathbf{G}}_{ao}) \rightarrow (X_a, \mathbf{f}_a, \underline{\mathbf{G}}_{ao'})$. Similarly, for all $m = M, M-1, \dots, 0$ and $(a, b) \in P^m$ we will choose a finite family $\underline{\mathbf{G}}_{abo}^m$ for $o \in O_{ab}^m$ of effective gauge-fixing data for $(X_{ab}^m, \mathbf{f}_{ab}^m)$, where $\text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ acts freely on O_{ab}^m , and lifts to isomorphisms between the $(X_{ab}^m, \mathbf{f}_{ab}^m, \underline{\mathbf{G}}_{abo}^m)$, and $\underline{\mathbf{G}}_{abo}^m|_{\partial X_{ab}^m}$ is determined in terms of the $\underline{\mathbf{G}}_{ab\bar{o}}^{m+1}$.

The construction involves a complex and difficult double induction on decreasing m . The underlying idea is that the families $(X_{ab}^m, \mathbf{f}_{ab}^m, \underline{\mathbf{G}}_{abo}^m)$ for $o \in O_{ab}^m$ have all the symmetries of $(X_a, \mathbf{f}_a, \mathbf{G}_a)$, so the symmetries are not broken, and we need not make arbitrary choices of isomorphisms (271). Using (270) and alternatives (A),(B) above we prove that

$$\partial(\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_{ao}]) = 0 \quad \text{in } KC_{k-1}^{\text{ef}}(Y; R). \quad (272)$$

We then set $\beta = [\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_{ao}]]$ in $KH_k^{\text{ef}}(Y; R)$, and show that $\Pi_{\text{ef}}^{\text{Kh}}(\beta) = \alpha$, so $\Pi_{\text{ef}}^{\text{Kh}}$ is *surjective*, as we want.

Step 5. Suppose $\beta \in KH_k^{\text{ef}}(Y; R)$ with $\Pi_{\text{ef}}^{\text{Kh}}(\beta) = 0$. We choose a cycle in $KC_k^{\text{ef}}(Y; R)$ representing β . As $KH_k^{\text{ef}}(Y; R) \cong H_k^{\text{si}}(Y; R)$ by Theorem 4.8 we can take this to be the image of a cycle $\sum_{s \in S} \zeta_s \tau_s$ in $C_k^{\text{si}}(Y; R)$, for S a finite indexing set, $\zeta_s \in R$ and $\tau_s : \Delta_k \rightarrow Y$ smooth. Thus β is represented by $\Pi_{\text{si}}^{\text{ef}}(\sum_{s \in S} \zeta_s \tau_s)$ in $KC_k^{\text{ef}}(Y; R)$. The image in $KC_k(Y; R)$ under $\Pi_{\text{ef}}^{\text{Kh}}$ is $\Pi_{\text{si}}^{\text{Kh}}(\sum_{s \in S} \zeta_s \tau_s)$, which represents $\Pi_{\text{ef}}^{\text{Kh}}(\beta) = 0$, and so is exact. Therefore as

for (221) there exists $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a] \in KC_{k+1}(Y; R)$ with

$$\partial(\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]) = \Pi_{\text{si}}^{\text{Kh}}(\sum_{s \in S} \zeta_s \tau_s) = \sum_{s \in S} \zeta_s [\Delta_k, \tau_s, \mathbf{G}_{\Delta_k}]. \quad (273)$$

We now apply Steps 1–4 to this $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$, replacing k -cycles by $(k+1)$ -chains. In Step 1 we use tent functions $\mathbf{T}_a$ to ‘cut’ each $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ into small pieces $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ for $c \in C_a$. When this process is applied to (273), the $(\Delta_k, \tau_s, \mathbf{G}_{\Delta_k})$ must also be cut into small pieces. To keep track of these pieces, it is convenient to use the method of Step 4 of §B in §B.4: we choose the $\mathbf{T}_a$ such that each $(\Delta_k, \tau_s, \mathbf{G}_{\Delta_k})$ is cut into the N^{th} barycentric subdivision of Δ_k , for some $N \gg 0$ independent of $s \in S$.

After slightly modifying the definitions of the functions G_{ac}^i in $\mathbf{G}_{ac}$, in Step 1 we construct a new chain $\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$ with all X_{ac} ‘arbitrarily small’, satisfying

$$\partial(\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]) = \Pi_{\text{si}}^{\text{Kh}} \circ \Upsilon^N(\sum_{s \in S} \zeta_s \tau_s) \quad (274)$$

for some $N \gg 0$, where $\Upsilon : C_k^{\text{si}}(Y; R) \rightarrow C_k^{\text{si}}(Y; R)$ is the barycentric subdivision operator of §B.4, so that $\Upsilon^N = \Upsilon \circ \dots \circ \Upsilon$ is the N^{th} barycentric subdivision operator. Note that we will not need the second part of Step 1, which constructs a homology $\sum_{a \in A} \rho_a [Z_{X_a, \mathbf{T}_a}, \mathbf{f}_a \circ \pi_a, \mathbf{H}_{X_a, \mathbf{T}_a}]$ between $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$ and $\sum_{a \in A} \sum_{c \in C_a} \rho_a [X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$, we only have to prove (274).

Going through Steps 2–4 with minor modifications then yields an effective Kuranishi chain $\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \mathbf{G}_{ao}]$ in $KC_{k+1}^{\text{ef}}(Y; R)$ satisfying

$$\partial\left(\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \mathbf{G}_{ao}]\right) = \Pi_{\text{si}}^{\text{ef}} \circ \Upsilon^N(\sum_{s \in S} \zeta_s \tau_s) \quad (275)$$

for some $N \gg 0$, where we change notation so that $A, \rho_a, X_a, \mathbf{f}_a, \mathbf{G}_a, N$ are different from the first versions. Since $\sum_{s \in S} \zeta_s \tau_s$ is a cycle in $C_k^{\text{si}}(Y; R)$, as in §B.4, $\Upsilon^N(\sum_{s \in S} \zeta_s \tau_s)$ is homologous to $\sum_{s \in S} \zeta_s \tau_s$, and thus $\Pi_{\text{si}}^{\text{ef}} \circ \Upsilon^N(\sum_{s \in S} \zeta_s \tau_s)$ is also a cycle in $KH_k^{\text{ef}}(Y; R)$ representing β . Therefore (275) implies that $\beta = 0$, so $\Pi_{\text{ef}}^{\text{Kh}} : KH_k^{\text{ef}}(Y; R) \rightarrow KH_k(Y; R)$ is *injective*.

Remark C.1. (a) We used relation Definition 4.3(iv) in Step 2 to lift to Kuranishi chains with trivial stabilizers, as in (260). But also, much of the work in Steps 1–3 is coping with problems caused by Definition 4.3(iv), and the equivalences between chains that it introduces.

We said in Remark 4.5(b) that Kuranishi homology $KH_*(Y; R)$ would still work, and Theorem 4.9 would still hold, if relation Definition 4.3(iv) were omitted in the definition of $KC_*(Y; R)$, that is, if we replaced $KC_*(Y; R)$ by $\widetilde{KC}_*(Y; R)$ in the notation of Step 1 above. We now briefly sketch how to prove Theorem 4.9 with this modified definition. This time there are four steps:

Step 1. Let $\alpha \in KH_*(Y; R)$. Represent α by a cycle $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$. Show that α can be represented by a cycle $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \tilde{\mathbf{G}}_a]$, in which we fix $X_a, \mathbf{f}_a$ but change to new gauge-fixing data $\tilde{\mathbf{G}}_a = (\tilde{\mathbf{I}}_a, \tilde{\eta}_a,$

$\tilde{G}_a^i : i \in \tilde{I}_a$) with $\tilde{I}_a = (\tilde{I}_a, (\tilde{V}_a^i, \dots, \tilde{\psi}_a^i), \tilde{f}_a^i : i \in \tilde{I}_a, \dots)$ in which each $\tilde{V}_a^i$ is an *effective* orbifold, although we do *not* require $(\tilde{V}_a^i, \dots, \tilde{\psi}_a^i)$ to be an effective Kuranishi neighbourhood, that is, the stabilizer groups of $\tilde{V}_a^i$ need not act trivially on the fibres of $\tilde{E}_a^i$.

Step 2. As in §A.2, we can use a tent function to cut an effective orbifold up into manifolds. Following Step 1 of the proof of Theorem 4.8 in Appendix B, we use the results of §A.3.5 to choose tent functions $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \tilde{\mathbf{G}}_a)$ for $a \in A$ which cut the $\tilde{V}_a^i$ up into manifolds $\tilde{V}_{ac}^i$, and the X_a up into Kuranishi spaces X_{ac} with trivial stabilizers, for $c \in C_a$. Thus we represent α by a cycle $\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \tilde{\mathbf{G}}_{ac}]$ in which the X_{ac} have trivial stabilizers, and the $\tilde{V}_{ac}^i$ in $\tilde{\mathbf{G}}_{ac}$ are manifolds.

Steps 3 and 4. At the end of the new Step 2, we are in exactly the same position as at the end of the old Step 3 above, noting that the new $KC_*(Y; R)$ is the old $\widehat{KC}_*(Y; R)$. We can now follow the old Steps 4 and 5 to complete the proof.

(b) In the proof of Theorem 4.8 we use the fact that R is a $\mathbb{Q}$ -algebra in two different ways: firstly, in (260) of Step 2 the $|\Gamma_{ac}|^{-1}$ must lie in R , and secondly, in (272) of Step 4 the $|O_a|^{-1}$ must lie in R , so for both we need $\mathbb{Q} \subseteq R$.

If as in (a) we omit relation Definition 4.3(iv) then we no longer need R to be a $\mathbb{Q}$ -algebra to define $KC_*(Y; R)$, $KH_*(Y; R)$, and in proving Theorem 4.9 by the method sketched in (a) we would no longer need R to be a $\mathbb{Q}$ -algebra in the first way. However, we would still need R to be a $\mathbb{Q}$ -algebra in the second way, in Step 4. If R were not a $\mathbb{Q}$ -algebra, the proof would fail, and in fact the author can show that $KH_*(Y; R) \cong H_*^{\text{sl}}(Y; R \otimes_{\mathbb{Z}} \mathbb{Q})$ in this case.

In the rest of this appendix we go through Steps 1–5.

C.1 Step 1: cutting the $X_a, \tilde{X}_d$ into small pieces $X_{ac}, \tilde{X}_{df}$

Let $k \in \mathbb{Z}$, $\alpha \in KH_k(Y; R)$, $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a] \in KC_k(Y; R)$ representing α , and $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d), \Gamma_d$ for $d \in D$ satisfying (255) be as in Step 1. Let the indexing sets A, D be disjoint, $A \cap D = \emptyset$. We now explain how to choose tent functions $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ for $a \in A$, and $\tilde{\mathbf{T}}_d$ for $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ for $d \in D$, satisfying conditions (a)–(c) of Step 1, such that $\mathbf{T}_a$ cuts X_a into arbitrarily small pieces X_{ac} for $c \in C_a$, and $\tilde{\mathbf{T}}_d$ cuts $\tilde{X}_d$ into arbitrarily small pieces $\tilde{X}_{df}$ for $f \in F_d$. This construction will be used in Steps 2 and 3. Choosing a tent function $\mathbf{T}$ to cut one $(X, \mathbf{f}, \mathbf{G})$ into arbitrarily small pieces was explained in §A.3.3 and §A.3.5. Our problem is to make these choices satisfy (a)–(c).

We introduce notation similar to $P^m, Q^m, \dots$ in Step 1 of Appendix B. By Lemma 3.7, for each $a \in A$ and $m \geq 0$ the triple $(\partial^m X_a, \mathbf{f}_a|_{\partial^m X_a}, \mathbf{G}_a|_{\partial^m X_a})$ splits as $\partial^m X_a = X_{a1}^m \amalg \dots \amalg X_{an_a^m}^m$ into *connected* triples $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) = (X_{ab}^m, \mathbf{f}_a|_{X_{ab}^m}, \mathbf{G}_a|_{X_{ab}^m})$ for $b = 1, \dots, n_a^m$. Similarly, for $d \in D$ and $m \geq 1$ the triple $(\partial^{m-1} \tilde{X}_d, \tilde{\mathbf{f}}_d|_{\partial^{m-1} \tilde{X}_d}, \tilde{\mathbf{G}}_d|_{\partial^{m-1} \tilde{X}_d})$ splits as $\partial^{m-1} \tilde{X}_d = X_{d1}^m \amalg \dots \amalg X_{dn_d^m}^m$

into connected triples $(X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m) = (X_{de}^m, \tilde{\mathbf{f}}_d|_{X_{de}^m}, \tilde{\mathbf{G}}_d|_{X_{de}^m})$ for $e = 1, \dots, \tilde{n}_d^m$. Also, for $d \in D$ and $m \geq 1$ the triple $(\partial^{m-1}(\tilde{X}_d/\Gamma_d), \pi_*(\tilde{\mathbf{f}}_d)|_{\partial^{m-1}(\tilde{X}_d/\Gamma_d)}, \pi_*(\tilde{\mathbf{G}}_d)|_{\partial^{m-1}(\tilde{X}_d/\Gamma_d)})$ splits as $\partial^{m-1}(\tilde{X}_d/\Gamma_d) = X_{d,-1}^m \amalg X_{d,-2}^m \amalg \dots \amalg X_{d,-\hat{n}_d^m}^m$ into connected $(X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m) = (X_{de}^m, \tilde{\mathbf{f}}_d|_{X_{de}^m}, \tilde{\mathbf{G}}_d|_{X_{de}^m})$ for $e = -1, -2, \dots, -\hat{n}_d^m$.

Note that this choice of notation may at first seem odd: in $X_{de}^m \subseteq \partial^{m-1}\tilde{X}_d$ and $X_{de}^m \subseteq \partial^{m-1}(\tilde{X}_d/\Gamma_d)$ we are changing from $m-1$ to m , and dropping accents ' $\sim$ '. The three sets of $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ and $(X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m)$ are distinguished as $A \cap D = \emptyset$, so the subscripts ' ab ' and ' de ' never coincide, and $X_{de} \subseteq \partial^{m-1}\tilde{X}_d$ have $e > 0$, and $X_{de}^m \subseteq \partial^{m-1}(\tilde{X}_d/\Gamma_d)$ have $e < 0$.

Define $\ddot{P}^m = \{(a, b) : a \in A, b = 1, \dots, n_a^m\} \cup \{(d, e) : d \in D, e = 1, \dots, \tilde{n}_d^m\} \cup \{(d, e) : d \in D, e = -1, -2, \dots, -\hat{n}_d^m\}$, omitting the second and third sets when $m = 0$. We write a general element of $\ddot{P}^m$ as (s, t) . Thus $(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$ can mean either $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$, a component of $\partial^m(X_a, \mathbf{f}_a, \mathbf{G}_a)$ for $a \in A$, or $(X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m)$, a component of $\partial^{m-1}(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ for $d \in D$ and $e > 0$ when $m \geq 1$, or $(X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m)$, a component of $\partial^{m-1}(\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d))$ for $d \in D$ and $e < 0$ when $m \geq 1$.

Since A, D are finite and $\partial^m X_a = \partial^{m-1}\tilde{X}_d = \emptyset$ for $m \gg 0$, $\ddot{P}^m$ is finite with $\ddot{P}^m = \emptyset$ for $m \gg 0$. Let $M \geq 0$ be largest with $\ddot{P}^M \neq \emptyset$. If $(s, t) \in \ddot{P}^m$ then $(\partial X_{st}^m, \mathbf{f}_{st}^m|_{\partial X_{st}^m}, \mathbf{G}_{st}^m|_{\partial X_{st}^m})$ is a union of connected components each of which is of the form $(X_{st'}^{m+1}, \mathbf{f}_{st'}^{m+1}, \mathbf{G}_{st'}^{m+1})$ for $(s, t') \in \ddot{P}^{m+1}$. Write $\ddot{B}_{st}^m$ for the set of such t' . Then $\partial X_{st}^m = \coprod_{t' \in \ddot{B}_{st}^m} X_{st'}^{m+1}$.

By induction on decreasing $m = M, M-1, \dots, 1, 0$ we will choose tent functions $\mathbf{T}_{st}^m$ for $(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$ for all $(s, t) \in \ddot{P}^m$, satisfying the conditions:

- (i) If $(s, t), (s', t') \in \ddot{P}^m$ and $(\mathbf{a}, \mathbf{b}) : (X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m) \rightarrow (X_{s't'}^m, \mathbf{f}_{s't'}^m, \mathbf{G}_{s't'}^m)$ is an isomorphism, not necessarily orientation-preserving, then $\mathbf{T}_{st}^m = \mathbf{b}^*(\mathbf{T}_{s't'}^m)$. That is, $T_{st}^{m,i} = T_{s't'}^{m,i} \circ b^i$ for all $i \in I_{st}^m$, where $b^i : V_{st}^{m,i} \rightarrow V_{s't'}^{m,i}$ is in $\mathbf{b}$ and $T_{st}^{m,i}, T_{s't'}^{m,i}$ are the tent functions on $V_{st}^{m,i}, V_{s't'}^{m,i}$ in $\mathbf{T}_{st}^m, \mathbf{T}_{s't'}^m$.
- (ii) Suppose $m \geq 1$, $d \in D$, $e = 1, \dots, \tilde{n}_d^m$ and $e = -1, -2, \dots, -\hat{n}_d^m$ with $X_{de'}^m = \pi^{m-1}(X_{de}^m)$. Then Γ_d acts on the indexing set $\{1, \dots, n_d^m\}$ of components of $\partial^{m-1}(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$. Write $\text{Stab}_{\Gamma_d}(e)$ for the stabilizer subgroup of $e \in \{1, \dots, n_d^m\}$ in Γ_d . Then $\text{Stab}_{\Gamma_d}(e)$ acts on $(X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m)$ by automorphisms, and the quotient $(X_{de}^m/\text{Stab}_{\Gamma_d}(e), \pi_*^m(\mathbf{f}_{de}^m), \pi_*^m(\mathbf{G}_{de}^m))$ is $(X_{de'}^m, \mathbf{f}_{de'}^m, \mathbf{G}_{de'}^m)$. Write $(\pi^m, \mathbf{p}^m) : (X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m) \rightarrow (X_{de'}^m, \mathbf{f}_{de'}^m, \mathbf{G}_{de'}^m)$ for the corresponding projection, so that $\pi^m : X_{de}^m \rightarrow X_{de'}^m$ is a strong submersion, and $p^{m,i} : V_{de}^{m,i} \rightarrow V_{de'}^{m,i}/\text{Stab}_{\Gamma_d}(e) = V_{de'}^{m,i}$ is the natural projection. Then $\mathbf{T}_{de}^m = (\mathbf{p}^m)^*(\mathbf{T}_{de'}^m)$, that is, $T_{de}^{m,i} = T_{de'}^{m,i} \circ p^{m,i}$ for all $i \in I_{de}^m$.
- (iii) If $(s, t) \in \ddot{P}^m$ then $(\partial X_{st}^m, \mathbf{f}_{st}^m|_{\partial X_{st}^m}, \mathbf{G}_{st}^m|_{\partial X_{st}^m}) = \coprod_{t' \in \ddot{B}_{st}^m} (X_{st'}^{m+1}, \mathbf{f}_{st'}^{m+1}, \mathbf{G}_{st'}^{m+1})$. We require $\min \mathbf{T}_{st}^m|_{\partial X_{st}^m} = \coprod_{t' \in \ddot{B}_{st}^m} \min \mathbf{T}_{st'}^{m+1}$ under this.
- (iv) Each $\mathbf{T}_{st}^m$ cuts $(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$ into 'arbitrarily small pieces'.

Note that taking $(s, t) = (s', t')$ in (i) implies that $\mathbf{T}_{st}^m$ is invariant under $\text{Aut}(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$ for all $(s, t) \in \ddot{P}^m$. Using this we will see later that (i), (ii)

imply (a),(b) in Step 1, and clearly (iv) implies (c).

We will carry out the induction to choose the $\mathbf{T}_{st}^m$ after Remark C.6. First we have more notation to define, and some work to do. Define an equivalence relation $\sim^m$ on $\check{P}^m$ by $(s, t) \sim^m (s', t')$ if there exists an isomorphism $(\mathbf{a}, \mathbf{b}) : (X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m) \rightarrow (X_{s't'}^m, \mathbf{f}_{s't'}^m, \mathbf{G}_{s't'}^m)$, where $\mathbf{a}$ need not identify orientations.

Define a second, weaker equivalence relation $\approx^m$ on $\check{P}^m$ to be the equivalence relation generated by $\sim^m$ and the additional relations when $m \geq 1$ that $(d, e) \approx^m (d, e')$ if $e > 0$ and $e' < 0$, so that $X_{de} \subseteq \partial^{m-1} \tilde{X}_d$ and $X_{de'}^m \subseteq \partial^{m-1}(\tilde{X}_d/\Gamma_d)$, and $X_{de'}^m = \pi^m(X_{de}^m)$, where $\pi^m : \partial^{m-1} \tilde{X}_d \rightarrow \partial^{m-1}(\tilde{X}_d/\Gamma_d)$ is the continuous map associated to $\pi^m : \partial^{m-1} \tilde{X}_d \rightarrow \partial^{m-1}(\tilde{X}_d/\Gamma_d)$ induced by the projection $\pi : \tilde{X}_d \rightarrow \tilde{X}_d/\Gamma_d$. For each (d, e) the image of $(X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m)$ under π is a connected component of $\partial^{m-1}(\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d))$, so for each $d \in D$ and $e = 1, \dots, \tilde{n}_d^m$ there is a unique $e' = -1, \dots, -\tilde{n}_d^m$ with $X_{de'}^m = \pi^m(X_{de}^m)$.

The relevance of $\sim^m, \approx^m$ is that if $(s, t) \sim^m (s', t')$ then (i) above implies that $\mathbf{T}_{st}^m = \mathbf{b}^*(\mathbf{T}_{s't'}^m)$. More generally, if $(s, t) \approx^m (s', t')$ then by a finite number of applications of (i),(ii) above we can determine $\mathbf{T}_{st}^m$ in terms of $\mathbf{T}_{s't'}^m$, and vice versa. Thus, the equivalence relations $\sim^m$ and $\approx^m$ correspond to the equivalences imposed on the $\mathbf{T}_{st}^m$ by (i) and by (i),(ii) respectively, *on the level of the indexing set $\check{P}^m$* . We can also define such equivalence relations on the levels of *topological spaces* $\coprod_{(s,t) \in \check{P}^m} X_{st}^m$, and *orbifolds* $\coprod_{(s,t) \in \check{P}^m : i \in I_{st}^m} V_{st}^{m,i}$.

Let $\sim_x^m$ be the equivalence relation on $\coprod_{(s,t) \in \check{P}^m} X_{st}^m$, as a set, given by $x \sim_x^m x'$ if $x \in X_{st}^m, x' \in X_{s't'}^m$, and there exists an isomorphism $(\mathbf{a}, \mathbf{b}) : (X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m) \rightarrow (X_{s't'}^m, \mathbf{f}_{s't'}^m, \mathbf{G}_{s't'}^m)$ with $a(x) = x'$, where $a : X_{st}^m \rightarrow X_{s't'}^m$ is the homeomorphism associated to $\mathbf{a}$. Let $\approx_x^m$ be the equivalence relation on $\coprod_{(s,t) \in \check{P}^m} X_{st}^m$ generated by $\sim_x^m$ and the additional relations when $m \geq 1$ and d, e, e' and $(\pi^m, \mathbf{p}^m) : (X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m) \rightarrow (X_{de'}^m, \mathbf{f}_{de'}^m, \mathbf{G}_{de'}^m)$ are as in (ii) above and $x \in X_{de}^m, x' \in X_{de'}^m$ with $\pi^m(x) = x'$, where $\pi^m : X_{de}^m \rightarrow X_{de'}^m$ is the continuous map associated to π^m , then $x \approx_x^m x'$. Define $\check{X}^m$ to be $(\coprod_{(s,t) \in \check{P}^m} X_{st}^m) / \approx_x^m$, the set of equivalence classes of $\approx_x^m$, with the quotient topology, that is, $\check{U} \subseteq \check{X}^m$ is open if its preimage $U \subseteq \coprod_{(s,t) \in \check{P}^m} X_{st}^m$ is open. Write $\pi_{\check{X}^m} : \coprod_{(s,t) \in \check{P}^m} X_{st}^m \rightarrow \check{X}^m$ for the natural, continuous projection.

Similarly, let $\sim_v^{m,i}$ be the equivalence relation on $\coprod_{(s,t) \in \check{P}^m : i \in I_{st}^m} V_{st}^{m,i}$ for each $i \in \mathbb{N}$ given by $v \sim_v^{m,i} v'$ if $v \in V_{st}^{m,i}, v' \in V_{s't'}^{m,i}$, and there exists an isomorphism $(\mathbf{a}, \mathbf{b}) : (X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m) \rightarrow (X_{s't'}^m, \mathbf{f}_{s't'}^m, \mathbf{G}_{s't'}^m)$ with $b^i(v) = v'$. Let $\approx_v^{m,i}$ be the equivalence relation on $\coprod_{(s,t) \in \check{P}^m : i \in I_{st}^m} V_{st}^{m,i}$ generated by $\sim_v^{m,i}$ and the additional relations when $m \geq 1$ and d, e, e' and $(\pi^m, \mathbf{p}^m) : (X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m) \rightarrow (X_{de'}^m, \mathbf{f}_{de'}^m, \mathbf{G}_{de'}^m)$ are as in (ii) above and $v \in V_{de}^{m,i}, v' \in V_{de'}^{m,i}$ with $p^{m,i}(v) = v'$, then $v \approx_v^{m,i} v'$. Define $\check{V}^{m,i} = (\coprod_{(s,t) \in \check{P}^m : i \in I_{st}^m} V_{st}^{m,i}) / \approx_v^{m,i}$, as a topological space. Write $\pi_{\check{V}^{m,i}} : \coprod_{(s,t) \in \check{P}^m : i \in I_{st}^m} V_{st}^{m,i} \rightarrow \check{V}^{m,i}$ for the projection.

In the same way, we can define $\approx_E^{m,i}$ on $\coprod_{(s,t) \in \check{P}^m : i \in I_{st}^m} E_{st}^{m,i}$, and $\check{E}^{m,i} = (\coprod_{(s,t) \in \check{P}^m : i \in I_{st}^m} E_{st}^{m,i}) / \approx_E^{m,i}$. Since the isomorphisms $(\mathbf{a}, \mathbf{b})$ and projections $(\pi^m, \mathbf{p}^m)$ used to generate $\approx_x^m, \approx_v^{m,i}, \approx_E^{m,i}$ are compatible with $s_{st}^{m,i}, \psi_{st}^{m,i}$ in Ku-

ranishi neighbourhoods $(V_{st}^{m,i}, \dots, \psi_{st}^{m,i})$ and coordinate changes between these neighbourhoods, and the data $\eta_i^{m,j}$ used to define open sets $\dot{V}^{m,i}, \dot{V}^{m,ij}$, all these descend to the topological spaces $\dot{X}^m, \dot{V}^{m,i}, \dot{E}^{m,i}$.

That is, $\dot{E}^{m,i} \rightarrow \dot{V}^{m,i}$ is a ‘bundle’ in some weak sense, with a well-defined ‘zero section’, and we have continuous section $\check{s}^{m,i} : \dot{V}^{m,i} \rightarrow \dot{E}^{m,i}$ which lifts to $s_{st}^{m,i} : V_{st}^{m,i} \rightarrow E_{st}^{m,i}$ for each $(s, t) \in \dot{P}^m$, and a continuous map $\check{\psi}^{m,i} : (\check{s}^{m,i})^{-1}(0) \rightarrow \dot{X}^m$ which is a homeomorphism with its image. There is an open set $\check{V}^{m,i} \subset \dot{V}^{m,i}$ which lifts to $\coprod_{(s,t) \in \dot{P}^m : i \in I_{st}^m} \dot{V}_{st}^{m,i} \subset \coprod_{(s,t) \in \dot{P}^m : i \in I_{st}^m} V_{st}^{m,i}$.

When $j \leq i$ we have an open set $\check{V}^{m,ij} \subset \check{V}^{m,j}$ which lifts to $V_{st}^{m,ij}$ in each $V_{st}^{m,j}$, and embeddings of topological spaces $\check{\phi}^{m,ij} : \check{V}^{m,ij} \rightarrow \check{V}^{m,i}, \hat{\phi}^{m,ij} : \dot{E}^{m,ij} \rightarrow \dot{E}^{m,i}$, which lift to $(V_{st}^{m,ij}, \phi_{st}^{m,ij}, \hat{\phi}_{st}^{m,ij})$ on $V_{st}^{m,j}$. Also we have an open set $\check{V}^{m,ij} \subset \check{V}^{m,i}$ which lifts to $V_{st}^{m,ij}$ in each $V_{st}^{m,i}$.

Having got this far, it seems obvious (though perhaps optimistic) to hope that $\dot{X}^m$ is a compact Kuranishi space, $\dot{V}^{m,i}, \dot{E}^{m,i}$ are orbifolds, $(\dot{V}^{m,i}, \dot{E}^{m,i}, \check{s}^{m,i}, \check{\psi}^{m,i})$ is a Kuranishi neighbourhood on $\dot{X}^m$, $(\check{\phi}^{m,ij}, \hat{\phi}^{m,ij})$ is a coordinate change between Kuranishi neighbourhoods, and all the other data in $\mathbf{f}_{st}^m, \mathbf{G}_{st}^m$ descends to $\dot{X}^m, \dot{V}^{m,i}, \dot{E}^{m,i}$ to define a triple $(\dot{X}^m, \check{\mathbf{f}}^m, \check{\mathbf{G}}^m)$ in Kuranishi homology. Then to choose data $\mathbf{T}_{st}^m$ satisfying (i), (ii) above for all $(s, t) \in \dot{P}^m$, we could choose a tent function $\check{\mathbf{T}}^m$ for $(\dot{X}^m, \check{\mathbf{f}}^m, \check{\mathbf{G}}^m)$ using the results of §A.3, and lift it to each $V_{st}^{m,i}$ using the projections $V_{st}^{m,i} \rightarrow \dot{V}^{m,i}$. The construction of $\approx_v^{m,i}$ would then imply (i), (ii).

It turns out that this does not quite work, for rather subtle reasons to do with the distinction between effective and non-effective orbifolds, that will be explained in Remark C.6. We can give $\dot{V}^{m,i}, \dot{E}^{m,i}$ the structure of effective orbifolds in a canonical way, but $\check{\pi}^{m,i} : \dot{E}^{m,i} \rightarrow \dot{V}^{m,i}$ may fail to be an orbibundle, and $\check{\phi}^{m,ij} : \dot{V}^{m,ij} \rightarrow \dot{V}^{m,i}$ may not be an isomorphism on stabilizer groups. So we cannot make $\dot{X}^m$ into a Kuranishi space. However, these problems will not matter when we apply the constructions of §A.3, so we can still choose $\check{\mathbf{T}}^m$ for $(\dot{X}^m, \check{\mathbf{f}}^m, \check{\mathbf{G}}^m)$ and lift to $\mathbf{T}_{st}^m$ for each $(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$.

We will need notions of (weak) *finite covers* for manifolds and orbifolds. Covering maps, fundamental group, and universal covers for topological spaces are discussed by Bredon [10, §III], and for orbifolds by Adem et al. [1, §2.2].

Definition C.2. Let V, W be manifolds of the same dimension, which may have boundary, corners, or g-corners, and $c : V \rightarrow W$ a smooth map. We call c a *finite cover* if it is a surjective, proper local diffeomorphism. Here c is *proper* means that if $S \subseteq W$ is compact in W then $c^{-1}(S)$ is compact in V , and c is a *local diffeomorphism* means that every $v \in V$ admits an open neighbourhood U in V such that $c(U)$ is open in W and $c|_U : U \rightarrow c(U)$ is a diffeomorphism.

Let $c : V \rightarrow W$ be a finite cover of *connected* manifolds V, W . Pick $v_0 \in V$, and set $w_0 = c(v_0) \in W$. We can form the *fundamental groups* $\pi_1(V, v_0)$, $\pi_1(W, w_0)$ and *universal covers* $\tilde{V}, \tilde{W}$ of V, W with base-points v_0, w_0 . Then $c : (V, v_0) \rightarrow (W, w_0)$ induces morphisms of fundamental groups $c_* : \pi_1(V, v_0) \rightarrow \pi_1(W, w_0)$, and $\tilde{c} : \tilde{V} \rightarrow \tilde{W}$ of universal covers. The important fact is that $c_* : \pi_1(V, v_0) \rightarrow \pi_1(W, w_0)$ is *injective*, and $\tilde{c} : \tilde{V} \rightarrow \tilde{W}$ is a *diffeomorphism*. Thus we may identify $\pi_1(V, v_0)$ with a subgroup of $\pi_1(W, w_0)$, and $\tilde{V}$

with $\tilde{W}$. So we have diffeomorphisms $W \cong \tilde{W}/\pi_1(W, w_0)$ and $V \cong \tilde{W}/\pi_1(V, v_0)$. One class of examples of finite covers of manifolds are *free finite group quotients*. Let V be a manifold and Γ a finite group acting freely on V by diffeomorphisms. Then $W = V/\Gamma$ is a manifold, and the projection $\pi : V \rightarrow V/\Gamma$ is a finite cover.

Definition C.3. Let V, W be orbifolds of the same dimension n , which may have boundary, corners, or g-corners, and $c : V \rightarrow W$ a smooth map. We call c a *finite cover* if the following condition holds. Suppose (U, Γ, ϕ) is an orbifold chart on W . Write $\pi : U \rightarrow U/\Gamma$ for the projection. Then $\phi \circ \pi : U \rightarrow W$ is a submersion, and $c : V \rightarrow W$ is a smooth map, so we can form the orbifold fibre product $U \times_{\phi \circ \pi, W, c} V$, which is an n -orbifold with smooth projection $\pi_U : U \times_{\phi \circ \pi, W, c} V \rightarrow U$. We require that for all such (U, Γ, ϕ) , $U \times_{\phi \circ \pi, W, c} V$ is an n -manifold, and $\pi_U : U \times_{\phi \circ \pi, W, c} V \rightarrow U$ is a finite cover of manifolds.

An important class of examples of finite covers of orbifolds are *finite group quotients*. Let V be an orbifold, and Γ a finite group acting on V by diffeomorphisms; note that Γ need not act freely, nor even effectively. Then $W = V/\Gamma$ is an orbifold, and the natural projection $\pi : V \rightarrow V/\Gamma$ is a finite cover of orbifolds.

As for finite covers of manifolds, one can also describe finite covers of connected orbifolds in terms of fundamental groups and universal covers. However, these are not the usual fundamental groups and universal covers of topological spaces [10, §III], but special orbifold versions, the *orbifold fundamental group* $\pi_1^{\text{orb}}(V, v_0)$ and *orbifold universal cover* $\tilde{V}^{\text{orb}}$ of a connected orbifold V . These are defined by Adem et al. [1, §2.2] using their language of groupoids, and the notion of classifying space of an orbifold.

One can also define $\pi_1^{\text{orb}}(V, v_0)$ in a similar way to the usual definition of $\pi_1(X)$ for a topological space X , in terms of isotopy classes of *smooth* paths $\gamma : [0, 1] \rightarrow V$ with $\gamma(0) = \gamma(1) = v_0$. However, at this point the discussion of Remark 2.12 becomes important. We must interpret $\gamma : [0, 1] \rightarrow V$, a smooth map of orbifolds, not in the weak sense of Satake, but as a *strong map* [57], *good map* [12] or *orbifold morphisms* [1]. Thus, γ carries with it some discrete extra data, which enables us to make natural choices of maps σ_{pq} in Definition 2.11. This discrete extra data makes the difference between $\pi_1^{\text{orb}}(V, v_0)$ and $\pi_1(V, v_0)$.

For example, let Γ be a finite subgroup of $\text{GL}(n, \mathbb{R})$. Then $\mathbb{R}^n/\Gamma$ is a connected orbifold. Choose 0Γ as the basepoint. As a topological space $\mathbb{R}^n/\Gamma$ is contractible, so $\pi_1(\mathbb{R}^n/\Gamma, 0\Gamma) = \{1\}$, and the universal cover is $\mathbb{R}^n/\Gamma$. But $\pi_1^{\text{orb}}(\mathbb{R}^n/\Gamma, 0\Gamma) = \Gamma$, with orbifold universal cover $(\widetilde{\mathbb{R}^n/\Gamma})^{\text{orb}} = \mathbb{R}^n$.

As for manifolds, $\tilde{V}^{\text{orb}}$ is a connected orbifold which is unique up to diffeomorphism with $\pi_1^{\text{orb}}(\tilde{V}^{\text{orb}}, \tilde{v}_0) = \{1\}$, and $\pi_1^{\text{orb}}(V, v_0)$ acts on $\tilde{V}^{\text{orb}}$ by diffeomorphisms (though not necessarily freely), with V naturally diffeomorphic to $\tilde{V}^{\text{orb}}/\pi_1^{\text{orb}}(V, v_0)$. If $c : V \rightarrow W$ is a finite cover of orbifolds with V, W connected (in the usual sense of topological spaces), $v_0 \in V$, and $w_0 = c(v_0) \in W$, then $c : (V, v_0) \rightarrow (W, w_0)$ induces an injective morphism of orbifold fundamental groups $c_* : \pi_1^{\text{orb}}(V, v_0) \rightarrow \pi_1^{\text{orb}}(W, w_0)$, and a diffeomorphism $\tilde{c} : \tilde{V}^{\text{orb}} \rightarrow \tilde{W}^{\text{orb}}$ of orbifold universal covers. Thus we may identify $\pi_1^{\text{orb}}(V, v_0)$ with a subgroup of $\pi_1^{\text{orb}}(W, w_0)$, and $\tilde{V}^{\text{orb}}$ with $\tilde{W}^{\text{orb}}$, and we have diffeomorphisms

$W \cong \tilde{W}^{\text{orb}}/\pi_1^{\text{orb}}(W, w_0)$ and $V \cong \tilde{W}^{\text{orb}}/\pi_1^{\text{orb}}(V, v_0)$.

The orbifold universal cover $\tilde{V}^{\text{orb}}$ of V is always an *effective* orbifold. Write $\text{Diff}(\tilde{V}^{\text{orb}})$ for the group of diffeomorphisms of $\tilde{V}^{\text{orb}}$. Then $\pi_1^{\text{orb}}(V, v_0)$ acts on $\tilde{V}^{\text{orb}}$ by a group morphism $\rho : \pi_1^{\text{orb}}(V, v_0) \rightarrow \text{Diff}(\tilde{V}^{\text{orb}})$. The kernel $\text{Ker } \rho$ is a finite group, isomorphic to the stabilizer group of a generic point of V . Now V is an *effective* orbifold if and only if generic points in V have trivial stabilizers. Thus, V is effective if and only if ρ is injective. Therefore, *if a connected orbifold V is effective, we can regard $\pi_1^{\text{orb}}(V, v_0)$ as a subgroup of $\text{Diff}(\tilde{V}^{\text{orb}})$, but if V is not effective, we cannot.* This will be important in Proposition C.5.

Definition C.4. Let V, W be orbifolds, and $c : V \rightarrow W$ a smooth map. Then as in Definition 2.11, for all $v \in V$ and $w = c(v)$ in W there exist orbifold charts (U, Γ, ϕ) on V and (U', Γ', ϕ') on W with $v \in \text{Im } \phi$ and $w \in \text{Im } \phi'$, a morphism of finite groups $\rho : \Gamma \rightarrow \Gamma'$, and a ρ -equivariant smooth map $\sigma : U \rightarrow U'$, such that $c \circ \phi(\Gamma u) = \phi'(\Gamma' \sigma(u))$ for all $u \in U$. We can also make U' smaller if necessary so that $c(\text{Im } \phi) = \text{Im } \phi'$.

We call c a *weak finite cover* if c is surjective and proper, and for all $v \in V$ we can choose $(U, \Gamma, \phi), (U', \Gamma', \phi'), \rho, \sigma$ above such that $\sigma : U \rightarrow U'$ is a finite cover of manifolds. If we also required $\rho : \Gamma \rightarrow \Gamma'$ to be injective then this would be equivalent to the definition of finite covers of orbifolds in Definition C.3. For weak finite covers we allow ρ not to be injective. As σ is ρ -equivariant, $\text{Ker } \rho$ must act trivially on U . Thus, if ρ is not injective then the nontrivial group $\text{Ker } \rho$ is contained in the stabilizer group of every point in the nonempty open set $\phi(U/\Gamma)$ in V , so V is a *non-effective* orbifold. Therefore a weak finite cover $c : V \rightarrow W$ is a finite cover if V is effective.

If $c : V \rightarrow W$ is a weak finite cover with V, W connected then choosing $v_0 \in V$ and setting $w_0 = c(v_0)$, we can form orbifold fundamental groups $\pi_1^{\text{orb}}(V, v_0), \pi_1^{\text{orb}}(W, w_0)$ and orbifold universal covers $\tilde{V}^{\text{orb}}, \tilde{W}^{\text{orb}}$, and we get a group morphism $c_* : \pi_1^{\text{orb}}(V, v_0) \rightarrow \pi_1^{\text{orb}}(W, w_0)$ and a smooth map $\tilde{c} : \tilde{V}^{\text{orb}} \rightarrow \tilde{W}^{\text{orb}}$. As for finite covers of orbifolds, it turns out that $\tilde{c}$ is a *diffeomorphism*. However, in the weak case, $c_* : \pi_1^{\text{orb}}(V, v_0) \rightarrow \pi_1^{\text{orb}}(W, w_0)$ *may not be injective*. Instead, if $v \in V$ is a generic point and $w = c(v)$ (which is generic in W) then $\text{Ker } c_*$ is isomorphic to the kernel of the induced morphism of stabilizer groups $\text{Stab}_V(v) \rightarrow \text{Stab}_W(w)$. Note that V is effective if and only if $\text{Stab}_V(v) = \{1\}$, and then $\text{Ker } c_*$ is trivial, and c is a finite cover.

We will show that the $\tilde{V}^{m,i}$ can be given the structure of effective orbifolds. The proof uses the finiteness properties of the gauge-fixing data $\mathbf{G}_{st}^m$.

Proposition C.5. *In the situation above, there is a unique way to give $\tilde{V}^{m,i}$ the structure of an effective i -orbifold, such that $\pi_{\tilde{V}^{m,i}} : \coprod_{(s,t) \in \tilde{P}^m : i \in I_{st}^m} V_{st}^{m,i} \rightarrow \tilde{V}^{m,i}$ lifts to a weak finite cover of orbifolds for each $i \in \mathbb{N}$. Furthermore:*

- (a) $\check{\phi}^{m,ij} : \tilde{V}^{m,ij} \rightarrow \tilde{V}^{m,i}$ lifts to a smooth, injective map of orbifolds for $j \leq i$ in $\mathbb{N}$, and $d\check{\phi}^{m,ij} : T\tilde{V}^{m,ij} \rightarrow (\check{\phi}^{m,ij})^*(T\tilde{V}^{m,i})$ is injective. However, $\check{\phi}^{m,ij}$ need not be an isomorphism on stabilizer groups, and so may not be an embedding in the sense of Definition 2.11. For $k \leq j \leq i$ we have $\check{\phi}^{m,ij} \circ \check{\phi}^{m,jk} = \check{\phi}^{m,ik}$ on $(\check{\phi}^{m,jk})^{-1}(\tilde{V}^{m,ij}) \cap \tilde{V}^{m,ik}$.

(b) We can also relate $\partial\check{V}^{m,i}$ to $\check{V}^{m+1,i-1}$. By definition we have

$$\partial[\coprod_{(s,t) \in \check{P}^m: i \in I_{st}^m} V_{st}^{m,i}] = \coprod_{(s',t') \in \check{P}^{m+1}: i-1 \in I_{s't'}^{m+1}} V_{s't'}^{m+1,i-1}. \quad (276)$$

The equivalence relation $\approx_V^{m,i}$ on $\coprod_{(s,t) \in \check{P}^m: i \in I_{st}^m} V_{st}^{m,i}$ induces $\approx_{\partial V}^{m,i}$ on the l.h.s. of (276), with homeomorphism $(\partial[\coprod_{(s,t) \in \check{P}^m: i \in I_{st}^m} V_{st}^{m,i}]) / \approx_{\partial V}^{m,i} \cong \partial\check{V}^{m,i}$, such that $\pi_{\check{V}^{m,i}}|_{\partial(\dots)} : \partial(\coprod_{(s,t) \in \check{P}^m: i \in I_{st}^m} V_{st}^{m,i}) \rightarrow \partial\check{V}^{m,i}$ is the projection from (276) to its quotient by $\approx_{\partial V}^{m,i}$.

The equivalence relation $\approx_V^{m+1,i-1}$ on the right hand side of (276) is stronger than $\approx_{\partial V}^{m,i}$, that is, $v \approx_{\partial V}^{m,i} v'$ implies $v \approx_V^{m+1,i-1} v'$. Thus, there is a natural, continuous, surjective projection

$$(\partial[\coprod_{\substack{(s,t) \in \check{P}^m \\ i \in I_{st}^m}} V_{st}^{m,i}]) / \approx_{\partial V}^{m,i} \longrightarrow (\coprod_{\substack{(s',t') \in \check{P}^{m+1} \\ i-1 \in I_{s't'}^{m+1}}} V_{s't'}^{m+1,i-1}) / \approx_V^{m+1,i-1},$$

that is, a projection $\partial\check{V}^{m,i} \rightarrow \check{V}^{m+1,i-1}$. This lifts to a weak finite cover of orbifolds $\tilde{\pi}_{\partial}^m : \partial\check{V}^{m,i} \rightarrow \check{V}^{m+1,i-1}$ with $\tilde{\pi}_{\partial}^m \circ \pi_{\check{V}^{m,i}}|_{\partial(\dots)} \equiv \pi_{\check{V}^{m+1,i-1}}$.

Proof. For each $(s,t) \in \check{P}^m$ and $i \in I_{st}^m$, the gauge-fixing data $\mathbf{G}_{st}^m$ includes a map $G_{st}^{m,i} : E_{st}^{m,i} \rightarrow P$, which is globally finite with constant $N_{st}^{m,i}$, say. Identifying $V_{st}^{m,i}$ with the zero section in $E_{st}^{m,i}$, we can regard $V_{st}^{m,i}$ as a suborbifold of $E_{st}^{m,i}$, and so restrict $G_{st}^{m,i}$ to $V_{st}^{m,i}$. Then $G_{st}^{m,i}|_{V_{st}^{m,i}}$ is also globally finite with constant $N_{st}^{m,i}$. Thus we can consider the map

$$\coprod_{(s,t) \in \check{P}^m: i \in I_{st}^m} G_{st}^{m,i}|_{V_{st}^{m,i}} : \coprod_{(s,t) \in \check{P}^m: i \in I_{st}^m} V_{st}^{m,i} \longrightarrow P. \quad (277)$$

It is a globally finite map, with constant $N = \sum_{(s,t) \in \check{P}^m: i \in I_{st}^m} N_{st}^{m,i}$.

Now the equivalence relation $\approx_V^{m,i}$ on $\coprod_{(s,t) \in \check{P}^m: i \in I_{st}^m} V_{st}^{m,i}$ is generated by maps $b^i : V_{st}^{m,i} \rightarrow V_{s't'}^{m,i}$ from isomorphisms $(\mathbf{a}, \mathbf{b})$, and maps $p^{m,i} : V_{de}^{m,i} \rightarrow V_{de'}^{m,i}$ from quotient maps $(\pi^m, \mathbf{p}^m)$. The definitions of $(\mathbf{a}, \mathbf{b})$ and $(\pi^m, \mathbf{p}^m)$ imply that $G_{s't'}^{m,i}|_{V_{s't'}^{m,i}} \circ b^i \equiv G_{st}^{m,i}|_{G_{st}^{m,i}}$ and $G_{de'}^{m,i}|_{V_{de'}^{m,i}} \circ p^{m,i} \equiv G_{de}^{m,i}|_{V_{de}^{m,i}}$. Hence the maps $G_{st}^{m,i}|_{G_{st}^{m,i}}$ are preserved by $\approx_V^{m,i}$, and descend to the quotient $\check{V}^{m,i}$. Thus, there exists $\check{G}^{m,i} : \check{V}^{m,i} \rightarrow P$ such that (277) is $\check{G}^{m,i} \circ \pi_{\check{V}^{m,i}}$. But as (277) is globally finite with constant N , it follows that $\pi_{\check{V}^{m,i}}$ is also globally finite with constant N . That is, each $\approx_V^{m,i}$ equivalence class in $\coprod_{(s,t) \in \check{P}^m: i \in I_{st}^m} V_{st}^{m,i}$ is of size at most N .

The maps $b^i, p^{m,i}$ generating $\approx_V^{m,i}$ behave well with respect to connected components: b^i is a diffeomorphism so identifies connected components, and $p^{m,i} : V_{de}^{m,i} \rightarrow V_{de'}^{m,i}$ is a finite quotient map and so takes each connected component of $V_{de}^{m,i}$ to a connected component of $V_{de'}^{m,i}$. It follows that $\pi_{\check{V}^{m,i}}$ behaves well with respect to connected components. That is, the image of each connected component of $V_{st}^{m,i}$ under $\pi_{\check{V}^{m,i}}$ is a connected component of $\check{V}^{m,i}$, and

conversely, the inverse image of any connected component of $\ddot{V}^{m,i}$ is the disjoint union of at most N connected components of $\coprod_{(s,t) \in \ddot{P}^m: i \in I_{st}^m} V_{st}^{m,i}$.

Fix $\ddot{v} \in \ddot{V}^{m,i}$. Let $\ddot{V}_{\ddot{v}}$ be the connected component of $\ddot{V}^{m,i}$ containing $\ddot{v}$. The preimage $\pi_{\ddot{V}^{m,i}}^{-1}(\{\ddot{v}\})$ is an $\approx_V^{m,i}$ equivalence class, and so is of size at most N . Write $\pi_{\ddot{V}^{m,i}}^{-1}(\{\ddot{v}\}) = \{v_1, v_2, \dots, v_k\}$, with $k \leq N$. Define V_a to be the connected component of $\coprod_{(s,t) \in \ddot{P}^m: i \in I_{st}^m} V_{st}^{m,i}$ containing v_a for $a = 1, \dots, k$. (Note that although $v_1, \dots, v_k$ are distinct, $V_1, \dots, V_k$ may not be distinct, there could be two or more v_a in the same connected component.) Then each v_a lies in $V_{st}^{m,i}$ for some $(s, t) \in \ddot{P}^m$, and V_a is the connected component of $V_{st}^{m,i}$ containing v_a . Write $\pi_1^{\text{orb}}(V_a, v_a)$ for the orbifold fundamental group, and $\tilde{V}_a^{\text{orb}}$ for the orbifold universal cover, of (V_a, v_a) . Then $V_a \cong \tilde{V}_a^{\text{orb}} / \pi_1^{\text{orb}}(V_a, v_a)$.

Now $\approx_V^{m,i}$ is generated by diffeomorphisms $b^i : V_{st}^{m,i} \rightarrow V_{s't'}^{m,i}$, and finite group quotients $p^{m,i} : V_{de}^{m,i} \rightarrow V_{de'}^{m,i}$. In both cases, these are *finite covers* of orbifolds. For each pair $a, b = 1, \dots, k$, define C_{ab} to be the set of smooth maps $c : V_a \rightarrow V_b$ with $c(v_a) = v_b$ that are the restrictions of b^i or $p^{m,i}$ above to connected components V_a . Then each C_{ab} is finite, as there are only finitely many possibilities for $(\mathbf{a}, \mathbf{b})$ and $(\boldsymbol{\pi}^m, \mathbf{p}^m)$. If $c : V_a \rightarrow V_b$ lies in C_{ab} then c is a finite cover of connected orbifolds V_a, V_b , and we have chosen base-points v_a, v_b with $c(v_a) = v_b$. Hence c induces an *injective group morphism* $c_* : \pi_1^{\text{orb}}(V_a, v_a) \rightarrow \pi_1^{\text{orb}}(V_b, v_b)$, and a *diffeomorphism* $\tilde{c} : \tilde{V}_a^{\text{orb}} \rightarrow \tilde{V}_b^{\text{orb}}$, as in Definition C.3.

It may be helpful to think of this data (V_a, v_a) for $a = 1, \dots, k$ and C_{ab} for $a, b = 1, \dots, k$ as defining a *quiver* Q , with vertices labelled by $1, \dots, k$, and directed edges C_{ab} from vertex a to vertex b for $a, b = 1, \dots, k$. Then Q is *connected*, since $\{v_1, \dots, v_k\}$ is an $\approx_V^{m,i}$ equivalence class, and equivalences are induced by the maps $c : V_a \rightarrow V_b$ in C_{ab} , so we can get between any two vertices v_d, v_e by a finite series of edges $c : V_a \rightarrow V_b$ or their inverses. However, Q need not be simply-connected. Choose a subquiver Q' of Q with the same set of vertices $\{1, \dots, k\}$ but a possibly smaller set of edges $C'_{ab} \subseteq C_{ab}$ for $a, b = 1, \dots, k$, such that Q' is connected and simply-connected.

As above, each $c \in C'_{ab}$ induces a diffeomorphism $\tilde{c} : \tilde{V}_a^{\text{orb}} \rightarrow \tilde{V}_b^{\text{orb}}$ of the universal covers. We use these diffeomorphisms for all $a, b = 1, \dots, k$ and $c \in C'_{ab}$, that is, all edges of Q' , to *identify* all the $\tilde{V}_a^{\text{orb}}$, so we write $\tilde{V}_a^{\text{orb}} \cong \tilde{V}^{\text{orb}}$ for all $a = 1, \dots, k$. This is well-defined as Q' is connected and simply-connected.

The action of $\pi_1^{\text{orb}}(V_a, v_a)$ on $\tilde{V}_a^{\text{orb}} \cong \tilde{V}^{\text{orb}}$ induces a group morphism $\rho_a : \pi_1^{\text{orb}}(V_a, v_a) \rightarrow \text{Diff}(\tilde{V}^{\text{orb}})$ for $a = 1, \dots, k$, which is injective if and only if V_a is effective. For all $a, b = 1, \dots, k$ and $c \in C_{ab}$, we have a diffeomorphism $\tilde{c} : \tilde{V}_a^{\text{orb}} \rightarrow \tilde{V}_b^{\text{orb}}$, which under the identifications $\tilde{V}_a^{\text{orb}} \cong \tilde{V}^{\text{orb}}$, $\tilde{V}_b^{\text{orb}} \cong \tilde{V}^{\text{orb}}$ becomes a diffeomorphism $\tilde{c} : \tilde{V}^{\text{orb}} \rightarrow \tilde{V}^{\text{orb}}$, so that $\tilde{c} \in \text{Diff}(\tilde{V}^{\text{orb}})$, which is the identity if $c \in C'_{ab}$. Define G to be the subgroup of $\text{Diff}(\tilde{V}^{\text{orb}})$ generated by the images $\rho_a(\pi_1^{\text{orb}}(V_a, v_a))$ for $a = 1, \dots, k$ and $\tilde{c} \in \text{Diff}(\tilde{V}^{\text{orb}})$ corresponding to $c \in C_{ab}$ for all $a, b = 1, \dots, k$.

For each $a = 1, \dots, k$, consider the composition

$$\tilde{V}^{\text{orb}} = \tilde{V}_a^{\text{orb}} \xrightarrow{\pi} \tilde{V}_a^{\text{orb}} / \pi_1^{\text{orb}}(V_a, v_a) \cong V_a \xrightarrow{\pi_{\ddot{V}^{m,i}}} \ddot{V}_{\ddot{v}}. \quad (278)$$

The projections $\pi_{\tilde{V}^{m,i}} : V_a \rightarrow \ddot{V}_{\tilde{v}}$ are quotients by the equivalence relation on $\coprod_{a=1}^k V_a$ induced by $c : V_a \rightarrow V_b$ for $c \in C_{ab}$ and $a, b = 1, \dots, k$. It is not difficult to see that the composition (278) is independent of $a = 1, \dots, k$, and factors as $\tilde{V}^{\text{orb}} \xrightarrow{\pi} \tilde{V}^{\text{orb}}/G \xrightarrow{i} \ddot{V}_{\tilde{v}}$, where π is the projection to the quotient, and i is a homeomorphism.

Thus, $\ddot{V}_{\tilde{v}}$ is naturally homeomorphic to $\tilde{V}^{\text{orb}}/G$. We claim that $\tilde{V}^{\text{orb}}/G$ is an *effective orbifold*. As G is a subgroup of $\text{Diff}(\tilde{V}^{\text{orb}})$, it acts effectively on $\tilde{V}^{\text{orb}}$ by definition, and as $\tilde{V}^{\text{orb}}$ is connected, the action is also locally effective. So it is enough to show that stabilizer group $\text{Stab}_G(\tilde{v})$ in G of any $\tilde{v}$ in $\tilde{V}^{\text{orb}}$ is finite. By factoring the projection $\tilde{V}^{\text{orb}} \rightarrow \tilde{V}^{\text{orb}}/G$ as

$$\tilde{V}^{\text{orb}} \xrightarrow{\pi} \tilde{V}^{\text{orb}}/\rho_1(\pi_1^{\text{orb}}(V_1, v_1)) \cong V'_1 \xrightarrow{\pi} \tilde{V}^{\text{orb}}/G \cong \ddot{V}_{\tilde{v}},$$

where V'_1 is the effective orbifold underlying V_1 , noting that the projection $\pi_{\tilde{V}^{m,i}} : V'_1 \rightarrow \ddot{V}_{\tilde{v}}$ is globally finite with constant N as V'_1, V_1 coincide as sets, considering a generic point $\tilde{v}'$ in $\tilde{V}^{\text{orb}}$, and using the local effectiveness of G , we see that $|G/\rho_1(\pi_1^{\text{orb}}(V_1, v_1))| \leq N$. Hence,

$$|\text{Stab}_G(\tilde{v})| \leq N |\text{Stab}_{\rho_1(\pi_1^{\text{orb}}(V_1, v_1))}(\tilde{v})| \leq N |\text{Stab}_{\pi_1^{\text{orb}}(V_1, v_1)}(\tilde{v})| \leq N |\text{Stab}_{V_1}(\pi(\tilde{v}))|,$$

which is finite. Therefore $\tilde{V}^{\text{orb}}/G$ is an effective orbifold, so $\ddot{V}_{\tilde{v}}$ is also an effective orbifold. As we can do this for each connected component $\ddot{V}_{\tilde{v}}$ of $\ddot{V}^{m,i}$, we have given $\ddot{V}^{m,i}$ the structure of an *effective orbifold*, as we have to prove.

By construction, if V is a connected component of $\coprod_{(s,t) \in \tilde{P}^m : i \in I_{st}^m} V_{st}^{m,i}$ and $v \in V$, then the restriction of $\pi_{\tilde{V}^{m,i}}$ to V may be identified with the projection $\tilde{V}^{\text{orb}}/\pi_1^{\text{orb}}(V, v) \rightarrow \tilde{V}^{\text{orb}}/G$, where G is a subgroup of $\text{Diff}(\tilde{V}^{\text{orb}})$ containing $\rho(\pi_1^{\text{orb}}(V, v))$ as a subgroup of finite index. By Definition C.4 it follows that $\tilde{V}^{\text{orb}}/\pi_1^{\text{orb}}(V, v) \rightarrow \tilde{V}^{\text{orb}}/G$ is a weak finite cover. Thus, the restriction of $\pi_{\tilde{V}^{m,i}} : \coprod_{(s,t) \in \tilde{P}^m : i \in I_{st}^m} V_{st}^{m,i} \rightarrow \ddot{V}^{m,i}$ to each connected component of $V_{st}^{m,i}$ is a weak finite cover of its image. Since $\pi_{\tilde{V}^{m,i}}$ is surjective and globally finite, it follows that $\pi_{\tilde{V}^{m,i}} : \coprod_{(s,t) \in \tilde{P}^m : i \in I_{st}^m} V_{st}^{m,i} \rightarrow \ddot{V}^{m,i}$ is a *weak finite cover*, as we want.

This completes the first paragraph of the proposition. Parts (a),(b) now follow from the construction. To define $\ddot{\phi}^{m,ij}$ as a map of orbifolds, for $(s, t) \in \tilde{P}^m$ consider $\phi_{st}^{m,ij} : V_{st}^{m,ij} \rightarrow V_{st}^{m,i}$, restrict to a connected component V^{ij} of $V_{st}^{m,ij}$ and the connected component V^i of $V_{st}^{m,i}$ containing $\phi_{st}^{m,ij}(V^{ij})$, write $\tilde{V}^{ij,\text{orb}}, \tilde{V}^{i,\text{orb}}$ for the orbifold universal covers of V^{ij}, V^i . Then $\phi_{st}^{m,ij}$ lifts to an embedding $\tilde{\phi}_{st}^{m,ij} : \tilde{V}^{ij,\text{orb}} \rightarrow \tilde{V}^{i,\text{orb}}$.

Now the image of V^{ij} in $\ddot{V}^{m,ij}$ is isomorphic as an orbifold to $\tilde{V}^{ij,\text{orb}}/G^{ij}$ for some $G^{ij} \subset \text{Diff}(\tilde{V}^{ij,\text{orb}})$, and the image of V^i in $\ddot{V}^{m,i}$ is isomorphic to $\tilde{V}^{i,\text{orb}}/G^i$ for some $G^i \subset \text{Diff}(\tilde{V}^{i,\text{orb}})$. Since $\phi_{st}^{m,ij}$ is compatible with the equivalences defining $\approx_v^{m,i}, \approx_v^{m,j}$, we see that $\tilde{\phi}_{st}^{m,ij}$ is equivariant under some morphism $\rho^{ij} : G^{ij} \rightarrow G^i$, and so descends to a morphism of orbifolds $\ddot{\phi}_{st}^{m,ij} : \tilde{V}^{ij,\text{orb}}/G^{ij} \rightarrow \tilde{V}^{i,\text{orb}}/G^i$. This lifts $\ddot{\phi}^{m,ij}$ from a continuous map to a map of orbifolds over the connected component $\pi_{\tilde{V}^{m,j}}(V^{ij})$ of $\ddot{V}^{m,ij}$, and proves (a).

For (b), to see that $\approx_v^{m+1,i-1}$ is stronger than $\approx_{\partial v}^{m,i}$, note that the morphisms $(a, b), (\pi^m, p^m)$ on $(X_{st}^m, f_{st}^m, G_{st}^m)$ used to define $\approx_v^{m,i}$ restrict on each con-

nected component $(X_{st'}^{m+1}, \mathbf{f}_{st'}^{m+1}, \mathbf{G}_{st'}^{m+1})$ of $(\partial X_{st}^m, \mathbf{f}_{st}^m|_{\partial X_{st}^m}, \mathbf{G}_{st}^m|_{\partial X_{st}^m})$ to morphisms $(\mathbf{a}', \mathbf{b}'), (\boldsymbol{\pi}^{m+1}, \mathbf{p}^{m+1})$ used to define $\approx_V^{m+1, i-1}$. But there may also be extra morphisms $(\mathbf{a}', \mathbf{b}'), (\boldsymbol{\pi}^{m+1}, \mathbf{p}^{m+1})$ on $(X_{st'}^{m+1}, \mathbf{f}_{st'}^{m+1}, \mathbf{G}_{st'}^{m+1})$ defining $\approx_V^{m+1, i-1}$, which are not the restriction of any $(\mathbf{a}, \mathbf{b}), (\boldsymbol{\pi}^m, \mathbf{p}^m)$ to the boundary of $(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$.

Thus, the set of morphisms $b^{i-1}, p^{m+1, i-1}$ used to define $\approx_V^{m+1, i-1}$ on the right hand side of (276) includes all the morphisms $b^i|_{\partial(\dots)}, p^{m, i}|_{\partial(\dots)}$ used to define $\approx_{\partial V}^{m, i}$ on the left hand side of (276), so $\approx_V^{m+1, i-1}$ is stronger than $\approx_{\partial V}^{m, i}$. The rest of (b) is straightforward. $\square$

Remark C.6. The proof of Proposition C.5 constructed an *effective* orbifold structure on $\tilde{V}^{m, i}$ from the orbifold structures of $V_{st}^{m, i}$. In doing so, it *discarded information* encoded in non-effective orbifold structures. If a connected component V_a of some $V_{st}^{m, i}$ is non-effective, then $\rho_a : \pi_1^{\text{orb}}(V_a, v_a) \rightarrow \text{Diff}(\tilde{V}_a^{\text{orb}})$ is not injective. We made the group G using only the image $\rho_a(\pi_1^{\text{orb}}(V_a, v_a))$, so all information about the kernel $\text{Ker } \rho_a$ is lost.

This loss of information means that in general $\tilde{E}^{m, i} \rightarrow \tilde{V}^{m, i}$ may not be an orbibundle, so that $(\tilde{V}^{m, i}, \tilde{E}^{m, i}, \tilde{s}^{m, i}, \tilde{\psi}^{m, i})$ may not be a Kuranishi neighbourhood; and $(\hat{\phi}^{m, ij}, \hat{\phi}^{m, ij})$ may not be a coordinate change between Kuranishi neighbourhoods; and $\tilde{X}^m$ may not be a Kuranishi space. It is easy to find examples of orbibundles $E \rightarrow V$ in which E is an effective orbifold, but V is a non-effective orbifold; for instance, $V = \{0\}/G$ and $E = \mathbb{R}^n/G$ for $G \subseteq \text{GL}(n, \mathbb{R})$ a nontrivial finite subgroup. The construction of Proposition C.5 would fix E , but replace V by its underlying effective orbifold, $\{0\}$ in our example. Then the projection $E \rightarrow V$, in our example $\mathbb{R}^n/G \rightarrow \{0\}$, is no longer an orbibundle.

Similarly, one can find examples of coordinate changes $(\phi^{ij}, \hat{\phi}^{ij})$ in which V^{ij} is a non-effective orbifold, but V^i is effective, and $\phi^{ij} : V^{ij} \rightarrow V^i$ maps V^{ij} to some orbifold stratum of V^i . The construction of Proposition C.5 would fix V^i , but replace V^{ij} by its underlying effective orbifold. Then $\phi^{ij} : V^{ij} \rightarrow V^i$ would no longer be an *embedding*, as required in Definition 2.18(a), since as in Definition 2.11 an embedding induces isomorphisms on stabilizer groups. So $(\phi^{ij}, \hat{\phi}^{ij})$ would no longer be a coordinate change.

One might expect that this is a solvable problem, and that there should be some way to modify the construction of Proposition C.5 to include the information in non-effective orbifolds, so that we can make $\tilde{X}^m$ into a compact Kuranishi space, $\tilde{\mathbf{f}}^m : \tilde{X}^m \rightarrow Y$ strongly smooth, and $\tilde{\mathbf{G}}^m$ gauge-fixing data including the $(\tilde{V}^{m, i}, \tilde{E}^{m, i}, \tilde{s}^{m, i}, \tilde{\psi}^{m, i})$. However, examples considered by the author show that *this really is not possible*. If we try to construct G not as a subgroup of $\text{Diff}(\tilde{V}^{\text{orb}})$, but as an abstract group with morphism $\rho : G \rightarrow \text{Diff}(\tilde{V}^{\text{orb}})$ containing each $\pi_1^{\text{orb}}(V_a, v_a)$ as a subgroup, there is no natural, functorial way to construct a candidate finite group $\text{Ker } \rho$ containing $\text{Ker } \rho_a$ for $a = 1, \dots, k$.

We now return to the problem of inductively constructing tent functions $\mathbf{T}_{st}^m$ for $(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$ satisfying (i)–(iv) at the beginning of §C.1. Firstly, by reverse induction on $m = M, M-1, M-2, \dots, 1, 0$ we choose $\tilde{\mathbf{T}}^m = (\tilde{T}^{m, i} : i \in \mathbb{N})$,

where $\tilde{T}^{m,i}$ is a tent function on $\check{V}^{m,i}$ which extends to a tent function on an open neighbourhood of the closure $\overline{\check{V}^{m,i}}$ of $\check{V}^{m,i}$ in $\check{V}^{m,i}$, satisfying:

- (A) whenever $j \leq i$ we have $\min \tilde{T}^{m,j}|_{\check{V}^{m,ij}} \equiv \min \tilde{T}^{m,i} \circ \dot{\phi}^{m,ij}|_{\check{V}^{m,ij}}$, and the submanifolds $S_{\{i_1, \dots, i_l\}}$ of Definition A.1 for $\tilde{T}^{m,i}$ intersect $\dot{\phi}^{ij}(\check{V}^{m,ij})$ transversely in $\check{V}^{m,i}$ wherever $t_{i_1}(u) = \min \tilde{T}^{m,i}(u)$.
- (B) for all i we have $\min \tilde{T}^{m,i}|_{\partial \check{V}^{m,i}} \equiv \min \tilde{T}^{m+1,i-1} \circ \tilde{\pi}_\partial^m$, where $\tilde{\pi}_\partial^m$ is as in Proposition C.5(b).
- (C) $\tilde{T}^{m,i}$ and its extension cut $\check{V}^{m,i}, \overline{\check{V}^{m,i}}$ into ‘arbitrarily small pieces’.

Here (A) is taken directly from Definition A.27, the definition of tent functions $\mathbf{T}$ for triples $(X, \mathbf{f}, \mathbf{G})$. Part (B) will imply (iii) at the beginning of §C.1, and also corresponds to the condition $\min(\mathbf{T}|_{\partial X}) = \min \mathbf{T}'$ in Theorem A.35. Part (C) will imply (iv) at the beginning of §C.1.

Suppose for the moment that we could treat $(\check{X}^m, \check{\mathbf{f}}^m, \check{\mathbf{G}}^m)$ as a triple in Kuranishi homology, and that the morphisms $\tilde{\pi}_\partial^m : \partial \check{V}^{m,i} \rightarrow \check{V}^{m+1,i-1}$ lift to $(\check{\mathbf{p}}_\partial^m, \check{\pi}_\partial^m) : (\partial \check{X}^m, \check{\mathbf{f}}^m|_{\partial \check{X}^m}, \check{\mathbf{G}}^m|_{\partial \check{X}^m}) \rightarrow (\check{X}^m, \check{\mathbf{f}}^m, \check{\mathbf{G}}^m)$. Then $\tilde{T}^m$ is a tent function for $(\check{X}^m, \check{\mathbf{f}}^m, \check{\mathbf{G}}^m)$ which cuts $(\check{X}^m, \check{\mathbf{f}}^m, \check{\mathbf{G}}^m)$ into arbitrarily small pieces, and (B) says that $\min(\tilde{T}^m|_{\partial \check{X}^m}) = \min \tilde{T}^{m+1} \circ \tilde{\pi}_\partial^m$. We could choose such $\tilde{T}^m$ by reverse induction on m using the results of §A.3.5.

Here is the important point. As in Remark C.6, we cannot make $(\check{X}^m, \check{\mathbf{f}}^m, \check{\mathbf{G}}^m)$ into a triple in Kuranishi homology. *But to apply the results of §A.3.5, this does not matter at all.* The ways in which $(\check{X}^m, \check{\mathbf{f}}^m, \check{\mathbf{G}}^m)$ fails to be a triple in Kuranishi homology — that $\dot{\phi}^{m,ij} : \check{V}^{m,ij} \rightarrow \check{V}^{m,i}$ may not be an embedding, since it may not be an isomorphism on stabilizer groups as in Proposition C.5(a), for instance — do not affect any of the proofs in §A.3. So we can still use the arguments of §A.3 to choose $\tilde{T}^m$ inductively, just as if $(\check{X}^m, \check{\mathbf{f}}^m, \check{\mathbf{G}}^m)$ were a triple in Kuranishi homology.

Therefore, by reverse induction on $m = M, M-1, \dots, 0$ we use Definition A.34 and Theorem A.35 to choose $\tilde{T}^m$. To apply these we must verify the assumption in Definition A.33 that the prescribed values $\min \tilde{T}^{m+1} \circ \tilde{\pi}_\partial^m$ for $\min(\tilde{T}^m|_{\partial \check{X}^m})$ restrict to σ -invariant data on $\partial^2 \check{X}^m$. That is, to be able to choose $\tilde{T}^{m,i}$, we must check that $(\min \tilde{T}^{m+1,i-1} \circ \tilde{\pi}_\partial^m)|_{\partial^2 \check{V}^{m,i}}$ is invariant under the natural involution $\sigma : \partial^2 \check{V}^{m,i} \rightarrow \partial^2 \check{V}^{m,i}$. But (B) for $(m+1, i-1)$ gives

$$(\min \tilde{T}^{m+1,i-1} \circ \tilde{\pi}_\partial^m)|_{\partial^2 \check{V}^{m,i}} \equiv \min \tilde{T}^{m+2,i-2} \circ \tilde{\pi}_\partial^{m+1} \circ (\tilde{\pi}_\partial^m|_{\partial^2 \check{V}^{m,i}}). \quad (279)$$

We claim that $\tilde{\pi}_\partial^{m+1} \circ (\tilde{\pi}_\partial^m|_{\partial^2 \check{V}^{m,i}}) \circ \sigma \equiv \tilde{\pi}_\partial^{m+1} \circ (\tilde{\pi}_\partial^m|_{\partial^2 \check{V}^{m,i}})$ as maps $\partial^2 \check{V}^{m,i} \rightarrow \check{V}^{m+2,i-2}$. Combining this with (279) shows that $(\min \tilde{T}^{m+1,i-1} \circ \tilde{\pi}_\partial^m)|_{\partial^2 \check{V}^{m,i}}$ is σ -invariant, as we have to show. To prove the claim, note that for each $(s, t) \in \check{P}^m$, we have a natural orientation-reversing involution $\sigma : \partial^2(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m) \rightarrow \partial^2(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$. The restriction of σ to each connected component $(X_{st'}^{m+2}, \mathbf{f}_{st'}^{m+2}, \mathbf{G}_{st'}^{m+2})$ of $\partial^2(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$ is one of the isomorphisms $(\mathbf{a}, \mathbf{b})$ in the definition of $\approx^{m+2}, \approx_X^{m+2}, \approx_V^{m+2,i-2}$.

Hence, $\sigma : \partial^2 V_{st}^{m,i} \rightarrow \partial^2 V_{st}^{m,i}$ is one of the diffeomorphisms in the definition of $\approx_V^{m+2,i-2}$. As $\sigma : \partial^2 \dot{V}^{m,i} \rightarrow \partial^2 \dot{V}^{m,i}$ lifts locally to $\sigma : \partial^2 V_{st}^{m,i} \rightarrow \partial^2 V_{st}^{m,i}$, which then projects to the identity on $\dot{V}^{m+2,i-2}$ by definition of $\approx_V^{m+2,i-2}$, the claim follows. Therefore by induction, we can use the results of §A.3.5 to choose $\tilde{T}^m$ satisfying (A)–(C) above for $m = M, M-1, \dots, 0$.

Now, for all $m = 0, \dots, M$, $(s, t) \in \dot{P}^m$ and $i \in I_{st}^m$ define $T_{st}^{m,i} : \dot{V}_{st}^{m,i} \rightarrow F([1, \infty))$ by $T_{st}^{m,i} = \tilde{T}^{m,i} \circ \pi_{\dot{V}^{m,i}}|_{\dot{V}_{st}^{m,i}}$. Since $\tilde{T}^{m,i} : \dot{V}^{m,i} \rightarrow F([1, \infty))$ is a tent function and $\pi_{\dot{V}^{m,i}}$ is a weak finite cover of orbifolds by Proposition C.5, it follows that $T_{st}^{m,i}$ is a *tent function* on $\dot{V}_{st}^{m,i}$, which extends to a tent function on an open neighbourhood of the closure $\overline{\dot{V}_{st}^{m,i}}$ of $\dot{V}_{st}^{m,i}$ in $V_{st}^{m,i}$, since $\tilde{T}^{m,i}$ extends to an open neighbourhood of $\overline{\dot{V}^{m,i}}$ in $\dot{V}^{m,i}$. Define $\mathbf{T}_{st}^m = (T_{st}^{m,i} : i \in I_{st}^m)$. Part (A) above implies that the $T_{st}^{m,i}$ for $i \in I_{st}^m$ satisfy the conditions of Definition A.27, and therefore $\mathbf{T}_{st}^m$ is a *tent function* for $(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$. We claim that these $\mathbf{T}_{st}^m$ for all $m = 0, \dots, M$ and $(s, t) \in \dot{P}^m$ satisfy conditions (i)–(iv) at the beginning of §C.1.

Parts (i),(ii) hold by construction of $\dot{V}^{m,i}$ using $\approx_V^{m,i}$. For (i), if $(s, t), (s', t')$ lie in $\dot{P}^m$, $(\mathbf{a}, \mathbf{b}) : (X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m) \rightarrow (X_{s't'}^m, \mathbf{f}_{s't'}^m, \mathbf{G}_{s't'}^m)$ is an isomorphism, and $i \in I_{st}^m$, then $b^i : V_{st}^{m,i} \rightarrow V_{s't'}^{m,i}$ is one of the maps generating the equivalence relation $\approx_V^{m,i}$ on $\coprod_{(s,t) \in \dot{P}^m, i \in I_{st}^m} V_{st}^{m,i}$. Since $\dot{V}^{m,i} = (\coprod_{(s,t) \in \dot{P}^m, i \in I_{st}^m} V_{st}^{m,i}) / \approx_V^{m,i}$, and $\pi_{\dot{V}^{m,i}} : \coprod_{(s,t) \in \dot{P}^m, i \in I_{st}^m} V_{st}^{m,i} \rightarrow \dot{V}^{m,i}$ is the natural projection, we see that $\pi_{\dot{V}^{m,i}}|_{\dot{V}_{st}^{m,i}} \equiv \pi_{\dot{V}^{m,i}}|_{\dot{V}_{s't'}^{m,i}} \circ b^i$, and composing with $\tilde{T}^{m,i}$ gives $T_{st}^{m,i} = T_{s't'}^{m,i} \circ b^i$. Part (ii) is the same, as the $p^{m,i}$ in (ii) are also maps generating $\approx_V^{m,i}$. Parts (iii),(iv) easily follow from (B),(C) above, respectively. This completes the construction of $\mathbf{T}_{st}^m$ for all m, s, t satisfying (i)–(iv) at the beginning of §C.1.

In the notation of the beginning of §C.1, for $a \in A$ and $d \in D$ we have

$$(X_a, \mathbf{f}_a, \mathbf{G}_a) = \coprod_{b=1}^{n_a^0} (X_{ab}^0, \mathbf{f}_{ab}^0, \mathbf{G}_{ab}^0), \quad (\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d) = \coprod_{e=1}^{\tilde{n}_d^1} (X_{de}^1, \mathbf{f}_{de}^1, \mathbf{G}_{de}^1)$$

$$\text{and} \quad (\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d)) = \coprod_{e=1}^{-\tilde{n}_d^1} (X_{de}^1, \mathbf{f}_{de}^1, \mathbf{G}_{de}^1).$$

Using these equations, define tent functions $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ and $\tilde{\mathbf{T}}_d$ for $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ and $\hat{\mathbf{T}}_d$ for $(\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d))$ by $\mathbf{T}_a = \coprod_{b=1}^{n_a^0} \mathbf{T}_{ab}^0$, and $\tilde{\mathbf{T}}_d = \coprod_{e=1}^{\tilde{n}_d^1} \mathbf{T}_{de}^1$, and $\hat{\mathbf{T}}_d = \coprod_{e=1}^{-\tilde{n}_d^1} \mathbf{T}_{de}^1$. We claim that these $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ satisfy conditions (a)–(c) of Step 1 of §C, shortly after equation (255).

We prove (a)–(c) using (i)–(iv) at the beginning of §C.1, which the $\mathbf{T}_{st}^m$ satisfy. For part (a), we have $\mathbf{T}_a|_{\partial X_a} = \coprod_{b=1}^{n_a^1} (X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1)$, so it follows from (i) that $\mathbf{T}_a|_{\partial X_a}$ is invariant under $\text{Aut}(\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a})$, and similarly $\tilde{\mathbf{T}}_d$ is invariant under $\text{Aut}(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$. Also (ii) implies that $\pi_*(\tilde{\mathbf{T}}_d) = \hat{\mathbf{T}}_d$, so again using (i) we see that $\pi_*(\tilde{\mathbf{T}}_d)$ is invariant under $\text{Aut}(\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d))$. Part (b) follows from (i) with $m = 1$, and (c) from (iv). This completes Step 1.

C.2 Step 2: lifting to $\dot{X}_{ac}$ with trivial stabilizers

We continue to use the notation of Step 1. For each $a \in A$, write $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ for $c \in C_a$ for the arbitrarily small pieces into which $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ is ‘cut’ using the tent function $\mathbf{T}_a$, in the notation of (211). We must show that we can choose the $\mathbf{T}_a$ so that $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}) \cong (\dot{X}_{ac}/\Gamma_{ac}, \pi_*(\dot{\mathbf{f}}_{ac}), \pi_*(\dot{\mathbf{G}}_{ac}))$ for each $c \in C_a$, where $(\dot{X}_{ac}, \dot{\mathbf{f}}_{ac}, \dot{\mathbf{G}}_{ac})$ is a triple and $\dot{X}_{ac}$ has *trivial stabilizers*.

Let $a \in A$ and $p \in X_a$, and write Γ_p for the stabilizer group $\text{Stab}(p)$. We first show that a small neighbourhood of p in X_a may be written, as a Kuranishi space, as a quotient $\dot{X}_p/\Gamma_p$, where $\dot{X}_p$ is a Kuranishi space with trivial stabilizers.

Let $(V_p, \dots, \psi_p)$ be a Kuranishi neighbourhood in the germ of X_a at p , and set $v = \psi_p^{-1}(p)$. As V_p is an orbifold, by §2.1 there exists an orbifold chart $(\dot{V}_p, \Gamma_p, \xi_p)$ for V_p near v , with group Γ_p . That is, $\dot{V}_p$ is a smooth manifold acted on by Γ_p , and $\xi_p : \dot{V}_p/\Gamma_p \rightarrow V_p$ is a diffeomorphism with an open neighbourhood of v in V_p . There is a unique $\dot{p} \in \dot{V}_p$ with $\xi_p(\Gamma_p \dot{p}) = v$, which is fixed by Γ_p . Making V_p smaller if necessary, we can assume $V_p = \xi_p(\dot{V}_p/\Gamma_p)$. Write $\pi_p : \dot{V}_p \rightarrow V_p$ for the corresponding projection, which is a submersion. Define $\dot{E}_p = \pi_p^*(E_p)$. Then $\dot{E}_p \rightarrow \dot{V}_p$ is a vector bundle, in the usual manifold sense. Define $\dot{s}_p = s_p \circ \pi_p$, so that $\dot{s}_p$ is a smooth section of $\dot{E}_p$.

Define a topological space $\dot{X}_p = \dot{s}_p^{-1}(0)$, and let $\dot{\psi}_p$ be the identity map on $\dot{X}_p$. Then $(\dot{V}_p, \dot{E}_p, \dot{s}_p, \dot{\psi}_p)$ is a Kuranishi neighbourhood covering $\dot{X}_p$. The action of Γ_p on $\dot{V}_p$ lifts naturally to an action on $\dot{E}_p$, and $\dot{s}_p$ is Γ_p -equivariant, so the subset $\dot{X}_p$ is preserved by Γ_p , and Γ_p acts on $\dot{X}_p$, with $\psi_p \circ \xi_p : \dot{X}_p/\Gamma_p \rightarrow X_a$ a homeomorphism with an open neighbourhood $\text{Im } \psi_p$ of p in X_a . Also $\psi_p \circ \pi_p : \dot{X}_p \rightarrow \text{Im } \psi_p \subset X_a$ is continuous.

We shall define a Kuranishi structure $\dot{\kappa}_p$ on $\dot{X}_p$. Let $\dot{q} \in \dot{X}_p$, and set $q = \psi_p \circ \pi_p(\dot{q})$. Then $q \in \text{Im } \psi_p$, so for all sufficiently small $(V_q, \dots, \psi_q)$ in the germ of X_a at q there is a coordinate change $(\phi_{pq}, \hat{\phi}_{pq}) : (V_q, \dots, \psi_q) \rightarrow (V_p, \dots, \psi_p)$. Define a Kuranishi neighbourhood $(\dot{V}_{\dot{q}}, \dots, \dot{\psi}_{\dot{q}})$ of $\dot{q}$ in $\dot{X}_p$ to be

$$(\dot{V}_{\dot{q}}, \dot{E}_{\dot{q}}, \dot{s}_{\dot{q}}, \dot{\psi}_{\dot{q}}) = (\dot{V}_p \times_{\pi_p, V_p, \phi_{pq}} V_q, \pi_{V_q}^*(E_q), s_q \circ \pi_{V_q}, \pi_{V_q}). \quad (280)$$

Since $\dot{V}_p$ is a manifold, π_p is a submersion, and ϕ_{pq} is an embedding, $\dot{V}_{\dot{q}}$ is a manifold, and $\dot{E}_{\dot{q}} \rightarrow \dot{V}_{\dot{q}}$ a vector bundle. Since $s_p \circ \phi_{pq} \equiv \hat{\phi}_{pq} \circ s_q$, ϕ_{pq} takes $s_q^{-1}(0)$ to $s_p^{-1}(0)$, and so $\dot{\psi}_{\dot{q}} = \pi_{V_q}$ takes $\dot{s}_{\dot{q}}^{-1}(0)$ to $\dot{s}_p^{-1}(0) = \dot{X}_p$, as we need.

Define the germ of Kuranishi neighbourhoods at $\dot{q}$ in $\dot{\kappa}_p$ to be the equivalence class of $(\dot{V}_{\dot{q}}, \dots, \dot{\psi}_{\dot{q}})$. This is independent of the choice of $(V_q, \dots, \psi_q)$, and so well-defined. It is now easy to show that the germ of coordinate changes between Kuranishi neighbourhoods on $\text{Im } \psi_p \subset X_a$ induces a germ of coordinate changes between Kuranishi neighbourhoods on $\dot{X}_p$, giving a Kuranishi structure $\dot{\kappa}_p$ on $\dot{X}_p$. Since the $\dot{V}_{\dot{q}}$ are all manifolds, $(\dot{X}_p, \dot{\kappa}_p)$ has *trivial stabilizers*. Furthermore, $\dot{\kappa}_p$ is invariant under the action of Γ_p on $\dot{X}_p$, and $\psi_p \circ \pi_p : \dot{X}_p \rightarrow \text{Im } \psi_p \subset X_a$ lifts

to an isomorphism of Kuranishi spaces between $(\dot{X}_p, \dot{\kappa}_p)/\Gamma_p$ and the restriction of the Kuranishi structure on X_a to $\text{Im } \psi_p$.

Thus, for all $a \in A$ and $p \in X_a$, we can find an open neighbourhood $\text{Im } \psi_p$ of p in X_a , and an isomorphism of Kuranishi spaces $\text{Im } \psi_p \cong \dot{X}_p/\Gamma_p$, where $\dot{X}_p$ has trivial stabilizers, and Γ_p is a finite group. Such $\text{Im } \psi_p$ cover X_a , so by Step 1 we can cut the X_a into pieces X_{ac} such that each X_{ac} is contained in some such subset $\text{Im } \psi_p$ in X_a .

Then we have $X_{ac} \cong \dot{X}_{ac}/\Gamma_{ac}$, where $\dot{X}_{ac} = \dot{X}_p \times_{\pi, X_a, \iota} X_{ac}$ is a compact Kuranishi space with trivial stabilizers, and $\Gamma_{ac} = \Gamma_p$ is a finite group acting on $\dot{X}_{ac}$ by strong diffeomorphisms. Here $\pi : \dot{X}_p \rightarrow \text{Im } \psi_p \subset X_a$ is the obvious projection, which is a strong submersion, and $\iota : X_{ac} \rightarrow X_a$ is the inclusion. Write $\pi_{ac} : \dot{X}_{ac} \rightarrow X_{ac}$ for the projection, which is a strong submersion, and $\dot{f}_{ac} = \dot{f}_{ac} \circ \pi_{ac} : \dot{X}_{ac} \rightarrow Y$, which is strongly smooth.

It remains to show that we can choose T_a for $a \in A$ and $X_{ac}, \dot{X}_{ac}$ for $c \in C_a$ so that G_{ac} on X_{ac} lifts to $\dot{G}_{ac}$ on $\dot{X}_{ac}$. In doing this, it is important that we can not only make the X_{ac} arbitrarily small, we can also make the subsets V_{ac}^i of V_a^i for $i \in I_a$ arbitrarily small. This is part (C) in §C.1, and can be seen from the construction in §A.3.3 used to define T_a .

In the situation above, let $q \in \text{Im } \psi_p$, and suppose $i \in I_a$ and $v \in (s_a^i)^{-1}(0) \subseteq V_a^i$ with $\psi_a^i(v) = q$. Let $(V_q, \dots, \psi_q)$ be a sufficiently small Kuranishi neighbourhood in the germ of X_a at q . Then as $(V_a^i, \dots, \psi_a^i)$ is a compatible Kuranishi neighbourhood on X_a , we are given a coordinate change $(\phi_q^i, \hat{\phi}_q^i) : (V_q, \dots, \psi_q) \rightarrow (V_a^i, \dots, \psi_a^i)$. From (280) we have a Kuranishi neighbourhood $(\dot{V}_q, \dot{E}_q, \dot{s}_q, \dot{\psi}_q)$ on $\dot{X}_p$ with $\dot{V}_q$ a manifold, a diffeomorphism $\xi_q : \dot{V}_q/\Gamma_p \rightarrow V_q$, and a submersion $\pi_q = \pi_{V_q} : \dot{V}_q \rightarrow V_q$.

Let W be an open neighbourhood of v in V_a^i . Since $\phi_q^i : V_q \rightarrow V_a^i$ is an embedding with $\phi_q^i(\psi_q^{-1}(q)) = v$, if W is sufficiently small we may extend the diffeomorphism $\xi_q : \dot{V}_q/\Gamma_p \rightarrow V_q$ to a diffeomorphism $\tilde{\xi}_q : \dot{W}/\Gamma_p \rightarrow W$, where $\dot{W}$ is a manifold with a Γ_p action, such that ϕ_q^i lifts over $(\phi_q^i)^{-1}(W)$ to a smooth Γ_p -equivariant map $\tilde{\phi}_q^i : (\phi_q^i \circ \pi_q)^{-1}(W) \rightarrow \dot{W}$ from the open subset $(\phi_q^i \circ \pi_q)^{-1}(W)$ in $\dot{V}_q$. Furthermore, if W is connected these $\dot{W}, \tilde{\xi}_q, \tilde{\phi}_q^i$ are unique up to canonical isomorphism.

As we can do this for any $q \in \text{Im } \psi_p \cap \text{Im } \psi_a^i$, and the diffeomorphisms $\xi_q : \dot{V}_q/\Gamma_p \rightarrow V_q$ all come from an isomorphism $\dot{X}_p/\Gamma_p \cong \text{Im } \psi_p \subset X_a$ and so are compatible under coordinate changes, and the $\dot{W}, \tilde{\xi}_q, \tilde{\phi}_q^i$ are unique up to *canonical* isomorphism above, we can patch all these local lifts together. This proves that if W is any sufficiently small open neighbourhood of some subset in $(\psi_a^i)^{-1}(\text{Im } \psi_p)$ in V_a^i , and every connected component of W meets $(\psi_a^i)^{-1}(\text{Im } \psi_p)$, then we can write $W \cong \dot{W}/\Gamma_p$ for some manifold $\dot{W}$, so that all embeddings $\phi_q^i : V_q \rightarrow W \subset V_a^i$ for $q \in \text{Im } \psi_p \cap \psi_a^i(W)$ lift canonically to embeddings $\tilde{\phi}_q^i : \dot{V}_q \rightarrow \dot{W}$.

Let us choose the T_a for $a \in A$ so that all V_{ac}^i are sufficiently small in this sense, which is possible by Step 1, and define $\dot{X}_{ac}, \Gamma_{ac} = \Gamma_p$ as above with

$X_{ac} \cong \dot{X}_{ac}/\Gamma_{ac}$. Then there exist a manifold $\dot{V}_{ac}^i$, an action of Γ_{ac} on $\dot{V}_{ac}^i$, and a diffeomorphism $\dot{V}_{ac}^i/\Gamma_{ac} \cong V_{ac}^i$, which are unique up to canonical isomorphism. To prove this, note that as $(V_{ac}^i, \dots, \psi_{ac}^i)$ is part of an excellent coordinate system, every connected component of V_{ac}^i intersects $(s_{ac}^i)^{-1}(0)$.

Define a Kuranishi neighbourhood $(\dot{V}_{ac}^i, \dot{E}_{ac}^i, \dot{s}_{ac}^i, \dot{\psi}_{ac}^i)$ on $\dot{X}_{ac}$, where $\dot{E}_{ac}^i = \pi_{V_{ac}^i}^*(E_{ac}^i)$ and $\dot{s}_{ac}^i \equiv s_{ac}^i \circ \pi_{V_{ac}^i}$, for $\pi_{V_{ac}^i} : \dot{V}_{ac}^i \rightarrow V_{ac}^i$ the projection, and $\dot{\psi}_{ac}^i$ is the natural lift of $\psi_{ac}^i \circ \pi_{V_{ac}^i} : (s_{ac}^i)^{-1}(0) \rightarrow X_{ac}$ to $\dot{X}_{ac}$. These extend to an *excellent coordinate system* $(\dot{I}_{ac}, \dot{\eta}_{ac}) = ((I_{ac}, (\dot{V}_{ac}^i, \dots, \dot{\psi}_{ac}^i), \dot{f}_{ac}^i : i \in I_{ac}, \dots), (\dot{\eta}_{i,ab}, \dot{\eta}_{i,ab}^j : i, j \in I_{ac}))$ for $(\dot{X}_{ac}, \dot{\mathbf{f}}_{ac})$, where $\dot{f}_{ac}^i = f_{ac}^i \circ \pi_{V_{ac}^i}$, $\dot{\eta}_{i,ab} = \eta_{i,ab} \circ \pi_{X_{ac}}$, and $\dot{\eta}_{i,ab}^j = \eta_{i,ab}^j \circ \pi_{V_{ac}^j}$. We also define $\dot{\mathbf{G}}_{ac} = (\dot{I}_{ac}, \dot{\eta}_{ac}, \dot{G}_{ac}^i : i \in \dot{I}_{ac})$, where $\dot{G}_{ac}^i : \dot{E}_{ac}^i \rightarrow P$ is given by $\dot{G}_{ac}^i = G_{ac}^i \circ \pi_{E_{ac}^i}$ for $\pi_{E_{ac}^i} : \dot{E}_{ac}^i \rightarrow E_{ac}^i$ the natural projection.

Then $(\dot{X}_{ac}, \dot{\mathbf{f}}_{ac}, \dot{\mathbf{G}}_{ac})$ is a triple in Kuranishi homology, and Γ_{ac} acts on $(\dot{X}_{ac}, \dot{\mathbf{f}}_{ac}, \dot{\mathbf{G}}_{ac})$ by automorphisms with $(\dot{X}_{ac}/\Gamma_{ac}, \pi_*(\dot{\mathbf{f}}_{ac}), \pi_*(\dot{\mathbf{G}}_{ac})) \cong (X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$, so Definition 4.3(iv) gives (260), and Step 2 follows.

C.3 Step 3: lifting to a cycle in $\widetilde{KC}_k(Y; R)$

As in Step 3 we now *change notation* from $\dot{X}_{ac}$ for $a \in A$ and $c \in C_a$ back to X_a for $a \in A$. That is, by Step 2 we can represent α by a cycle $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$ in $KC_k(Y; R)$, where each X_a is a compact, oriented Kuranishi space with *trivial stabilizers*, and each V_a^i in $\mathbf{G}_a$ is a *manifold*. Equation (255) holds for $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ for $d \in D$. These $A, X_a, \mathbf{f}_a, \mathbf{G}_a, \rho_a, D, \tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d, \Gamma_d, \eta_d$ are different from those in Steps 1 and 2.

We now repeat Step 1, choosing tent functions $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ for $a \in A$, and $\tilde{\mathbf{T}}_d$ for $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ for $d \in D$, satisfying conditions (a)–(c) of Step 1, such that $\mathbf{T}_a$ cuts $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ into small pieces $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ for $c \in C_a$, and $\tilde{\mathbf{T}}_d$ cuts $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ into small pieces $(\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df})$ for $f \in F_d$. Equations (262)–(267) follow as in Step 3.

We must prove that provided the $X_{ac}, \tilde{X}_{df}$ are sufficiently small, the right hand side of (267) is zero. To do this, we recall the notation of §C.1, fixing $m = 1$. For $a \in A$ and $d \in D$ we have

$$\begin{aligned} (\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a}) &= \coprod_{b=1}^{n_a^1} (X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1), \\ (\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d) &= \coprod_{e=1}^{\tilde{n}_d^1} (X_{de}^1, \mathbf{f}_{de}^1, \mathbf{G}_{de}^1), \quad \text{and} \\ (\tilde{X}_d/\Gamma_d, \pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d)) &= \coprod_{e=1}^{-\tilde{n}_d^1} (X_{de}^1, \mathbf{f}_{de}^1, \mathbf{G}_{de}^1). \end{aligned} \tag{281}$$

Write $(X^1, \mathbf{f}^1, \mathbf{G}^1)$ for the disjoint union of the triples on the left hand sides of equations (281) over all $a \in A$ and $d \in D$, and $(V^{1,i}, \dots, \psi^{1,i})$ for $i \in I^1$ for the Kuranishi neighbourhoods in $\mathbf{G}^1$.

Write $\mathbf{T}^1$ for the tent function for $(X^1, \mathbf{f}^1, \mathbf{G}^1)$ which is the disjoint union of the tent functions $\mathbf{T}_a|_{\partial X_a}, \tilde{\mathbf{T}}_d$ and $\pi_*(\tilde{\mathbf{T}}_d)$ on the triples in (281) chosen in Step 1. Write $(X_c^1, \mathbf{f}_c^1, \mathbf{G}_c^1)$ for $c \in C^1$ for the pieces into which $(X^1, \mathbf{f}^1, \mathbf{G}^1)$ is ‘cut’

by $\mathbf{T}^1$, as in (211). These $(X_c^1, \mathbf{f}_c^1, \mathbf{G}_c^1)$ consist of triples $(\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df})$ and $(\tilde{X}_{df}/\text{Stab}_{\Gamma_d}(f), \pi_*(\tilde{\mathbf{f}}_{df}), \pi_*(\tilde{\mathbf{G}}_{df}))$ which occur in (267), and also the decomposition of $(\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a})$ using $\mathbf{T}_a|_{\partial X_a}$ for $a \in A$. Write $\pi : X_c^1 \rightarrow X^1$ for the natural projection.

Now $\tilde{P}^1$ is the disjoint union of all $(a, b), (d, e)$ occurring in (281). Hence

$$\coprod_{(s,t) \in \tilde{P}^1} X_{st}^1 = \coprod_{a \in A} \partial X_a \amalg \coprod_{d \in D} \tilde{X}_d \amalg \coprod_{d \in D} \tilde{X}_d/\Gamma_d = X^1. \quad (282)$$

Also $\approx_X^1$ is an equivalence relation on (282) with $\tilde{X}^1 = X^1/\approx_X^1$, and projection $\pi_{\tilde{X}^1} : X^1 \rightarrow \tilde{X}^1$, which is continuous, surjective, and globally finite, with $\tilde{X}^1$ a compact topological space. We shall prove Step 3 by working locally on $\tilde{X}^1$. Fix some $\tilde{p} \in \tilde{X}^1$. As $\pi_{\tilde{X}^1}$ is surjective and globally finite, $\pi_{\tilde{X}^1}^{-1}(\tilde{p})$ is finitely many points $p_{(1)}, \dots, p_{(n)}$ in X^1 .

We now use the ideas of Step 2. For each $j = 1, \dots, n$, there exists an open neighbourhood $U_{(j)}$ of $p_{(j)}$ in X^1 such that as a Kuranishi space we may write $U_{(j)} \cong \tilde{U}_{(j)}/\Gamma_{(j)}$, where $\tilde{U}_{(j)}$ has *trivial stabilizers*, and $\Gamma_{(j)} = \text{Stab}_{X^1}(p_{(j)})$ is a finite group acting on $\tilde{U}_{(j)}$ by strong diffeomorphisms, fixing the pullback $\dot{p}_{(j)}$ of $p_{(j)}$ in $\tilde{U}_{(j)}$. Making the $U_{(j)}$ smaller if necessary, we suppose $U_{(1)}, \dots, U_{(n)}$ are *disjoint*. Similarly, if $i \in \mathbb{N}$ with $p_{(j)} \in \text{Im } \psi^{1,i}$, so that $p_{(j)} = \psi^{1,i}(v_{(j)}^i)$ for some unique $v_{(j)}^i \in V^{1,i}$, we choose disjoint open neighbourhoods $W_{(j)}^i$ of $v_{(j)}^i$ in $V^{1,i}$ for $j = 1, \dots, n$ with $W_{(j)}^i \cong \tilde{W}_{(j)}^i/\Gamma_{(j)}$, where $\tilde{W}_{(j)}^i$ is a *manifold*, and $\Gamma_{(j)}$ is as before, since $\text{Stab}_{X^1}(p_{(j)}) \cong \text{Stab}_{V^{1,i}}(v_{(j)}^i)$.

For $j = 1, \dots, n$, define $(X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)})$ to be the disjoint union of those components $(X_c^1, \mathbf{f}_c^1, \mathbf{G}_c^1)$ for $c \in C^1$ with $p_{(j)} \in \pi(X_c^1)$. There is exactly one such component if $p_{(j)}$ lies in the interior of $\pi(X_{(j)})$, but if $p_{(j)}$ lies in a codimension k corner for $k \geq 1$, there may be more than one. Write $(V_{(j)}^i, \dots, \psi_{(j)}^i)$ for $i \in I_{(j)}$ for the Kuranishi neighbourhoods in $\mathbf{G}_{(j)}$, and $\pi : V_{(j)}^i \rightarrow V^{1,i}$ for the natural projections.

Suppose now that the $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ in Step 1 have been chosen such that $\pi(X_{(j)}) \subseteq U_{(j)}$ and $\pi(V_{(j)}^i) \subseteq W_{(j)}^i$ for all $j = 1, \dots, n$ and $i \in I_{(j)}$. This is possible as Step 1 allows us to cut $X^1, V^{1,i}$ into ‘arbitrarily small pieces’, and so to make $\pi(X_{(j)})$ and $\pi(V_{(j)}^i)$ arbitrarily small. Then the proof of Step 2 gives a triple $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$, an action of $\Gamma_{(j)}$ on $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$ by isomorphisms, and an isomorphism $(\dot{X}_{(j)}/\Gamma_{(j)}, \pi_*(\dot{\mathbf{f}}_{(j)}), \pi_*(\dot{\mathbf{G}}_{(j)})) \cong (X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)})$, for each j .

Making the pieces smaller if necessary, we see that $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$ is *unique up to isomorphism*, that is, there are no arbitrary choices in the construction of $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$ from $(X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)})$ and $p_{(j)}$. We can think of $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$ as like a kind of ‘Kuranishi space universal cover’ of $(X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)})$, by analogy with orbifold universal covers in §C.1.

We now come to one of the crucial steps in the proof. We claim that *the triples $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$ for $j = 1, \dots, n$ are all isomorphic*, through either orientation-preserving or orientation-reversing isomorphisms. To prove this, note that the equivalence relation $\approx_X^1$ on X^1 is generated by (possibly

orientation-reversing) isomorphisms of components of $(X^1, \mathbf{f}^1, \mathbf{G}^1)$, and finite group quotients between components of $(X^1, \mathbf{f}^1, \mathbf{G}^1)$. Since $\{p_{(1)}, \dots, p_{(n)}\}$ is an $\approx_x^1$ -equivalence class, any two $p_{(j)}, p_{(k)}$ are connected by a finite chain of such isomorphisms or finite group quotients. Thus, to show the $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$ are all isomorphic for $j = 1, \dots, n$, it is enough to prove that if $p_{(j)}$ and $p_{(k)}$ are identified by an isomorphism or finite group quotient of their components, then $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$ and $(\dot{X}_{(k)}, \dot{\mathbf{f}}_{(k)}, \dot{\mathbf{G}}_{(k)})$ are isomorphic.

The case of isomorphisms is easy: if $(\mathbf{a}, \mathbf{b}) : (X_{(j)}^1, \mathbf{f}_{(j)}^1, \mathbf{G}_{(j)}^1) \rightarrow (X_{(k)}^1, \mathbf{f}_{(k)}^1, \mathbf{G}_{(k)}^1)$ is a (possibly orientation-reversing) isomorphism between the components of $(X^1, \mathbf{f}^1, \mathbf{G}^1)$ containing $p_{(j)}, p_{(k)}$ with $a(p_{(j)}) = p_{(k)}$, then condition (i) in §C.1 implies that $\mathbf{T}^1|_{X_{(j)}^1} = \mathbf{b}^*(\mathbf{T}^1|_{X_{(k)}^1})$, so $(\mathbf{a}, \mathbf{b})$ lifts to a (possibly orientation-reversing) isomorphism between $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$ and $(\dot{X}_{(k)}, \dot{\mathbf{f}}_{(k)}, \dot{\mathbf{G}}_{(k)})$.

Suppose $p_{(j)}$ and $p_{(k)}$ are related by one of the finite group quotients used to define $\approx_x^1$. Then in §C.1(ii) with $m = 1$, we have $(X_{(j)}^1, \mathbf{f}_{(j)}^1, \mathbf{G}_{(j)}^1) = (X_{de}^1, \mathbf{f}_{de}^1, \mathbf{G}_{de}^1)$, and $\Gamma = \text{Stab}_{\Gamma_d}(e)$, and $(X_{(k)}^1, \mathbf{f}_{(k)}^1, \mathbf{G}_{(k)}^1) = (X_{(j)}^1/\Gamma, \pi_*(\mathbf{f}_{(j)}^1), \pi_*(\mathbf{G}_{(j)}^1))$ with $p_{(k)} = p_{(j)}\Gamma$, where $(\pi, \mathbf{p}) : (X_{(j)}^1, \mathbf{f}_{(j)}^1, \mathbf{G}_{(j)}^1) \rightarrow (X_{(k)}^1, \mathbf{f}_{(k)}^1, \mathbf{G}_{(k)}^1)$ is the projection to the quotient. Part (ii) of §C.1 gives $\mathbf{T}^1|_{X_{(j)}^1} = \mathbf{p}^*(\mathbf{T}^1|_{X_{(k)}^1})$.

It is now easy to see that the action of Γ on $(X_{(j)}^1, \mathbf{f}_{(j)}^1, \mathbf{G}_{(j)}^1)$ lifts to $(X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)})$, and $(X_{(k)}, \mathbf{f}_{(k)}, \mathbf{G}_{(k)}) \cong (X_{(j)}/\Gamma, \pi_*(\mathbf{f}_{(j)}), \pi_*(\mathbf{G}_{(j)}))$. Using the obvious shorthand notation, we now have isomorphisms

$$\begin{aligned} (X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)}) &\cong (\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})/\Gamma_{(j)}, & \Gamma_{(j)} &= \text{Stab}(p_{(j)}), \\ (X_{(k)}, \mathbf{f}_{(k)}, \mathbf{G}_{(k)}) &\cong (\dot{X}_{(k)}, \dot{\mathbf{f}}_{(k)}, \dot{\mathbf{G}}_{(k)})/\Gamma_{(k)}, & \Gamma_{(k)} &= \text{Stab}(p_{(k)}), \\ \text{and} & & & \\ & (X_{(k)}, \mathbf{f}_{(k)}, \mathbf{G}_{(k)}) \cong (X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)})/\Gamma. \end{aligned} \quad (283)$$

As the $U_{(j)}$ are disjoint, $p_{(j)}$ is the only point in its $\approx_x^1$ -equivalence class in $X_{(j)}$, so the action of Γ on $X_{(j)}$ must fix $p_{(j)}$. The equalities $X_{(k)} = X_{(j)}/\Gamma$, $p_{(k)} = p_{(j)}\Gamma$ now imply that $\text{Stab}_{X^1}(p_{(j)})$ is a normal subgroup of $\text{Stab}_{X^1}(p_{(k)})$ with $\Gamma \cong \text{Stab}_{X^1}(p_{(k)})/\text{Stab}_{X^1}(p_{(j)})$. So by (283) we see that $\Gamma_{(j)}$ is normal in $\Gamma_{(k)}$ with $\Gamma \cong \Gamma_{(k)}/\Gamma_{(j)}$. Considering the group quotients in (283), and using uniqueness up to isomorphism of $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$, $(\dot{X}_{(k)}, \dot{\mathbf{f}}_{(k)}, \dot{\mathbf{G}}_{(k)})$, we see that $(\dot{X}_{(k)}, \dot{\mathbf{f}}_{(k)}, \dot{\mathbf{G}}_{(k)})/\Gamma_{(j)} \cong (X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)})$, and deduce that $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)}) \cong (\dot{X}_{(k)}, \dot{\mathbf{f}}_{(k)}, \dot{\mathbf{G}}_{(k)})$, in this case with an isomorphism-preserving isomorphism.

This proves that the $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$ are all isomorphic for all j , possibly with orientation changes. Choose one and write it as $(\dot{X}, \dot{\mathbf{f}}, \dot{\mathbf{G}})$, and then identify $(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$ with $(\epsilon_{(j)}\dot{X}, \dot{\mathbf{f}}, \dot{\mathbf{G}})$ for $j = 1, \dots, n$, where $\epsilon_{(j)} = \pm 1$ is to change orientations if necessary.

We are now ready to prove Step 3. We first derive a modified version of equation (267). Fix $a \in A$, and consider the term $\sum_{c \in C_a} \partial[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$ on the left hand side of (267). Since X_a is ‘cut into pieces’ X_{ac} for $c \in C_a$, we can divide the boundaries ∂X_{ac} into ‘external’ components, which are part of ∂X_a , and ‘internal’ components, which are new boundary components created

by cutting the interior of X_a along some hypersurface. When we cut X_a along a hypersurface, the two sides of the cut yield identical boundary components with opposite orientations. Therefore the contributions of the ‘internal’ components to $\sum_{c \in C_a} \partial[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$ cancel out.

This leaves only the ‘external’ components, which are the pieces into which $(\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a})$ is ‘cut’ using the tent function $\mathbf{T}_a|_{\partial X_a}$. Write these as $(X_{ac'}^\partial, \mathbf{f}_{ac'}^\partial, \mathbf{G}_{ac'}^\partial)$ for $c' \in C_a^\partial$. Then in $\widehat{KC}_{k-1}(Y; R)$ we have

$$\sum_{c \in C_a} \partial[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}] = \sum_{c' \in C_a^\partial} [X_{ac'}^\partial, \mathbf{f}_{ac'}^\partial, \mathbf{G}_{ac'}^\partial]. \quad (284)$$

Substituting (284) into the left hand of (267) yields

$$\begin{aligned} \sum_{a \in A} \sum_{c' \in C_a^\partial} \rho_a [X_{ac'}^\partial, \mathbf{f}_{ac'}^\partial, \mathbf{G}_{ac'}^\partial] = \\ \sum_{d \in D} \eta_d \left(\sum_{f \Gamma_d \in F_d / \Gamma_d} [\tilde{X}_{df} / \text{Stab}_{\Gamma_d}(f), \pi_*(\tilde{\mathbf{f}}_{df}), \pi_*(\tilde{\mathbf{G}}_{df})] \right. \\ \left. - |\Gamma_d|^{-1} \sum_{f \in F_d} [\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df}] \right). \end{aligned} \quad (285)$$

We must prove that both sides of (285) are zero. Note that as X_a for $a \in A$ have trivial stabilizers, so does ∂X_a , and so all $X_{ac'}^\partial$ on the left hand side of (285) have *trivial stabilizers*.

Our approach is to work locally in $\ddot{X}^1$. Now each term $X_{ac'}^\partial, \tilde{X}_{df} / \text{Stab}_{\Gamma_d}(f), \tilde{X}_{df}$ in (285) has a natural projection π to X^1 , and so to $\ddot{X}^1$ via $\pi_{\ddot{X}^1} : X^1 \rightarrow \ddot{X}^1$. (This did not hold for (267), as the ‘internal’ boundary components of ∂X_a do not project to X^1 .) Going through the proof of (285), one can show that it also holds locally in $\ddot{X}^1$. Thus, if we fix $\ddot{p} \in \ddot{X}^1$ and restrict to those terms in (285) whose image in $\ddot{X}^1$ contains $\ddot{p}$, the resulting equation still holds. We will show that each side of this resulting equation is zero.

For $a \in A$, the sum of terms in $\sum_{c' \in C_a^\partial} [X_{ac'}^\partial, \mathbf{f}_{ac'}^\partial, \mathbf{G}_{ac'}^\partial]$ whose image in $\ddot{X}^1$ contain $\ddot{p}$ is

$$\sum_{j=1, \dots, n: p_{(j)} \in \partial X_a} [X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)}]. \quad (286)$$

Now if $p_{(j)} \in \partial X_a$ then $\text{Stab}(p_{(j)}) = \{1\}$ as ∂X_a has trivial stabilizers, so $\Gamma_{(j)} = \{1\}$ by (283), giving $(X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)}) \cong \epsilon_{(j)}(\dot{X}_{(j)}, \dot{\mathbf{f}}_{(j)}, \dot{\mathbf{G}}_{(j)})$. So another way to write (286) is

$$\sum_{j=1, \dots, n: p_{(j)} \in \partial X_a} \epsilon_{(j)}[\dot{X}, \dot{\mathbf{f}}, \dot{\mathbf{G}}]. \quad (287)$$

Similarly, for $d \in D$ the sum of terms in $\sum_{f \in F_d} [\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df}]$ whose image in $\ddot{X}^1$ contains $\ddot{p}$ is

$$\sum_{j=1, \dots, n: p_{(j)} \in \tilde{X}_d} [X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)}]. \quad (288)$$

The sum of terms in $\sum_{f \Gamma_d \in F_d / \Gamma_d} [\tilde{X}_{df} / \text{Stab}_{\Gamma_d}(f), \pi_*(\tilde{\mathbf{f}}_{df}), \pi_*(\tilde{\mathbf{G}}_{df})]$ whose images contain $\ddot{p}$ is the quotient of (288) by Γ_d acting on $\coprod_{j=1, \dots, n: p_{(j)} \in \tilde{X}_d} X_{(j)}$. This action of Γ_d factors through an action on $\{j = 1, \dots, n : p_{(j)} \in \tilde{X}_d\}$.

Hence we may write the sum of terms in $[\tilde{X}_{df}/\text{Stab}_{\Gamma_d}(f), \pi_*(\tilde{\mathbf{f}}_{df}), \pi_*(\tilde{\mathbf{G}}_{df})]$ whose images contain $\check{p}$ as

$$\sum_{j \in \Gamma_d \in \{j=1, \dots, n: p_{(j)} \in \tilde{X}_d\}/\Gamma_d} [X_{(j)}/\text{Stab}_{\Gamma_d}(j), \pi_*(\mathbf{f}_{(j)}), \pi_*(\mathbf{G}_{(j)})]. \quad (289)$$

Here $(X_{(j)}/\text{Stab}_{\Gamma_d}(j), \pi_*(\mathbf{f}_{(j)}), \pi_*(\mathbf{G}_{(j)}))$ is one of the quotients $(X_{(k)}^1, \mathbf{f}_{(k)}^1, \mathbf{G}_{(k)}^1) = (X_{(j)}^1/\Gamma, \pi_*(\mathbf{f}_{(j)}^1), \pi_*(\mathbf{G}_{(j)}^1))$ above.

Combining (285)–(289), replacing $[X_{(j)}, \mathbf{f}_{(j)}, \mathbf{G}_{(j)}]$ by $\epsilon_{(j)}[\dot{X}/\Gamma_{(j)}, \pi_*(\dot{\mathbf{f}}), \pi_*(\dot{\mathbf{G}})]$, and restricting to terms in (285) whose images contain $\check{p}$ yields

$$\begin{aligned} & \sum_{a \in A} \sum_{j=1, \dots, n: p_{(j)} \in \partial X_a} \rho_a \epsilon_{(j)} [\dot{X}, \dot{\mathbf{f}}, \dot{\mathbf{G}}] = \\ & \sum_{d \in D} \eta_d \left(\sum_{j \in \Gamma_d \in \{j=1, \dots, n: p_{(j)} \in \tilde{X}_d\}/\Gamma_d} \epsilon_{(j)} [\dot{X}/\text{Stab}_{\Gamma_d}(j) \times \Gamma_{(j)}, \pi_*(\dot{\mathbf{f}}), \pi_*(\dot{\mathbf{G}})] \right. \\ & \quad \left. - |\Gamma_d|^{-1} \sum_{j=1, \dots, n: p_{(j)} \in \tilde{X}_d} \epsilon_{(j)} [\dot{X}/\Gamma_{(j)}, \pi_*(\dot{\mathbf{f}}), \pi_*(\dot{\mathbf{G}})] \right). \end{aligned} \quad (290)$$

Equation (290) holds in $\widetilde{KC}_{k-1}(Y; R)$. Projecting to $KC_{k-1}(Y; R)$, each term $(\dots)$ on the right hand side vanishes by relation Definition 4.3(iv), which holds in $KC_{k-1}(Y; R)$ but not in $\widetilde{KC}_{k-1}(Y; R)$. Thus

$$\sum_{a \in A} \sum_{j=1, \dots, n: p_{(j)} \in \partial X_a} \rho_a \epsilon_{(j)} [\dot{X}, \dot{\mathbf{f}}, \dot{\mathbf{G}}] = 0 \quad \text{in } KC_{k-1}(Y; R). \quad (291)$$

There are now two possibilities: either there exists an orientation-reversing isomorphism (a, b) of $(\dot{X}, \dot{\mathbf{f}}, \dot{\mathbf{G}})$, so that $[\dot{X}, \dot{\mathbf{f}}, \dot{\mathbf{G}}] = 0$ in both $KC_{k-1}(Y; R)$ and $\widetilde{KC}_{k-1}(Y; R)$ by Definition 4.3(ii), or the coefficient of $[\dot{X}, \dot{\mathbf{f}}, \dot{\mathbf{G}}]$ is zero in (291). (More complicated scenarios of having to use several applications of relations in $KC_{k-1}(Y; R)$ to prove $[\dot{X}, \dot{\mathbf{f}}, \dot{\mathbf{G}}] = 0$ can be excluded, because of the general form of (290).) In both cases, (291) also holds in $\widetilde{KC}_{k-1}(Y; R)$. Hence both sides of (290) are zero in $\widetilde{KC}_{k-1}(Y; R)$.

We have shown that for a given $\check{p} \in \check{X}^1$, if the $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ are chosen to make the pieces $X_{ac}, \tilde{X}_{df}$ sufficiently small near the preimage of $\check{p}$, then on each side of (285), the sum of terms whose image in $\check{X}^1$ contains $\check{p}$ is zero. On $\check{X}^1$, we can write the ‘sufficiently small’ condition as requiring that $\pi(X_{ac}), \pi(\tilde{X}_{df}) \subseteq \check{U}_{\check{p}}$ for any $X_{ac}, \tilde{X}_{df}$ whose image contains $\check{p}$, where $\check{U}_{\check{p}}$ is the intersection in $\check{X}^1$ of the images of $U_{(j)}$ for $j = 1, \dots, n$, so that $\check{U}_{\check{p}}$ is an open neighbourhood of $\check{p}$ in $\check{X}^1$. Such open sets $\check{U}_{\check{p}}$ for all $\check{p}$ in $\check{X}^1$ form an open cover of $\check{X}^1$, which is compact. So we can choose a finite subset $\check{p}^1, \dots, \check{p}^l$ with $\check{X}^1 = \check{U}_{\check{p}^1} \cup \dots \cup \check{U}_{\check{p}^l}$.

We can then choose the $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ to make the $X_{ac}^\partial, \tilde{X}_{df}$ small enough that each $\pi(X_{ac}^\partial)$ or $\pi(\tilde{X}_{df})$ lies in $\check{U}_{\check{p}^i}$ for some $i = 1, \dots, l$, and with the corresponding smallness conditions in $V^{1,i}$, so that each $(X_{ac'}^\partial, \mathbf{f}_{ac'}^\partial, \mathbf{G}_{ac'}^\partial)$ and $(\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df})$ can be written naturally as a quotient of a triple with trivial stabilizers by a finite group, which is the stabilizer group of a preimage of some $\check{p}^i$ in $\check{X}^1$.

This method does not necessarily ensure that every $\pi(X_{ac'}^\partial)$ or $\pi(\tilde{X}_{df})$ contains some $\check{p}^i$, only that it lies in some $\check{U}_{\check{p}^i}$. But this does not matter, as we can rewrite the argument above, replacing restricting to terms in (285) whose image in $\check{X}^1$ contains $\check{p}^i$ for some $i = 1, \dots, l$ by restricting to terms in (285) whose image in $\check{X}^1$ lies in $\check{U}_{\check{p}^i}$ for some $i = 1, \dots, l$.

Thus, with the $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ chosen in this way, equation (285) has the following properties. Firstly, for every term in (285), the projection of $X_{ac'}^\partial$ or $\tilde{X}_{df}/\text{Stab}_{\Gamma_d}(f)$ or $\tilde{X}_{df}$ to $\check{X}^1$ lies in $\check{U}_{\check{p}^i}$ for at least one $i = 1, \dots, l$. Secondly, for each $i = 1, \dots, l$, if we restrict to the terms in (285) whose projection to $\check{X}^1$ lies in $\check{U}_{\check{p}^i}$, the resulting equation is true, and also, both sides are zero. Thirdly, an easy generalization of the second property shows that for any subset S of $\{1, \dots, l\}$, if we restrict to the terms in (285) whose projection to $\check{X}^1$ lies in $\check{U}_{\check{p}^i}$ for all $i \in S$, the resulting equation is true, and also, both sides are zero.

By the first property, we see that summing over all $\emptyset \neq S \subseteq \{1, \dots, l\}$ of the restriction to the terms in (285) whose projection to $\check{X}^1$ lies in $\check{U}_{\check{p}^i}$ for all $i \in S$, multiplied by $(-1)^{|S|-1}$, yields (285). But the third property says that each of these restrictions has both sides zero, so (285) has both sides zero. Therefore both sides of (267) are zero. This proves (268), and the first part of Step 3.

It remains to show that if the X_{ac} are chosen sufficiently small, for all $a \in A$, $c \in C_a$ and $m \geq 2$, each component of $(\partial^m X_{ac}, \mathbf{f}_{ac}|_{\partial^m X_{ac}}, \mathbf{G}_{ac}|_{\partial^m X_{ac}})$ occurs as the intersection of m *distinct* components of $(\partial X_{ac}, \mathbf{f}_{ac}|_{\partial X_{ac}}, \mathbf{G}_{ac}|_{\partial X_{ac}})$. We use notation $(X_{acd}^m, \mathbf{f}_{acd}^m, \mathbf{G}_{acd}^m)$ for $d = 1, \dots, n_{ac}^m$, $V_{acd}^{m, i-m} \subseteq \partial^m V_{ac}^i$ and $(v, B_1, \dots, B_m)$ for points in $\partial^m V_{ac}^i$ as in Step 3.

Recall that $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ is a piece in the decomposition of $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ using $\mathbf{T}_a = (T_a^i : i \in I_a)$, where $T_a^i : \dot{V}_a^i \rightarrow F([1, \infty))$ is a tent function on the manifold $\dot{V}_a^i$. By Definition A.1, each $v \in \dot{V}_a^i$ has an open neighbourhood U in $\dot{V}_a^i$ such that $T_a^i|_U$ may be written $T_a^i(u) = \{t_p(u) : p = 1, \dots, N, u \in U_p\}$, for some open sets $U_1, \dots, U_N$ in U . Considering the constructions of tent functions in §A, we see that how small this open set U needs to be can be made to depend only on $\dot{V}_a^i$ and other data in $(X_a, \mathbf{f}_a, \mathbf{G}_a)$, not on the choices in $\mathbf{T}_a$ which make the $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ small.

Therefore by choosing $\mathbf{T}_a$ to make the pieces $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ sufficiently small, we can arrange that $\pi(V_{ac}^i)$ for each $c \in C_a$ is contained in some such open set U in $\dot{V}_a^i$. Then V_{ac}^i is a piece in the decomposition of U using the tent function $T_a^i|_U$ which has the *global* definition $T_a^i(u) = \{t_p(u) : p = 1, \dots, N, u \in U_p\}$. Consider the boundary ∂V_{ac}^i . It is a disjoint union of two types of components: (a) the lift of part of $\partial \dot{V}_a^i$, a piece in the decomposition of $\partial \dot{V}_a^i$ using $T_a^i|_{\partial \dot{V}_a^i}$, and (b) part of the hypersurface $t_p(u) = t_q(u)$ in U , for $p, q = 1, \dots, N$, $p \neq q$. To specify the order of p, q we require that $t_p - t_q$ should be increasing in the direction of the outward normal to the boundary component of X_{ac} .

Let $d = 1, \dots, n_{ac}^1$ and $i-1 \in I_{acd}^1$, so that $\emptyset \neq V_{abd}^{1, i-1} \subseteq \partial V_{ac}^i$. Then for each $(v, B) \in V_{abd}^{1, i-1}$, the classification of B into either (a) or (b) with (p, q) depends only on d . That is, if $(v, B), (v', B')$ lie in $V_{abd}^{1, i-1}$ then B is a hypersurface $t_p = t_q$ if and only if B' is a hypersurface $t_p = t_q$, with the same (p, q) . This

holds even though $V_{abd}^{1,i-1}$ may not be connected, because of the global definition of the T_a^j near $\pi(V_{ac}^j)$, as $(X_{acd}^1, \mathbf{f}_{acd}^1, \mathbf{G}_{acd}^1)$ is connected, and since the way in which we make T_a^i, T_a^j compatible under coordinate changes $(\phi_a^{ij}, \hat{\phi}_a^{ij})$ ensures that the hypersurface $t_p = t_q$ in $U^j \cap \dot{V}_a^{ij} \subseteq \dot{V}_a^j$ maps to a hypersurface $t_{p'} = t_{q'}$ in $U^i \subseteq \dot{V}_a^i$ under ϕ_a^{ij} , for p', q' depending only on p, q, i, j .

Now let $m \geq 2$ and $d = 1, \dots, n_{ac}^m$, so that $(X_{acd}^m, \mathbf{f}_{acd}^m, \mathbf{G}_{acd}^m)$ is a component of $(\partial^m X_{ac}, \mathbf{f}_{ac}|_{\partial^m X_{ac}}, \mathbf{G}_{ac}|_{\partial^m X_{ac}})$. Suppose $i - m \in I_{acd}^m$, so that $\emptyset \neq V_{acd}^{m,i-m} \subseteq \partial^m V_{ac}^i$, let $(v, B_1, \dots, B_m) \in V_{acd}^{m,i-m}$, and let $(v, B_l) \in V_{acd_l}^{1,i-1}$ for $l = 1, \dots, m$ and $d_l = 1, \dots, n_{ac}^1$. We must show that $d_1, \dots, d_m$ are *distinct*. Suppose $d_l = d_{l'}$ for $l \neq l'$. As V_a^i is a manifold, $B_1, \dots, B_m$ are distinct, so $B_l \neq B_{l'}$. As above, $B_l, B_{l'}$ are classified into either (a) or (b) with (p, q) , and this classification depends only on $d_l = d_{l'}$. Thus, if B_l is the hypersurface $t_p = t_q$ near v then so is $B_{l'}$, contradicting $B_l \neq B_{l'}$. So the only possibility is that $B_l, B_{l'}$ are of type (a), that is, they are local boundary components of V_a^i .

Since V_a^i is a manifold, we can exclude this possibility by making the X_{ac} sufficiently small, as it would mean that at $v \in V_a^i$ we have two distinct local boundary components $B_l, B_{l'}$ which are connected via some path in $\partial V_a^i \cap \pi(V_{ac}^i)$, or by some more complicated path involving coordinate changes $(\phi_a^{ij}, \hat{\phi}_a^{ij})$. This can only happen if some nontrivial global topology in V_a^i happens in the image $\pi(V_{ac}^i)$, which is ruled out by making the V_{ac}^i small. Therefore $d_1, \dots, d_m$ are distinct, as we have to prove. This completes Step 3.

C.4 Step 4: lifting to a cycle in $KC_k^{\text{ef}}(Y; R)$

As in Step 4, we again change notation from X_{ac} back to X_a , and introduce new notation $n_a^m, \dots, B_{ab}^m$. Also, for $a \in A$ and $0 \leq m \leq m+l \leq M$ define $\theta_a^{ml} : \{1, \dots, n_a^{m+l}\} \rightarrow \{1, \dots, n_a^m\}$ by $\theta_a^{ml}(\bar{b}) = b$ if $X_{ab}^{m+l} \subseteq \partial^l X_{ab}^m$. Then θ_a^{m0} is the identity, and

$$(\partial^l X_{ab}^m, \mathbf{f}_{ab}^m|_{\partial^l X_{ab}^m}, \mathbf{G}_{ab}^m|_{\partial^l X_{ab}^m}) = \coprod_{\bar{b}=1, \dots, n_a^{m+l} : \theta_a^{ml}(\bar{b})=b} (X_{ab}^{m+l}, \mathbf{f}_{ab}^{m+l}, \mathbf{G}_{ab}^{m+l}) \quad (292)$$

is the decomposition of $\partial^l X_{ab}^m$ given by Lemma 3.7.

We represent α by a cycle $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$ in $\widetilde{KC}_k(Y; R)$ satisfying (269), where each X_a has trivial stabilizers, each V_a^i is a manifold, each component of $(\partial^m X_a, \mathbf{f}_a|_{\partial^m X_a}, \mathbf{G}_a|_{\partial^m X_a})$ occurs as the intersection of m distinct components of $(\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a})$, and (270) holds with each $(\bar{a}, \bar{b}) \in R^1$ satisfying (A) or (B) in Step 4. Our goal is to choose finite families $\underline{\mathbf{G}}_{ao}$ for $o \in O_a$ of effective gauge-fixing data for $(X_a, \mathbf{f}_a)$, such that $\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_{ao}]$ is an effective Kuranishi cycle in class $\beta \in KH_k^{\text{ef}}(Y; R)$ with $\Pi_{\text{ef}}^{\text{Kh}}(\beta) = \alpha$.

Here is our inductive hypothesis:

Hypothesis C.7. Let $N \geq 0$ be given. Suppose that for all $m = N, N+1, \dots, M$ and all $(a, b) \in R^m$, we have chosen the following data:

- (i) A nonempty finite set O_{ab}^m ;

- (ii) A free left action of the finite group $\text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ on O_{ab}^m ;
(iii) For all $o \in O_{ab}^m$ and $l \geq 0$, a map

$$f_{abo}^{ml} : \{\bar{b} = 1, \dots, n_a^{m+l} : \theta_a^{ml}(\bar{b}) = b\} \longrightarrow \coprod_{\substack{\bar{b}=1, \dots, n_a^{m+l} \\ \theta_a^{ml}(\bar{b})=b}} O_{\phi^{m+l}(a, \bar{b})}^{m+l} \quad (293)$$

with $f_{abo}^{ml}(\bar{b}) \in O_{\phi^{m+l}(a, \bar{b})}^{m+l}$ for all $\bar{b}$; and

- (iv) For each $o \in O_{ab}^m$, a family of maps $\underline{\mathbf{G}}_{abo}^{m,i} : E_{ab}^{m,i} \rightarrow \underline{P}$ for $i \in I_{ab}^m$ such that $\underline{\mathbf{G}}_{abo}^m = (I_{ab}^m, \eta_{ab}^m, \underline{\mathbf{G}}_{abo}^{m,i} : i \in I_{ab}^m)$ is effective gauge-fixing data for $(X_{ab}^m, \mathbf{f}_{ab}^m)$. Thus $\underline{\mathbf{G}}_{abo}^{m,i}$ must satisfy the injectivity condition Definition 3.36(b).

These should satisfy the following conditions:

- (a) If $o, o' \in O_{ab}^m$ and $\underline{\mathbf{G}}_{abo}^{m,i} \equiv \underline{\mathbf{G}}_{abo'}^{m,i}$ for all $i \in I_{ab}^m$, then $o = o'$.
(b) Let $o \in O_{ab}^m$ and $(\mathbf{u}, \mathbf{v}) \in \text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$, and set $o' = (\mathbf{u}, \mathbf{v}) \cdot o$ using the action in (ii). Then for all $i \in I_{ab}^m$ we have diffeomorphisms $\hat{v}^i : E_{ab}^{m,i} \rightarrow E_{ab}^{m,i}$, and we require that $\underline{\mathbf{G}}_{abo}^{m,i} \equiv \underline{\mathbf{G}}_{abo'}^{m,i} \circ \hat{v}^i$.
(c) Let $(a, b) \in R^m$, $i \in I_{ab}^m$, $o \in O_{ab}^m$ and $l \geq 0$. Then (292) implies that

$$\begin{aligned} \partial^l E_{ab}^{m,i} &= \coprod_{\bar{b}=1, \dots, n_a^{m+l} : \theta_a^{ml}(\bar{b})=b} E_{a\bar{b}}^{m+l, i-l} \\ &\stackrel{\cong}{\simeq} \coprod_{\bar{b}=1, \dots, n_a^{m+l} : \theta_a^{ml}(\bar{b})=b} E_{\phi^{m+l}(a, \bar{b})}^{m+l, i-l}, \end{aligned} \quad (294)$$

where in the notation of Step 4 we have an isomorphism $(\mathbf{a}, \mathbf{b})_{a\bar{b}}^{m+l} : (X_{a\bar{b}}^{m+l}, \mathbf{f}_{a\bar{b}}^{m+l}, \mathbf{G}_{a\bar{b}}^{m+l}) \rightarrow (X_{\phi^{m+l}(a, \bar{b})}^{m+l}, \mathbf{f}_{\phi^{m+l}(a, \bar{b})}^{m+l}, \mathbf{G}_{\phi^{m+l}(a, \bar{b})}^{m+l})$ which includes a diffeomorphism $\hat{b}_{a\bar{b}}^{m+l, i-l} : E_{a\bar{b}}^{m+l, i-l} \rightarrow E_{\phi^{m+l}(a, \bar{b})}^{m+l, i-l}$, and we use these diffeomorphisms to make the identification in the second line of (294).

We have functions $\underline{\mathbf{G}}_{abo}^{m,i} : E_{ab}^{m,i} \rightarrow \underline{P}$ and $\underline{\mathbf{G}}_{\phi^{m+l}(a, \bar{b}) f_{abo}^{ml}(\bar{b})}^{m+l, i-l} : E_{\phi^{m+l}(a, \bar{b})}^{m+l, i-l} \rightarrow \underline{P}$ for each $\bar{b}$ with $\theta_a^{ml}(\bar{b}) = b$. We require that

$$\underline{\mathbf{G}}_{abo}^{m,i} \big|_{\partial^l E_{ab}^{m,i}} \equiv \coprod_{\bar{b}=1, \dots, n_a^{m+l} : \theta_a^{ml}(\bar{b})=b} \underline{\mathbf{G}}_{\phi^{m+l}(a, \bar{b}) f_{abo}^{ml}(\bar{b})}^{m+l, i-l} \quad (295)$$

under the identification (294).

When $l = 0$, θ_a^{m0} is the identity, so the domain of f_{abo}^{m0} is $\{b\}$, and if $f_{abo}^{m0}(b) = o'$ then (294) reduces to $E_{ab}^{m,i} = E_{ab}^{m,i}$ and (295) to $\underline{\mathbf{G}}_{abo}^{m,i} \equiv \underline{\mathbf{G}}_{abo'}^{m,i}$. Thus (a) implies that $f_{abo}^{m0}(b) = o$.

Before giving the two remaining conditions, we define some notation. Fix $m \geq 0$, and suppose that we have chosen data $O_{ab}^{m'}, f_{abo}^{m'}, \underline{\mathbf{G}}_{abo}^{m'}$ for all $m' > m$ and all a, b, o, i, l satisfying (i)–(iv) and (a)–(c) above. Then in an inductive step we will wish to choose $O_{ab}^m, f_{abo}^m, \underline{\mathbf{G}}_{abo}^m$ for all a, b, o, i, l . For $l > 0$, consider

which maps $f_{ab}^{ml} : \{\bar{b} = 1, \dots, n_a^{m+l} : \theta_a^{ml}(\bar{b}) = b\} \rightarrow \coprod_{\bar{b}} O_{\phi^{m+l}(a, \bar{b})}^{m+l}$ could be f_{abo}^{ml} in (293) for some $o \in O_{ab}^m$. We will derive a necessary condition on such f_{ab}^{ml} .

Suppose that for some $o \in O_{ab}^m$ we have chosen maps f_{abo}^{ml} for $l \geq 0$ and $\underline{G}_{abo}^{m,i}$ for $i \in I_{ab}^m$ satisfying (iii),(iv),(c) above. Let $l > 0$. Then by (c) for a, b, m and (a),(c) for $m' = m + l > m$, we see that f_{abo}^{ml} uniquely determines and is determined by the restrictions $\underline{G}_{abo}^{m,i}|_{\partial^l E_{ab}^{m,i}}$ for $i \in I_{ab}^m$. Since these determine $\underline{G}_{abo}^{m,i}|_{\partial^{l'} E_{ab}^{m,i}}$ for $l' > l$, it follows that f_{abo}^{ml} determines $f_{abo}^{ml'}$ for all $l' > l$.

Observe that there is a natural action of the symmetric group S_l on $\partial^l E_{ab}^{m,i}$ generalizing the involution $\sigma : \partial^2 E_{ab}^{m,i} \rightarrow \partial^2 E_{ab}^{m,i}$ discussed in Definitions 2.7 and 2.9. The restriction (pullback) $\underline{G}_{abo}^{m,i}|_{\partial^l E_{ab}^{m,i}}$ is invariant under this S_l action on $\partial^l E_{ab}^{m,i}$. This gives rise to a consistency condition on f_{abo}^{ml} . Suppose f_{ab}^{ml} is some map as in (293), with $f_{ab}^{ml}(\bar{b}) \in O_{\phi^{m+l}(a, \bar{b})}^{m+l}$ for all $\bar{b}$. Then (c) determines functions $\partial^l E_{ab}^{m,i} \rightarrow \underline{P}$ for $i \in I_{ab}^m$, by the r.h.s. of (295) with f_{ab}^{ml} in place of f_{abo}^{ml} . To be of the form $\underline{G}_{abo}^{m,i}|_{\partial^l E_{ab}^{m,i}}$, these functions must be S_l -invariant.

We say f_{ab}^{ml} is S_l -invariant if the functions it determines are S_l -invariant in this way. We say f_{ab}^{ml} is allowable if it is S_l -invariant, and also each map $f_{ab}^{ml'}$ for $l' > l$ determined by f_{ab}^{ml} is $S_{l'}$ -invariant. Write D_{ab}^{ml} for the set of allowable maps f_{ab}^{ml} . When $l = 0$, each f_{ab}^{m0} maps $\{b\} \rightarrow O_{ab}^m$, and for each $o \in O_{ab}^m$ we have $f_{abo}^{m0} : b \rightarrow o$, which is allowable. Thus D_{ab}^{m0} is the set of maps $\{b\} \rightarrow O_{ab}^m$, and $f_{ab}^{m0} \mapsto f_{ab}^{m0}(b)$ induces a bijection $D_{ab}^{m0} \rightarrow O_{ab}^m$.

When $l' > l \geq 0$, write $\Pi_{ab}^{ml'} : D_{ab}^{ml} \rightarrow D_{ab}^{ml'}$ for the natural projection taking $\Pi_{ab}^{ml'} : f_{abo}^{ml} \mapsto f_{abo}^{ml'}$, defined by restriction from $\partial^l E_{ab}^{m,i}$ to $\partial^{l'} E_{ab}^{m,i}$ as above.

Suppose $\tilde{m} > m \geq 0$ and $l > \tilde{l} \geq 0$ with $m + l = \tilde{m} + \tilde{l}$, and that $\hat{b} = 1, \dots, n_a^{\tilde{m}}$ with $\theta_a^{m(\tilde{m}-m)}(\hat{b}) = b$. Set $(\tilde{a}, \tilde{b}) = \phi^{\tilde{m}}(a, \hat{b})$. Then we have an isomorphism $(a, b)_{ab}^{\tilde{m}} : (X_{ab}^{\tilde{m}}, \mathbf{f}_{ab}^{\tilde{m}}, \mathbf{G}_{ab}^{\tilde{m}}) \rightarrow (X_{\tilde{a}\tilde{b}}^{\tilde{m}}, \mathbf{f}_{\tilde{a}\tilde{b}}^{\tilde{m}}, \mathbf{G}_{\tilde{a}\tilde{b}}^{\tilde{m}})$. Composing this with the inclusion $X_{ab}^{\tilde{m}} \subseteq \partial^{\tilde{m}-m} X_{ab}^m$ and applying $\partial^{\tilde{l}}$ to both sides induces a morphism $(a_{ab}^{\tilde{m}})^{-1} : \partial^{\tilde{l}} X_{ab}^{\tilde{m}} \rightarrow \partial^l X_{ab}^m$ which is an isomorphism with a union of components of $\partial^l X_{ab}^m$.

Since the components of $\partial^{\tilde{l}} X_{\tilde{a}\tilde{b}}^{\tilde{m}}$ are $X_{\tilde{a}\tilde{b}}^{\tilde{m}+\tilde{l}}$ for $\tilde{b} = 1, \dots, n_a^{\tilde{m}+\tilde{l}}$ with $\theta_a^{\tilde{m}\tilde{l}}(\tilde{b}) = \tilde{b}$, and the components of $\partial^l X_{ab}^m$ are $X_{ab'}^{m+l}$ for $b' = 1, \dots, n_a^{m+l}$ with $\theta_a^{ml}(b') = b$, there is a natural, injective map $I_{abb}^{ml\tilde{m}\tilde{l}} : \{\tilde{b} = 1, \dots, n_a^{\tilde{m}+\tilde{l}} : \theta_a^{\tilde{m}\tilde{l}}(\tilde{b}) = \tilde{b}\} \rightarrow \{b' = 1, \dots, n_a^{m+l} : \theta_a^{ml}(b') = b\}$ such that $(a_{ab}^{\tilde{m}})^{-1}(X_{\tilde{a}\tilde{b}}^{\tilde{m}+\tilde{l}}) = X_{ab'}^{m+l}$ when $I_{abb}^{ml\tilde{m}\tilde{l}}(\tilde{b}) = b'$. Let $f_{ab}^{ml} \in D_{ab}^{ml}$, and define $\tilde{f}_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}} = f_{ab}^{ml} \circ I_{abb}^{ml\tilde{m}\tilde{l}}$. Then the conditions for f_{ab}^{ml} to be allowable restrict to imply that $\tilde{f}_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$ is allowable, so $\tilde{f}_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}} \in D_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$. Thus we define a projection $P_{abb}^{ml\tilde{m}\tilde{l}} : D_{ab}^{ml} \rightarrow D_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$ mapping $P_{abb}^{ml\tilde{m}\tilde{l}} : f_{ab}^{ml} \mapsto \tilde{f}_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}} \circ I_{abb}^{ml\tilde{m}\tilde{l}}$.

Here are the final conditions:

- (d) When $l' > l \geq 0$ the projection $\Pi_{ab}^{ml'} : D_{ab}^{ml} \rightarrow D_{ab}^{ml'}$ is surjective, and the preimage of each point is the same number $|D_{ab}^{ml}|/|D_{ab}^{ml'}|$ of points.
- (e) For all $m, \tilde{m}, l, \tilde{l}, a, b, \hat{b}$ as above, $P_{abb}^{ml\tilde{m}\tilde{l}} : D_{ab}^{ml} \rightarrow D_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$ is surjective, and

the preimage of each point is $|D_{ab}^{ml}|/|D_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}|$ points.

Here in (ii), $\text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ is finite by Theorem 3.20. We shall prove by reverse induction on N that Hypothesis C.7 holds for all $N \geq 0$. For $m > M$ we have $R^m = \emptyset$, so Hypothesis C.7 is vacuous for $N > M$, the initial step. For the inductive step, we shall suppose Hypothesis C.7 holds for $N = n + 1$, and prove it holds for $N = n \geq 0$.

To do this, we will first fix $(a, b) \in R^n$ and construct the sets D_{ab}^{nl} of allowable maps f_{ab}^{nl} for $l > 0$, and prove by reverse induction on l that (d),(e) hold. Then setting $l = 1$, the maps f_{ab}^{n1} in D_{ab}^{n1} uniquely determine the possible values of $\underline{\mathbf{G}}_{abo}^{n,i}|_{\partial E_{ab}^{n,i}}$ for $i \in I_{ab}^n$. We shall choose O_{ab}^n and $\underline{\mathbf{G}}_{abo}^{n,i}$ for $o \in O_{ab}^n$ and $i \in I_{ab}^n$ with these boundary values, and prove that (a)–(e) hold.

Since the sets D_{ab}^{nl} of allowable maps f_{ab}^{nl} for $l > 0$ depend only on data O_{ab}^m and $(\underline{\mathbf{G}}_{abo}^{m,i})_{i \in I_{ab}^m}$ for $m > n$ which we have already chosen, they are already well-defined without making further choices. When $l > M - n$ we have $\partial^l X_{ab}^n = \emptyset$, so that $\{\bar{b} = 1, \dots, n_a^{n+l} : \theta_a^{nl}(\bar{b}) = b\} = \emptyset$, and the only possible map (293) is the trivial map $\emptyset \rightarrow \emptyset$, which we write $\emptyset$. It is allowable, so $D_{ab}^{nl} = \{\emptyset\}$, that is, the one point set with element $\emptyset$. Thus for $l' > l > M - n$ part (d) is immediate, and for $l > M - n$ part (e) is trivial as there are no such $\hat{b}$.

Suppose by induction that for some $p > 0$, (d) holds for $m = n$ and all $l' > l > p$, and (e) holds for $m = n$ and all $l > p$. This holds for $p \geq M - n$ from above. We shall prove that (d) holds for all $m = n$ and $l' > l = p$, and (e) holds for $m = n$ and $l = p$. Then by induction (d),(e) hold for all $l', l > 0$. For (d) it is sufficient to prove the case $l = p$ and $l' = p + 1$, as the case $l' > p + 1$ follows by composing projections $D_{ab}^{np} \rightarrow D_{ab}^{n(p+1)} \rightarrow D_{ab}^{nl'}$ and using the inductive hypothesis.

For the first part of (d) we must show that the projection $\Pi_{ab}^{np(p+1)} : D_{ab}^{np} \rightarrow D_{ab}^{n(p+1)}$ is surjective. Fix $f_{ab}^{n(p+1)}$ in $D_{ab}^{n(p+1)}$. We will construct f_{ab}^{np} in D_{ab}^{np} which restricts to $f_{ab}^{n(p+1)}$. Let $\hat{b}$ lie in the domain of f_{ab}^{np} , that is, $\hat{b} = 1, \dots, n_a^{n+p}$ with $\theta_a^{np}(\hat{b}) = b$. Set $(a', b') = \phi^{n+p}(a, \hat{b})$. We shall construct the set of possible values of $f_{ab}^{np}(\hat{b})$ in $O_{a'b'}^{n+p}$. Consider the diagram:

$$\begin{array}{ccccc}
 D_{ab}^{n(p+1)} & \xleftarrow{\Pi_{ab}^{np(p+1)}} & D_{ab}^{np} & & \\
 \downarrow P_{ab\hat{b}}^{n(p+1)(n+p)1} & & \downarrow P_{ab\hat{b}}^{np(n+p)0} & \searrow f_{ab}^{np} \mapsto f_{ab}^{np}(\hat{b}) & \\
 D_{a'b'}^{(n+p)1} & \xleftarrow{\Pi_{a'b'}^{(n+p)01}} & D_{a'b'}^{(n+p)0} & \xrightarrow[\cong]{f_{a'b'}^{(n+p)0} \mapsto f_{a'b'}^{(n+p)0}(b')} & O_{a'b'}^{n+p}
 \end{array} \tag{296}$$

It is easy to show from the definitions that (296) commutes. Thus a necessary condition for $f_{ab}^{np} \in D_{ab}^{np}$ to satisfy $\Pi_{ab}^{np(p+1)}(f_{ab}^{np}) = f_{ab}^{n(p+1)}$, as in (d), is that if $f_{ab}^{np}(\hat{b}) = o' \in O_{a'b'}^{n+1}$ then $\Pi_{a'b'}^{(n+p)01}(\{b'\} \mapsto o') = P_{ab\hat{b}}^{n(p+1)(n+p)1}(f_{ab}^{n(p+1)})$, where $\{b'\} \mapsto o'$ in $D_{a'b'}^{(n+p)0}$ is the map $\{b'\} \rightarrow O_{a'b'}^{n+1}$ taking $b' \mapsto o'$.

Since $p > 0$, Hypothesis C.7(d) with $m = n + p > n$ implies that $\Pi_{a'b'}^{(n+p)01}$ is surjective, and the preimage of each point is $|D_{a'b'}^{(n+p)0}|/|D_{a'b'}^{(n+p)1}|$ points. Thus,

for each $\hat{b}$ in the domain of f_{ab}^{np} there is a nonempty set $S_{\hat{b}} \subseteq O_{a'b'}^{n+1}$ of possible values o' for $f_{ab}^{np}(\hat{b})$ making (296) commute, with $|S_{\hat{b}}| = |D_{a'b'}^{(n+p)0}|/|D_{a'b'}^{(n+p)1}|$. For f_{ab}^{np} to lie in D_{ab}^{np} it is necessary that $f_{ab}^{np}(\hat{b}) \in S_{\hat{b}}$ for all $\hat{b}$ in the domain. Any such map f_{ab}^{np} induces the fixed map $f_{ab}^{n(p+1)}$ in D_{ab}^{np} , by construction. As $f_{ab}^{n(p+1)}$ is allowable, any $f_{ab}^{n'l'}$ induced from f_{ab}^{np} for $l' > p$ is induced from $f_{ab}^{n(p+1)}$, and so is $S_{l'}$ -invariant. Therefore such an f_{ab}^{np} is allowable if and only if it is S_p -invariant.

There are natural S_p -actions on the domain $\{\hat{b} = 1, \dots, n_a^{n+p} : \theta_a^{np}(\hat{b}) = b\}$ and range $\coprod_{\hat{b}: \theta_a^{np}(\hat{b})=b} O_{\phi^{m+l}(a, \hat{b})}^{m+l}$ of f_{ab}^{np} , and f_{ab}^{np} is S_p -invariant if and only if it is equivariant w.r.t. these actions. Here is the crux of this part of the argument: by the last part of Step 3, each point of $\partial^p X_{ab}^n \subseteq \partial^{n+p} X_a$ occurs at an ordered intersection of $n+p$ *distinct* pieces X_{ad}^1 for $d = 1, \dots, n_a^1$. Thus, to each point in $\partial^p X_{ab}^n$ we can associate an ordered $(n+p)$ -tuple $(d_1, \dots, d_{n+p})$, where $d_1, \dots, d_{n+p}$ are distinct elements of $\{1, \dots, n_a^1\}$. On a connected component $X_{ab}^{n+p} \subset \partial^p X_{ab}^n$ this $n+p$ -tuple is constant. Hence, $(d_1, \dots, d_{n+p})$ depends only on $\hat{b}$ in $\{\hat{b} = 1, \dots, n_a^{n+p} : \theta_a^{np}(\hat{b}) = b\}$, where X_{ab}^{n+p} is the component of $\partial^p X_{ab}^n$ containing the chosen point.

Now S_p acts on $\partial^p X_{ab}^n$ by permuting p out of the $n+p$ pieces X_{ad}^1 meeting at the point. Thus, the S_p -action is compatible with an action of S_p on $n+p$ -tuples $(d_1, \dots, d_{n+p})$, by permuting p of d_i . Since $d_1, \dots, d_{n+p}$ are *distinct*, this action of S_p on $n+p$ -tuples $(d_1, \dots, d_{n+p})$ is free. Since $(d_1, \dots, d_{n+p})$ depends only on $\hat{b}$, and the action of S_p on the $(d_1, \dots, d_{n+p})$ is free, it follows that the action of S_p on the domain $\{\hat{b} = 1, \dots, n_a^{n+p} : \theta_a^{np}(\hat{b}) = b\}$ of f_{ab}^{np} is *free*.

We can now show that $\Pi_{ab}^{np(p+1)} : D_{ab}^{np} \rightarrow D_{ab}^{n(p+1)}$ is *surjective*. From above, for fixed $f_{ab}^{n(p+1)} \in D_{ab}^{n(p+1)}$, the preimage of $f_{ab}^{n(p+1)}$ is the set of S_p -invariant f_{ab}^{np} with $f_{ab}^{np}(\hat{b}) \in S_{\hat{b}}$ for all $\hat{b}$. To construct such f_{ab}^{np} , we choose one point $\hat{b}$ in each S_p -orbit in the domain, and choose $f_{ab}^{np}(\hat{b}) \in S_{\hat{b}}$ arbitrarily at each such $\hat{b}$. This is possible as $S_{\hat{b}} \neq \emptyset$. Then we extend f_{ab}^{np} in an S_p -invariant way. This is possible, uniquely, as S_p acts freely on the domain. So there exist such f_{ab}^{np} , and $\Pi_{ab}^{np(p+1)}$ is surjective.

This proof also gives us the size of $|(\Pi_{ab}^{np(p+1)})^{-1}(f_{ab}^{n(p+1)})|$: it is the product of $|S_{\hat{b}}|$ over representatives $\hat{b}$ of each S_p -orbit. But $|S_{\hat{b}}| = |D_{a'b'}^{(n+p)0}|/|D_{a'b'}^{(n+p)1}|$, which is independent of $f_{ab}^{n(p+1)}$, so the preimage of each $f_{ab}^{n(p+1)}$ is the same number of points, which must be $|D_{ab}^{np}|/|D_{ab}^{n(p+1)}|$. If $l' > p$ then either $l' = p+1$, when $\Pi_{ab}^{npl'}$ is surjective from above, or $l' > p+1$, when $\Pi_{ab}^{npl'} = \Pi_{ab}^{n(p+1)l'} \circ \Pi_{ab}^{np(p+1)}$. But $\Pi_{ab}^{n(p+1)l'}$ is surjective by induction, and $\Pi_{ab}^{np(p+1)}$ is surjective from above. Hence $\Pi_{ab}^{npl'}$ is surjective for all $l' > p$, which proves the first part of (d). The second part also follows, since $\Pi_{ab}^{n(p+1)l'}$ and $\Pi_{ab}^{np(p+1)}$ both have the same number of points in the preimage of each point.

Next we prove (e) holds. Let $m, l, \tilde{m}, \tilde{l}, a, b, \hat{b}, \tilde{a}, \tilde{b}$ be as in (e) with $m = n$

and $l = p$. We must show $P_{abb}^{mp\tilde{m}\tilde{l}} : D_{ab}^{np} \rightarrow D_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$ is surjective. Consider the diagram

$$\begin{array}{ccc} D_{ab}^{np} & \xrightarrow{\quad} & D_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}} \\ \downarrow \Pi_{ab}^{np(p+1)} & \begin{array}{c} P_{abb}^{np\tilde{m}\tilde{l}} \\ P_{abb}^{n(p+1)\tilde{m}(\tilde{l}+1)} \end{array} & \downarrow \Pi_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}(\tilde{l}+1)} \\ D_{ab}^{n(p+1)} & \xrightarrow{\quad} & D_{\tilde{a}\tilde{b}}^{\tilde{m}(\tilde{l}+1)}. \end{array} \quad (297)$$

It is easy to show from the definitions that (297) commutes. Fix $f_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$ in $D_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$. By (e) and induction, $P_{abb}^{n(p+1)\tilde{m}(\tilde{l}+1)}$ is surjective, so we may choose $f_{ab}^{n(p+1)}$ in $D_{ab}^{n(p+1)}$ with $P_{abb}^{n(p+1)\tilde{m}(\tilde{l}+1)}(f_{ab}^{n(p+1)}) = \Pi_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}(\tilde{l}+1)}(f_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}})$.

We now claim that there exists f_{ab}^{np} in D_{ab}^{np} with $\Pi_{ab}^{np(p+1)}(f_{ab}^{np}) = f_{ab}^{n(p+1)}$ and $P_{abb}^{np\tilde{m}\tilde{l}}(f_{ab}^{np}) = f_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$. From above, the set of f_{ab}^{np} with $\Pi_{ab}^{np(p+1)}(f_{ab}^{np}) = f_{ab}^{n(p+1)}$ is the set of S_p -invariant maps f_{ab}^{np} with $f_{ab}^{np}(\hat{b}) \in S_{\hat{b}}$ for all $\hat{b}$. By definition, $P_{abb}^{np\tilde{m}\tilde{l}}(f_{ab}^{np}) = f_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$ holds if and only if f_{ab}^{np} takes prescribed values on $\text{Im } I_{abb}^{np\tilde{m}\tilde{l}}$. Since $f_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$ is allowable, $\text{Im } I_{abb}^{np\tilde{m}\tilde{l}}$ is an S_p -invariant subset of the domain of f_{ab}^{np} , and the prescribed values are S_p -equivariant. As $P_{abb}^{n(p+1)\tilde{m}(\tilde{l}+1)}(f_{ab}^{n(p+1)}) = \Pi_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}(\tilde{l}+1)}(f_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}})$, the prescribed values on $\text{Im } I_{abb}^{np\tilde{m}\tilde{l}}$ lie in $S_{\hat{b}}$ for each $\hat{b}$.

Thus, we take f_{ab}^{np} to have the prescribed values on $\text{Im } I_{abb}^{np\tilde{m}\tilde{l}}$, and then for each S_p -orbit in the complement of $\text{Im } I_{abb}^{np\tilde{m}\tilde{l}}$ in the domain, we fix a representative $\hat{b}$, choose $f_{ab}^{np}(\hat{b}) \in S_{\hat{b}} \neq \emptyset$ arbitrarily, and extend f_{ab}^{np} uniquely to be S_p -invariant. This f_{ab}^{np} lies in D_{ab}^{np} with $P_{abb}^{np\tilde{m}\tilde{l}}(f_{ab}^{np}) = f_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$, so $P_{abb}^{np\tilde{m}\tilde{l}}$ is *surjective*. The previous argument shows that the preimage of each $f_{\tilde{a}\tilde{b}}^{\tilde{m}\tilde{l}}$ is the same number of points, proving (e).

This concludes the inductive step. Thus, we have shown by induction that (d) holds for $m = n$ and all $l' > l > 0$, and (e) holds for $m = n$ and all $l > 0$. Since D_{ab}^{nl} is the one point set $\{\emptyset\}$ for $l > M - n$, and $\Pi_{ab}^{n1l} : D_{ab}^{n1} \rightarrow D_{ab}^{nl}$ is surjective by (d), we see that when $l = 1$, the set D_{ab}^{n1} is nonempty. But functions f_{ab}^{n1} in D_{ab}^{n1} prescribe allowable boundary conditions for the data $(\underline{G}_{abo}^{n,i})_{i \in I_{ab}^n}$ for $o \in O_{ab}^n$ that we must construct. Therefore, in the proof so far we have shown that the set of allowable, consistent boundary conditions for $(\underline{G}_{abo}^{n,i})_{i \in I_{ab}^n}$ is nonempty.

We shall now choose the data of Hypothesis C.7(i)–(iv) when $m = n$. For part (i), define $O_{ab}^n = \text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n) \times D_{ab}^{n1}$ for each $(a, b) \in R^n$. Since $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)$ is a finite group by Theorem 3.20 and D_{ab}^{n1} is a nonempty finite set from above, O_{ab}^n is a nonempty finite set, so Hypothesis C.7(i) holds.

For (ii), define a left action of $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)$ on O_{ab}^n to be the product of the left action of $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)$ on itself with the natural action of $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)$ on D_{ab}^{n1} , combining the morphism $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n) \rightarrow \text{Aut}(\partial X_{ab}^n, \mathbf{f}_{ab}^n|_{\partial X_{ab}^n}, \mathbf{G}_{ab}^n|_{\partial X_{ab}^n})$ with the interpretation of D_{ab}^{n1} as a set of choices of data on $(\partial X_{ab}^n, \mathbf{f}_{ab}^n|_{\partial X_{ab}^n}, \mathbf{G}_{ab}^n|_{\partial X_{ab}^n})$. This action of $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)$ on O_{ab}^n

is *free*, since $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)$ acts freely on itself, so part (ii) holds.

For (iii), we must define maps f_{abo}^{nl} in (293) for all $o \in O_{ab}^n$ and $l \geq 0$. When $l = 0$, as in the last part of (c) we define $f_{abo}^{n0}(b) = o$ for all $o \in O_{ab}^n$. When $l = 1$, if $o \in O_{ab}^n$ then $o = ((\mathbf{u}, \mathbf{v}), f_{ab}^{n1})$ for $(\mathbf{u}, \mathbf{v}) \in \text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)$ and $f_{ab}^{n1} \in D_{ab}^{n1}$. As above, $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)$ acts on D_{ab}^{n1} . Define $\tilde{f}_{ab}^{n1} = (\mathbf{u}, \mathbf{v}) \cdot f_{ab}^{n1}$ using this action, so that $\tilde{f}_{ab}^{n1} \in D_{ab}^{n1}$. Now define $f_{abo}^{n1} = \tilde{f}_{ab}^{n1}$. This is valid as by construction, D_{ab}^{n1} is the set of possibilities for f_{abo}^{n1} . When $l > 1$ we have a map $\Pi_{ab}^{n1l} : D_{ab}^{n1} \rightarrow D_{ab}^{nl}$ and we define $f_{abo}^{nl} = \Pi_{ab}^{n1l}(f_{abo}^{n1})$. The definitions imply that $f_{abo}^{nl}(\bar{b}) \in O_{\phi^{n+l}(a, \bar{b})}^{n+l}$ for all $\bar{b}$, completing Hypothesis C.7(iii).

For (iv), write $(\mathbf{1}, \mathbf{1})$ for the identity in $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)$. We first choose the $(\underline{\mathbf{G}}_{abo}^{n,i})_{i \in I_{ab}^n}$ for each $(a, b) \in R^n$ and $o \in O_{ab}^n$ of the form $((\mathbf{1}, \mathbf{1}), f_{ab}^{n1})$ for $f_{ab}^{n1} \in D_{ab}^{n1}$. Via Hypothesis C.7(c), this f_{ab}^{n1} determines the restrictions $\underline{\mathbf{G}}_{abo}^{n,i}|_{\partial E_{ab}^{n,i}}$. That is, f_{ab}^{n1} determines maps $\underline{\mathbf{H}}_{abo}^{n,i} : \partial E_{ab}^{n,i} \rightarrow \underline{P}$ for all $i \in I_{ab}^n$, and we seek $\underline{\mathbf{G}}_{abo}^{n,i} : E_{ab}^{n,i} \rightarrow \underline{P}$ for $i \in I_{ab}^n$ with $\underline{\mathbf{G}}_{abo}^{n,i}|_{\partial E_{ab}^{n,i}} \equiv \underline{\mathbf{H}}_{abo}^{n,i}$, satisfying the conditions of Definition 3.36, so that $\underline{\mathbf{G}}_{abo}^n = (\mathbf{I}_{ab}^n, \boldsymbol{\eta}_{ab}^n, \underline{\mathbf{G}}_{abo}^{n,i} : i \in I_{ab}^n)$ is *effective gauge-fixing data* for $(X_{ab}^n, \mathbf{f}_{ab}^n)$, as in Hypothesis C.7(iv).

Since f_{ab}^{n1} is allowable, the induced f_{ab}^{n2} is invariant under $S_2 \cong \mathbb{Z}_2$. Now f_{ab}^{n2} is the restriction of f_{ab}^{n1} to $\partial^2 X_{ab}^n$, and S_2 -invariance means $\underline{\mathbf{H}}_{abo}^{n,i}|_{\partial^2 E_{ab}^{n,i}}$ is invariant under the natural involution $\sigma : \partial^2 E_{ab}^{n,i} \rightarrow \partial^2 E_{ab}^{n,i}$, for all $i \in I_{ab}^n$. But this is exactly the necessary and sufficient condition for $\underline{\mathbf{H}}_{abo}^{n,i}$ to extend to a map $\underline{\mathbf{G}}_{abo}^{n,i} : E_{ab}^{n,i} \rightarrow \underline{P}$ for $i \in I_{ab}^n$ with $\underline{\mathbf{G}}_{abo}^{n,i}|_{\partial E_{ab}^{n,i}} \equiv \underline{\mathbf{H}}_{abo}^{n,i}$, by Principle 2.8(a).

Thus, $\underline{\mathbf{H}}_{abo}^{n,i}$ prescribes consistent values for $\underline{\mathbf{G}}_{abo}^{n,i}$ on $E_{ab}^{n,i} \setminus (E_{ab}^{n,i})^\circ$, and it remains to specify $\underline{\mathbf{G}}_{abo}^{n,i}$ on $(E_{ab}^{n,i})^\circ$. Choose some $k_o \geq 0$, and maps $\underline{\mathbf{G}}_{abo}^{n,i}|_{(E_{ab}^{n,i})^\circ} : (E_{ab}^{n,i})^\circ \rightarrow \mathbb{R}^{k_o} \subset \underline{P}$ for $i \in I_{ab}^n$ such that $\coprod_{i \in I_{ab}^n} \underline{\mathbf{G}}_{abo}^{n,i}|_{(E_{ab}^{n,i})^\circ} : \coprod_{i \in I_{ab}^n} (E_{ab}^{n,i})^\circ \rightarrow \underline{P}$ is *injective*. This is possible if $k_o \gg 0$, and we can take $\underline{\mathbf{G}}_{abo}^{n,i}|_{(E_{ab}^{n,i})^\circ}$ smooth. In addition, we require that the choices of integers k_o for all $o = ((\mathbf{1}, \mathbf{1}), f_{ab}^{n1})$ for $f_{ab}^{n1} \in D_{ab}^{n1}$ should be *distinct*.

We now claim that $\underline{\mathbf{G}}_{abo}^n$ satisfies Definition 3.36. Part (a) holds as the $V_{ab}^{n,i}$ are manifolds, so $(V_{ab}^{n,i}, \dots, \psi_{ab}^{n,i})$ is automatically an effective Kuranishi neighbourhood for all $i \in I_{ab}^n$. Part (b) when $l = 0$ holds as $\coprod_{i \in I_{ab}^n} \underline{\mathbf{G}}_{abo}^{n,i}|_{(E_{ab}^{n,i})^\circ}$ is injective, and when $l > 0$ holds by induction and Hypothesis C.7(iv) for $m = n + l$, since by construction $\underline{\mathbf{G}}_{abo}^n|_{\partial^l X_{ab}^n}$ is a disjoint union of effective gauge-fixing data $\underline{\mathbf{G}}_{a'b'o'}^{n+l}$, so Definition 3.36(b) with $l = 0$ for these $\underline{\mathbf{G}}_{a'b'o'}^{n+l}$ implies Definition 3.36(b) for $\underline{\mathbf{G}}_{abo}^n$ with $l > 0$. Hence $\underline{\mathbf{G}}_{abo}^n$ is effective gauge-fixing data.

This defines the data $(\underline{\mathbf{G}}_{abo}^{n,i})_{i \in I_{ab}^n}$ for each $(a, b) \in R^n$ and $o \in O_{ab}^n$ of the form $((\mathbf{1}, \mathbf{1}), f_{ab}^{n1})$. For general $o \in O_{ab}^n$, write $o = ((\mathbf{u}, \mathbf{v}), f_{ab}^{n1})$ and $\dot{o} = (\mathbf{u}, \mathbf{v})^{-1} \cdot o = ((\mathbf{1}, \mathbf{1}), \dot{f}_{ab}^{n1})$, where $\dot{f}_{ab}^{n1} = (\mathbf{u}, \mathbf{v})^{-1} \cdot f_{ab}^{n1}$, so that we have already defined $\underline{\mathbf{G}}_{ab\dot{o}}^n$, and set $\underline{\mathbf{G}}_{abo}^n = (\mathbf{u}, \mathbf{v})_*(\underline{\mathbf{G}}_{ab\dot{o}}^n)$. That is, we define $\underline{\mathbf{G}}_{abo}^{n,i} \equiv (\hat{v}_{ab}^{n,i})_*(\underline{\mathbf{G}}_{ab\dot{o}}^{n,i}) \equiv \underline{\mathbf{G}}_{ab\dot{o}}^{n,i} \circ (\hat{v}_{ab}^{n,i})^{-1}$ for all $i \in I_{ab}^n$, where $\hat{v}_{ab}^{n,i} : E_{ab}^{n,i} \rightarrow E_{ab}^{n,i}$ is the diffeomorphism in $\mathbf{v}$. Then $\underline{\mathbf{G}}_{abo}^n$ is effective gauge-fixing data for all $o \in O_{ab}^n$, giving Hypothesis C.7(iv).

We must now prove Hypothesis C.7(a)–(e) hold when $m = n$. For (a), suppose $o = ((\mathbf{u}, \mathbf{v}), f_{ab}^{n1})$ and $\tilde{o} = ((\tilde{\mathbf{u}}, \tilde{\mathbf{v}}), \tilde{f}_{ab}^{n1})$ are distinct elements of O_{ab}^n . We must prove that $\underline{\mathbf{G}}_{abo}^n \neq \underline{\mathbf{G}}_{ab\tilde{o}}^n$. Write $f_{ab}^{n1} = (\mathbf{u}, \mathbf{v})^{-1} \cdot f_{ab}^{n1}$, $\dot{o} = ((\mathbf{1}, \mathbf{1}), \dot{f}_{ab}^{n1})$ and $\tilde{f}_{ab}^{n1} = (\tilde{\mathbf{u}}, \tilde{\mathbf{v}})^{-1} \cdot \tilde{f}_{ab}^{n1}$, $\ddot{o} = ((\mathbf{1}, \mathbf{1}), \ddot{f}_{ab}^{n1})$. Then $\underline{\mathbf{G}}_{abo}^n = (\mathbf{u}, \mathbf{v})_*(\underline{\mathbf{G}}_{ab\dot{o}}^n)$ and $\underline{\mathbf{G}}_{ab\tilde{o}}^n = (\tilde{\mathbf{u}}, \tilde{\mathbf{v}})_*(\underline{\mathbf{G}}_{ab\ddot{o}}^n)$. Since the $\underline{\mathbf{G}}_{ab\dot{o}}^{n,i}$ map $(E_{ab}^{n,i})^\circ \rightarrow \mathbb{R}^{k_{\dot{o}}}$, it follows that the $\underline{\mathbf{G}}_{abo}^{n,i}$ map $(E_{ab}^{n,i})^\circ \rightarrow \mathbb{R}^{k_{\dot{o}}}$, and similarly the $\underline{\mathbf{G}}_{ab\tilde{o}}^{n,i}$ map $(E_{ab}^{n,i})^\circ \rightarrow \mathbb{R}^{k_{\ddot{o}}}$.

If $\dot{o} \neq \ddot{o}$ then $k_{\dot{o}} \neq k_{\ddot{o}}$, as from above the k_o for all $o = ((\mathbf{1}, \mathbf{1}), f_{ab}^{n1})$ are distinct, so $\underline{\mathbf{G}}_{abo}^{n,i}|_{(E_{ab}^{n,i})^\circ}$ and $\underline{\mathbf{G}}_{ab\tilde{o}}^{n,i}|_{(E_{ab}^{n,i})^\circ}$ map to different spaces, giving $\underline{\mathbf{G}}_{abo}^{n,i} \neq \underline{\mathbf{G}}_{ab\tilde{o}}^{n,i}$ and $\underline{\mathbf{G}}_{abo}^n \neq \underline{\mathbf{G}}_{ab\tilde{o}}^n$ as we want. So suppose $\dot{o} = \ddot{o}$. Then $\underline{\mathbf{G}}_{abo}^n = (\mathbf{u}, \mathbf{v})_*(\underline{\mathbf{G}}_{ab\dot{o}}^n)$ and $\underline{\mathbf{G}}_{ab\tilde{o}}^n = (\tilde{\mathbf{u}}, \tilde{\mathbf{v}})_*(\underline{\mathbf{G}}_{ab\dot{o}}^n)$, so that $\underline{\mathbf{G}}_{ab\tilde{o}}^n = (\tilde{\mathbf{u}}, \tilde{\mathbf{v}})_*(\underline{\mathbf{G}}_{abo}^n)$, where $(\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) = (\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) \circ (\mathbf{u}, \mathbf{v})^{-1}$ in $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)$. As $o \neq \tilde{o}$ we have $(\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) \neq (\mathbf{1}, \mathbf{1})$. Since $\underline{\mathbf{G}}_{abo}^n$ is effective gauge-fixing data and $(X_{ab}^n, \mathbf{f}_{ab}^n, \underline{\mathbf{G}}_{abo}^n)$ is connected we have $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \underline{\mathbf{G}}_{abo}^n) = \{1\}$ by Theorem 3.39(b). Hence $(\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) \notin \text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \underline{\mathbf{G}}_{abo}^n)$, so $\underline{\mathbf{G}}_{abo}^n \neq (\tilde{\mathbf{u}}, \tilde{\mathbf{v}})_*(\underline{\mathbf{G}}_{abo}^n)$, and $\underline{\mathbf{G}}_{abo}^n \neq \underline{\mathbf{G}}_{ab\tilde{o}}^n$.

This proves (a). Part (b) is immediate from $\underline{\mathbf{G}}_{abo}^n = (\mathbf{u}, \mathbf{v})_*(\underline{\mathbf{G}}_{ab\dot{o}}^n)$ when $o = ((\mathbf{u}, \mathbf{v}), f_{ab}^{n1})$ and $\dot{o} = (\mathbf{u}, \mathbf{v})^{-1} \cdot o$. Part (c) when $l = 0$ is trivial, as we defined $f_{abo}^{n0}(b) = o$. For $o = ((\mathbf{1}, \mathbf{1}), f_{ab}^{n1})$ we have $f_{abo}^{n1} = f_{ab}^{n1}$, and part (c) for $m = n$, $l = 1$ and this o holds by construction, since we used (c) for $l = 1$ to choose the values of $\underline{\mathbf{G}}_{abo}^{n,i}|_{\partial E_{ab}^{n,i}}$ for $i \in I_{ab}^n$. Part (c) for $m = n$ and $l > 1$ follows from (c) for $m = n$, $l = 1$, since the corresponding f_{ab}^{nl} are induced by f_{ab}^{n1} . This proves (c) for all o of the form $o = ((\mathbf{1}, \mathbf{1}), f_{ab}^{n1})$. Part (c) for general $o \in O_{ab}^n$ follows by equivariance of the construction under $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \underline{\mathbf{G}}_{abo}^n)$.

It remains to prove (d),(e). We have already done this for $m = n$ and $l > 0$ above, so only the case $l = 0$ remains. In fact it is enough to prove (d) in the case $m = n$, $l = 0$ and $l' = 1$, since we can then deduce (d) with $l = 0$ and $l' > 1$ and (e) when $l = 0$ as in the inductive proof above. Thus, we have to prove that $\Pi_{ab}^{n01} : D_{ab}^{n0} \rightarrow D_{ab}^{n1}$ is surjective, and the preimage of each point is the same number of points. But as above each f_{ab}^{n0} in D_{ab}^{n0} maps $\{b\} \rightarrow O_{ab}^n$, and $f_{ab}^{n0} \mapsto f_{ab}^{n0}(b)$ induces a bijection $D_{ab}^{n0} \rightarrow O_{ab}^n$. Identifying D_{ab}^{n0} with $O_{ab}^n = \text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n) \times D_{ab}^{n1}$, we see that Π_{ab}^{n01} maps $\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n) \times D_{ab}^{n1} \rightarrow D_{ab}^{n1}$, and is given explicitly by $\Pi_{ab}^{n01} : ((\mathbf{u}, \mathbf{v}), f_{ab}^{n1}) \mapsto f_{ab}^{n1}$. Hence Π_{ab}^{n01} is surjective, and the preimage of each point is the same number of points $|\text{Aut}(X_{ab}^n, \mathbf{f}_{ab}^n, \mathbf{G}_{ab}^n)|$. This completes (d),(e), and the inductive step in the proof of Hypothesis C.7. Therefore by induction, Hypothesis C.7 holds for all $N \geq 0$.

We can now at last complete Step 4. Consider the data of Hypothesis C.7 when $m = 0$. Since each $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ is connected by Step 3, we have $n_a^0 = 1$, with $X_{a1}^0 = X_a$. Setting $m = 0$ and $b = 1$, define $O_a = O_{a1}^0$ and $\underline{\mathbf{G}}_{ao} = \underline{\mathbf{G}}_{a1o}^0$ for all $a \in A$ and $o \in O_a$. Using the notation introduced in Step 4 we have

$$\begin{aligned} \partial[X_a, \mathbf{f}_a, \mathbf{G}_a] &= \sum_{b=1}^{n_a^1} [X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1] = \sum_{b=1}^{n_a^1} \epsilon_{ab}^1 [X_{\phi^1(a,b)}^1, \mathbf{f}_{\phi^1(a,b)}^1, \mathbf{G}_{\phi^1(a,b)}^1] \\ &= \sum_{(\bar{a}, \bar{b}) \in R^1} \left\{ \sum_{\substack{b=1, \dots, n_a^1: \\ \phi^1((a,b)) = (\bar{a}, \bar{b})}} \epsilon_{ab}^1 \right\} [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1] \end{aligned} \quad (298)$$

in $\widetilde{KC}_{k-1}(Y; R)$, where in the second step we use the isomorphism $(\mathbf{a}, \mathbf{b})_{ab}^1 : (X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1) \rightarrow (X_{\phi^1(a,b)}^1, \mathbf{f}_{\phi^1(a,b)}^1, \mathbf{G}_{\phi^1(a,b)}^1)$, which multiplies orientation by ϵ_{ab}^1 . We need an analogue of (298) with $\mathbf{G}_a$ replaced by $\mathbf{G}_{ao}$ for $o \in O_a$. Now Hypothesis C.7(c) with $m = 0$, $b = 1$ and $l = 1$ implies that $(\mathbf{a}, \mathbf{b})_{ab}^1$ identifies $(X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ao}|_{X_{ab}^1})$ with $(X_{\phi^1(a,b)}^1, \mathbf{f}_{\phi^1(a,b)}^1, \mathbf{G}_{\phi^1(a,b)\bar{o}}^1)$, where $\bar{o} \in O_{\phi^1(a,b)}^1$ is given by $\bar{o} = f_{a1o}^{01}(b)$. Thus following (298), in $KC_{k-1}^{\text{ef}}(Y; R)$ we have

$$\partial[X_a, \mathbf{f}_a, \mathbf{G}_{ao}] = \sum_{\substack{(\bar{a}, \bar{b}) \in R^1, \\ \bar{o} \in O_{\bar{a}\bar{b}}^1}} \left\{ \sum_{\substack{b=1, \dots, n_a^1: \\ \phi^1((a,b))=(\bar{a}, \bar{b}), \bar{o}=f_{a1o}^{01}(b)}} \epsilon_{ab}^1 \right\} [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}\bar{o}}^1]. \quad (299)$$

Consider the composition $O_a = O_a^0 \cong D_{a1}^{00} \xrightarrow{\Pi_{a1}^{001}} D_{a1}^{01} \xrightarrow{P_{a1b}^{0110}} D_{ab}^{10} \cong O_{ab}^1$ for $a \in A$ and $b = 1, \dots, n_a^1$. This maps o to $\bar{o} = f_{a1o}^{01}(b)$. Now Hypothesis C.7(d), (e) say that Π_{a1}^{001} and P_{a1b}^{0110} are surjective and the preimages of each point is the same size. Hence for fixed $a \in A$, $b = 1, \dots, n_a^1$ and $(\bar{a}, \bar{b}) = \phi^1(a, b)$, the map $O_a \rightarrow O_{\bar{a}\bar{b}}^1$ taking $o \mapsto \bar{o} = f_{a1o}^{01}(b)$ is surjective, and the preimage of each point is $|O_a|/|O_{\bar{a}\bar{b}}^1|$ points. Therefore, if we average equation (299) over all $o \in O_a$, on the right hand side each $O_{\bar{a}\bar{b}}^1$ occurs equally often, yielding

$$\begin{aligned} \partial(\sum_{o \in O_a} |O_a|^{-1} [X_a, \mathbf{f}_a, \mathbf{G}_{ao}]) = \\ \sum_{(\bar{a}, \bar{b}) \in R^1} \left\{ \sum_{\substack{b=1, \dots, n_a^1: \\ \phi^1((a,b))=(\bar{a}, \bar{b})}} \epsilon_{ab}^1 \right\} \sum_{\bar{o} \in O_{\bar{a}\bar{b}}^1} |O_{\bar{a}\bar{b}}^1|^{-1} [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}\bar{o}}^1]. \end{aligned} \quad (300)$$

Multiplying (300) by ρ_a and summing over $a \in A$ gives

$$\begin{aligned} \partial(\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \mathbf{G}_{ao}]) = \\ \sum_{(\bar{a}, \bar{b}) \in R^1} \left\{ \sum_{\substack{a \in A, b=1, \dots, n_a^1: \\ \phi^1((a,b))=(\bar{a}, \bar{b})}} \rho_a \epsilon_{ab}^1 \right\} \sum_{\bar{o} \in O_{\bar{a}\bar{b}}^1} |O_{\bar{a}\bar{b}}^1|^{-1} [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}\bar{o}}^1]. \end{aligned} \quad (301)$$

Now in Step 4 of §C we showed that for each $(\bar{a}, \bar{b}) \in R^1$, either (A) $(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)$ admits an orientation-reversing automorphism, or (B) the coefficient $\{\dots\}$ on the second line of (301) is zero in R . We claim that in case (A), the final sum $\sum_{\bar{o} \in O_{\bar{a}\bar{b}}^1} |O_{\bar{a}\bar{b}}^1|^{-1} [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}\bar{o}}^1]$ in (301) is zero. Thus in both cases (A) and (B), the product on the second line of (301) is zero, so both sides of (301) are zero, proving equation (272).

To prove the claim, suppose $(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)$ admits an orientation-reversing automorphism. There is a natural group morphism $\gamma : \text{Aut}(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1) \rightarrow \{\pm 1\}$ with $\gamma(\mathbf{u}, \mathbf{v}) = 1$ if $\mathbf{u}$ is orientation-preserving, and $\gamma(\mathbf{u}, \mathbf{v}) = -1$ if $\mathbf{u}$ is orientation-reversing. As there exists an orientation-reversing automorphism of $(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)$, this γ is surjective.

By Hypothesis C.7(ii), $\text{Aut}(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)$ acts freely on $O_{\bar{a}\bar{b}}^1$. Choose a subset $\tilde{O}_{\bar{a}\bar{b}}^1$ in $O_{\bar{a}\bar{b}}^1$ containing one point from each orbit of $\text{Aut}(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)$.

Then there is a bijection $\text{Aut}(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1) \times \tilde{O}_{\bar{a}\bar{b}}^1 \rightarrow O_{\bar{a}\bar{b}}^1$ mapping $((\mathbf{u}, \mathbf{v}), o) \mapsto (\mathbf{u}, \mathbf{v}) \cdot o$. Thus we have

$$\begin{aligned} \sum_{\bar{o} \in \tilde{O}_{\bar{a}\bar{b}}^1} [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \underline{\mathbf{G}}_{\bar{a}\bar{b}\bar{o}}^1] &= \sum_{\bar{o} \in \tilde{O}_{\bar{a}\bar{b}}^1} \sum_{(\mathbf{u}, \mathbf{v}) \in \text{Aut}(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)} [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \underline{\mathbf{G}}_{\bar{a}\bar{b}(\mathbf{u}, \mathbf{v})\bar{o}}^1] \\ &= \sum_{\bar{o} \in \tilde{O}_{\bar{a}\bar{b}}^1} \sum_{(\mathbf{u}, \mathbf{v}) \in \text{Aut}(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)} \gamma(\mathbf{u}, \mathbf{v}) [X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \underline{\mathbf{G}}_{\bar{a}\bar{b}\bar{o}}^1] = 0, \end{aligned} \quad (302)$$

where in the second step we use the isomorphism $(\mathbf{u}, \mathbf{v}) : (X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \underline{\mathbf{G}}_{\bar{a}\bar{b}\bar{o}}^1) \rightarrow (X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \underline{\mathbf{G}}_{\bar{a}\bar{b}(\mathbf{u}, \mathbf{v})\bar{o}}^1)$, which follows from Hypothesis C.7(b) and multiplies orientations by $\gamma(\mathbf{u}, \mathbf{v})$, and in the third step we note that $\sum_{(\mathbf{u}, \mathbf{v}) \in \text{Aut}(X_{\bar{a}\bar{b}}^1, \mathbf{f}_{\bar{a}\bar{b}}^1, \mathbf{G}_{\bar{a}\bar{b}}^1)} \gamma(\mathbf{u}, \mathbf{v}) = 0$ as γ is surjective, so exactly half of $(\mathbf{u}, \mathbf{v})$ have $\gamma(\mathbf{u}, \mathbf{v}) = 1$, and half have $\gamma(\mathbf{u}, \mathbf{v}) = -1$. Equation (302) proves the claim, and (272) follows.

Thus $\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_{ao}]$ is a cycle in $KC_k^{\text{ef}}(Y; R)$, so $\beta = [\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_{ao}]]$ is well-defined in $KH_k^{\text{ef}}(Y; R)$. Finally, we must prove that $\Pi_{\text{ef}}^{\text{Kh}}(\beta) = \alpha$. By definition of $\Pi_{\text{ef}}^{\text{Kh}}$, $\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \Pi(\underline{\mathbf{G}}_{ao})]$ is a cycle in $KC_k(Y; R)$ representing $\Pi_{\text{ef}}^{\text{Kh}}(\beta)$. Hence it is sufficient to construct a homology between $\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \Pi(\underline{\mathbf{G}}_{ao})]$ and $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$.

For each $a \in A$, consider $[0, 1] \times X_a$ as a compact oriented Kuranishi space with the product orientation, and define $\mathbf{g}_a = \mathbf{f}_a \circ \pi_{X_a} : [0, 1] \times X_a \rightarrow Y$, a strongly smooth map. Set $J_a = \{i + 1 : i \in I_a\}$. For $i \in I_a$, define a Kuranishi neighbourhood on $[0, 1] \times X_a$ by $(W_a^{i+1}, F_a^{i+1}, t_a^{i+1}, \chi_a^{i+1}) = ([0, 1] \times V_a^i, [0, 1] \times E_a^i, s_a^i \circ \pi_{V_a^i}, \text{id}_{[0, 1]} \times \psi^i)$, define $g_a^{i+1} : W_a^{i+1} \rightarrow Y$ by $g_a^{i+1} = f_a^i \circ \pi_{V_a^i}$, and $\zeta_{i+1, a} : [0, 1] \times X_a \rightarrow [0, 1]$ by $\zeta_{i+1, a} = \eta_{i, a} \circ \pi_{X_a}$. For $i, j \in I_a$, define $W_a^{(i+1)(j+1)} = [0, 1] \times V_a^{ij}$, $\psi_a^{(i+1)(j+1)} = \text{id}_{[0, 1]} \times \phi_a^{ij}$, $\hat{\psi}_a^{(i+1)(j+1)} = \text{id}_{[0, 1]} \times \hat{\phi}_a^{ij}$ and $\zeta_{i+1, a}^{j+1} : W_a^{j+1} \rightarrow [0, 1]$ by $\zeta_{i+1, a}^{j+1} = \eta_{i, a}^j \circ \pi_{V_a^j}$. Write $\mathbf{J}_a = (J_a, (W_a^j, \dots, \chi_a^j), g_a^j : j \in J, \dots)$, and $\boldsymbol{\zeta}_a = (\zeta_{j, a} : j \in J, \zeta_{j, a}^k : j, k \in J)$. It is easy to check that $(\mathbf{J}_a, \boldsymbol{\zeta}_a)$ is an *excellent coordinate system* for $([0, 1] \times X_a, \mathbf{g}_a)$.

For $o \in O_a$ and $i \in I_a$, define $H_{ao}^{i+1} : F_a^{i+1} \rightarrow P$ by, for $u \in [0, 1]$ and $e \in E_a^i$,

$$H_{ao}^{i+1}(u, e) = \begin{cases} G_a^i(e), & u = 0, \\ \mu(u, G_a^i(e)), & u \in (0, 1), \\ \Pi \circ \underline{G}_{ao}^i(e), & u = 1. \end{cases} \quad (303)$$

Write $\mathbf{H}_{ao} = (\mathbf{J}_a, \boldsymbol{\zeta}_a, H_{ao}^j : j \in J_a)$. As in the proofs of Theorem 4.15 and Proposition 4.32, one can show $\mathbf{H}_{ao}$ is *gauge-fixing data* for $([0, 1] \times X_a, \mathbf{g}_a)$.

We have $\partial([0, 1] \times X_a) = (\{1\} \times X_a) \Pi - (\{0\} \times X_a) \Pi - ([0, 1] \times \partial X_a)$ in oriented Kuranishi spaces. The construction of $(\mathbf{J}_a, \boldsymbol{\zeta}_a)$ and the cases $u = 0$, $u = 1$ in (303) ensure that $\mathbf{H}_{ao}|_{\{0\} \times X_a} = \mathbf{G}_a$ and $\mathbf{H}_{ao}|_{\{1\} \times X_a} = \Pi(\underline{\mathbf{G}}_{ao})$, identifying $\{0\} \times X_a \cong X_a \cong \{1\} \times X_a$. Thus in $KC_k(Y; R)$ we have

$$\begin{aligned} \partial([0, 1] \times X_a, \mathbf{g}_a, \mathbf{H}_{ao}) &= [X_a, \mathbf{f}_a, \Pi(\underline{\mathbf{G}}_{ao})] - [X_a, \mathbf{f}_a, \mathbf{G}_a] \\ &\quad - [[0, 1] \times \partial X_a, \mathbf{g}_a|_{[0, 1] \times \partial X_a}, \mathbf{H}_{ao}|_{[0, 1] \times \partial X_a}]. \end{aligned} \quad (304)$$

Following the proof of (272) in equations (299)–(302) above, but replacing $X_a, \mathbf{f}_a, \underline{\mathbf{G}}_{ao}$ by $[0, 1] \times X_a, \mathbf{g}_a, \mathbf{H}_{ao}$, and X_{ab}^1 by $[0, 1] \times X_{ab}^1$, and so on, yields

$$\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [[0, 1] \times \partial X_a, \mathbf{g}_a|_{[0, 1] \times \partial X_a}, \mathbf{H}_{ao}|_{[0, 1] \times \partial X_a}] = 0. \quad (305)$$

Thus, multiplying (304) by $\rho_a |O_a|^{-1}$, summing over all $a \in A$ and $o \in O_a$, and using (305) to cancel out the contribution of the final term in (304) gives

$$\begin{aligned} & \partial \left(\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [[0, 1] \times X_a, \mathbf{g}_a, \mathbf{H}_{ao}] \right) = \\ & \sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \Pi(\underline{\mathbf{G}}_{ao})] - \sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]. \end{aligned}$$

Therefore $\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \Pi(\underline{\mathbf{G}}_{ao})]$ is homologous to $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$, and $\Pi_{\text{ef}}^{\text{Kh}}(\beta) = \alpha$. Hence for each $\alpha \in KH_k(Y; R)$ we can construct $\beta \in KH_k^{\text{ef}}(Y; R)$ with $\Pi_{\text{ef}}^{\text{Kh}}(\beta) = \alpha$, and $\Pi_{\text{ef}}^{\text{Kh}} : KH_*^{\text{ef}}(Y; R) \rightarrow KH_*(Y; R)$ is *surjective*. This completes Step 4.

C.5 Step 5: $\Pi_{\text{ef}}^{\text{Kh}} : KH_*^{\text{ef}}(Y; R) \rightarrow KH_*(Y; R)$ is injective

As in Step 5, suppose $\beta \in KH_k^{\text{ef}}(Y; R)$ with $\Pi_{\text{ef}}^{\text{Kh}}(\beta) = 0$, represent β by the image of a cycle $\sum_{s \in S} \zeta_s \tau_s$ in singular homology, and choose $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$ in $KC_{k+1}(Y; R)$ satisfying (273). We now explain how to apply Steps 1–4 to this $\sum_{a \in A} \rho_a [X_a, \mathbf{f}_a, \mathbf{G}_a]$, replacing k -cycles by $(k+1)$ -chains, to construct $\sum_{a \in A} \sum_{o \in O_a} \rho_a |O_a|^{-1} [X_a, \mathbf{f}_a, \underline{\mathbf{G}}_{ao}]$ in $KC_{k+1}^{\text{ef}}(Y; R)$ satisfying (275). This will imply that $\beta = 0$, so $\Pi_{\text{ef}}^{\text{Kh}} : KH_k^{\text{ef}}(Y; R) \rightarrow KH_k(Y; R)$ is *injective*.

First we discuss Step 1. Introducing connected $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ and Γ_d, η_d for $d \in D$ as in Step 1, generalizing (255), the lift of (273) to $KC_k(Y; R)$ is

$$\begin{aligned} \sum_{a \in A} \rho_a [\partial X_a, \mathbf{f}_a|_{\partial X_a}, \mathbf{G}_a|_{\partial X_a}] &= \sum_{s \in S} \zeta_s [\Delta_k, \tau_s, \mathbf{G}_{\Delta_k}] + \\ \sum_{d \in D} \eta_d ([\tilde{X}_d / \Gamma_d, \pi_*(\tilde{\mathbf{f}}_d), \pi_*(\tilde{\mathbf{G}}_d)] - \frac{1}{|\Gamma_d|} [\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d]). \end{aligned} \quad (306)$$

Write $\partial(X_a, \mathbf{f}_a, \mathbf{G}_a) = \coprod_{b=1}^{n_a^1} (X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1)$ as usual. Then (306) implies that some of the components $(X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1)$ or $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ must be isomorphic to $(\Delta_k, \tau_s, \mathbf{G}_{\Delta_k})$ for $s \in S$.

In Step 1 we choose tent functions $\mathbf{T}_a$ for $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ for $a \in A$ and $\tilde{\mathbf{T}}_d$ for $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ for $d \in D$ to cut $X_a, \tilde{X}_d$ into ‘arbitrarily small pieces’ $X_{ac}, \tilde{X}_{df}$ for $c \in C_a, f \in F_d$. The new issue we must deal with here is that when $(X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1)$ or $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ is isomorphic to $(\Delta_k, \tau_s, \mathbf{G}_{\Delta_k})$, the tent function $\mathbf{T}_a$ or $\tilde{\mathbf{T}}_d$ also cuts $(\Delta_k, \tau_s, \mathbf{G}_{\Delta_k})$ into pieces, so $\sum_{s \in S} \zeta_s [\Delta_k, \tau_s, \mathbf{G}_{\Delta_k}]$ is modified. We must keep track of these modifications, and in particular ensure that the modified version of $\sum_{s \in S} \zeta_s [\Delta_k, \tau_s, \mathbf{G}_{\Delta_k}]$ is still the image of a cycle in $KC_k^{\text{ef}}(Y; R)$ representing β .

We do this using an idea from Step 4 of §B in §B.4. We fix some $N \gg 0$, and choose the tent functions $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ such that if $(X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1)$ or $(\tilde{X}_d, \tilde{\mathbf{f}}_d, \tilde{\mathbf{G}}_d)$ is isomorphic to $(\Delta_k, \tau_s, \mathbf{G}_{\Delta_k})$, then $\mathbf{T}_a|_{X_{ab}^1}$ or $\tilde{\mathbf{T}}_d$ cuts Δ_k into the N^{th} barycentric subdivision of Δ_k . Since we construct the $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ in §B.1 by first choosing

tent functions $\mathbf{T}_{st}^m$ for $(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m)$ by induction on decreasing $m = M, M-1, \dots, 0$, we do this by inserting an extra condition: if for $0 < m \leq M$ we have $(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m) \cong (\Delta_{k+1-m}, \tau, \mathbf{G}_{\Delta_{k+1-m}})$ for some smooth $\tau : \Delta_{k+1-m} \rightarrow Y$, then we choose $\mathbf{T}_{st}^m$ to be a standard tent function on Δ_{k+1-m} which cuts Δ_{k+1-m} into its N^{th} barycentric subdivision.

We must check this extra condition is compatible with conditions (i)–(iv) near the beginning of §C.1. Part (i) is immediate since our condition is expressed using isomorphism classes. For (ii), as Δ_{k+1-m} is a simply-connected manifold we cannot write $(\Delta_{k+1-m}, \tau, \mathbf{G}_{\Delta_{k+1-m}})$ as a quotient $(X_{de}^m / \text{Stab}_{\Gamma_d}(e), \pi_*^m(\mathbf{f}_{de}^m))$ for $(X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m)$ connected and $\text{Stab}_{\Gamma_d}(e) \neq \{1\}$, and if $(X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m) \cong (\Delta_{k+1-m}, \tau, \mathbf{G}_{\Delta_{k+1-m}})$ then $\text{Stab}_{\Gamma_d}(e)$ must act trivially on $(X_{de}^m, \mathbf{f}_{de}^m, \mathbf{G}_{de}^m)$ as $\text{Aut}(\Delta_{k+1-m}, \tau, \mathbf{G}_{\Delta_{k+1-m}}) = \{1\}$. Thus $X_{de}^m \cong \Delta_{k+1-m} \times (\{0\} / \text{Stab}_{\Gamma_d}(e))$, so we deal with (ii) by extending the extra condition to say that if $(X_{st}^m, \mathbf{f}_{st}^m, \mathbf{G}_{st}^m) \cong (\Delta_{k+1-m} \times (\{0\} / \Gamma), \tau, \mathbf{G}_{\Delta_{k+1-m}})$ for Γ a finite group, then we again choose $\mathbf{T}_{st}^m$ to be a standard tent function on $\Delta_{k+1-m} \times (\{0\} / \Gamma)$ which cuts Δ_{k+1-m} into its N^{th} barycentric subdivision.

Part (iii) holds automatically since ∂X_{st}^m is the disjoint union of $k+2-m$ copies of Δ_{k-m} , and on each of these we have chosen $\mathbf{T}_{st'}^{m+1}$ to be a standard tent function inducing the N^{th} barycentric subdivision of Δ_{k-m} in the previous inductive step. Part (iv) is satisfied provided $N \gg 0$ is large enough, since then the N^{th} barycentric subdivision cuts $X_{st}^m \cong \Delta_{k+1-m}$ into arbitrarily small pieces. Thus we can carry out Step 1 in such a way that the $\mathbf{T}_a, \tilde{\mathbf{T}}_d$ cut each $(\Delta_k, \tau_s, \mathbf{G}_{\Delta_k})$ into the N^{th} barycentric subdivision of Δ_k for some fixed $N \gg 0$.

We also make one additional change to the definition of the gauge-fixing data $\mathbf{G}_{ac}$ for $(X_{ac}, \mathbf{f}_{ac})$ for $c \in C_a$. Here $(X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac})$ was defined in Proposition A.29 using the triple $(Z_{X_a, \mathbf{T}_a}, \mathbf{f}_a \circ \pi, \mathbf{H}_{X_a, \mathbf{T}_a})$ constructed from $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ and $\mathbf{T}_a$ in Definition A.28. The issue is that when we use a standard tent function on Δ_k to cut $(\Delta_k, \tau_s, \mathbf{G}_{\Delta_k})$ into its N^{th} barycentric subdivision, we obtain $(k!)^N$ small pieces $(\Delta_{kl}, \tau_s|_{\Delta_{kl}}, \mathbf{G}_{\Delta_{kl}})$ for l in an indexing set $L_{k,N}$, where each Δ_{kl} is a small k -simplex with a natural affine diffeomorphism $\Delta_{kl} \cong \Delta_k$. However, because of the way the function H^{i+1} is defined in Definition A.28, the diffeomorphism $\Delta_{kl} \cong \Delta_k$ does not identify $\mathbf{G}_{\Delta_{kl}}$ with $\mathbf{G}_{\Delta_k}$, as the functions $G_{\Delta_{kl}}^k : \Delta_{kl} \rightarrow P$ and $G_{\Delta_k}^k : \Delta_k \rightarrow P$ in $\mathbf{G}_{\Delta_{kl}}, \mathbf{G}_{\Delta_k}$ are not identified.

We deal this by modifying the definition of the functions $H_a^{k+2} : W_a^{k+2} \rightarrow P$ in $\mathbf{H}_{X_a, \mathbf{T}_a}$, by hand, which then modifies the functions $G_{ac}^{k+1} : V_{ac}^{k+1} \rightarrow P$ in $\mathbf{G}_{ac}$, to identify each $\mathbf{G}_{\Delta_{kl}}$ with $\mathbf{G}_{\Delta_k}$, and similarly for $\tilde{X}_d, \tilde{\mathbf{T}}_d$. That is, whenever $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ is a component of $\partial^m(X_a, \mathbf{f}_a, \mathbf{G}_a)$ isomorphic to some $(\Delta_{k+1-m}, \tau, \mathbf{G}_{\Delta_{k+1-m}})$, so that $\mathbf{T}_{ab}^m$ is a standard tent function on $X_{ab}^m \cong \Delta_{k+1-m}$ inducing the N^{th} subdivision, then on the image in $Z_{X_a, \mathbf{T}_a}$ (and hence each X_{ac}) of each piece $X_{abl}^m \cong \Delta_{(k+1-m)l}$ for $l \in L_{k+1-m, N}$, we define the function H_a^{k+2} to be identified with $G_{\Delta_{k+1-m}}^{k+1-m} : \Delta_{k+1-m} \rightarrow P$ under the identification $X_{abl}^m \cong \Delta_{(k+1-m)l} \cong \Delta_{k+1-m}$.

This prescription is compatible on overlaps between the images of different simplices $X_{abl}^m, X_{ab'l'}^m$, as two such simplices must intersect in a face of each, and the functions $G_{\Delta_k}^k$ restrict to $G_{\Delta_l}^l$ on any face $\Delta \cong \Delta_l$ of Δ_k . With these

changes, following the proof in Step 1 that $\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]$ is homologous to $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]$ proves equation (274), as we want. This completes our discussion of the changes to Step 1.

Step 2 requires essentially no changes. For Step 3, the analogue of (267) is

$$\begin{aligned} & \partial(\sum_{a \in A} \sum_{c \in C_a} \rho_a[X_{ac}, \mathbf{f}_{ac}, \mathbf{G}_{ac}]) - \Pi_{\text{si}}^{\text{Kh}} \circ \Upsilon^N(\sum_{s \in S} \zeta_s \tau_s) = \\ & \sum_{d \in D} \eta_d \left(\sum_{f \in F_d / \Gamma_d} [\tilde{X}_{df} / \text{Stab}_{\Gamma_d}(f), \pi_*(\tilde{\mathbf{f}}_{df}), \pi_*(\tilde{\mathbf{G}}_{df})] \right. \\ & \quad \left. - |\Gamma_d|^{-1} \sum_{f \in F_d} [\tilde{X}_{df}, \tilde{\mathbf{f}}_{df}, \tilde{\mathbf{G}}_{df}] \right). \end{aligned} \quad (307)$$

which is a lift of (274) to $\widetilde{KC}_k(Y; R)$. As for (267), all terms on the l.h.s. of (307) have *trivial stabilizers*, since the X_a which we started with at the beginning of Step 3 (after changing notation) do, so the X_{ac} do, and the Kuranishi spaces in $\Pi_{\text{si}}^{\text{Kh}} \circ \Upsilon^N(\sum_{s \in S} \zeta_s \tau_s)$ are k -simplices Δ_k , and so are manifolds.

In Step 3, we proved that both sides of (267) are zero in $\widetilde{KC}_{k-1}(Y; R)$ provided the $X_{ac}, \tilde{X}_{df}$ are taken sufficiently small, because all terms on the l.h.s. of (267) have trivial stabilizers. Since this also holds for the l.h.s. of (307), the same argument shows that both sides of (307) are zero in $\widetilde{KC}_k(Y; R)$ for $X_{ac}, \tilde{X}_{df}$ sufficiently small. Therefore (274) holds in $\widetilde{KC}_k(Y; R)$, as well as $KC_k(Y; R)$. This completes the changes to Step 3.

At the beginning of Step 4 we have a chain $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]$ satisfying

$$\partial(\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]) = \Pi_{\text{si}}^{\text{Kh}} \circ \Upsilon^N(\sum_{s \in S} \zeta_s \tau_s) \quad (308)$$

in $\widetilde{KC}_k(Y; R)$ for $N \gg 0$, where $\sum_{s \in S} \zeta_s \tau_s$ is the same as it has been all the way through, but $\sum_{a \in A} \rho_a[X_a, \mathbf{f}_a, \mathbf{G}_a]$ has been changed several times, and now the X_a have trivial stabilizers, the V_a^i are manifolds, the $(X_a, \mathbf{f}_a, \mathbf{G}_a)$ are connected, and each component of $\partial^m X_a$ occurs as the intersection of m distinct components of ∂X_a . Our goal is to choose finite sets O_a and maps $\underline{\mathbf{G}}_a^i : V_a^i \rightarrow \underline{P}$ for $a \in A$ and $i \in I_a$ such that (308) lifts to equation (275) in $KH_k^{\text{ef}}(Y; R)$.

The issue is this: we have to choose these maps $\underline{\mathbf{G}}_a^i$ such that whenever some component $(X_{ab}^1, \mathbf{f}_{ab}^1, \mathbf{G}_{ab}^1)$ of $\partial(X_a, \mathbf{f}_a, \mathbf{G}_a)$ is isomorphic to $\Pi_{\text{si}}^{\text{Kh}}(v)$ for some $v : \Delta_k \rightarrow Y$ occurring in the N^{th} barycentric subdivision of some $\tau_s : \Delta_k \rightarrow Y$, then the corresponding component of $\partial(X_a, \mathbf{f}_a, \underline{\mathbf{G}}_{ao})$ must be isomorphic to $\Pi_{\text{si}}^{\text{ef}}(v)$ for all $o \in O_a$. That is, $E_{ab}^{1,k}$ in $\mathbf{G}_{ab}^1$ is a component of ∂E_a^{k+1} in $\mathbf{G}_a$, and the maps $\underline{\mathbf{G}}_{ao}^{k+1} | E_{ab}^{1,k} : E_{ab}^{1,k} \rightarrow \underline{P}$ and $\underline{\mathbf{G}}_{\Delta_k}^k : \Delta_k \rightarrow \underline{P}$ should agree under the given diffeomorphism $E_{ab}^{1,k} \cong \Delta_k$. If we do this, then all terms on the l.h.s. of (308) isomorphic to terms on the r.h.s. of (308) lift to terms on the l.h.s. of (275) isomorphic to terms on the r.h.s. of (275), and so (275) holds.

We ensure this by adding an extra condition in the inductive proof of Hypothesis C.7 in §C.4, when for part (iv) we chose maps $\underline{\mathbf{G}}_{abo}^{m,i} : E_{ab}^{m,i} \rightarrow \underline{P}$ for $i \in I_{ab}^m$ such that $\underline{\mathbf{G}}_{abo}^m = (\mathbf{I}_{ab}^m, \boldsymbol{\eta}_{ab}^m, \underline{\mathbf{G}}_{abo}^{m,i} : i \in I_{ab}^m)$ is effective gauge-fixing data for $(X_{ab}^m, \mathbf{f}_{ab}^m)$. We require that if any $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ is isomorphic to $(\Delta_{k+1-m}, \tau, \mathbf{G}_{\Delta_{k+1-m}})$ for some smooth $\tau : \Delta_{k+1-m} \rightarrow Y$, then the corresponding maps $\underline{\mathbf{G}}_{abo}^{m,k+1-m} : E_{ab}^{m,k+1-m} \rightarrow \underline{P}$ for $o \in O_{ab}^m$ are identified with $\underline{\mathbf{G}}_{\Delta_{k+1-m}}^{k+1-m} :$

$\Delta_{k+1-m} \rightarrow \underline{P}$ under the given diffeomorphism $E_{ab}^{m,k+1-m} \cong \Delta_{k+1-m}$. Then $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{abo}^m)$ is isomorphic to $(\Delta_{k+1-m}, \tau, \mathbf{G}_{\Delta_{k+1-m}})$.

We must check that prescribing these particular choices are compatible with conditions (a)–(e), so that the inductive proof that we can choose data satisfying Hypothesis C.7 is still valid with this extra condition. For part (a), we claim that if $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ is isomorphic to some $(\Delta_{k+1-m}, \tau, \mathbf{G}_{\Delta_{k+1-m}})$ then O_{ab}^m is a single point. We prove this by reverse induction on $m = k+1, k, \dots, 0$. For the inductive step, note that in §C.4 we defined $O_{ab}^m = \text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) \times D_{ab}^{m1}$. As $X_{ab}^m \cong \Delta_{k+1-m}$, it has $k+2-m$ boundary faces $X_{ab'}^{m+1}$, each of which is a copy of Δ_{k-m} , and D_{ab}^{m1} consists of a choice of point in $O_{\phi^{m+1}(a,b')}^{m+1}$ satisfying some consistency conditions.

By induction, each such $O_{\phi^{m+1}(a,b')}^{m+1}$ is a point, as $(X_{ab'}^{m+1}, \mathbf{f}_{ab'}^{m+1}, \mathbf{G}_{ab'}^{m+1}) \cong (\Delta_{k-m}, v, \mathbf{G}_{\Delta_{k-m}})$, where $v = \tau \circ F_j^{k+1-m}$ for some $j = 0, \dots, k+1-m$, in the notation of §4.1. Therefore D_{ab}^{m1} is one point, as there is only one possible choice in $O_{\phi^{m+1}(a,b')}^{m+1}$ for each b' . Also $\text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) = \{1\}$ as $\text{Aut}(\Delta_{k+1-m}, \tau, \mathbf{G}_{\Delta_{k+1-m}}) = \{1\}$. Hence $O_{ab}^m = \text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) \times D_{ab}^{m1}$ is one point, proving the inductive step.

Since O_{ab}^m is a single point, (a) is trivial as there do not exist distinct o, o' in O_{ab}^m . Part (b) is also trivial, as $\text{Aut}(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) = \{1\}$. Part (c) holds since as above each boundary component $(X_{ab'}^{m+1}, \mathbf{f}_{ab'}^{m+1}, \mathbf{G}_{ab'}^{m+1})$ of $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m)$ is isomorphic to $(\Delta_{k-m}, v, \mathbf{G}_{\Delta_{k-m}})$ for some v , so in the previous inductive step we chose $\mathbf{G}_{ab'o'}^{m+1} \cong \mathbf{G}_{\Delta_{k-m}}$ for $o' \in O_{ab'}^{m+1}$, and in §4.3 we defined $\mathbf{G}_{\Delta_{k+1-m}}$ so that it restricted to $\mathbf{G}_{\Delta_{k-m}}$ on each Δ_{k-m} face of Δ_{k+1-m} . And (d),(e) are not affected as they do not concern choice of the $\mathbf{G}_{abo}^{m,i}$.

Therefore the inductive proof goes through, and we can choose data satisfying Hypothesis C.7 with this extra condition when $(X_{ab}^m, \mathbf{f}_{ab}^m, \mathbf{G}_{ab}^m) \cong (\Delta_{k+1-m}, \tau, \mathbf{G}_{\Delta_{k+1-m}})$. The proof of (272) using equations (298)–(302) then generalizes to give (275), and Step 5 follows. This completes the proof of Theorem 4.9.

D Identification of products $\cup, \cap, \bullet$

We now prove Theorem 4.34, that is, we shall show that $\Pi_{cs}^{\text{Kch}}, \Pi_{cs}^{\text{ec}}, \Pi_{si}^{\text{Kh}}, \Pi_{si}^{\text{ef}}$ identify the cup, cap and intersection products $\cup, \cap, \bullet$ on $H_{cs}^*, H_{*}^{\text{si}}(Y; R)$ with those on $KH^*, KH_*(Y; R)$ and $KH_{ec}^*, KH_{*}^{\text{ef}}(Y; R)$. Proving this is tricky since we do not have a good description of $\Pi_{cs}^{\text{Kch}}, \Pi_{cs}^{\text{ec}}$ at the cochain level. Our approach is to first suppose that Y is oriented, and prove that $\Pi_{si}^{\text{Kh}} : H_{*}^{\text{si}}(Y; R) \rightarrow KH_*(Y; R)$, $\Pi_{si}^{\text{ef}} : H_{*}^{\text{si}}(Y; R) \rightarrow KH_{*}^{\text{ef}}(Y; R)$ identify the intersection products $\bullet$ on $H_{*}^{\text{si}}(Y; R)$ and $KH_*, KH_{*}^{\text{ef}}(Y; R)$, using chain level descriptions of $\bullet$ on $C_{*}^{\text{si}}(Y; R)$ and $KC_*, KC_{*}^{\text{ef}}(Y; R)$. Then since Poincaré duality identifies $\bullet$ with $\cup, \cap$, we deduce that $\Pi_{cs}^{\text{Kch}}, \Pi_{cs}^{\text{ec}}, \Pi_{si}^{\text{Kh}}, \Pi_{si}^{\text{ef}}$ also identify cup and cap products.

Chain-level definitions of $\bullet$ on $KC_*, KC_{*}^{\text{ef}}(Y; R)$ are given in (111) and (114). Describing $\bullet$ on singular chains $C_{*}^{\text{si}}(Y; R)$ is more complex. Two good references are Bredon [10, VI.11], and Lefschetz [46, §IV], which is the original source for much of the theory of the intersection product on singular homology of

manifolds. As these do not discuss orbifolds, we assume Y is a *manifold*.

Let Y be an oriented n -manifold without boundary and $\alpha \in H_k^{\text{si}}(Y; R)$, $\beta \in H_l^{\text{si}}(Y; R)$, so that $\alpha \bullet \beta \in H_{k+l-n}^{\text{si}}(Y; R)$. For Theorem 4.34(c) we must prove that $\Pi_{\text{si}}^*(\alpha \bullet \beta) = \Pi_{\text{si}}^*(\alpha) \bullet \Pi_{\text{si}}^*(\beta)$ in $KH_{k+l-n}(Y; R)$ or $KH_{k+l-n}^{\text{ef}}(Y; R)$. First consider the case in which $\alpha = [K]$ and $\beta = [L]$, where K, L are compact, oriented, immersed submanifolds of Y without boundary intersecting transversely in Y , with immersions $\iota_K : K \rightarrow Y$ and $\iota_L : L \rightarrow Y$.

Choose effective gauge-fixing data $\underline{\mathbf{G}}_K, \underline{\mathbf{G}}_L$ for $(K, \iota_K), (L, \iota_L)$, which is possible by Theorem 3.39(a), and set $\mathbf{G}_K = \Pi(\underline{\mathbf{G}}_K)$, $\mathbf{G}_L = \Pi(\underline{\mathbf{G}}_L)$. Then $[K, \iota_K, \mathbf{G}_K] \in KC_k(Y; R)$ with $\partial[K, \iota_K, \mathbf{G}_K] = 0$ as $\partial K = \emptyset$, so $[K, \iota_K, \mathbf{G}_K] \in KH_k(Y; R)$. Similarly $[K, \iota_K, \underline{\mathbf{G}}_K] \in KH_k^{\text{ef}}(Y; R)$, $[L, \iota_L, \mathbf{G}_L] \in KH_l(Y; R)$, and $[L, \iota_L, \underline{\mathbf{G}}_L] \in KH_l^{\text{ef}}(Y; R)$. Considering Step 3 in Appendix B, when we construct an inverse $(\Pi_{\text{si}}^{\text{ef}})^{-1}$ for $\Pi_{\text{si}}^{\text{ef}} : H_k^{\text{si}}(Y; R) \rightarrow KH_k^{\text{ef}}(Y; R)$, we see that $(\Pi_{\text{si}}^{\text{ef}})^{-1}([K, \iota_K, \underline{\mathbf{G}}_K])$ is represented, as a singular chain, by a triangulation of K by k -simplices. But this is how we define the class $[K]$ of K in $H_k^{\text{si}}(Y; R)$. Since $\alpha = [K]$, this gives $[K, \iota_K, \underline{\mathbf{G}}_K] = \Pi_{\text{si}}^{\text{ef}}(\alpha)$. As $[K, \iota_K, \mathbf{G}_K] = \Pi_{\text{ef}}^{\text{Kh}}([K, \iota_K, \underline{\mathbf{G}}_K])$ and $\Pi_{\text{si}}^{\text{Kh}} = \Pi_{\text{ef}}^{\text{Kh}} \circ \Pi_{\text{si}}^{\text{ef}}$, we have $[K, \iota_K, \mathbf{G}_K] = \Pi_{\text{si}}^{\text{Kh}}(\alpha)$. Similarly $[L, \iota_L, \underline{\mathbf{G}}_L] = \Pi_{\text{si}}^{\text{ef}}(\beta)$ and $[L, \iota_L, \mathbf{G}_L] = \Pi_{\text{si}}^{\text{Kh}}(\beta)$.

Therefore $\Pi_{\text{si}}^{\text{Kh}}(\alpha), \Pi_{\text{si}}^{\text{Kh}}(\beta)$ are represented in $KC_*(Y; R)$ by $[K, \iota_K, \mathbf{G}_K], [L, \iota_L, \mathbf{G}_L]$, so by (111), $\Pi_{\text{si}}^{\text{Kh}}(\alpha) \bullet \Pi_{\text{si}}^{\text{Kh}}(\beta)$ is represented in $KC_{k+l-n}(Y; R)$ by

$$\begin{aligned} & \Pi_{\text{Kch}}^{\text{Kh}}(\Pi_{\text{Kh}}^{\text{Kh}}([K, \iota_K, \mathbf{G}_K]) \cup \Pi_{\text{Kh}}^{\text{Kh}}[L, \iota_L, \mathbf{G}_L]) \\ &= \Pi_{\text{Kch}}^{\text{Kh}}([K^Y, \iota_K^Y, \mathbf{C}_{\mathbf{G}_K}^Y] \cup [L^Y, \iota_L^Y, \mathbf{C}_{\mathbf{G}_L}^Y]) \\ &= \Pi_{\text{Kch}}^{\text{Kh}}([K^Y \times_Y L^Y, \pi_Y, \mathbf{C}_{\mathbf{G}_K}^Y \times_Y \mathbf{C}_{\mathbf{G}_L}^Y]) \\ &= [K^Y \times_Y L^Y, \pi_Y, \mathbf{G}_{\mathbf{C}_{\mathbf{G}_K}^Y \times_Y \mathbf{C}_{\mathbf{G}_L}^Y}], \end{aligned} \tag{309}$$

using the definitions of $\Pi_{\text{Kh}}^{\text{Kh}}, \Pi_{\text{Kch}}^{\text{Kh}}, \cup$ in Definitions 4.14 and 4.29.

Since K, L intersect transversely in Y , their intersection $K \cap L$ is a compact immersed $(k+l-n)$ -submanifold of Y , which can also be considered as a fibre product $K \times_{\iota_K, Y, \iota_L} L$. Following Bredon [10, p. 375] we define an *orientation* on $K \cap L$, using the orientations on K, L, Y . Then [10, Th. VI.11.9] shows that $[K] \bullet [L] = [K \cap L]$ in $H_{k+l-n}^{\text{si}}(Y; R)$. Thus, by the argument above, $\Pi_{\text{si}}^{\text{Kh}}(\alpha \bullet \beta)$ is represented in $KC_{k+l-n}(Y; R)$ by $[K \cap L, \iota_{K \cap L}, \mathbf{G}_{K \cap L}]$, for some choice of gauge-fixing data $\mathbf{G}_{K \cap L}$ for $(K \cap L, \iota_{K \cap L})$.

Hence, to show that $\Pi_{\text{si}}^{\text{Kh}}(\alpha \bullet \beta) = \Pi_{\text{si}}^{\text{Kh}}(\alpha) \bullet \Pi_{\text{si}}^{\text{Kh}}(\beta)$, by (309) it is enough to prove that $[K \cap L, \iota_{K \cap L}, \mathbf{G}_{K \cap L}]$ is homologous to $[K^Y \times_Y L^Y, \pi_Y, \mathbf{G}_{\mathbf{C}_{\mathbf{G}_K}^Y \times_Y \mathbf{C}_{\mathbf{G}_L}^Y}]$ in $KC_{k+l-n}(Y; R)$. Now the topological space underlying $K^Y \times_Y L^Y$ is just $K \cap L$, so $K^Y \times_Y L^Y$ is just $K \cap L$ with a different Kuranishi structure. Examining the definitions shows that $K^Y \times_Y L^Y$ is isomorphic as a Kuranishi space to $((K \cap L)^Y)^Y$, in the notation of Definition 4.14, and that π_Y agrees with $(\iota_{K \cap L}^Y)^Y$. Comparing our orientation conventions in §2.7 with Bredon's shows that the

orientation conventions of $K^Y \times_Y L^Y$ also agree. Now

$$\begin{aligned} & \Pi_{K \cap L}^{\text{Kh}} \circ \Pi_{K \cap L}^{\text{Kh}} \circ \Pi_{K \cap L}^{\text{Kh}} \circ \Pi_{K \cap L}^{\text{Kh}} ([K \cap L, \iota_{K \cap L}, \mathbf{G}_{K \cap L}]) \\ &= [((K \cap L)^Y)^Y, (\iota_{K \cap L}^Y)^Y, \mathbf{G}_{\mathcal{C}_{\mathbf{G}_{K \cap L}}^Y}]. \end{aligned} \quad (310)$$

The proof of Theorem 4.15 shows that (310) is homologous to $[K \cap L, \iota_{K \cap L}, \mathbf{G}_{K \cap L}]$. But the right hand side is $[K^Y \times_Y L^Y, \pi_Y, \mathbf{G}_{\mathcal{C}_{\mathbf{G}_K}^Y \times_Y \mathcal{C}_{\mathbf{G}_L}^Y}]$ with an alternative choice of gauge-fixing data. As $\partial(K^Y \times_Y L^Y) = \emptyset$, the homology class does not depend on the gauge-fixing data. Therefore $[K \cap L, \iota_{K \cap L}, \mathbf{G}_{K \cap L}]$ is homologous to $[K^Y \times_Y L^Y, \pi_Y, \mathbf{G}_{\mathcal{C}_{\mathbf{G}_K}^Y \times_Y \mathcal{C}_{\mathbf{G}_L}^Y}]$, so $\Pi_{\text{si}}^{\text{Kh}}(\alpha \bullet \beta) = \Pi_{\text{si}}^{\text{Kh}}(\alpha) \bullet \Pi_{\text{si}}^{\text{Kh}}(\beta)$. Similarly $\Pi_{\text{si}}^{\text{ef}}(\alpha \bullet \beta) = \Pi_{\text{si}}^{\text{ef}}(\alpha) \bullet \Pi_{\text{si}}^{\text{ef}}(\beta)$.

This proves Theorem 4.34(c) in the case when α, β can be represented by compact oriented submanifolds K, L intersecting transversely. For the general case we can use ideas in Lefschetz [46], particularly [46, §IV], which is the original source for much of the theory of the intersection product on singular homology of manifolds. Lefschetz shows that we can choose a triangulation $\mathcal{T}$ of Y by convex polyhedra, with a dual triangulation $\mathcal{T}^*$, such that if P, P^* are polyhedra in $\mathcal{T}, \mathcal{T}^*$ of dimensions k, l then P, P^* intersect transversely in a convex polyhedron of dimension $k + l - n$, or $P \cap P^* = \emptyset$.

We may represent α by an R -linear combination of k -dimensional polyhedra P in $\mathcal{T}$; to make a singular chain, we replace each P by its barycentric subdivision into k -simplices. Similarly, we represent β by an R -linear combination of l -dimensional polyhedra P^* in $\mathcal{T}^*$. Then $\alpha \bullet \beta$ is represented by the corresponding R -bilinear sum of transverse intersections $P \cap P^*$. We then use the argument above with K, L replaced by R -linear combinations of polygons P, P^* , ensuring compatibility at boundaries $\partial P, \partial P^*$ using the methods of Appendix B.

This proves Theorem 4.34(c). Parts (a),(b) when Y is oriented follow using Poincaré duality and Theorems 4.15 and 4.26, since Poincaré duality identifies $\cup, \cap, \bullet$ on both $H_*^{\text{si}}(Y; R), H_{\text{cs}}^*(Y; R)$ and $KH_*(Y; R), KH^*(Y; R)$ and $KH_*^{\text{ef}}(Y; R), KH_{\text{ec}}^*(Y; R)$. When Y is not orientable we deduce (a),(b) using *orientation bundles*, as in Definition 4.16. Part (a) then implies (d), since the identities in $H_{\text{cs}}^0(Y; R), KH^0(Y; R), KH_{\text{ec}}^0(Y; R)$ are characterized uniquely by the cup products $\cup$. This completes the proof of Theorem 4.34.

References

- [1] A. Adem, J. Leida and Y. Ruan, *Orbifolds and Stringy Topology*, Cambridge Tracts in Math. 171, Cambridge University Press, Cambridge, 2007.
- [2] M. Aganagic, A. Klemm and C. Vafa, *Disc instantons, mirror symmetry, and the duality web*, Z. Naturforsch. A57 (2002), 1–28. hep-th/0105045.
- [3] M. Aganagic and C. Vafa, *Mirror Symmetry, D-branes and counting holomorphic discs*, hep-th/0012041, 2000.

- [4] M. Akaho and D. Joyce, *Lagrangian Floer theory using Kuranishi cohomology*, in preparation, 2009.
- [5] M.F. Atiyah, *Bordism and cobordism*, Proc. Camb. Phil. Soc. 57 (1961), 200–208.
- [6] K. Behrend, *Gromov–Witten invariants in algebraic geometry*, Invent. Math. 127 (1997), 601–617. alg-geom/9601011.
- [7] K. Behrend, *Cohomology of stacks*, pages 249–294 in E. Arbarello, G. Ellingsrud and L. Goettsche, editors, *Intersection theory and moduli*, ICTP Lect. Notes volume XIX, Trieste, 2004.
Available online at <http://publications.ictp.it/>.
- [8] K. Behrend and B. Fantechi, *The intrinsic normal cone*, Invent. Math. 128 (1997), 45–88. alg-geom/9601010.
- [9] J. Bryan and R. Pandharipande, *BPS states of curves in Calabi–Yau 3-folds*, Geometry & Topology 5 (2001) 287–318. math.AG/0009025.
- [10] G.E. Bredon, *Topology and Geometry*, Graduate Texts in Math. 139, Springer-Verlag, New York, 1993.
- [11] M. Chas and D. Sullivan, *String Topology*, math.DG/9911159, 1999. To appear in Annals of Mathematics.
- [12] W. Chen and Y. Ruan, *Orbifold Gromov–Witten theory*, pages 25–86 in A. Adem, J. Morava and Y. Ruan, editors, *Orbifolds in mathematics and physics*, Cont. Math. 310, A.M.S., Providence, RI, 2002. math.AG/0103156.
- [13] W. Chen and Y. Ruan, *A new cohomology theory for orbifold*, Comm. Math. Phys. 248 (2004), 1–31. math.AG/0004129.
- [14] C.-H. Cho, *Counting real pseudo-holomorphic discs and spheres in dimension four and six*, math.SG/0604501, 2006.
- [15] C.-H. Cho, *On the counting of holomorphic discs in toric Fano manifolds*, math.SG/0604502, 2006.
- [16] P.E. Connor, *Differentiable Periodic Maps*, second edition, Springer Lecture Notes in Mathematics 738, Springer-Verlag, Berlin, 1979.
- [17] P.E. Connor and E.E. Floyd, *The relation of cobordism to K-theories*, Springer Lecture Notes in Mathematics 28, Springer-Verlag, Berlin, 1966.
- [18] P.E. Connor and E.E. Floyd, *Torsion in SU-bordism*, Memoirs of the A.M.S. 60, A.M.S., Providence, RI, 1966.
- [19] D.A. Cox and S. Katz, *Mirror Symmetry and Algebraic Geometry*, Mathematical Surveys and Monographs 68, A.M.S., Providence, RI, 1999.

- [20] T. Ekholm, J. Etnyre and M. Sullivan, *The contact homology of Legendrian submanifolds in $\mathbb{R}^{2n+1}$* , J. Diff. Geom. 71 (2005), 177–305.
- [21] Y. Eliashberg, A. Givental and H. Hofer, *Introduction to Symplectic Field Theory*, Geom. Funct. Anal. 2000, Special Volume, Part II, 560–673. math.SG/0010059.
- [22] K. Fukaya, *Floer homology and Mirror Symmetry II*, pages 31–127 in K. Fukaya, S. Nishikawa and J. Spruck, editors, *Minimal surfaces, geometric analysis and symplectic geometry*, Advanced Studies in Pure Math. 34, Mathematical Society of Japan, Tokyo, 2002.
- [23] K. Fukaya, *Application of Floer homology of Lagrangian submanifolds to symplectic topology*, pages 231–276 in P. Biran, O. Cornea and F. Lalonde, editors, *Morse theoretic methods in nonlinear analysis and in symplectic topology*, NATO Sci. Ser. II Math. Phys. Chem., 217, Springer, 2006.
- [24] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian intersection Floer theory – anomaly and obstruction*, preprint, final(?) version, 2008. 1385 pages.
- [25] K. Fukaya and K. Ono, *Arnold Conjecture and Gromov–Witten invariant*, Topology 38 (1999), 933–1048.
- [26] K. Fukaya and K. Ono, *Floer homology and Gromov–Witten invariant over integer of general symplectic manifolds — summary*, pages 75–91 in M. Maruyama and T. Sunada, editors, *Taniguchi Conference on Mathematics, Nara '98*, Advanced Studies in Pure Math. 31, Mathematical Society of Japan, Tokyo, 2001.
- [27] W. Fulton and R. MacPherson, *Categorical framework for the study of singular spaces*, Mem. A.M.S. 243, A.M.S., Providence, RI, 1981.
- [28] R. Gopakumar and C. Vafa, *M-theory and topological strings. I*, hep-th/9809187, 1998.
- [29] R. Gopakumar and C. Vafa, *M-theory and topological strings. II*, hep-th/9812127, 1998.
- [30] T. Graber and E. Zaslow, *Open-string Gromov–Witten invariants: calculations and a mirror ‘theorem’*, pages 107–121 in A. Adem, J. Morava and Y. Ruan, editors, *Orbifolds in mathematics and physics*, Cont. Math. 310, A.M.S., Providence, RI, 2002. hep-th/0109075.
- [31] M. Gromov, *Pseudoholomorphic curves in symplectic manifolds*, Invent. math. 82 (1985), 307–347.
- [32] J. Harris and I. Morrison, *Moduli of curves*, Graduate Texts in Math. 187, Springer-Verlag, New York, 1998.

- [33] H. Hofer, *A general Fredholm theory and applications*, math.SG/0509366, 2005.
- [34] H. Hofer, *Lectures on polyfolds and applications I*, preprint available from <http://www.cims.nyu.edu/~hofer/>, 2008.
- [35] H. Hofer, K. Wysocki and E. Zehnder, *A general Fredholm theory I: A splicing-based differential geometry*, J. Eur. Math. Soc. 9 (2007), 841–876. math.FA/0612604.
- [36] H. Hofer, K. Wysocki and E. Zehnder, *A general Fredholm theory II: implicit function theorems*, arXiv:0705.1310, 2007.
- [37] H. Hofer, K. Wysocki and E. Zehnder, *Integration theory for zero sets of polyfold Fredholm sections*, arXiv:0711.0781, 2007.
- [38] D. Joyce, *Kuranishi homology and Kuranishi cohomology: a User's Guide*, math.SG/0710.5634, version 2, 2008. 29 pages.
- [39] D. Joyce, *A theory of open Gromov–Witten invariants*, in preparation, 2008.
- [40] D. Joyce, *Kuranishi (co)homology and String Topology*, in preparation, 2008.
- [41] D. Joyce, *Towards a symplectic proof of the integrality conjecture for Gopakumar–Vafa invariants*, in preparation, 2009.
- [42] S. Katz and C.-C. M. Liu, *Enumerative geometry of stable maps with Lagrangian boundary conditions and multiple covers of the disc*, Adv. Theor. Math. Phys. 5 (2001), 1–49. math.AG/0103074.
- [43] A. Klemm and R. Pandharipande, *Enumerative geometry of Calabi–Yau 4-folds*, math.AG/0702189.
- [44] M. Kontsevich, *Homological Algebra of Mirror Symmetry*, pages 120–139 in *Proceedings of the International Congress of Mathematicians, (Zürich, 1994)*, vol. I, Birkhäuser, Basel, 1995. alg-geom/9411018.
- [45] M. Kontsevich and Y. Manin, *Gromov–Witten classes, quantum cohomology, and enumerative geometry*, Comm. Math. Phys. 164 (1994), 525–562. hep-th/9402147.
- [46] S. Lefschetz, *Topology*, A.M.S., New York, 1930.
- [47] J. Li and Y.S. Song, *Open string instantons and relative stable morphisms*, Adv. Theor. Math. Phys. 5 (2001), 67–91. hep-th/0103100.
- [48] J. Li and G. Tian, *Virtual moduli cycles and Gromov–Witten invariants of algebraic varieties*, J. A.M.S. 11 (1998), 119–174. alg-geom/9602007.

- [49] J. Li and G. Tian, *Virtual moduli cycles and Gromov–Witten invariants of general symplectic manifolds*, pages 47–83 in R.J. Stern, editor, *Topics in symplectic 4-manifolds (Irvine, CA, 1996)*, International Press, Cambridge, MA, 1998. alg-geom/9608032.
- [50] J. Li and G. Tian, *Comparison of the algebraic and symplectic Gromov–Witten invariants*, Asian J. Math. 3 (1999), 689–728. alg-geom/9712035.
- [51] C.-C.M. Liu, *Moduli of J -holomorphic curves with Lagrangian boundary conditions and open Gromov–Witten invariants for an S^1 -Equivariant Pair*, math.SG/0210257, 2002.
- [52] G. Liu and G. Tian, *Floer homology and Arnold conjecture*, J. Diff. Geom. 49 (1998), 1–74.
- [53] G. Lu and G. Tian, *Constructing virtual Euler cycles and classes*, International Mathematics Research Surveys 2007. math.SG/0605290.
- [54] E. Lupercio and B. Uribe, *Gerbés over orbifolds and twisted K -theory*, Comm. Math. Phys. 245 (2004), 449–489. math.AT/01050393.
- [55] D. McDuff, *The virtual moduli cycle*, pages 73–102 in Y. Eliashberg, D. Fuchs, T. Ratiu and A. Weinstein, editors, *Northern California Symplectic Geometry Seminar*, A.M.S. Translations 196, A.M.S., Providence, RI, 1999.
- [56] D. McDuff and D. Salamon, *J -holomorphic curves and symplectic topology*, Colloquium Publications 52, A.M.S., Providence, R.I., 2004.
- [57] I. Moerdijk and D.A. Pronk, *Orbifolds, sheaves and groupoids*, K-theory 12 (1997), 3–21.
- [58] H. Ooguri and C. Vafa, *Knot invariants and topological strings*, Nucl. Phys. B577 (2000), 69–108. hep-th/9912123.
- [59] R. Pandharipande, *Hodge integrals and degenerate contributions*, Comm. Math. Phys. 208 (1999), 489506. math.AG/9811140.
- [60] R. Pandharipande, *Three questions in Gromov–Witten theory*, pages 503–512 in vol. II of *Proceedings of the International Congress of Mathematicians, (Beijing, 2002)*, Higher Education Press, Beijing, 2002. math.AG/0302077.
- [61] R. Pandharipande, J. Solomon and J. Walcher, *Disk enumeration on the quintic 3-fold*, math.SG/0610901, 2006.
- [62] R. Pandharipande and R. Thomas, *Curve counting via stable pairs in the derived category*, arXiv:0707.2348, 2007.
- [63] R. Pandharipande and R. Thomas, *Stable pairs and BPS invariants*, arXiv:0711.3899, 2007.

- [64] R. Pandharipande and A. Zinger, *Enumerative geometry of Calabi–Yau 5-folds*, arXiv:0803.1640, 2008.
- [65] Y. Ruan, *Virtual neighbourhoods and pseudo-holomorphic curves*, Turkish J. Math. 23 (1999), 161–231. alg-geom/96110121.
- [66] I. Satake, *On a generalization of the notion of manifold*, Proc. Nat. Acad. Sci. U.S.A. 42 (1956), 359–363.
- [67] P. Seidel, *Fukaya categories and deformations*, pages 351–360 in *Proceedings of the International Congress of Mathematicians (Beijing, 2002)*, vol. II, Higher Ed. Press, Beijing, 2002. math.SG/0206155.
- [68] P. Seidel, *A biased view of symplectic cohomology*, arXiv:0704.2055, 2007.
- [69] P. Seidel, *Fukaya categories and Picard–Lefschetz theory*, Zürich Lectures in Advanced Mathematics, European Mathematical Society, Zürich, 2008.
- [70] B. Siebert, *Gromov–Witten invariants of general symplectic manifolds*, dg-ga/9608005, 1996.
- [71] B. Siebert, *Symplectic Gromov–Witten invariants*, pages 375–424 in K. Hulek, F. Catanese, C. Peters and M. Reid, editors, *New trends in algebraic geometry (Warwick, 1996)*, L.M.S. Lecture Notes 264, C.U.P. 1999.
- [72] B. Siebert, *Algebraic and symplectic Gromov–Witten invariants coincide*, Ann. Inst. Fourier (Grenoble) 49 (1999), 1743–1795. dg-ga/9804108.
- [73] J. Solomon, *Intersection theory of the moduli space of holomorphic curves with Lagrangian boundary conditions*, math.SG/0606429, 2006.
- [74] R.E. Stong, *Notes on cobordism theory*, Princeton University Press, Princeton, NJ, 1968.
- [75] G.W. Whitehead, *Generalized homology theories*, Trans. A.M.S. 102 (1962), 292–311.
- [76] J.-Y. Welschinger, *Invariants of real symplectic 4-manifolds and lower bounds in real enumerative geometry*, math.AG/0303145, 2003.
- [77] J.-Y. Welschinger, *Enumerative invariants of strongly semipositive real symplectic six-manifolds*, math.AG/0509121, 2005.

THE MATHEMATICAL INSTITUTE, 24-29 ST. GILES, OXFORD, OX1 3LB, U.K.
 E-MAIL: joyce@maths.ox.ac.uk